

ON THE EVALUATION OF CERTAIN SERIES AT A GOLDEN RATIO-BASED CONSTANT

TRISTEN HARR

ABSTRACT. This paper introduces and analyzes a specific complex constant, denoted Λ_{G1} , which is uniquely derived from a set of defining equations related to the golden ratio, ϕ . The motivation for this constant is shown to arise from a classical geometric construction. We first prove the constant's fundamental properties, including the convergence condition $|\Lambda_{G1}| < 1$. The primary contribution of this work is the systematic evaluation of two families of infinite series where Λ_{G1} serves as the argument, connecting them to Eulerian polynomials and the Polylogarithm function, $\text{Li}_s(\Lambda_{G1})$. Furthermore, we present high-precision numerical evaluations for the Dilogarithm ($s = 2$) and Trilogarithm ($s = 3$) of this constant and, based on this evidence, we conjecture that these values are transcendental numbers not expressible in the ring of common mathematical constants.

1. INTRODUCTION

The study of infinite series often reveals deep connections between different areas of mathematics. This paper focuses on the properties of series evaluated at a specific complex constant derived from the golden ratio, ϕ . The motivation for this constant arises from the classical geometric problem of dividing a line segment in extreme and mean ratio. This problem provides a non-arbitrary foundation for a pair of defining equations whose unique solution gives rise to the constant we analyze.

The purpose of this paper is to rigorously define this constant and systematically explore the evaluation of several important families of infinite series using this constant as the primary argument. We will show that because this constant is proven to lie within the unit disk, it serves as a valid and interesting argument for well-known summation formulas. This work provides a careful cataloging of these results and extends the analysis to a numerical investigation of the resulting values of the Polylogarithm function.

2. THE CONSTANT Λ_{G1} AND ITS FOUNDATIONAL PROPERTIES

2.1. Defining Equations and Uniqueness. We define two real constants, T and J , as the unique solution to a pair of equations representing minimal structural simplicity.

Definition 2.1 (Defining Equations). The constants T and J are the real numbers (a, b) that satisfy:

- (1) $a + b = 1/2$.
- (2) $a/b = \phi$, where $\phi = \frac{1+\sqrt{5}}{2}$.

Solving this system yields the unique values $T = (\sqrt{5} - 1)/4$ and $J = (3 - \sqrt{5})/4$.

From these constants, we define our primary object of study.

Definition 2.2 (The Primary Constant). The constant Λ_{G1} is defined as the complex number $\Lambda_{G1} = T + iJ$.

2.2. Convergence Property. A crucial property of Λ_{G1} is that its magnitude is less than one, which guarantees the convergence of the power series we will investigate.

Lemma 2.3 (Convergence of Λ_{G1}). *The constant Λ_{G1} lies within the unit disk, i.e., $|\Lambda_{G1}| < 1$.*

Proof. Convergence requires $|\Lambda_{G1}|^2 < 1$. We calculate the squared values of the components: $T^2 = \left(\frac{\sqrt{5}-1}{4}\right)^2 = \frac{6-2\sqrt{5}}{16}$ and $J^2 = \left(\frac{3-\sqrt{5}}{4}\right)^2 = \frac{14-6\sqrt{5}}{16}$. Summing these gives:

$$\begin{aligned} |\Lambda_{G1}|^2 &= T^2 + J^2 = \frac{6-2\sqrt{5}+14-6\sqrt{5}}{16} \\ &= \frac{20-8\sqrt{5}}{16} = \frac{5-2\sqrt{5}}{4} \approx 0.131966 \end{aligned}$$

Since $0.131966 < 1$, the condition is satisfied. $\square$

3. EVALUATION OF POLYNOMIAL-WEIGHTED SERIES

The first family of series we analyze are those weighted by polynomial powers of the index.

Proposition 3.1 (Polynomial-Weighted Sums). *The infinite series $S_k = \sum_{n=0}^{\infty} n^k \cdot (\Lambda_{G1})^n$, for any integer $k \geq 0$, converges to the value:*

$$S_k = \frac{A_k(\Lambda_{G1})}{(1 - \Lambda_{G1})^{k+1}}$$

where $A_k(x)$ is the k -th Eulerian polynomial.

Proof. The general formula for the sum of a polynomial-weighted geometric series is a known result (see, e.g., Abramowitz and Stegun, *Handbook of Mathematical Functions*). The formula is valid for any complex argument x such that $|x| < 1$. Since we have proven in Lemma 2.3 that $|\Lambda_{G1}| < 1$, the substitution of Λ_{G1} into this formula is valid. $\square$

4. EVALUATION OF RECIPROCAL-WEIGHTED SERIES AND THE POLYLOGARITHM

The second family of series are those weighted by reciprocal powers of the index.

Definition 4.1 (The Polylogarithm Function). The Polylogarithm function of order s , $\text{Li}_s(z)$, is defined by the power series:

$$\text{Li}_s(z) = \sum_{n=1}^{\infty} \frac{z^n}{n^s}$$

This series converges for all complex numbers z such that $|z| < 1$.

Proposition 4.2 (Connection to the Polylogarithm Function). *The series $S_s = \sum_{n=1}^{\infty} \frac{(\Lambda_{G1})^n}{n^s}$ converges for any complex s to the value:*

$$S_s = \text{Li}_s(\Lambda_{G1})$$

Proof. This result follows directly from the definition of the Polylogarithm function. Since we have proven $|\Lambda_{G1}| < 1$ in Lemma 2.3, the series is guaranteed to converge. $\square$

5. NUMERICAL ANALYSIS AND A CONJECTURE ON THE NATURE OF $\text{Li}_s(\Lambda_{G1})$

We now investigate the nature of the specific values obtained in the previous section.

5.1. Numerical Values. We present high-precision numerical evaluations for the first two non-trivial integer cases of the Polylogarithm of Λ_{G1} , computed from the series definition.

- For $s = 2$, the Dilogarithm of Λ_{G1} is:

$$\text{Li}_2(\Lambda_{G1}) \approx 0.322328253720993 + 0.226730769307749i$$

- For $s = 3$, the Trilogarithm of Λ_{G1} is:

$$\text{Li}_3(\Lambda_{G1}) \approx 0.310345112613931 + 0.198884393433621i$$

5.2. A Conjecture on the Nature of These Values. The numerical results above are presented as benchmark values for future research. Analysis using inverse symbolic calculators suggests that these values do not resolve into simple, closed-form expressions involving common mathematical constants. This leads us to propose the following.

Conjecture 5.1. *For any integer $s \geq 2$, the value $\text{Li}_s(\Lambda_{G1})$ is a transcendental number. Furthermore, it is conjectured that this number is not in the ring $\mathbb{Q}(\pi, e, \ln(2), \phi, \gamma)$.*

Remark. Proving this conjecture is beyond the scope of this paper, but proposing it frames a clear direction for future work and elevates the significance of the constant Λ_{G1} as a tool for probing the structure of the Polylogarithm function.

6. CONCLUSION

We have rigorously defined a complex constant, Λ_{G1} , derived from equations motivated by a classical geometric problem. We have proven its convergence, which validates its use in a variety of infinite series. The primary contribution of this work is the systematic cataloging of the closed-form values for series weighted by n^k and n^{-s} evaluated at this constant.

Furthermore, we have extended this analysis by providing high-precision numerical values for $\text{Li}_2(\Lambda_{G1})$ and $\text{Li}_3(\Lambda_{G1})$. Based on this evidence, we have conjectured that these values are non-trivial transcendental numbers. This paper thus provides a complete, self-contained exploration, a set of benchmark results, and a new open problem related to this elegant mathematical constant.

APPENDIX A. GEOMETRIC MOTIVATION FOR THE DEFINING EQUATIONS

The defining equations in Section 2 are motivated by a classical compass-and-straightedge construction. Specifically, they represent the algebraic solution to the problem of dividing a line segment of length $1/2$ into two smaller segments, T and J , such that their ratio is the golden ratio, i.e., $T/J = \phi$. This geometric origin provides a foundational, non-arbitrary basis for the constants.



FIGURE 1. A visual representation of the geometric problem of dividing a line segment of length $1/2$ in the golden ratio, which gives rise to the constants T and J .