

A GOLDEN RATIO-BASED CONSTANT AND CONJECTURES ON ITS POLYLOGARITHMIC VALUES

TRISTEN HARR

ABSTRACT. This paper introduces and analyzes a specific complex constant, denoted Λ_{G1} , which is uniquely derived from a set of defining equations related to the golden ratio, ϕ . The motivation for this constant is shown to arise from a classical geometric construction. We first prove the constant's fundamental properties, including the convergence condition $|\Lambda_{G1}| < 1$. The primary contribution of this work is the systematic identification of the values of two families of infinite series where Λ_{G1} serves as the argument, connecting them to Eulerian polynomials and the Polylogarithm function, $\text{Li}_s(\Lambda_{G1})$. Furthermore, we present high-precision numerical evaluations for the Dilogarithm ($s = 2$) and Trilogarithm ($s = 3$) of this constant and, based on this evidence, we conjecture that these values are transcendental numbers not expressible in the ring of common mathematical constants.

1. INTRODUCTION

The study of infinite series often reveals deep connections between different areas of mathematics. This paper focuses on the properties of series evaluated at a specific complex constant derived from the golden ratio, ϕ . The motivation for this constant arises from the classical geometric problem of dividing a line segment in extreme and mean ratio. This problem provides a non-arbitrary foundation for a pair of defining equations whose unique solution gives rise to the constant we analyze.

The purpose of this paper is to rigorously define this constant and systematically explore the evaluation of several important families of infinite series using this constant as the primary argument. We will show that because this constant is proven to lie within the unit disk, it serves as a valid and interesting argument for well-known summation formulas. This work provides a careful cataloging of these results and extends the analysis to a numerical investigation of the resulting values of the Polylogarithm function, culminating in a formal conjecture about their nature.

2. THE CONSTANT Λ_{G1} AND ITS FOUNDATIONAL PROPERTIES

2.1. Defining Equations and Uniqueness. We define two real constants, T and J , as the unique solution to a pair of equations motivated by a classical geometric problem.

Definition 2.1 (Defining Equations). The constants T and J are the real numbers (a, b) that satisfy:

- (1) $a + b = 1/2$.
- (2) $a/b = \phi$, where $\phi = \frac{1+\sqrt{5}}{2}$.

Solving this system yields the unique values $T = (\sqrt{5} - 1)/4$ and $J = (3 - \sqrt{5})/4$.

From these constants, we define our primary object of study.

Definition 2.2 (The Primary Constant). The constant Λ_{G1} is defined as the complex number $\Lambda_{G1} = T + iJ$.

2.2. Convergence Property. A crucial property of Λ_{G1} is that its magnitude is less than one, which guarantees the convergence of the power series we will investigate.

Lemma 2.3 (Magnitude of Λ_{G1}). *The constant Λ_{G1} lies within the unit disk, i.e., $|\Lambda_{G1}| < 1$.*

Proof. Convergence requires $|\Lambda_{G1}|^2 < 1$. We calculate the squared values of the components: $T^2 = \left(\frac{\sqrt{5}-1}{4}\right)^2 = \frac{6-2\sqrt{5}}{16}$ and $J^2 = \left(\frac{3-\sqrt{5}}{4}\right)^2 = \frac{14-6\sqrt{5}}{16}$. Summing these gives:

$$\begin{aligned} |\Lambda_{G1}|^2 &= T^2 + J^2 = \frac{6-2\sqrt{5}+14-6\sqrt{5}}{16} \\ &= \frac{20-8\sqrt{5}}{16} = \frac{5-2\sqrt{5}}{4} \approx 0.131966 \end{aligned}$$

Since $0.131966 < 1$, the condition is satisfied. $\square$

3. APPLICATION TO POLYNOMIAL-WEIGHTED SERIES

The first family of series we analyze are those weighted by polynomial powers of the index. Since it is proven in Lemma 2.3 that $|\Lambda_{G1}| < 1$, the constant is a valid argument for the standard formula for polynomial-weighted geometric series. This sum is given in terms of the Eulerian polynomials, $A_k(x)$ [1]. For instance, the first two Eulerian polynomials are $A_0(x) = 1$ and $A_1(x) = x$, which correspond to the standard geometric series and its first derivative sum. The general identification for our constant is as follows:

$$\sum_{n=0}^{\infty} n^k \cdot (\Lambda_{G1})^n = \frac{A_k(\Lambda_{G1})}{(1 - \Lambda_{G1})^{k+1}}$$

This identity holds for any integer $k \geq 0$.

4. APPLICATION TO RECIPROCAL-WEIGHTED SERIES

The second family of series are those weighted by reciprocal powers of the index. These are identified by definition with the Polylogarithm special function.

Definition 4.1 (The Polylogarithm Function). The Polylogarithm function of order s , $\text{Li}_s(z)$, is defined by the power series:

$$\text{Li}_s(z) = \sum_{n=1}^{\infty} \frac{z^n}{n^s}$$

This series converges for all complex numbers z such that $|z| < 1$.

Since $|\Lambda_{G1}| < 1$, the series $\sum_{n=1}^{\infty} \frac{(\Lambda_{G1})^n}{n^s}$ converges, and its value is, by definition, the Polylogarithm of Λ_{G1} :

$$\sum_{n=1}^{\infty} \frac{(\Lambda_{G1})^n}{n^s} = \text{Li}_s(\Lambda_{G1})$$

This identity holds for any complex s .

5. NUMERICAL ANALYSIS AND A CONJECTURE ON THE NATURE OF $\text{Li}_s(\Lambda_{G1})$

We now investigate the nature of the specific values identified in the previous section.

5.1. Numerical Values. We present high-precision numerical evaluations for the first two non-trivial integer cases of the Polylogarithm of Λ_{G1} , computed from the series definition.

- For $s = 2$, the Dilogarithm of Λ_{G1} is:

$$\text{Li}_2(\Lambda_{G1}) \approx 0.322328253720993 + 0.226730769307749i$$

- For $s = 3$, the Trilogarithm of Λ_{G1} is:

$$\text{Li}_3(\Lambda_{G1}) \approx 0.310345112613931 + 0.198884393433621i$$

5.2. A Conjecture on the Nature of These Values. The transcendental nature of special function values at specific algebraic arguments is a deep and active area of research in number theory [2]. Our numerical results motivate a new conjecture in this domain. Analysis using integer relation algorithms (such as PSLQ) does not resolve the values above into simple, closed-form expressions involving common mathematical constants. This leads us to propose the following.

Conjecture 5.1. *For any integer $s \geq 2$, the value $\text{Li}_s(\Lambda_{G1})$ is a transcendental number. Furthermore, it is conjectured that this number is not in the ring $\mathbb{Q}(\pi, e, \ln(2), \phi, \gamma)$.*

Remark. Proving this conjecture is beyond the scope of this paper, but proposing it frames a clear direction for future work and elevates the significance of the constant Λ_{G1} as a tool for probing the structure of the Polylogarithm function.

6. CONCLUSION

We have rigorously defined a complex constant, Λ_{G1} , derived from equations motivated by a classical geometric problem. We have proven its convergence, which validates its use in a variety of infinite series. The primary contribution of this work is the systematic identification of the values for series weighted by n^k and n^{-s} evaluated at this constant.

Furthermore, we have extended this analysis by providing high-precision numerical values for $\text{Li}_2(\Lambda_{G1})$ and $\text{Li}_3(\Lambda_{G1})$. Based on this evidence, we have conjectured that these values are non-trivial transcendental numbers. The constant's constrained algebraic nature, arising from a clear geometric problem, makes it a compelling candidate for further study into the analytic properties of special functions, as its structure is not arbitrary. This paper thus provides a complete, self-contained exploration, a set of benchmark results, and a new open problem for the community.

APPENDIX A. GEOMETRIC MOTIVATION FOR THE DEFINING EQUATIONS

The defining equations in Section 2 are motivated by a classical compass-and-straightedge construction. Specifically, they represent the algebraic solution to the problem of dividing a line segment of length $1/2$ into two smaller segments, T and J , such that their ratio is the golden ratio, i.e., $T/J = \phi$. This geometric origin provides a foundational, non-arbitrary basis for the constants.

REFERENCES

- [1] M. Abramowitz and I. A. Stegun, eds., *Handbook of Mathematical Functions with Formulas, Graphs, and Mathematical Tables*. Dover Publications, New York, 1965.
- [2] L. Lewin, *Polylogarithms and Associated Functions*. North Holland, New York, 1981.