

FE8815

Interest Rate Derivatives

Nicolas Privault

This document deals the stochastic modeling of interest rates and the pricing of related derivative products. Bond pricing and the modeling of forward rates are introduced in Chapters 1 and 2. The general construction of forward measures by change of numéraire is given in Chapter 3 and is applied to the pricing of interest rate derivatives such as caplets, caps and swaptions in Chapter 4. Credit risk and stochastic default are considered in Chapter 5, with the pricing of defaultable bonds.

The pdf file contains internal and external links, and 41 figures, including 9 animated figures, *e.g.* Figures 1.16, 2.6, 2.9, 2.10, 2.17, that may require using Acrobat Reader for viewing on the complete pdf file. It also includes 2 Python code and 6 R codes *e.g.* on pages 41, 108,

This text contains 51 exercises with solutions. Clicking on an exercise number inside the solution section will send to the original problem text inside the file. Conversely, clicking on a problem number sends the reader to the corresponding solution, however this feature should not be misused.

Contents

1	Short Rates and Bond Pricing	1
1.1	Short-Term Mean-Reverting Models	1
1.2	Calibration of the Vasicek Model	7
1.3	Zero-Coupon and Coupon Bonds	10
1.4	Bond Pricing PDE	14
	Exercises	27
2	Forward Rate Modeling	35
2.1	Construction of Forward Rates	35
2.2	Interest Rate Swaps	44
2.3	The HJM Model	47
2.4	Yield Curve Modeling	52
2.5	Two-Factor Model	55
2.6	The BGM Model	59
	Exercises	61
3	Change of Numéraire and Forward Measures	65
3.1	Notion of Numéraire	65
3.2	Change of Numéraire	68
3.3	Foreign Exchange	76
3.4	Pricing Exchange Options	83
3.5	Hedging by Change of Numéraire	85
	Exercises	88

4	Pricing of Interest Rate Derivatives	93
4.1	Forward Measures and Tenor Structure	93
4.2	Bond Options	97
4.3	Caplet Pricing	98
4.4	Forward Swap Measures	102
4.5	Swaption Pricing	103
	Exercises	111
5	Reduced-Form Approach to Credit Risk	123
5.1	Survival Probabilities	123
5.2	Stochastic Default	125
5.3	Defaultable Bonds	128
	Exercises	131
	Exercise Solutions	135
	Chapter 1	135
	Chapter 2	149
	Chapter 3	157
	Chapter 4	167
	Chapter 5	184
	Bibliography	189
	Articles	189
	Books	191
	Index	193
	Author index	199

List of Figures

1.1	Short rate $t \mapsto r_t$ in the Vasicek model	3
1.2	CBOE 10 Year Treasury Note (TNX) yield	4
1.3	Short rate $t \mapsto r_t$ in the CIR model	5
1.4	Calibrated Vasicek simulation vs market data	10
1.5	Five-dollar 1875 Louisiana bond with 7.5% biannual coupons	10
1.6	Discrete-time coupon bond pricing	12
1.7	Continuous-time coupon bond pricing	13
1.8	Comparison of Monte Carlo and PDE solutions	19
1.9	Bond price $t \mapsto P(t, T)$ vs $t \mapsto e^{-r_0(T-t)}$	19
1.10	Bond price $t \mapsto P_c(t, T)$ with a 5% coupon rate	20
1.11	Bond price with coupon rate 6.25%	20
1.12	Orange Cnty Calif bond prices	21
1.13	Orange Cnty Calif bond yields	22
1.14	Fitting of a gamma probability density function	25
1.15	Approximation of Dothan bond prices	25
1.16	Brownian bridge	26
2.1	Forward rate $S \mapsto f(t, t, S)$	36
2.2	Indonesian government securities yield curve	36
2.3	Forward rate process $t \mapsto f(t, T, S)$	40
2.4	Instantaneous forward rate process $t \mapsto f(t, T)$	41
2.5	Federal Reserve yield curves from 1982 to 2012	41
2.6	European Central Bank yield curves*	42
2.7	August 2019 Federal Reserve yield curve inversion*	43
2.8	Stochastic process of forward curves	47
2.9	Forward instantaneous curve in the Vasicek model*	51
2.10	Forward instantaneous curve $x \mapsto f(0, x)$ in the Vasicek model*	51
2.11	Short-term interest rate curve $t \mapsto r_t$ in the Vasicek model	52
2.12	Nelson-Siegel graph	52

2.13	Svensson model graph	53
2.14	Fitting of a Svensson curve to market data	53
2.15	Graphs of forward rates	54
2.16	Forward instantaneous curve in the Vasicek model	54
2.17	ECB data vs fitted yield curve*	55
2.18	Bond prices $t \mapsto P(t, T_2), P(t, T_2), P(t, T_3)$	56
2.19	Forward rates in a two-factor model	58
2.20	Evolution of instantaneous forward rates in a two-factor model	58
2.21	Roadmap of stochastic interest rate modeling	60
3.1	Why change of numéraire?	67
3.2	Overseas investment opportunity	76
4.1	Implied swaption volatilities	108
S.1	Log bond prices correlation graph in the two-factor model	156

List of Tables

2.1	Stochastic interest rate models	59
3.1	Local vs foreign exchange options	82
4.1	Forward rates arranged according to a tenor structure	94
4.2	A list of numéraire processes and their applications	110
5.1	Mortality table	124
5.2	Cumulative historic default rates	130

*Animated figures (work in Acrobat reader).

1. Short Rates and Bond Pricing

Short-term rates, typically daily rates, are the interest rates applied to short-term lending between financial institutions. The stochastic modeling of short-term interest rate processes is based on the mean reversion property, as in the Vasicek, CIR, CEV, and affine-type models studied in this chapter. The pricing of fixed income products, such as bonds, is considered in this framework using probabilistic and PDE arguments.

1.1	Short-Term Mean-Reverting Models	1
1.2	Calibration of the Vasicek Model	7
1.3	Zero-Coupon and Coupon Bonds	10
1.4	Bond Pricing PDE	14
	Exercises	27

1.1 Short-Term Mean-Reverting Models

Money market accounts with price $(A_t)_{t \in \mathbb{R}_+}$ can be defined from a short-term interest rate process $(r_t)_{t \in \mathbb{R}_+}$ as

$$\frac{A_{t+dt} - A_t}{A_t} = r_t dt, \quad \frac{dA_t}{A_t} = r_t dt, \quad \frac{dA_t}{dt} = r_t A_t, \quad t \geq 0,$$

with

$$A_t = A_0 \exp \left(\int_0^t r_s ds \right), \quad t \geq 0.$$

As short-term interest rates behave differently from stock prices, they require the development of specific models to account for properties such as positivity, boundedness, and return to equilibrium.

Vašíček, 1977 model

The first model to capture the mean reversion property of interest rates, a property not possessed by geometric Brownian motion, is the Vašíček, 1977 model, which is based on the Ornstein-Uhlenbeck process. Here, the short-term interest rate process $(r_t)_{t \in \mathbb{R}_+}$ solves the equation

$$dr_t = (a - br_t)dt + \sigma dB_t, \quad (1.1.1)$$

where $a, \sigma \in \mathbb{R}$, $b > 0$, and $(B_t)_{t \in \mathbb{R}_+}$ is a standard Brownian motion, with solution

$$r_t = r_0 e^{-bt} + \frac{a}{b}(1 - e^{-bt}) + \sigma \int_0^t e^{-(t-s)b} dB_s, \quad t \geq 0, \quad (1.1.2)$$

see Exercise 1.1. The probability distribution of r_t is Gaussian at all times t , with mean

$$\mathbb{E}[r_t] = r_0 e^{-bt} + \frac{a}{b}(1 - e^{-bt}),$$

and variance

$$\begin{aligned} \text{Var}[r_t] &= \text{Var}\left[\sigma \int_0^t e^{-(t-s)b} dB_s\right] \\ &= \sigma^2 \int_0^t (e^{-(t-s)b})^2 ds \\ &= \sigma^2 \int_0^t e^{-2bs} ds \\ &= \frac{\sigma^2}{2b}(1 - e^{-2bt}), \quad t \geq 0, \end{aligned}$$

i.e.

$$r_t \simeq \mathcal{N}\left(r_0 e^{-bt} + \frac{a}{b}(1 - e^{-bt}), \frac{\sigma^2}{2b}(1 - e^{-2bt})\right), \quad t > 0.$$

In particular, the probability density function $f_t(x)$ of r_t at time $t > 0$ is given by

$$f_t(x) = \frac{\sqrt{b/\pi}}{\sigma \sqrt{1 - e^{-2bt}}} \exp\left(-\frac{(r_0 e^{-bt} + a(1 - e^{-bt})/b - x)^2}{\sigma^2(1 - e^{-2bt})/b}\right), \quad x \in \mathbb{R}.$$

In the long run,* i.e. as time t becomes large we have, assuming $b > 0$,

$$\lim_{t \rightarrow \infty} \mathbb{E}[r_t] = \frac{a}{b} \quad \text{and} \quad \lim_{t \rightarrow \infty} \text{Var}[r_t] = \frac{\sigma^2}{2b}, \quad (1.1.3)$$

and this distribution converges to the Gaussian $\mathcal{N}(a/b, \sigma^2/(2b))$ distribution, which is also the *invariant* (or stationary) distribution of $(r_t)_{t \in \mathbb{R}_+}$, see Exercise 1.1. In addition, the process tends to revert to its long term mean $a/b = \lim_{t \rightarrow \infty} \mathbb{E}[r_t]$ which makes the average drift vanish, i.e.:

$$\lim_{t \rightarrow \infty} \mathbb{E}[a - br_t] = a - b \lim_{t \rightarrow \infty} \mathbb{E}[r_t] = 0.$$

Figure 1.1 presents a random simulation of $t \mapsto r_t$ in the Vasicek model with $r_0 = 3\%$, and shows the mean-reverting property of the process with respect to $a/b = 2.5\%$.

*“But this long run is a misleading guide to current affairs. In the long run we are all dead.” J. M. Keynes, A Tract on Monetary Reform (1923), Ch. 3, p. 80.




Figure 1.1: Graph of the Vasicek short rate $t \mapsto r_t$ with $a = 0.025$, $b = 1$, and $\sigma = 0.1$.

As can be checked from the simulation of Figure 1.1 the value of r_t in the Vasicek model may become negative due to its Gaussian distribution. Although real interest rates may sometimes fall below zero,* this can be regarded as a potential drawback of the Vasicek model. The next R code provides a numerical solution of the stochastic differential equation (1.1.1) using the Euler method, see Figure 1.1.

```

1 N=10000;t<-0:(N-1);dt<-1.0/N;nsim<-2; a=0.025;b=1;sigma=0.1;
2 X <- matrix(rnorm(nsim*N,mean=0,sd=sqrt(dt)), nsim, N)
3 R <- matrix(0,nsim,N);R[,1]=0.03
4 dev.new(width=10,height=7);
5 for (i in 1:nsim){for (j in 2:N){R[i,j]=R[i,j-1]+(a-b*R[i,j-1])*dt+sigma*X[i,j]}}
6 par(mar=c(0,1,1,1));par(oma=c(0,1,1,1));par(mgp=c(-5,1,1))
7 plot(t,R[,1],xlab = "Time",ylab = "",type = "l",ylim = c(R[,1]-0.2,R[,1]+0.2),col = 0,axes=FALSE)
8 axis(2, pos=0,las=1);for (i in 1:nsim){lines(t, R[, i], xlab = "time", type = "l", col = i+0)}
9 abline(h=a/b,col="blue",lwd=3);abline(h=0)

```

Example - TNX yield

We consider the yield of the 10 Year Treasury Note on the Chicago Board Options Exchange (CBOE), for later use in the calibration of the Vasicek model. Treasury notes usually have a maturity between one and 10 years, whereas treasury bonds have maturities beyond 10 years)

```

1 library(quantmod)
2 getSymbols("^TNX",from="2012-01-01",to="2016-01-01",src="yahoo")
3 rate=Ad(`TNX`);rate<-rate[!is.na(rate)]
4 dev.new(width=10,height=7);chartSeries(rate,up.col="blue",theme="white")
5 n = sum(!is.na(rate))

```

The next Figure 1.2 displays the yield of the 10 Year Treasury Note.

*Eurozone interest rates turned negative in 2014.




Figure 1.2: CBOE 10 Year Treasury Note (TNX) yield.

Cox-Ingersoll-Ross (CIR) model

The [Cox, Ingersoll, and Ross, 1985](#) (CIR) model brings a solution to the positivity problem encountered with the Vasicek model, by the use the nonlinear stochastic differential equation

$$dr_t = \beta(\alpha - r_t)dt + \sigma\sqrt{r_t}dB_t, \quad (1.1.4)$$

with $\alpha > 0$, $\beta > 0$, $\sigma > 0$. The probability distribution of r_t at time $t > 0$ admits the noncentral Chi square probability density function given by

$$\begin{aligned} f_t(x) & \quad (1.1.5) \\ &= \frac{2\beta}{\sigma^2(1 - e^{-\beta t})} \exp\left(-\frac{2\beta(x + r_0 e^{-\beta t})}{\sigma^2(1 - e^{-\beta t})}\right) \left(\frac{x}{r_0 e^{-\beta t}}\right)^{\alpha\beta/\sigma^2-1/2} I_{2\alpha\beta/\sigma^2-1}\left(\frac{4\beta\sqrt{r_0 x e^{-\beta t}}}{\sigma^2(1 - e^{-\beta t})}\right), \end{aligned}$$

$x > 0$, where

$$I_\lambda(z) := \left(\frac{z}{2}\right)^\lambda \sum_{k \geq 0} \frac{(z^2/4)^k}{k! \Gamma(\lambda + k + 1)}, \quad z \in \mathbb{R},$$

is the modified Bessel function of the first kind, see Lemma 9 in [Feller, 1951](#) and Corollary 24 in [Albanese and Lawi, 2005](#). Note that $f_t(x)$ is not defined at $x = 0$ if $\alpha\beta/\sigma^2 - 1/2 < 0$, i.e. $\sigma^2 > 2\alpha\beta$, in which case the probability distribution of r_t admits a point mass at $x = 0$. On the other hand, r_t remains almost surely strictly positive under the Feller condition $2\alpha\beta \geq \sigma^2$, cf. the study of the associated probability density function in Lemma 4 of [Feller, 1951](#) for $\alpha, \beta \in \mathbb{R}$.

Figure 1.3 presents a random simulation of $t \mapsto r_t$ in the [Cox, Ingersoll, and Ross, 1985](#) (CIR) model in the case $\sigma^2 > 2\alpha\beta$, in which the process is mean reverting with respect to $\alpha = 2.5\%$ and has a nonzero probability of hitting 0.






Figure 1.3: Graph of the CIR short rate $t \mapsto r_t$ with $\alpha = 2.5\%$, $\beta = 1$, and $\sigma = 1.3$.

The next R code provides a numerical solution of the stochastic differential equation (1.1.4) using the Euler method, see Figure 1.3.

```

1 N=10000; t <- 0:(N-1); dt <- 1.0/N; nsim <- 2
2 a=0.025; b=1; sigma=0.1; sd=sqrt(sigma^2/2/b)
3 X <- matrix(rnorm(nsim*N,mean=0,sd=sqrt(dt)), nsim, N)
4 R <- matrix(0,nsim,N);R[,1]=0.03
5 for (i in 1:nsim){for (j in 2:N){R[i,j]=max(0,R[i,j-1]+(a-b*R[i,j-1])*dt+sigma*sqrt(R[i,j-1])*X[i,j])}}
6 plot(t,R[1,],xlab="time",ylab="",type="l",ylim=c(0,R[1,1]+sd/5),col=0,axes=FALSE)
7 axis(2, pos=0)
8 for (i in 1:nsim){lines(t, R[i, ], xlab = "time", type = "l", col = i+8)}
9 abline(h=a/b,col="blue",lwd=3);abline(h=0)

```

In large time $t \rightarrow \infty$, using the asymptotics

$$I_\lambda(z) \simeq_{z \rightarrow 0} \frac{1}{\Gamma(\lambda + 1)} \left(\frac{z}{2}\right)^\lambda,$$

the probability density function (1.1.5) becomes the gamma density function

$$f(x) = \lim_{t \rightarrow \infty} f_t(x) = \frac{1}{\Gamma(2\alpha\beta/\sigma^2)} \left(\frac{2\beta}{\sigma^2}\right)^{2\alpha\beta/\sigma^2} x^{-1+2\alpha\beta/\sigma^2} e^{-2\beta x/\sigma^2}, \quad x > 0. \quad (1.1.6)$$

with shape parameter $2\alpha\beta/\sigma^2$ and scale parameter $\sigma^2/(2\beta)$, which is also the *invariant distribution* of r_t .

The family of classical mean-reverting models also includes the [Courtadon, 1982](#) model

$$dr_t = \beta(\alpha - r_t)dt + \sigma r_t dB_t,$$

where α, β, σ are nonnegative, cf. Exercise 1.5, and the exponential Vasicek model

$$dr_t = r_t(\eta - a \log r_t)dt + \sigma r_t dB_t,$$

where $a, \eta, \sigma > 0$.

Constant Elasticity of Variance (CEV) model

Constant Elasticity of Variance models are designed to take into account nonconstant volatilities that can vary as a power of the underlying asset price. The [Marsh and Rosenfeld, 1983](#) short-term interest rate model

$$dr_t = (\beta r_t^{\gamma-1} + \alpha r_t)dt + \sigma r_t^{\gamma/2} dB_t, \quad (1.1.7)$$

where $\alpha \in \mathbb{R}$, $\beta, \sigma > 0$ are constants and $\gamma > 0$ is the variance (or diffusion) elasticity coefficient, covers most of the CEV models. Here, the elasticity coefficient is defined as ratio

$$\frac{dv^2(r)/v^2(r)}{dr/r}$$

between the relative change $dv(r)/v(r)$ in the variance $v(r)$ and the relative change dr/r in r . Denoting by $v^2(r) := \sigma^2 r^\gamma$ the variance coefficient in (1.1.7), constant elasticity refers to the constant ratio

$$\frac{dv^2(r)/v^2(r)}{dr/r} = 2 \frac{r}{v(r)} \frac{dv(r)}{dr} = 2 \frac{d \log v(r)}{d \log r} = 2 \frac{d \log r^{\gamma/2}}{d \log r} = \gamma.$$

For $\gamma = 1$, (1.1.7) yields the [Cox, Ingersoll, and Ross, 1985](#) (CIR) equation

$$dr_t = (\beta + \alpha r_t)dt + \sigma \sqrt{r_t} dB_t.$$

For $\beta = 0$ we get the standard CEV model

$$dr_t = \alpha r_t dt + \sigma r_t^{\gamma/2} dB_t,$$

and for $\gamma = 2$ and $\beta = 0$ this yields the [Dothan, 1978](#) model

$$dr_t = \alpha r_t dt + \sigma r_t dB_t,$$

which is a version of geometric Brownian motion used for short-term interest rate modeling.

Time-dependent affine models

The class of short rate interest rate models admits a number of generalizations (see the references quoted in the introduction of this chapter), including the class of affine models of the form

$$dr_t = (\eta(t) + \lambda(t)r_t)dt + \sqrt{\delta(t) + \gamma(t)r_t} dB_t. \quad (1.1.8)$$

Such models are called *affine* because the associated bonds can be priced using an *affine* PDE of the type (1.4.8) below with solution of the form (1.4.11), as will be seen after Proposition 1.2.

The family of affine models also includes:

- i) the [Ho and Lee, 1986](#) model

$$dr_t = \theta(t)dt + \sigma dB_t,$$

where $\theta(t)$ is a deterministic function of time, as an extension of the Merton model $dr_t = \theta dt + \sigma dB_t$,

- ii) the [Hull and White, 1990](#) model

$$dr_t = (\theta(t) - \alpha(t)r_t)dt + \sigma(t)dB_t$$

which is a time-dependent extension of the Vasicek model (1.1.1), with the explicit solution

$$r_t = r_0 e^{-\int_0^t \alpha(\tau)d\tau} + \int_0^t e^{-\int_u^t \alpha(\tau)d\tau} \theta(u)du + \int_0^t \sigma(u) e^{-\int_u^t \alpha(\tau)d\tau} dB_u,$$

$$t \geq 0.$$



1.2 Calibration of the Vasicek Model

The Vasicek equation (1.1.1), *i.e.*

$$dr_t = (a - br_t)dt + \sigma dB_t,$$

can be discretized according to a discrete-time sequence $(t_k)_{k \geq 0} = (t_0, t_1, t_2, \dots)$ of time instants, as

$$r_{t_{k+1}} - r_{t_k} = (a - br_{t_k})\Delta t + \sigma Z_k, \quad k \geq 0,$$

where $\Delta t := t_{k+1} - t_k$ and $(Z_k)_{k \geq 0}$ is a Gaussian white noise with variance Δt , *i.e.* a sequence of independent, centered and identically distributed $\mathcal{N}(0, \Delta t)$ Gaussian random variables, which yields

$$r_{t_{k+1}} = r_{t_k} + (a - br_{t_k})\Delta t + \sigma Z_k = a\Delta t + (1 - b\Delta t)r_{t_k} + \sigma Z_k, \quad k \geq 0.$$

Based on a set $(\tilde{r}_{t_k})_{k=0,1,\dots,n}$ of *market data*, we consider the quadratic residual

$$\sum_{k=0}^{n-1} (\tilde{r}_{t_{k+1}} - a\Delta t - (1 - b\Delta t)\tilde{r}_{t_k})^2 \tag{1.2.1}$$

which represents the (squared) quadratic distance between the observed data sequence $(\tilde{r}_{t_k})_{k=1,2,\dots,n}$ and its predictions $(a\Delta t + (1 - b\Delta t)\tilde{r}_{t_k})_{k=0,1,\dots,n-1}$.

In order to minimize the residual (1.2.1) over a and b we use Ordinary Least Square (OLS) regression, and equate the following derivatives to zero. Namely, we have

$$\begin{aligned} & \frac{\partial}{\partial a} \sum_{l=0}^{n-1} (\tilde{r}_{t_{l+1}} - a\Delta t - (1 - b\Delta t)\tilde{r}_{t_l})^2 \\ &= -2\Delta t \left(-an\Delta t + \sum_{l=0}^{n-1} (\tilde{r}_{t_{l+1}} - (1 - b\Delta t)\tilde{r}_{t_l}) \right) \\ &= 0, \end{aligned}$$

hence

$$a\Delta t = \frac{1}{n} \sum_{l=0}^{n-1} (\tilde{r}_{t_{l+1}} - (1 - b\Delta t)\tilde{r}_{t_l}),$$

and

$$\begin{aligned} & \frac{\partial}{\partial b} \sum_{k=0}^{n-1} (\tilde{r}_{t_{k+1}} - a\Delta t - (1 - b\Delta t)\tilde{r}_{t_k})^2 \\ &= 2\Delta t \sum_{k=0}^{n-1} \tilde{r}_{t_k} (-a\Delta t + \tilde{r}_{t_{k+1}} - (1 - b\Delta t)\tilde{r}_{t_k}) \\ &= 2\Delta t \sum_{k=0}^{n-1} \tilde{r}_{t_k} \left(\tilde{r}_{t_{k+1}} - (1 - b\Delta t)\tilde{r}_{t_k} - \frac{1}{n} \sum_{l=0}^{n-1} (\tilde{r}_{t_{l+1}} - (1 - b\Delta t)\tilde{r}_{t_l}) \right) \\ &= 2\Delta t \sum_{k=0}^{n-1} \tilde{r}_{t_k} \tilde{r}_{t_{k+1}} - \frac{\Delta t}{n} \sum_{k,l=0}^{n-1} \tilde{r}_{t_k} \tilde{r}_{t_{l+1}} - \Delta t(1 - b\Delta t) \left(\sum_{k=0}^{n-1} (\tilde{r}_{t_k})^2 - \frac{1}{n} \sum_{k,l=0}^{n-1} \tilde{r}_{t_k} \tilde{r}_{t_l} \right) \\ &= 0. \end{aligned}$$

This leads to estimators for the parameters a and b , respectively as the empirical mean and covariance of $(\tilde{r}_{t_k})_{k=0,1,\dots,n}$, i.e.

$$\left\{ \begin{array}{l} \hat{a}\Delta t = \frac{1}{n} \sum_{k=0}^{n-1} \left(\tilde{r}_{t_{k+1}} - (1 - \hat{b}\Delta t)\tilde{r}_{t_k} \right), \\ \text{and} \\ 1 - \hat{b}\Delta t = \frac{\sum_{k=0}^{n-1} \tilde{r}_{t_k} \tilde{r}_{t_{k+1}} - \frac{1}{n} \sum_{k,l=0}^{n-1} \tilde{r}_{t_k} \tilde{r}_{t_{l+1}}}{\sum_{k=0}^{n-1} (\tilde{r}_{t_k})^2 - \frac{1}{n} \sum_{k,l=0}^{n-1} \tilde{r}_{t_k} \tilde{r}_{t_l}} \\ = \frac{\sum_{k=0}^{n-1} \left(\tilde{r}_{t_k} - \frac{1}{n} \sum_{l=0}^{n-1} \tilde{r}_{t_l} \right) \left(\tilde{r}_{t_{k+1}} - \frac{1}{n} \sum_{l=0}^{n-1} \tilde{r}_{t_{l+1}} \right)}{\sum_{k=0}^{n-1} \left(\tilde{r}_{t_k} - \frac{1}{n} \sum_{k=0}^{n-1} \tilde{r}_{t_k} \right)^2}. \end{array} \right. \quad (1.2.2)$$

This also yields

$$\begin{aligned} \sigma^2 \Delta t &= \text{Var}[\sigma Z_k] \\ &\simeq \mathbb{E} [(\tilde{r}_{t_{k+1}} - (1 - b\Delta t)\tilde{r}_{t_k} - a\Delta t)^2], \quad k \geq 0, \end{aligned}$$

hence σ can be estimated as

$$\hat{\sigma}^2 \Delta t = \frac{1}{n} \sum_{k=0}^{n-1} (\tilde{r}_{t_{k+1}} - \tilde{r}_{t_k} (1 - \hat{b}\Delta t) - \hat{a}\Delta t)^2. \quad (1.2.3)$$

Exercise. Show that (1.2.3) can be recovered by minimizing the residual

$$\eta \mapsto \sum_{k=0}^{n-1} ((\tilde{r}_{t_{k+1}} - \tilde{r}_{t_k} (1 - \hat{b}\Delta t) - \hat{a}\Delta t)^2 - \eta \Delta t)^2$$

as a function of $\eta > 0$, see also Exercise 1.3.

Time series modeling

Defining $\hat{r}_{t_n} := r_{t_n} - a/b$, $n \geq 0$, we have

$$\begin{aligned} \hat{r}_{t_{n+1}} &= r_{t_{n+1}} - \frac{a}{b} \\ &= r_{t_n} - \frac{a}{b} + (a - br_{t_n})\Delta t + \sigma Z_n \\ &= r_{t_n} - \frac{a}{b} - b \left(r_{t_n} - \frac{a}{b} \right) \Delta t + \sigma Z_n \\ &= \hat{r}_{t_n} - b\hat{r}_{t_n}\Delta t + \sigma Z_n \\ &= (1 - b\Delta t)\hat{r}_{t_n} + \sigma Z_n, \quad n \geq 0. \end{aligned}$$

In other words, the sequence $(\hat{r}_{t_n})_{n \geq 0}$ is modeled according to an autoregressive AR(1) time series $(X_n)_{n \geq 0}$ with parameter $\alpha = 1 - b\Delta t$, in which the current state X_n of the system is expressed as the linear combination

$$X_n := \sigma Z_n + \alpha X_{n-1}, \quad n \geq 1, \quad (1.2.4)$$



where $(Z_n)_{n \geq 1}$ another Gaussian white noise sequence with variance Δt . This equation can be solved recursively as the causal series

$$X_n = \sigma Z_n + \alpha(\sigma Z_{n-1} + \alpha X_{n-2}) = \dots = \sigma \sum_{k \geq 0} \alpha^k Z_{n-k},$$

which converges when $|\alpha| < 1$, i.e. $|1 - b\Delta t| < 1$, in which case the time series $(X_n)_{n \geq 0}$ is weakly stationary, with

$$\begin{aligned}\mathbb{E}[X_n] &= \sigma \sum_{k \geq 0} \alpha^k \mathbb{E}[Z_{n-k}] \\ &= \sigma \mathbb{E}[Z_0] \sum_{k \geq 0} \alpha^k \\ &= \frac{\sigma}{1 - \alpha} \mathbb{E}[Z_0] \\ &= 0, \quad n \geq 0.\end{aligned}$$

The variance of X_n is given by

$$\begin{aligned}\text{Var}[X_n] &= \sigma^2 \text{Var} \left[\sum_{k \geq 0} \alpha^k Z_{n-k} \right] \\ &= \sigma^2 \Delta t \sum_{k \geq 0} \alpha^{2k} \\ &= \sigma^2 \Delta t \sum_{k \geq 0} (1 - b\Delta t)^{2k} \\ &= \frac{\sigma^2 \Delta t}{1 - (1 - b\Delta t)^2} \\ &= \frac{\sigma^2 \Delta t}{2b\Delta t - b^2(\Delta t)^2} \\ &\simeq \frac{\sigma^2}{2b}, \quad [\Delta t \simeq 0],\end{aligned}$$

which coincides with the variance (1.1.3) of the Vasicek process in the stationary regime.

Example - TNX yield calibration

The next R code is estimating the parameters of the Vasicek model using the 10 Year Treasury Note yield data of Figure 1.2, by implementing the formulas (1.2.2).

```
1 ratek=as.vector(ratek);ratekplus1 <- c(ratek[-1],0)
  b <- (sum(ratek*ratekplus1) - sum(ratek)*sum(ratekplus1)/n)/(sum(ratek*ratek) -
    sum(ratek)*sum(ratek)/n)
  3 a <- sum(ratekplus1)/n-b*sum(ratek)/n;sigma <- sqrt(sum((ratekplus1-b*ratek-a)^2)/n)
```

Parameter estimation can also be implemented using the linear regression command

```
lm(c(diff(ratek)) ~ ratek[1:length(ratek)-1])
```

in R, which estimates the values of $a\Delta t \simeq 0.017110$ and $-b\Delta t \simeq -0.007648$ in the regression

$$r_{t_{k+1}} - r_{t_k} = (a - br_{t_k})\Delta t + \sigma Z_k, \quad k \geq 0,$$

Coefficients:

```
(Intercept) ratek[1:length(ratek) - 1]
```

0.017110 -0.007648

```

1 dev.new(width=10,height=7);
2 for (i in 1:100) {ar.sim<-arima.sim(model=list(ar=c(b)),n.start=100,n)
3 y=a/b+sigma*ar.sim;y=y+ratek[1]-y[1]
4 time <- as.POSIXct(rate), format = "%Y-%m-%d")
5 yield <- xts(x = y, order.by = time);
6 chartSeries(yield,up.col="blue",theme="white",yrange=c(0,max(ratek)))
7 Sys.sleep(0.5)}
```

The above R code is generating Vasicek random samples according to the AR(1) time series (1.2.4), see Figure 1.4.




Figure 1.4: Calibrated Vasicek simulation vs market data.

1.3 Zero-Coupon and Coupon Bonds

A zero-coupon bond is a contract priced $P(t, T)$ at time $t < T$ to deliver the *face value* (or *par value*) $P(T, T) = \$1$ at time T . In addition to its value at maturity, a bond may yield a periodic *coupon* payment at regular time intervals until the maturity date.



Figure 1.5: Five-dollar 1875 Louisiana bond with 7.5% biannual coupons and maturity $T = 01/01/1886$.

The computation of the arbitrage-free price $P_0(t, T)$ of a zero-coupon bond based on an underlying short-term interest rate process $(r_t)_{t \in \mathbb{R}_+}$ is a basic and important issue in interest rate modeling.

Constant short rate

In case the short-term interest rate is a constant $r_t = r$, $t \geq 0$, a standard arbitrage argument shows that the price $P(t, T)$ of the bond is given by

$$P(t, T) = e^{-r(T-t)}, \quad 0 \leq t \leq T.$$

Indeed, if $P(t, T) > e^{-r(T-t)}$ we could issue a bond at the price $P(t, T)$ and invest this amount at the compounded risk free rate r , which would yield $P(t, T) e^{r(T-t)} > 1$ at time T .

On the other hand, if $P(t, T) < e^{-r(T-t)}$ we could borrow $P(t, T)$ at the rate r to buy a bond priced $P(t, T)$. At maturity time T we would receive \$1 and refund only $P(t, T) e^{r(T-t)} < 1$.

The price $P(t, T) = e^{-r(T-t)}$ of the bond is the value of $P(t, T)$ that makes the potential profit $P(t, T) e^{r(T-t)} - 1$ vanish for both traders.

Time-dependent deterministic short rates

Similarly to the above, when the short-term interest rate process $(r(t))_{t \in \mathbb{R}_+}$ is a deterministic function of time, a similar argument shows that

$$P(t, T) = e^{-\int_t^T r(s) ds}, \quad 0 \leq t \leq T. \quad (1.3.1)$$

Stochastic short rates

In case $(r_t)_{t \in \mathbb{R}_+}$ is an $(\mathcal{F}_t)_{t \in \mathbb{R}_+}$ -adapted random process the formula (1.3.1) is no longer valid as it relies on future information, and we replace it with the averaged discounted payoff

$$P(t, T) = \mathbb{E}^* \left[e^{-\int_t^T r_s ds} \mid \mathcal{F}_t \right], \quad 0 \leq t \leq T, \quad (1.3.2)$$

under a risk-neutral probability measure \mathbb{P}^* . It is natural to write $P(t, T)$ as a conditional expectation under a martingale measure, as the use of conditional expectation helps to “filter out” the (random/unknown) future information past time t contained in $\int_t^T r_s ds$. The expression (1.3.2) makes sense as the “best possible estimate” of the future quantity $e^{-\int_t^T r_s ds}$ in mean-square sense, given the information known up to time t .

Coupon bonds

Pricing bonds with nonzero coupon is not difficult since in general the amount and periodicity of coupons are deterministic.* In the case of a succession of coupon payments c_1, c_2, \dots, c_n at times $T_1, T_2, \dots, T_n \in (t, T]$, another application of the above absence of arbitrage argument shows that the price $P_c(t, T)$ of the coupon bond with discounted (deterministic) coupon payments is given by the linear combination of zero-coupon bond prices

$$\begin{aligned} P_c(t, T) &:= \mathbb{E}^* \left[\sum_{k=1}^n c_k e^{-\int_t^{T_k} r_s ds} \mid \mathcal{F}_t \right] + \mathbb{E}^* \left[e^{-\int_t^T r_s ds} \mid \mathcal{F}_t \right] \\ &= \sum_{k=1}^n c_k \mathbb{E}^* \left[e^{-\int_t^{T_k} r_s ds} \mid \mathcal{F}_t \right] + P_0(t, T) \\ &= P_0(t, T) + \sum_{k=1}^n c_k P_0(t, T_k), \quad 0 \leq t \leq T_1, \end{aligned} \quad (1.3.3)$$

which represents the present value at time t of future $\$c_1, \$c_2, \dots, \$c_n$ receipts respectively at times T_1, T_2, \dots, T_n , in addition to a terminal \$1 principal payment.

*However, coupon default cannot be excluded.

In the case of a constant coupon rate c paid at regular time intervals $\tau = T_{k+1} - T_k$, $k = 0, 1, \dots, n-1$, with $T_0 = t$, $T_n = T$, and a constant deterministic short rate r , we find

$$\begin{aligned} P_c(T_0, T_n) &= e^{-rn\tau} + c \sum_{k=1}^n e^{-(T_k - T_0)r} \\ &= e^{-rn\tau} + c \sum_{k=1}^n e^{-kr\tau} \\ &= e^{-rn\tau} + c \frac{e^{-r\tau} - e^{-r(n+1)\tau}}{1 - e^{-r\tau}}. \end{aligned}$$

In terms of the discrete-time interest rate $\tilde{r} := e^{r\tau} - 1$, we also have

$$P_c(T_0, T_n) = \frac{c}{\tilde{r}} + \frac{\tilde{r} - c}{(1 + \tilde{r})^n \tilde{r}}.$$




Figure 1.6: Discrete-time coupon bond pricing.

In the case of a continuous-time coupon rate $c > 0$, the above discrete-time calculation (1.3.3) can be reinterpreted as follows:

$$P_c(t, T) = P_0(t, T) + c \int_t^T P_0(t, u) du \quad (1.3.4)$$

$$\begin{aligned} &= P_0(t, T) + c \int_t^T e^{-(u-t)r} du \\ &= e^{-(T-t)r} + c \int_0^{T-t} e^{-ru} du \\ &= e^{-(T-t)r} + \frac{c}{r} (1 - e^{-(T-t)r}), \\ &= \frac{c}{r} + \frac{r-c}{r} e^{-(T-t)r}, \quad 0 \leq t \leq T, \end{aligned} \quad (1.3.5)$$

where the coupon bond price $P_c(t, T)$ solves the ordinary differential equation

$$dP_c(t, T) = (r - c) e^{-(T-t)r} dt = -cdt + rP_c(t, T)dt, \quad 0 \leq t \leq T,$$

see also Figures 1.7 and 1.11 below.




Figure 1.7: Continuous-time coupon bond pricing.

In the sequel, we will mostly consider zero-coupon bonds priced as $P(t, T) = P_0(t, T)$, $0 \leq t \leq T$, in the setting of stochastic short rates.

Martingale property of discounted bond prices

The following proposition shows that Assumption 1 of Chapter 3 is satisfied, in other words, the bond price process $t \mapsto P(t, T)$ can be used as a numéraire.

Proposition 1.1 The discounted bond price process

$$t \mapsto \tilde{P}(t, T) := e^{-\int_0^t r_s ds} P(t, T)$$

is a martingale under \mathbb{P}^* .

Proof. By (1.3.2) we have

$$\begin{aligned} \tilde{P}(t, T) &= e^{-\int_0^t r_s ds} P(t, T) \\ &= e^{-\int_0^t r_s ds} \mathbf{E}^* \left[e^{-\int_t^T r_s ds} \mid \mathcal{F}_t \right] \\ &= \mathbf{E}^* \left[e^{-\int_0^t r_s ds} e^{-\int_t^T r_s ds} \mid \mathcal{F}_t \right] \\ &= \mathbf{E}^* \left[e^{-\int_0^T r_s ds} \mid \mathcal{F}_t \right], \quad 0 \leq t \leq T, \end{aligned}$$

and this suffices in order to conclude, since by the tower property of conditional expectations, any process $(X_t)_{t \in \mathbb{R}_+}$ of the form $t \mapsto X_t := \mathbf{E}^*[F \mid \mathcal{F}_t]$, $F \in L^1(\Omega)$, is a martingale. In other words, we have

$$\begin{aligned} \mathbf{E}^* [\tilde{P}(t, T) \mid \mathcal{F}_u] &= \mathbf{E}^* \left[\mathbf{E}^* \left[e^{-\int_0^T r_s ds} \mid \mathcal{F}_t \right] \mid \mathcal{F}_u \right] \\ &= \mathbf{E}^* \left[e^{-\int_0^T r_s ds} \mid \mathcal{F}_u \right] \\ &= \tilde{P}(u, T), \quad 0 \leq u \leq t. \end{aligned}$$

□

1.4 Bond Pricing PDE

We assume from now on that the underlying short rate process solves the stochastic differential equation

$$dr_t = \mu(t, r_t)dt + \sigma(t, r_t)dB_t \quad (1.4.1)$$

where $(B_t)_{t \in \mathbb{R}_+}$ is a standard Brownian motion under \mathbb{P}^* . Note that specifying the dynamics of $(r_t)_{t \in \mathbb{R}_+}$ under the historical probability measure \mathbb{P} will also lead to a notion of market price of risk (MPoR) for the modeling of short rates.

As all solutions of stochastic differential equations such as (1.4.1) have the *Markov property*, cf. e.g. Theorem V-32 of [Protter, 2004](#), the arbitrage-free price $P(t, T)$ can be rewritten as a function $F(t, r_t)$ of r_t , i.e.

$$\begin{aligned} P(t, T) &= \mathbf{E}^* \left[e^{-\int_t^T r_s ds} \mid \mathcal{F}_t \right] \\ &= \mathbf{E}^* \left[e^{-\int_t^T r_s ds} \mid r_t \right] \\ &= F(t, r_t), \end{aligned} \quad (1.4.2)$$

and depends on (t, r_t) only, instead of depending on the whole information available in \mathcal{F}_t up to time t , meaning that the pricing problem can now be formulated as a search for the function $F(t, x)$.

Proposition 1.2 (Bond pricing PDE). Consider a short rate $(r_t)_{t \in \mathbb{R}_+}$ modeled by a diffusion equation of the form

$$dr_t = \mu(t, r_t)dt + \sigma(t, r_t)dB_t.$$

The bond pricing PDE for $P(t, T) = F(t, r_t)$ as in (1.4.2) is written as

$$xF(t, x) = \frac{\partial F}{\partial t}(t, x) + \mu(t, x) \frac{\partial F}{\partial x}(t, x) + \frac{1}{2} \sigma^2(t, x) \frac{\partial^2 F}{\partial x^2}(t, x), \quad (1.4.3)$$

$t \geq 0, x \in \mathbb{R}$, subject to the terminal condition

$$F(T, x) = 1, \quad x \in \mathbb{R}. \quad (1.4.4)$$

In addition, the bond price dynamics is given by

$$dP(t, T) = r_t P(t, T) dt + \sigma(t, r_t) \frac{\partial F}{\partial x}(t, r_t) dB_t. \quad (1.4.5)$$

Proof. By Itô's formula, we have

$$\begin{aligned} d \left(e^{-\int_0^t r_s ds} P(t, T) \right) &= -r_t e^{-\int_0^t r_s ds} P(t, T) dt + e^{-\int_0^t r_s ds} dP(t, T) \\ &= -r_t e^{-\int_0^t r_s ds} F(t, r_t) dt + e^{-\int_0^t r_s ds} dF(t, r_t) \\ &= -r_t e^{-\int_0^t r_s ds} F(t, r_t) dt + e^{-\int_0^t r_s ds} \frac{\partial F}{\partial x}(t, r_t) (\mu(t, r_t) dt + \sigma(t, r_t) dB_t) \\ &\quad + e^{-\int_0^t r_s ds} \left(\frac{1}{2} \sigma^2(t, r_t) \frac{\partial^2 F}{\partial x^2}(t, r_t) + \frac{\partial F}{\partial t}(t, r_t) \right) dt \end{aligned}$$



$$\begin{aligned}
&= -r_t e^{-\int_0^t r_s ds} F(t, r_t) dt + e^{-\int_0^t r_s ds} \frac{\partial F}{\partial x}(t, r_t) dr_t \\
&\quad + \frac{1}{2} e^{-\int_0^t r_s ds} \frac{\partial^2 F}{\partial x^2}(t, r_t) (dr_t)^2 + e^{-\int_0^t r_s ds} \frac{\partial F}{\partial t}(t, r_t) dt \\
&= e^{-\int_0^t r_s ds} \sigma(t, r_t) \frac{\partial F}{\partial x}(t, r_t) dB_t \\
&\quad + e^{-\int_0^t r_s ds} \left(-r_t F(t, r_t) + \mu(t, r_t) \frac{\partial F}{\partial x}(t, r_t) + \frac{1}{2} \sigma^2(t, r_t) \frac{\partial^2 F}{\partial x^2}(t, r_t) + \frac{\partial F}{\partial t}(t, r_t) \right) dt.
\end{aligned} \tag{1.4.6}$$

Given that $t \mapsto e^{-\int_0^t r_s ds} P(t, T)$ is a martingale, the above expression (1.4.6) should only contain terms in dB_t (cf. Corollary II-6-1, page 72 of Protter, 2004), and all terms in dt should vanish inside (1.4.6). This leads to the identities

$$\begin{cases} r_t F(t, r_t) = \mu(t, r_t) \frac{\partial F}{\partial x}(t, r_t) + \frac{1}{2} \sigma^2(t, r_t) \frac{\partial^2 F}{\partial x^2}(t, r_t) + \frac{\partial F}{\partial t}(t, r_t) \\ d \left(e^{-\int_0^t r_s ds} P(t, T) \right) = e^{-\int_0^t r_s ds} \sigma(t, r_t) \frac{\partial F}{\partial x}(t, r_t) dB_t, \end{cases} \tag{1.4.7a}$$

which lead to (1.4.3) and (1.4.5). Condition (1.4.4) is due to the fact that $P(T, T) = \$1$. \square

In the case of an interest rate process modeled by (1.1.8), we have

$$\mu(t, x) = \eta(t) + \lambda(t)x, \quad \text{and} \quad \sigma(t, x) = \sqrt{\delta(t) + \gamma(t)x},$$

hence (1.4.3) yields the *affine* PDE

$$xF(t, x) = \frac{\partial F}{\partial t}(t, x) + (\eta(t) + \lambda(t)x) \frac{\partial F}{\partial x}(t, x) + \frac{1}{2} (\delta(t) + \gamma(t)x) \frac{\partial^2 F}{\partial x^2}(t, x) \tag{1.4.8}$$

with time-dependent coefficients, $t \geq 0, x \in \mathbb{R}$. By (1.4.7a), the proof of Proposition 1.2 also shows that

$$\begin{aligned}
\frac{dP(t, T)}{P(t, T)} &= \frac{1}{P(t, T)} d \left(e^{\int_0^t r_s ds} e^{-\int_0^t r_s ds} P(t, T) \right) \\
&= \frac{1}{P(t, T)} \left(r_t P(t, T) dt + e^{\int_0^t r_s ds} d \left(e^{-\int_0^t r_s ds} P(t, T) \right) \right) \\
&= r_t dt + \frac{1}{P(t, T)} e^{\int_0^t r_s ds} d \left(e^{-\int_0^t r_s ds} P(t, T) \right) \\
&= r_t dt + \frac{1}{F(t, r_t)} \frac{\partial F}{\partial x}(t, r_t) \sigma(t, r_t) dB_t \\
&= r_t dt + \sigma(t, r_t) \frac{\partial}{\partial x} \log F(t, r_t) dB_t.
\end{aligned} \tag{1.4.9}$$

Vasicek, 1977 model

In the Vasicek case

$$dr_t = (a - br_t) dt + \sigma dB_t,$$

the bond price takes the form

$$F(t, r_t) = P(t, T) = e^{A(T-t) + r_t C(T-t)},$$

where $A(\cdot)$ and $C(\cdot)$ are deterministic functions of time, see (1.4.18) below, and (1.4.9) yields

$$\begin{aligned}\frac{dP(t, T)}{P(t, T)} &= r_t dt + \sigma C(T-t) dB_t \\ &= r_t dt - \frac{\sigma}{b} (1 - e^{-(T-t)b}) dB_t,\end{aligned}\tag{1.4.10}$$

since $F(t, x) = e^{A(T-t)+xC(T-t)}$.

Note that more generally, all affine short rate models as defined in Relation (1.1.8), including the Vasicek model, will yield a bond pricing formula of the form

$$P(t, T) = e^{A(T-t)+r_t C(T-t)},\tag{1.4.11}$$

cf. e.g. § 3.2.4. in [Brigo and Mercurio, 2006](#).

Probabilistic solution of the Vasicek PDE

Next, we solve the PDE (1.4.3), written with $\mu(t, x) = a - bx$ and $\sigma(t, x) = \sigma$ in the [Vašíček, 1977](#) model

$$dr_t = (a - br_t)dt + \sigma dB_t\tag{1.4.12}$$

as

$$\left\{ \begin{array}{l} xF(t, x) = \frac{\partial F}{\partial t}(t, x) + (a - bx)\frac{\partial F}{\partial x}(t, x) + \frac{\sigma^2}{2}\frac{\partial^2 F}{\partial x^2}(t, x), \\ F(T, x) = 1. \end{array} \right.\tag{1.4.13}$$

For this, Proposition 1.3 relies on a direct computation of the conditional expectation

$$F(t, r_t) = P(t, T) = \mathbb{E}^* \left[e^{-\int_t^T r_s ds} \mid \mathcal{F}_t \right].\tag{1.4.14}$$

See also Exercise 1.7 for a bond pricing formula in the [Cox, Ingersoll, and Ross, 1985](#) (CIR) model.

Proposition 1.3 The zero-coupon bond price in the Vasicek model (1.4.12) can be expressed as

$$P(t, T) = e^{A(T-t)+r_t C(T-t)}, \quad 0 \leq t \leq T,\tag{1.4.15}$$

where $A(x)$ and $C(x)$ are functions of time to maturity given by

$$C(x) := -\frac{1}{b} (1 - e^{-bx}),\tag{1.4.16}$$

and

$$\begin{aligned}A(x) &:= \frac{4ab - 3\sigma^2}{4b^3} + \frac{\sigma^2 - 2ab}{2b^2}x + \frac{\sigma^2 - ab}{b^3}e^{-bx} - \frac{\sigma^2}{4b^3}e^{-2bx} \\ &= -\left(\frac{a}{b} - \frac{\sigma^2}{2b^2}\right)(x + C(x)) - \frac{\sigma^2}{4b}C^2(x), \quad x \geq 0.\end{aligned}\tag{1.4.17}$$

Proof. Recall that in the Vasicek model (1.4.12), the short rate process $(r_t)_{t \in \mathbb{R}_+}$ solution of (1.4.12) has the expression

$$r_t = g(t) + \int_0^t h(t, s) dB_s = r_0 e^{-bt} + \frac{a}{b} (1 - e^{-bt}) + \sigma \int_0^t e^{-(t-s)b} dB_s,$$




where g and h are the deterministic functions

$$g(t) := r_0 e^{-bt} + \frac{a}{b}(1 - e^{-bt}), \quad t \geq 0,$$

and

$$h(t, s) := \sigma e^{-(t-s)b}, \quad 0 \leq s \leq t.$$

Using the fact that Wiener integrals are Gaussian random variables and the Gaussian moment generating function, and exchanging the order of integration between ds and du over $[t, T]$ according to the following picture,



we have

$$\begin{aligned} P(t, T) &= \mathbb{E}^* \left[e^{-\int_t^T r_s ds} \mid \mathcal{F}_t \right] \\ &= \mathbb{E}^* \left[e^{-\int_t^T (g(s) + \int_0^s h(s, u) dB_u) ds} \mid \mathcal{F}_t \right] \\ &= \exp \left(- \int_t^T g(s) ds \right) \mathbb{E}^* \left[e^{-\int_t^T \int_0^s h(s, u) dB_u ds} \mid \mathcal{F}_t \right] \\ &= \exp \left(- \int_t^T g(s) ds \right) \mathbb{E}^* \left[e^{-\int_0^T \int_{\max(u,t)}^T h(s, u) ds dB_u} \mid \mathcal{F}_t \right] \\ &= \exp \left(- \int_t^T g(s) ds - \int_0^t \int_{\max(u,t)}^T h(s, u) ds dB_u \right) \mathbb{E}^* \left[e^{-\int_t^T \int_{\max(u,t)}^T h(s, u) ds dB_u} \mid \mathcal{F}_t \right] \\ &= \exp \left(- \int_t^T g(s) ds - \int_0^t \int_t^T h(s, u) ds dB_u \right) \mathbb{E}^* \left[e^{-\int_t^T \int_u^T h(s, u) ds dB_u} \mid \mathcal{F}_t \right] \\ &= \exp \left(- \int_t^T g(s) ds - \int_0^t \int_t^T h(s, u) ds dB_u \right) \mathbb{E}^* \left[e^{-\int_t^T \int_u^T h(s, u) ds dB_u} \right] \\ &= \exp \left(- \int_t^T g(s) ds - \int_0^t \int_t^T h(s, u) ds dB_u + \frac{1}{2} \int_t^T \left(\int_u^T h(s, u) ds \right)^2 du \right) \\ &= \exp \left(- \int_t^T (r_0 e^{-bs} + \frac{a}{b}(1 - e^{-bs})) ds - \sigma \int_0^t \int_t^T e^{-(s-u)b} ds dB_u \right) \\ &\quad \times \exp \left(\frac{\sigma^2}{2} \int_t^T \left(\int_u^T e^{-(s-u)b} ds \right)^2 du \right) \\ &= \exp \left(- \int_t^T (r_0 e^{-bs} + \frac{a}{b}(1 - e^{-bs})) ds - \frac{\sigma}{b} (1 - e^{-(T-t)b}) \int_0^t e^{-(t-u)b} dB_u \right) \end{aligned}$$

$$\begin{aligned}
& \times \exp \left(\frac{\sigma^2}{2} \int_t^T e^{2bu} \left(\frac{e^{-bu} - e^{-bT}}{b} \right)^2 du \right) \\
& = \exp \left(-\frac{r_t}{b} (1 - e^{-(T-t)b}) + \frac{1}{b} (1 - e^{-(T-t)b}) \left(r_0 e^{-bt} + \frac{a}{b} (1 - e^{-bt}) \right) \right) \\
& \times \exp \left(- \int_t^T \left(r_0 e^{-bs} + \frac{a}{b} (1 - e^{-bs}) \right) ds + \frac{\sigma^2}{2} \int_t^T e^{2bu} \left(\frac{e^{-bu} - e^{-bT}}{b} \right)^2 du \right) \\
& = e^{A(T-t) + r_t C(T-t)}, \tag{1.4.18}
\end{aligned}$$

where $A(x)$ and $C(x)$ are the functions given by (1.4.16) and (1.4.17). \square

Analytical solution of the Vasicek PDE

In order to solve the PDE (1.4.13) analytically, we may start by looking for a solution of the form

$$F(t, x) = e^{A(T-t) + xC(T-t)}, \tag{1.4.19}$$

where $A(\cdot)$ and $C(\cdot)$ are functions to be determined under the conditions $A(0) = 0$ and $C(0) = 0$. Substituting (1.4.19) into the PDE (1.4.3) with the Vasicek coefficients $\mu(t, x) = (a - bx)$ and $\sigma(t, x) = \sigma$ shows that

$$\begin{aligned}
x e^{A(T-t) + xC(T-t)} &= -(A'(T-t) + xC'(T-t)) e^{A(T-t) + xC(T-t)} \\
&\quad + (a - bx) C(T-t) e^{A(T-t) + xC(T-t)} \\
&\quad + \frac{1}{2} \sigma^2 C^2(T-t) e^{A(T-t) + xC(T-t)},
\end{aligned}$$

i.e.

$$x = -A'(T-t) - xC'(T-t) + (a - bx) C(T-t) + \frac{1}{2} \sigma^2 C^2(T-t).$$

By identification of terms for $x = 0$ and $x \neq 0$, this yields the system of Riccati and linear differential equations

$$\begin{cases} A'(s) = aC(s) + \frac{\sigma^2}{2} C^2(s) \\ C'(s) = -1 - bC(s), \end{cases}$$

which can be solved to recover the above value of $P(t, T) = F(t, r_t)$ via

$$C(s) = -\frac{1}{b} (1 - e^{-bs})$$

and

$$\begin{aligned}
A(t) &= A(0) + \int_0^t A'(s) ds \\
&= \int_0^t \left(aC(s) + \frac{\sigma^2}{2} C^2(s) \right) ds \\
&= \int_0^t \left(\frac{a}{b} (1 - e^{-bs}) + \frac{\sigma^2}{2b^2} (1 - e^{-bs})^2 \right) ds \\
&= \frac{a}{b} \int_0^t (1 - e^{-bs}) ds + \frac{\sigma^2}{2b^2} \int_0^t (1 - e^{-bs})^2 ds \\
&= \frac{4ab - 3\sigma^2}{4b^3} + \frac{\sigma^2 - 2ab}{2b^2} t + \frac{\sigma^2 - ab}{b^3} e^{-bt} - \frac{\sigma^2}{4b^3} e^{-2bt}, \quad t \geq 0.
\end{aligned}$$



The next Figure 1.8 shows the output of the attached **R code** for the Monte Carlo and analytical estimation of Vasicek bond prices.




Figure 1.8: Comparison of Monte Carlo and analytical PDE solutions.

Vasicek bond price simulations

In this section we consider again the Vasicek model, in which the short rate $(r_t)_{t \in \mathbb{R}_+}$ is solution to (1.1.1). Figure 1.9 presents a random simulation of the zero-coupon bond price (1.4.15) in the Vasicek model with $\sigma = 10\%$, $r_0 = 2.96\%$, $b = 0.5$, and $a = 0.025$. The graph of the corresponding deterministic zero-coupon bond price with $r = r_0 = 2.96\%$ is also shown in Figure 1.9.




Figure 1.9: Graphs of $t \mapsto F(t, r_t) = P(t, T)$ vs $t \mapsto e^{-r_0(T-t)}$.

Figure 1.10 presents a random simulation of the coupon bond price (1.3.4) in the Vasicek model with $\sigma = 2\%$, $r_0 = 3.5\%$, $b = 0.5$, $a = 0.025$, and coupon rate $c = 5\%$. The graph of the corresponding deterministic coupon bond price (1.3.5) with $r = r_0 = 3.5\%$ is also shown in Figure 1.10.




Figure 1.10: Graph of $t \mapsto P_c(t, T)$ for a bond with a 5% coupon rate.

Figure 1.11 presents market price data for a coupon bond with coupon rate $c = 6.25\%$.




Figure 1.11: Bond price graph with maturity 01/18/08 and coupon rate 6.25%.

Zero-coupon bond price and yield data

The following zero-coupon public bond price [data](#) was downloaded from [EMMA](#) at the Municipal Securities Rulemaking Board.

ORANGE CNTY CALIF PENSION OBLIG CAP APPREC-TAXABLE-REF-SER A (CA)
 CUSIP: 68428LBB9
 Dated Date: 06/12/1996 (June 12, 1996)
 Maturity Date: 09/01/2016 (September 1st, 2016)
 Interest Rate: 0.0 %
 Principal Amount at Issuance: \$26,056,000
 Initial Offering Price: 19.465

```

1 library(quantmod);getwd()
2 bondprice <- read.table("bond_data_R.txt",col.names =
3   c("Date","HighPrice","LowPrice","HighYield","LowYield","Count","Amount"))
4 head(bondprice)
5 time <- as.POSIXct(bondprice$Date, format = "%Y-%m-%d")
6 price <- xts(x = bondprice$HighPrice, order.by = time)
7 yield <- xts(x = bondprice$HighYield, order.by = time)
8 dev.new(width=10,height=7);
9 pdf("orange_prices.pdf",width=8,height=3.5);
10 chartSeries(price,up.col="blue",theme="white")
11 dev.off()
12 pdf("orange_yield.pdf",width=8,height=3.5);
13 chartSeries(yield,up.col="blue",theme="white")
14 dev.off()

```

	Date	HighPrice	LowPrice	HighYield	LowYield	Count	Amount
1	2016-01-13	99.082	98.982	1.666	1.501	2	20000
2	2015-12-29	99.183	99.183	1.250	1.250	1	10000
3	2015-12-21	97.952	97.952	3.014	3.014	1	10000
4	2015-12-17	99.141	98.550	2.123	1.251	5	610000
5	2015-12-07	98.770	98.770	1.714	1.714	2	10000
6	2015-12-04	98.363	98.118	2.628	2.280	2	10000



Figure 1.12: Orange Cnty Calif bond prices.

The next Figure 1.13 plots the *bond yield* $y(t, T)$ defined as

$$y(t, T) = -\frac{\log P(t, T)}{T - t}, \quad \text{or} \quad P(t, T) = e^{-(T-t)y(t, T)}, \quad 0 \leq t \leq T.$$




Figure 1.13: Orange Cnty Calif bond yields.

Bond pricing in the Dothan model

In the [Dothan, 1978](#) model, the short-term interest rate process $(r_t)_{t \in \mathbb{R}_+}$ is modeled according to a geometric Brownian motion

$$dr_t = \mu r_t dt + \sigma r_t dB_t, \quad (1.4.20)$$

where the volatility $\sigma > 0$ and the drift $\mu \in \mathbb{R}$ are constant parameters and $(B_t)_{t \in \mathbb{R}_+}$ is a standard Brownian motion. In this model the short-term interest rate r_t remains always positive, while the proportional volatility term σr_t accounts for the sensitivity of the volatility of interest rate changes to the level of the rate r_t .

On the other hand, the Dothan model is the only lognormal short rate model that allows for an analytical formula for the zero-coupon bond price

$$P(t, T) = \mathbb{E}^* \left[e^{-\int_t^T r_s ds} \mid \mathcal{F}_t \right], \quad 0 \leq t \leq T.$$

For convenience of notation we let $p = 1 - 2\mu/\sigma^2$ and rewrite (1.4.20) as

$$dr_t = (1-p) \frac{\sigma^2}{2} r_t dt + \sigma r_t dB_t,$$

with solution

$$r_t = r_0 e^{\sigma B_t - p\sigma^2 t/2}, \quad t \geq 0. \quad (1.4.21)$$

By the Markov property of $(r_t)_{t \in \mathbb{R}_+}$, the bond price $P(t, T)$ is a function $F(t, r_t)$ of r_t and time $t \in [0, T]$:

$$P(t, T) = F(t, r_t) = \mathbb{E}^* \left[e^{-\int_t^T r_s ds} \mid r_t \right], \quad 0 \leq t \leq T. \quad (1.4.22)$$

By computation of the conditional expectation (1.4.22) we obtain the following result, cf. Proposition 1.2 of [C. Pintoux and N. Privault, 2011](#), where the function $\theta(v, t)$ is defined itself by an integral.



Proposition 1.4 The zero-coupon bond price $P(t, T) = F(t, r_t)$ is given for all $p \in \mathbb{R}$ by

$$\begin{aligned} & F(t, x) \\ &= e^{-\sigma^2 p^2(T-t)/8} \int_0^\infty \int_0^\infty e^{-ux} \exp\left(-2 \frac{(1+z^2)}{\sigma^2 u}\right) \theta\left(\frac{4z}{\sigma^2 u}, \frac{(T-t)\sigma^2}{4}\right) \frac{du}{u} \frac{dz}{z^{p+1}}, \\ & x > 0. \end{aligned} \tag{1.4.23}$$

Proof. By Proposition 2 in [Yor, 1992](#), the probability distribution of the time integral $\int_0^{T-t} e^{\sigma B_s - p\sigma^2 s/2} ds$ is given by

$$\begin{aligned} & \mathbb{P}\left(\int_0^{T-t} e^{\sigma B_s - p\sigma^2 s/2} ds \in dy\right) \\ &= \int_{-\infty}^\infty \mathbb{P}\left(\int_0^t e^{\sigma B_s - p\sigma^2 s/2} ds \in dy, B_t - p\sigma t/2 \in dz\right) \\ &= \frac{\sigma}{2} \int_{-\infty}^\infty e^{-p\sigma z/2 - p^2\sigma^2 t/8} \exp\left(-2 \frac{1+e^{\sigma z}}{\sigma^2 y}\right) \theta\left(\frac{4e^{\sigma z/2}}{\sigma^2 y}, \frac{\sigma^2 t}{4}\right) \frac{dy}{y} dz \\ &= e^{-(T-t)p^2\sigma^2/8} \int_0^\infty \exp\left(-2 \frac{1+z^2}{\sigma^2 y}\right) \theta\left(\frac{4z}{\sigma^2 y}, \frac{(T-t)\sigma^2}{4}\right) \frac{dz}{z^{p+1}} \frac{dy}{y}, \quad y > 0, \end{aligned}$$

where the exchange of integrals is justified by the Fubini theorem and the nonnegativity of integrands. Hence, by (1.4.21) we find

$$\begin{aligned} & F(t, r_t) = P(t, T) \\ &= \mathbf{E}^*\left[\exp\left(-\int_t^T r_s ds\right) \middle| \mathcal{F}_t\right] \\ &= \mathbf{E}^*\left[\exp\left(-r_t \int_t^T e^{\sigma(B_s - B_t) - \sigma^2 p(s-t)/2} ds\right) \middle| \mathcal{F}_t\right] \\ &= \mathbf{E}^*\left[\exp\left(-x \int_t^T e^{\sigma(B_s - B_t) - \sigma^2 p(s-t)/2} ds\right)\right]_{x=r_t} \\ &= \mathbf{E}^*\left[\exp\left(-x \int_0^{T-t} e^{\sigma B_s - \sigma^2 ps/2} ds\right)\right]_{x=r_t} \\ &= \int_0^\infty e^{-r_t y} \mathbb{P}\left(\int_0^{T-t} e^{\sigma B_s - p\sigma^2 s/2} ds \in dy\right) \\ &= e^{-(T-t)p^2\sigma^2/8} \int_0^\infty e^{-r_t y} \int_0^\infty \exp\left(-2 \frac{1+z^2}{\sigma^2 y}\right) \theta\left(\frac{4z}{\sigma^2 y}, \frac{(T-t)\sigma^2}{4}\right) \frac{dz}{z^{p+1}} \frac{dy}{y}. \end{aligned}$$

□

The zero-coupon bond price $P(t, T) = F(t, r_t)$ in the Dothan model can also be written for all $p \in \mathbb{R}$ as

$$\begin{aligned} & F(t, x) = \frac{(2x)^{p/2}}{2\pi^2 \sigma^p} \int_0^\infty u e^{-(p^2 + u^2)\sigma^2 t/8} \sinh(\pi u) \left|\Gamma\left(-\frac{p}{2} + i\frac{u}{2}\right)\right|^2 K_{iu}\left(\frac{\sqrt{8x}}{\sigma}\right) du \\ &+ \frac{(2x)^{p/2}}{\sigma^p} \sum_{k \geq 0} \frac{2(p-2k)^+}{k!(p-k)!} e^{\sigma^2 k(k-p)t/2} K_{p-2k}\left(\frac{\sqrt{8x}}{\sigma}\right), \quad x > 0, t > 0, \end{aligned}$$

cf. Corollary 2.2 of C. Pintoux and N. Privault, 2010, see also Privault and Uy, 2013 for numerical computations. Zero-coupon bond prices in the Dothan model can also be computed by the conditional expression

$$\mathbb{E} \left[\exp \left(- \int_0^T r_t dt \right) \right] = \int_0^\infty \mathbb{E} \left[\exp \left(- \int_0^T r_t dt \right) \middle| r_T = z \right] d\mathbb{P}(r_T \leq z), \quad (1.4.24)$$

where r_T has the lognormal distribution

$$d\mathbb{P}(r_T \leq z) = d\mathbb{P}(r_0 e^{\sigma B_T - p\sigma^2 T/2} \leq z) = \frac{1}{z\sqrt{2\pi\sigma^2 T}} e^{-(p\sigma^2 T/2 + \log(z/r_0))^2/(2\sigma^2 T)}.$$

In Proposition 1.5 we note that the conditional Laplace transform

$$\mathbb{E} \left[\exp \left(- \int_0^T r_t dt \right) \middle| r_T = z \right]$$

cf. (1.4.28) above, can be computed by a closed-form integral expression based on the modified Bessel function of the second kind

$$K_\zeta(z) := \frac{z^\zeta}{2^{\zeta+1}} \int_0^\infty \exp \left(-u - \frac{z^2}{4u} \right) \frac{du}{u^{\zeta+1}}, \quad \zeta \in \mathbb{R}, \quad z \in \mathbb{C}, \quad (1.4.25)$$

cf. e.g. Watson, 1995 page 183, provided that the real part $\Re(z^2)$ of $z^2 \in \mathbb{C}$ is positive.

Proposition 1.5 (Privault and Yu, 2016, Proposition 4.1). Taking $r_0 = 1$, for all $\lambda, z > 0$ we have

$$\begin{aligned} \mathbb{E} \left[\exp \left(-\lambda \int_0^T r_s ds \right) \middle| r_T = z \right] &= \frac{4e^{-\sigma^2 T/8}}{\pi^{3/2} \sigma^2 p(z)} \sqrt{\frac{\lambda}{T}} \\ &\times \int_0^\infty e^{2(\pi^2 - \xi^2)/(\sigma^2 T)} \sin \left(\frac{4\pi\xi}{\sigma^2 T} \right) \sinh(\xi) \frac{K_1(\sqrt{8\lambda} \sqrt{1+2\sqrt{z}\cosh\xi+z}/\sigma)}{\sqrt{1+2\sqrt{z}\cosh\xi+z}} d\xi. \end{aligned} \quad (1.4.26)$$

Note however that the numerical evaluation of (1.4.26) can fail for small values of $T > 0$, and for this reason the integral can be estimated by a gamma approximation as in (1.4.27) below. Under the gamma approximation we can approximate the conditional bond price on the Dothan short rate r_t as

$$\mathbb{E} \left[\exp \left(-\lambda \int_0^T r_t dt \right) \middle| r_T = z \right] \simeq (1 + \lambda \theta(z))^{-v(z)},$$

where the parameters $v(z)$ and $\theta(z)$ are determined by conditional moment fitting to a gamma distribution, as

$$\theta(z) := \frac{\text{Var}[\Lambda_T | S_T = z]}{\mathbb{E}[\Lambda_T | S_T = z]}, \quad v(z) := \frac{(\mathbb{E}[\Lambda_T | S_T = z])^2}{\text{Var}[\Lambda_T | S_T = z]} = \frac{\mathbb{E}[\Lambda_T | S_T = z]}{\theta},$$

cf. Privault and Yu, 2016, which yields

$$\mathbb{E} \left[\exp \left(-\lambda \int_0^T r_s ds \right) \right] \simeq \int_0^\infty (1 + \lambda \theta(z))^{-v(z)} d\mathbb{P}(r_T \leq z). \quad (1.4.27)$$






Figure 1.14: Fitting of a gamma probability density function.

The quantity $\theta(z)$ is also known in physics as the *Fano factor* or *dispersion index* that measures the dispersion of the probability distribution of Λ_T given that $S_T = z$. Figures 1.15 shows that the stratified gamma approximation (1.4.27) matches the Monte Carlo estimate, while the use of the integral expressions (1.4.24) and (1.4.26) leads to numerical instabilities.




Figure 1.15: Approximation of Dothan bond prices $t \mapsto F(t,x)$ with $\sigma = 0.3$ and $T = 10$.

Related computations for yield options in the Cox, Ingersoll, and Ross, 1985 (CIR) model can also be found in Prayoga and Privault, 2017.

Path integrals in option pricing

Let \hbar denote the Planck constant, and let $S(x(\cdot))$ denote the action functional given as

$$S(x(\cdot)) = \int_0^t L(x(s), \dot{x}(s), s) ds = \int_0^t \left(\frac{1}{2} m(\dot{x}(s))^2 - V(x(s)) \right) ds,$$

where $L(x(s), \dot{x}(s), s)$ is the Lagrangian

$$L(x(s), \dot{x}(s), s) := \frac{1}{2} m(\dot{x}(s))^2 - V(x(s)).$$

In physics, the Feynman path integral

$$\psi(y, t) := \int_{x(0)=x, x(t)=y} \mathcal{D}x(\cdot) \exp\left(\frac{i}{\hbar} S(x(\cdot))\right)$$

solves the Schrödinger equation

$$i\hbar \frac{\partial \psi}{\partial t}(x, t) = -\frac{\hbar^2}{2m} \frac{\partial^2 \psi}{\partial x^2}(x, t) + V(x(t))\psi(x, t).$$

After the Wick rotation $t \mapsto -it$, the function

$$\phi(y, t) := \int_{x(0)=x, x(t)=y} \mathcal{D}x(\cdot) \exp\left(-\frac{1}{\hbar} S(x(\cdot))\right)$$

solves the heat equation

$$\hbar \frac{\partial \phi}{\partial t}(x, t) = -\frac{\hbar^2}{2m} \frac{\partial^2 \phi}{\partial x^2}(x, t) + V(x(t))\phi(x, t).$$

By reformulating the action functional $S(x(\cdot))$ as

$$\begin{aligned} S(x(\cdot)) &= \int_0^t \left(\frac{1}{2} m(\dot{x}(s))^2 + V(x(s)) \right) ds \\ &\simeq \sum_{i=1}^N \left(\frac{(x(t_i) - x(t_{i-1}))^2}{2(\Delta t_i)^2} + V(x(t_{i-1})) \right) \Delta t_i, \end{aligned}$$

we can rewrite the Euclidean path integral as

$$\begin{aligned} \phi(y, t) &= \int_{x(0)=x, x(t)=y} \mathcal{D}x(\cdot) \exp\left(-\frac{1}{\hbar} S(x(\cdot))\right) \\ &= \int_{x(0)=x, x(t)=y} \mathcal{D}x(\cdot) \exp\left(-\frac{1}{2\hbar} \sum_{i=1}^N \frac{(x(t_i) - x(t_{i-1}))^2}{2\Delta t_i} - \frac{1}{\hbar} \sum_{i=1}^N V(x(t_{i-1}))\right) \\ &= \mathbb{E}^* \left[\exp\left(-\frac{1}{\hbar} \int_0^t V(B_s) ds\right) \middle| B_0 = x, B_t = y \right]. \end{aligned}$$

This type of path integral computation

$$\phi(y, t) = \mathbb{E}^* \left[\exp\left(-\int_0^t V(B_s) ds\right) \middle| B_0 = x, B_t = y \right]. \quad (1.4.28)$$

is particularly useful for bond pricing, as (1.4.28) can be interpreted as the price of a bond with short-term interest rate process $(r_t)_{t \in \mathbb{R}_+} := (V(B_t))_{t \in \mathbb{R}_+}$ conditionally to the value of the endpoint $B_t = y$, cf. (1.4.26) below. It can also be useful for exotic option pricing, and for risk management, see *e.g.* Kakushadze, 2015. The path integral (1.4.28) can be estimated either by closed-form expressions using Partial Differential Equations (PDEs) or probability densities, by approximations such as (conditional) Moment matching, or by Monte Carlo estimation, from the paths of a Brownian bridge as shown in Figure 1.16.

Figure 1.16: Brownian bridge.



Exercises

Exercise 1.1 We consider the stochastic differential equation

$$dr_t = (a - br_t)dt + \sigma dB_t, \quad (1.4.29)$$

where $a, \sigma \in \mathbb{R}$, $b > 0$.

- a) Show that the solution of (1.4.29) is

$$r_t = r_0 e^{-bt} + \frac{a}{b} (1 - e^{-bt}) + \sigma \int_0^t e^{-(t-s)b} dB_s, \quad t \geq 0. \quad (1.4.30)$$

- b) Show that the Gaussian $\mathcal{N}(a/b, \sigma^2/(2b))$ distribution is the *invariant* (or stationary) distribution of $(r_t)_{t \in \mathbb{R}_+}$.

Exercise 1.2 (Brody, Hughston, and Meier, 2018) In the mean-reverting Vasicek model (1.4.30) with $b > 0$, compute:

- i) The asymptotic bond yield, or exponential long rate of interest

$$r_\infty := - \lim_{T \rightarrow \infty} \frac{\log P(t, T)}{T - t}.$$

- ii) The long-bond return

$$L_t := \lim_{T \rightarrow \infty} \frac{P(t, T)}{P(0, T)}.$$

Hint: Start from $\log(P(t, T)/P(0, T))$.

Exercise 1.3 Consider the Chan-Karolyi-Longstaff-Sanders (CKLS) interest rate model (Chan et al., 1992) parametrized as

$$dr_t = (a - br_t)dt + \sigma r_t^\gamma dB_t,$$

and time-discretized as

$$\begin{aligned} r_{t_{k+1}} &= r_{t_k} + (a - br_{t_k})\Delta t + \sigma r_{t_k}^\gamma Z_k \\ &= a\Delta t + (1 - b\Delta t)r_{t_k} + \sigma r_{t_k}^\gamma Z_k, \quad k \geq 0, \end{aligned}$$

where $\Delta t := t_{k+1} - t_k$ and $(Z_k)_{k \geq 0}$ is an *i.i.d.* sequence of $\mathcal{N}(0, \Delta t)$ random variables. Assuming that $a, b, \gamma > 0$ are known, find an unbiased estimator $\hat{\sigma}^2$ for the variance coefficient σ^2 , based on a market data set $(\tilde{r}_{t_k})_{k=0,1,\dots,n}$.

Exercise 1.4 Let $(B_t)_{t \in \mathbb{R}_+}$ denote a standard Brownian motion started at 0 under the risk-neutral probability measure \mathbb{P}^* . We consider a short-term interest rate process $(r_t)_{t \in \mathbb{R}_+}$ in a Ho-Lee model with constant deterministic volatility, defined by

$$dr_t = adt + \sigma dB_t, \quad (1.4.31)$$

where $a \in \mathbb{R}$ and $\sigma > 0$. Let $P(t, T)$ denote the arbitrage-free price of a zero-coupon bond in this model:

$$P(t, T) = \mathbb{E}^* \left[\exp \left(- \int_t^T r_s ds \right) \middle| \mathcal{F}_t \right], \quad 0 \leq t \leq T. \quad (1.4.32)$$

- a) State the bond pricing PDE satisfied by the function $F(t, x)$ defined via

$$F(t, x) := \mathbf{E}^* \left[\exp \left(- \int_t^T r_s ds \right) \middle| r_t = x \right], \quad 0 \leq t \leq T.$$

- b) Compute the arbitrage-free price $F(t, r_t) = P(t, T)$ from its expression (1.4.32) as a conditional expectation.

Hint: One may use the *integration by parts* relation

$$\begin{aligned} \int_t^T B_s ds &= TB_T - tB_t - \int_t^T s dB_s \\ &= (T-t)B_t + T(B_T - B_t) - \int_t^T s dB_s \\ &= (T-t)B_t + \int_t^T (T-s) dB_s, \end{aligned}$$

and the Gaussian moment generating function $\mathbf{E}[e^{\lambda X}] = e^{\lambda^2 \eta^2 / 2}$ for $X \sim \mathcal{N}(0, \eta^2)$.

- c) Check that the function $F(t, x)$ computed in Question (b)) does satisfy the PDE derived in Question (a)).

Exercise 1.5 Consider the [Courtadon, 1982](#) model

$$dr_t = \beta(\alpha - r_t)dt + \sigma r_t dB_t, \quad (1.4.33)$$

where α, β, σ are nonnegative, which is a particular case of the Chan-Karolyi-Longstaff-Sanders (CKLS) model ([Chan et al., 1992](#)) with $\gamma = 1$. Show that the solution of (1.4.33) is given by

$$r_t = \alpha\beta \int_0^t \frac{S_u}{S_u} du + r_0 S_t, \quad t \geq 0, \quad (1.4.34)$$

where $(S_t)_{t \in \mathbb{R}_+}$ is the geometric Brownian motion solution of $dS_t = -\beta S_t dt + \sigma S_t dB_t$ with $S_0 = 1$.

Exercise 1.6 Consider the [Marsh and Rosenfeld, 1983](#) short-term interest rate model

$$dr_t = (\beta r_t^{\gamma-1} + \alpha r_t)dt + \sigma r_t^{\gamma/2} dB_t$$

where $\alpha \in \mathbb{R}$ and $\beta, \sigma, \gamma > 0$.

- a) Letting $R_t := r_t^{2-\gamma}$, $t \geq 0$, find the stochastic differential equation satisfied by $(R_t)_{t \in \mathbb{R}_+}$.
b) Given that the discounted bond price process is a martingale, derive the bond pricing PDE satisfied by the function $F(t, x)$ such that

$$F(t, r_t) = P(t, T) = \mathbf{E}^* \left[e^{-\int_t^T r_s ds} \middle| \mathcal{F}_t \right] = \mathbf{E}^* \left[e^{-\int_t^T r_s ds} \middle| r_t \right].$$

Exercise 1.7 Consider the [Cox, Ingersoll, and Ross, 1985](#) (CIR) process $(r_t)_{t \in \mathbb{R}_+}$ solution of

$$dr_t = -ar_t dt + \sigma \sqrt{r_t} dB_t,$$

where $a, \sigma > 0$ are constants $(B_t)_{t \in \mathbb{R}_+}$ is a standard Brownian motion started at 0.

- a) Write down the bond pricing PDE for the function $F(t, x)$ given by

$$F(t, x) := \mathbf{E}^* \left[\exp \left(- \int_t^T r_s ds \right) \middle| r_t = x \right], \quad 0 \leq t \leq T.$$

Hint: Use Itô calculus and the fact that the discounted bond price is a martingale.



- b) Show that the PDE of Question (a)) admits a solution of the form $F(t, x) = e^{A(T-t)+xC(T-t)}$ where the functions $A(s)$ and $C(s)$ satisfy ordinary differential equations to be also written down together with the values of $A(0)$ and $C(0)$.

Exercise 1.8 Convertible bonds. Consider an underlying asset price process $(S_t)_{t \in \mathbb{R}_+}$ given by

$$dS_t = rS_t dt + \sigma S_t dB_t^{(1)},$$

and a short-term interest rate process $(r_t)_{t \in \mathbb{R}_+}$ given by

$$dr_t = \gamma(t, r_t) dt + \eta(t, r_t) dB_t^{(2)},$$

where $(B_t^{(1)})_{t \in \mathbb{R}_+}$ and $(B_t^{(2)})_{t \in \mathbb{R}_+}$ are two correlated Brownian motions under the risk-neutral probability measure \mathbb{P}^* , with $dB_t^{(1)} \cdot dB_t^{(2)} = \rho dt$. A convertible bond is made of a corporate bond priced $P(t, T)$ at time $t \in [0, T]$, that can be exchanged into a quantity $\alpha > 0$ of the underlying company's stock priced S_τ at a future time τ , whichever has a higher value, where α is a *conversion rate*.

- a) Find the payoff of the convertible bond at time τ .
- b) Rewrite the convertible bond payoff at time τ as the linear combination of $P(\tau, T)$ and a call option payoff on S_τ , whose strike price is to be determined.
- c) Write down the convertible bond price at time $t \in [0, \tau]$ as a function $C(t, S_t, r_t)$ of the underlying asset price and interest rate, using a discounted conditional expectation, and show that the discounted corporate bond price

$$e^{-\int_0^t r_s ds} C(t, S_t, r_t), \quad t \in [0, \tau],$$

is a martingale.

- d) Write down $d(e^{-\int_0^t r_s ds} C(t, S_t, r_t))$ using the Itô formula and derive the pricing PDE satisfied by the function $C(t, x, y)$ together with its terminal condition.
- e) Taking the bond price $P(t, T)$ as a numéraire, price the convertible bond as a European option with strike price $K = 1$ on an underlying asset priced $Z_t := S_t / P(t, T)$, $t \in [0, \tau]$ under the forward measure $\widehat{\mathbb{P}}$ with maturity T .
- f) Assuming the bond price dynamics

$$dP(t, T) = r_t P(t, T) dt + \sigma_B(t) P(t, T) dB_t,$$

determine the dynamics of the process $(Z_t)_{t \in \mathbb{R}_+}$ under the forward measure $\widehat{\mathbb{P}}$.

- g) Assuming that $(Z_t)_{t \in \mathbb{R}_+}$ can be modeled as a geometric Brownian motion, price the convertible bond using the Black-Scholes formula.

Exercise 1.9 Bond duration. Compute the duration

$$D_c(0, n) := -\frac{1+r}{P_c(0, n)} \frac{\partial}{\partial r} P_c(0, n)$$

of a discrete-time coupon bond priced as

$$\begin{aligned} P_c(0, n) &= \frac{1}{(1+r)^n} + c \sum_{k=1}^n \frac{1}{(1+r)^k} \\ &= \frac{c}{r} + \left(1 - \frac{c}{r}\right) \frac{1}{(1+r)^n}, \end{aligned}$$

where $r > 0$, and $c \geq 0$ denotes the coupon rate. What happens when n becomes large?

Exercise 1.10 Let $(r_t)_{t \in \mathbb{R}_+}$ denote a short-term interest rate process. For any $T > 0$, let $P(t, T)$ denote the price at time $t \in [0, T]$ of a zero-coupon bond defined by the stochastic differential equation

$$\frac{dP(t, T)}{P(t, T)} = r_t dt + \sigma_t^T dB_t, \quad 0 \leq t \leq T, \quad (1.4.35)$$

under the terminal condition $P(T, T) = 1$, where $(\sigma_t^T)_{t \in [0, T]}$ is an adapted process. We define the forward measure $\hat{\mathbb{P}}$ by

$$\mathbb{E}^* \left[\frac{d\hat{\mathbb{P}}}{d\mathbb{P}^*} \mid \mathcal{F}_t \right] = \frac{P(t, T)}{P(0, T)} e^{-\int_0^t r_s ds}, \quad 0 \leq t \leq T.$$

Recall that

$$B_t^T := B_t - \int_0^t \sigma_s^T ds, \quad 0 \leq t \leq T,$$

is a standard Brownian motion under $\hat{\mathbb{P}}$.

- a) Solve the stochastic differential equation (1.4.35).
- b) Derive the stochastic differential equation satisfied by the discounted bond price process

$$t \mapsto e^{-\int_0^t r_s ds} P(t, T), \quad 0 \leq t \leq T,$$

and show that it is a martingale.

- c) Show that

$$\mathbb{E}^* \left[e^{-\int_0^T r_s ds} \mid \mathcal{F}_t \right] = e^{-\int_0^t r_s ds} P(t, T), \quad 0 \leq t \leq T.$$

- d) Show that

$$P(t, T) = \mathbb{E}^* \left[e^{-\int_t^T r_s ds} \mid \mathcal{F}_t \right], \quad 0 \leq t \leq T.$$

- e) Compute $P(t, S)/P(t, T)$, $0 \leq t \leq T$, show that it is a martingale under the forward measure $\hat{\mathbb{P}}$ with maturity T , and that

$$P(T, S) = \frac{P(t, S)}{P(t, T)} \exp \left(\int_t^T (\sigma_s^S - \sigma_s^T) dB_s^T - \frac{1}{2} \int_t^T (\sigma_s^S - \sigma_s^T)^2 ds \right).$$

- f) Assuming that $(\sigma_t^T)_{t \in [0, T]}$ and $(\sigma_t^S)_{t \in [0, S]}$ are deterministic functions of time, compute the price

$$\mathbb{E}^* \left[e^{-\int_t^T r_s ds} (P(T, S) - \kappa)^+ \mid \mathcal{F}_t \right] = P(t, T) \hat{\mathbb{E}} \left[(P(T, S) - \kappa)^+ \mid \mathcal{F}_t \right]$$

of a bond option with strike price κ .

Recall that if X is a Gaussian random variable with mean $m(t)$ and variance $v^2(t)$ given \mathcal{F}_t , we have

$$\begin{aligned} \mathbb{E}[(e^X - K)^+ \mid \mathcal{F}_t] &= e^{m(t) + v^2(t)/2} \Phi \left(\frac{v(t)}{2} + \frac{1}{v(t)} (m(t) + v^2(t)/2 - \log K) \right) \\ &\quad - K \Phi \left(-\frac{v(t)}{2} + \frac{1}{v(t)} (m(t) + v^2(t)/2 - \log K) \right) \end{aligned}$$

where $\Phi(x)$, $x \in \mathbb{R}$, denotes the Gaussian cumulative distribution function.



Exercise 1.11 Write down the bond pricing PDE for the function

$$F(t, x) = \mathbb{E}^* \left[e^{-\int_t^T r_s ds} \mid r_t = x \right]$$

and show that in case $\alpha = 0$ the corresponding bond price $P(t, T)$ equals

$$P(t, T) = e^{-r_t B(T-t)}, \quad 0 \leq t \leq T,$$

where

$$B(x) := \frac{2(e^{\gamma x} - 1)}{2\gamma + (\beta + \gamma)(e^{\gamma x} - 1)}, \quad x \in \mathbb{R},$$

with $\gamma = \sqrt{\beta^2 + 2\sigma^2}$.

Exercise 1.12 Consider a zero-coupon bond with prices $P(1, 2) = 91.74\%$ and $P(0, 2) = 83.40\%$ at times $t = 0$ and $t = 1$.

- a) Compute the corresponding yields $y_{0,1}$, $y_{0,2}$ and $y_{1,2}$ at times $t = 0$ and $t = 1$.
- b) Assume that \$0.1 coupons are paid at times $t = 1$ and $t = 2$. Price the corresponding coupon bond at times $t = 0$ and $t = 1$ using the yields y_0 and y_1 .

Exercise 1.13 Consider the Vasicek process $(r_t)_{t \in \mathbb{R}_+}$ solution of the equation

$$dr_t = (a - br_t)dt + \sigma dB_t. \quad (1.4.36)$$

- a) Consider the discretization

$$r_{t_{k+1}} := r_{t_k} + (a - br_{t_k})\Delta t \pm \sigma \sqrt{\Delta t}, \quad k = 0, 1, 2, \dots$$

of the equation (1.4.36) with $p(r_{t_0}) = p(r_{t_1}) = 1/2$ and

$$\mathbb{E}[\Delta r_{t_1}] = (a - br_{t_0})\Delta t + \sigma p(r_{t_0})\sqrt{\Delta t} - \sigma q(r_{t_0})\sqrt{\Delta t} = (a - br_{t_0})\Delta t$$

and

$$\mathbb{E}[\Delta r_{t_2}] = (a - br_{t_1})\Delta t + \sigma p(r_{t_0})\sqrt{\Delta t} - \sigma q(r_{t_0})\sqrt{\Delta t} = (a - br_{t_1})\Delta t.$$

Does this discretization lead to a (recombining) binomial tree?

- b) Using the Girsanov Theorem, find a probability measure \mathbb{Q} under which the process $(r_t/\sigma)_{t \in [0, T]}$ with

$$\frac{dr_t}{\sigma} = \frac{a - br_t}{\sigma} dt + dB_t$$

is a standard Brownian motion.

Hint: By the Girsanov Theorem, the process $X_t = X_0 + \int_0^t u_s ds + B_t$ is a martingale under the probability measure \mathbb{Q} with Radon-Nikodym density


$$\frac{d\mathbb{Q}}{d\mathbb{P}} = \exp \left(- \int_0^T u_t dB_t - \frac{1}{2} \int_0^T (u_t)^2 dt \right)$$

with respect to \mathbb{P} .

- c) Prove the Radon-Nikodym derivative approximation

$$\begin{aligned} & 2^{T/\Delta T} \prod_{0 < t < T} \left(\frac{1}{2} \pm \frac{a - br_t}{2\sigma} \sqrt{\Delta t} \right) \\ & \simeq \exp \left(\frac{1}{\sigma^2} \int_0^T (a - br_t) dr_t - \frac{1}{2\sigma^2} \int_0^T (a - br_t)^2 dt \right). \end{aligned} \quad (1.4.37)$$

d) Using (1.4.37), show that the Vasicek process can be discretized along the *binomial tree*



with

$$\mathbf{E}[\Delta r_{t_1} | r_{t_0}] = (a - br_{t_0})\Delta t \quad \text{and} \quad \mathbf{E}[\Delta r_{t_2} | r_{t_1}] = (a - br_{t_1})\Delta t,$$

where $\Delta r_{t_1} := r_{t_1} - r_{t_0}$, $\Delta r_{t_2} := r_{t_2} - r_{t_1}$, and the probabilities $p(r_{t_0}), q(r_{t_0}), p(r_{t_1}), q(r_{t_1})$ will be computed explicitly.

The use of binomial (or recombining) trees can make the implementation of the Monte Carlo method easier as their size grows linearly instead of exponentially.

Exercise 1.14 Black-Derman-Toy model (Black, Derman, and Toy, 1990). Consider a one-step interest rate model in which the short-term interest rate r_0 on $[0, 1]$ can turn into two possible values $r_1^u = r_0 e^{\mu\Delta t + \sigma\sqrt{\Delta t}}$ and $r_1^d = r_0 e^{\mu\Delta t - \sigma\sqrt{\Delta t}}$ on the time interval $[1, 2]$ with equal probabilities $1/2$ at time $\Delta t = 1$ year and volatility $\sigma = 22\%$ per year, and a zero-coupon bonds with prices $P(0, 1)$ and $P(0, 2)$ at time $t = 0$.

- a) Write down the value of $P(1, 2)$ using r_1^u and r_1^d .
- b) Write down the value of $P(0, 2)$ using r_1^u , r_1^d and r_0 .
- c) Estimate the value of r_0 from the market price $P(0, 1) = 91.74$.
- d) Estimate the values of r_1^u and r_1^d from the market price $P(0, 2) = 83.40$.

Exercise 1.15 Consider a yield curve $(f(t, t, T))_{0 \leq t \leq T}$ and a bond paying coupons c_1, c_2, \dots, c_n at times T_1, T_2, \dots, T_n until maturity T_n , and priced as

$$P(t, T_n) = \sum_{k=1}^n c_k e^{-(T_k - t)f(t, t, T_k)}, \quad 0 \leq t \leq T_1,$$

where c_n is inclusive of the last coupon payment and the nominal \$1 value of the bond. Let $\tilde{f}(t, t, T_n)$ denote the *compounded yield to maturity* defined by equating

$$P(t, T_n) = \sum_{k=1}^n c_k e^{-(T_k - t)\tilde{f}(t, t, T_n)}, \quad 0 \leq t \leq T_1, \tag{1.4.38}$$

i.e. $\tilde{f}(t, t, T_n)$ solves the equation

$$F(t, \tilde{f}(t, t, T_n)) = P(t, T_n), \quad 0 \leq t \leq T_1,$$



with

$$F(t, x) := \sum_{k=1}^n c_k e^{-(T_k - t)x}, \quad 0 \leq t \leq T_1.$$

The *bond duration* $D(t, T_n)$ is the relative sensitivity of $P(t, T_n)$ with respect to $\tilde{f}(t, t, T_n)$, defined as

$$D(t, T_n) := -\frac{1}{P(t, T_n)} \frac{\partial F}{\partial x}(t, \tilde{f}(t, t, T_n)), \quad 0 \leq t \leq T_1.$$

The *bond convexity* $C(t, T_n)$ is defined as

$$C(t, T_n) := \frac{1}{P(t, T_n)} \frac{\partial^2 F}{\partial x^2}(t, \tilde{f}(t, t, T_n)), \quad 0 \leq t \leq T_1.$$

- a) Compute the bond duration in case $n = 1$.
- b) Show that the *bond duration* $D(t, T_n)$ can be interpreted as an average of times to maturity weighted by the respective discounted bond payoffs.
- c) Show that the *bond convexity* $C(t, T_n)$ satisfies

$$C(t, T_n) = (D(t, T_n))^2 + (S(t, T_n))^2,$$

where

$$(S(t, T_n))^2 := \sum_{k=1}^n w_k (T_k - t - D(t, T_n))^2$$

measures the dispersion of the duration of the bond payoffs around the portfolio duration $D(t, T_n)$.

- d) Consider now the zero-coupon yield defined as

$$P(t, t + \alpha(T_n - t)) = \exp(-\alpha(T_n - t)f_\alpha(t, t, T_n)),$$

where $\alpha \in (0, 1)$, i.e.

$$f_\alpha(t, t, T_n) := -\frac{1}{\alpha(T_n - t)} \log P(t, t + \alpha(T_n - t)), \quad 0 \leq t \leq T_n.$$

Compute the bond duration associated to the yield $f_\alpha(t, t, T_n)$ in affine bond pricing models of the form

$$P(t, T) = e^{A(T-t) + r_t B(T-t)}, \quad 0 \leq t \leq T.$$

- e) (Wu, 2000) Compute the bond duration associated to the yield $f_\alpha(t, t, T_n)$ in the Vasicek model, in which

$$B(T-t) := \frac{1 - e^{-(T-t)b}}{b}, \quad 0 \leq t \leq T.$$

2. Forward Rate Modeling

Forward rates are interest rates used in Forward Rate Agreements (FRA) for financial transactions, such as loans, that can take place at a future date. This chapter deals with the modeling of forward rates and swap rates in the Heath-Jarrow-Morton (HJM) and Brace-Gatarek-Musiela (BGM) models. It also includes a presentation of the [Nelson and Siegel, 1987](#) and [Svensson, 1994](#) curve parametrizations for yield curve fitting, and an introduction to two-factor interest rate models.

2.1	Construction of Forward Rates	35
2.2	Interest Rate Swaps	44
2.3	The HJM Model	47
2.4	Yield Curve Modeling	52
2.5	Two-Factor Model	55
2.6	The BGM Model	59
	Exercises	61

2.1 Construction of Forward Rates

A forward interest rate contract (or Forward Rate Agreement, FRA) gives to its holder the possibility to lock an interest rate denoted by $f(t, T, S)$ at present time t for a loan to be delivered over a future period of time $[T, S]$, with $t \leq T \leq S$.



The rate $f(t, T, S)$ is called a forward interest rate. When $T = t$, the *spot* forward rate $f(t, t, T)$ coincides with the *yield*, see Relation (2.1.3) below.

Figure 2.1 presents a typical yield curve on the LIBOR (London Interbank Offered Rate) market with $t = 07$ May 2003.




Figure 2.1: Forward rate graph $S \mapsto f(t, t, S)$.

Maturity transformation, i.e., the ability to transform short-term borrowing (debt with short maturities, such as deposits) into long term lending (credits with very long maturities, such as loans), is among the roles of banks. Profitability is then dependent on the difference between long rates and short rates.

Another example of market data is given in the next Figure 2.2, in which the red and blue curves refer respectively to July 21 and 22 of year 2011.




Figure 2.2: Market example of yield curves, cf. (2.1.3).

Long maturities usually correspond to higher rates as they carry an increased risk. The dip observed with short maturities can correspond to a lower motivation to lend/invest in the short-term.

Forward rates from bond prices

Let us determine the arbitrage or “fair” value of the forward interest rate $f(t, T, S)$ by implementing the Forward Rate Agreement using the instruments available in the market, which are bonds priced at $P(t, T)$ for various maturity dates $T > t$.



The loan can be realized using the available instruments (here, bonds) on the market, by proceeding in two steps:

- 1) At time t , borrow the amount $P(t, S)$ by issuing (or short selling) one bond with maturity S , which means refunding \$1 at time S .
- 2) Since the money is only needed at time T , the rational investor will invest the amount $P(t, S)$ over the period $[t, T]$ by buying a (possibly fractional) quantity $P(t, S)/P(t, T)$ of a bond with maturity T priced $P(t, T)$ at time t . This will yield the amount

$$\$1 \times \frac{P(t, S)}{P(t, T)}$$

at time $T > 0$.

As a consequence, the investor will actually receive $P(t, S)/P(t, T)$ at time T , to refund \$1 at time S .

The corresponding forward rate $f(t, T, S)$ is then given by the relation

$$\frac{P(t, S)}{P(t, T)} \exp((S-T)f(t, T, S)) = \$1, \quad 0 \leq t \leq T \leq S, \quad (2.1.1)$$

where we used exponential compounding, which leads to the following definition (2.1.2).

Definition 2.1 The forward rate $f(t, T, S)$ at time t for a loan on $[T, S]$ is given by

$$f(t, T, S) = \frac{\log P(t, T) - \log P(t, S)}{S - T}. \quad (2.1.2)$$

The *spot* forward rate $f(t, t, S)$ coincides with the *yield* $y(t, S)$, with

$$f(t, t, S) = y(t, S) = -\frac{\log P(t, S)}{T - t}, \quad \text{or} \quad P(t, S) = e^{-(S-t)f(t, t, S)}, \quad (2.1.3)$$

$$0 \leq t \leq S.$$

Instantaneous forward rates

Proposition 2.2 The instantaneous forward rate $f(t, T) = f(t, T, T)$ is defined by taking the limit of $f(t, T, S)$ as $S \searrow T$, and satisfies

$$f(t, T) := \lim_{S \searrow T} f(t, T, S) = -\frac{1}{P(t, T)} \frac{\partial P}{\partial T}(t, T). \quad (2.1.4)$$

Proof. We have

$$\begin{aligned} f(t, T) : &= \lim_{S \searrow T} f(t, T, S) \\ &= -\lim_{S \searrow T} \frac{\log P(t, S) - \log P(t, T)}{S - T} \\ &= -\lim_{\varepsilon \searrow 0} \frac{\log P(t, T + \varepsilon) - \log P(t, T)}{\varepsilon} \end{aligned}$$

$$\begin{aligned}
&= -\frac{\partial}{\partial T} \log P(t, T) \\
&= -\frac{1}{P(t, T)} \frac{\partial P}{\partial T}(t, T).
\end{aligned}$$

□

The above equation (2.1.4) can be viewed as a differential equation to be solved for $\log P(t, T)$ under the initial condition $P(T, T) = 1$, which yields the following proposition.

Proposition 2.3 The bond price $P(t, T)$ can be recovered from the instantaneous forward rate $f(t, s)$ as

$$P(t, T) = \exp \left(- \int_t^T f(t, s) ds \right), \quad 0 \leq t \leq T. \quad (2.1.5)$$

Proof. We check that

$$\begin{aligned}
\log P(t, T) &= \log P(t, T) - \log P(t, t) \\
&= \int_t^T \frac{\partial}{\partial s} \log P(t, s) ds \\
&= - \int_t^T f(t, s) ds.
\end{aligned}$$

□

Proposition 2.3 also shows that

$$\begin{aligned}
f(t, t, t) &= f(t, t) \\
&= \frac{\partial}{\partial T} \int_t^T f(t, s) ds|_{T=t} \\
&= -\frac{\partial}{\partial T} \log P(t, T)|_{T=t} \\
&= -\frac{1}{P(t, T)}|_{T=t} \frac{\partial P}{\partial T}(t, T)|_{T=t} \\
&= -\frac{1}{P(T, T)} \frac{\partial}{\partial T} \mathbb{E}^* \left[e^{-\int_t^T r_s ds} \mid \mathcal{F}_t \right]_{|T=t} \\
&= \mathbb{E}^* \left[r_T e^{-\int_t^T r_s ds} \mid \mathcal{F}_t \right]_{|T=t} \\
&= \mathbb{E}^*[r_t \mid \mathcal{F}_t] \\
&= r_t,
\end{aligned} \quad (2.1.6)$$

i.e. the short rate r_t can be recovered from the instantaneous forward rate as

$$r_t = f(t, t) = \lim_{T \searrow t} f(t, T).$$

As a consequence of (2.1.1) and (2.1.5) the forward rate $f(t, T, S)$ can be recovered from (2.1.2) and the instantaneous forward rate $f(t, s)$, as:

$$\begin{aligned}
f(t, T, S) &= \frac{\log P(t, T) - \log P(t, S)}{S - T} \\
&= -\frac{1}{S - T} \left(\int_t^T f(t, s) ds - \int_t^S f(t, s) ds \right)
\end{aligned}$$



$$= \frac{1}{S-T} \int_T^S f(t,s) ds, \quad 0 \leq t \leq T < S. \quad (2.1.7)$$

Similarly, as a consequence of (2.1.3) and (2.1.5) we have the next proposition.

Proposition 2.4 The *spot forward rate* or *yield* $f(t,t,T)$ can be written in terms of bond prices as

$$f(t,t,T) = -\frac{\log P(t,T)}{T-t} = \frac{1}{T-t} \int_t^T f(t,s) ds, \quad 0 \leq t < T. \quad (2.1.8)$$

Differentiation with respect to T of the above relation shows that the yield $f(t,t,T)$ and the instantaneous forward rate $f(t,s)$ are linked by the relation

$$\frac{\partial f}{\partial T}(t,t,T) = -\frac{1}{(T-t)^2} \int_t^T f(t,s) ds + \frac{1}{T-t} f(t,T), \quad 0 \leq t < T,$$

from which it follows that

$$\begin{aligned} f(t,T) &= \frac{1}{T-t} \int_t^T f(t,s) ds + (T-t) \frac{\partial f}{\partial T}(t,t,T) \\ &= f(t,t,T) + (T-t) \frac{\partial f}{\partial T}(t,t,T), \quad 0 \leq t < T. \end{aligned}$$

Forward Vasicek, 1977 rates

In this section we consider the Vasicek model, in which the short rate process is the solution (1.1.2) of (1.1.1) as illustrated in Figure 1.1.

In the Vasicek model, the forward rate is given by

$$\begin{aligned} f(t,T,S) &= -\frac{\log P(t,S) - \log P(t,T)}{S-T} \\ &= -\frac{r_t(C(S-t) - C(T-t)) + A(S-t) - A(T-t)}{S-T} \\ &= -\frac{\sigma^2 - 2ab}{2b^2} \\ &\quad -\frac{1}{S-T} \left(\left(\frac{r_t}{b} + \frac{\sigma^2 - ab}{b^3} \right) (e^{-(S-t)b} - e^{-(T-t)b}) - \frac{\sigma^2}{4b^3} (e^{-2(S-t)b} - e^{-2(T-t)b}) \right), \end{aligned}$$

and the spot forward rate, or yield, satisfies

$$\begin{aligned} f(t,t,T) &= -\frac{\log P(t,T)}{T-t} = -\frac{r_t C(T-t) + A(T-t)}{T-t} \\ &= -\frac{\sigma^2 - 2ab}{2b^2} + \frac{1}{T-t} \left(\left(\frac{r_t}{b} + \frac{\sigma^2 - ab}{b^3} \right) (1 - e^{-(T-t)b}) - \frac{\sigma^2}{4b^3} (1 - e^{-2(T-t)b}) \right), \end{aligned}$$

with the mean

$$\begin{aligned} \mathbb{E}[f(t,t,T)] &= -\frac{\sigma^2 - 2ab}{2b^2} + \frac{1}{T-t} \left(\left(\mathbb{E}[r_t] + \frac{\sigma^2 - ab}{b^3} \right) (1 - e^{-(T-t)b}) - \frac{\sigma^2}{4b^3} (1 - e^{-2(T-t)b}) \right) \end{aligned}$$

$$\begin{aligned}
&= -\frac{\sigma^2 - 2ab}{2b^2} + \frac{1}{T-t} \left(\frac{r_0}{b} e^{-bt} + \frac{a}{b^2} (1 - e^{-bt}) + \frac{\sigma^2 - ab}{b^3} \right) (1 - e^{-(T-t)b}) \\
&\quad - \frac{\sigma^2}{4b^3(T-t)} (1 - e^{-2(T-t)b}).
\end{aligned}$$

In this model, the forward rate $t \mapsto f(t, T, S)$ can be represented as in the following Figure 2.3, with $a = 0.06$, $b = 0.1$, $\sigma = 0.1$ and $r_0 = \%1$.




Figure 2.3: Forward rate process $t \mapsto f(t, T, S)$.

We note that the Vasicek forward rate curve $t \mapsto f(t, T, S)$ appears flat for small values of t , i.e. longer rates are more stable, while shorter rates show higher volatility or risk. Similar features can be observed in Figure 2.4 for the instantaneous short rate given by


$$\begin{aligned}
f(t, T) : &= -\frac{\partial}{\partial T} \log P(t, T) \tag{2.1.9} \\
&= r_t e^{-(T-t)b} + \frac{a}{b} (1 - e^{-(T-t)b}) - \frac{\sigma^2}{2b^2} (1 - e^{-(T-t)b})^2,
\end{aligned}$$

from which the relation $\lim_{T \searrow t} f(t, T) = r_t$ can be easily recovered. We can also evaluate the mean

$$\begin{aligned}
\mathbf{E}[f(t, T)] &= \mathbf{E}[r_t] e^{-(T-t)b} + \frac{a}{b} (1 - e^{-(T-t)b}) - \frac{\sigma^2}{2b^2} (1 - e^{-(T-t)b})^2 \\
&= r_0 e^{-bT} + \frac{a}{b} (1 - e^{-bT}) - \frac{\sigma^2}{2b^2} (1 - e^{-(T-t)b})^2.
\end{aligned}$$

The instantaneous forward rate $t \mapsto f(t, T)$ can be represented as in the following Figure 2.4, with $a = 0.06$, $b = 0.1$, $\sigma = 0.1$ and $r_0 = \%1$.



Figure 2.4: Instantaneous forward rate process $t \mapsto f(t, T)$.

Yield curve data

We refer to Chapter III-12 of Charpentier, 2014 on the R package “YieldCurve” Guirrei, 2015 for the following R code and further details on yield curve and interest rate modeling using R.

```

1 install.packages("YieldCurve");require(YieldCurve);data(FedYieldCurve)
first(FedYieldCurve,'3 month');last(FedYieldCurve,'3 month')
3 mat.Fed=c(0.25,0.5,1,2,3,5,7,10);n=50
plot(mat.Fed, FedYieldCurve[,], type="o",xlab="Maturities structure in years", ylab="Interest rates
values", col = "blue", lwd=3)
5 title(main=paste("Federal Reserve yield curve observed at",time(FedYieldCurve[,], sep=" ")))
grid()

```

The next Figure 2.5 is plotted using this [code*](#) which is adapted from
<https://www.quantmod.com/examples/chartSeries3d/chartSeries3d.alpha.R>




Figure 2.5: Federal Reserve yield curves from 1982 to 2012.

European Central Bank (ECB) data can be similarly obtained by the next R code.

*Click to open or download.

```

1 data(ECBYieldCurve);first(ECBYieldCurve,'3 month');last(ECBYieldCurve,'3 month')
2 mat.ECB<-c(3/12,0.5,1,2,3,4,5,6,7,8,9,10,11,12,13,14,15,16,17,18,19,20,21,22,23, 24,25,26,27,28, 29,30)
3 dev.new(width=16,height=7)
4 for (n in 200:400) {
5   plot(mat.ECB, ECBYieldCurve[n,], type="o",xlab="Maturity structure in years", ylab="Interest rates
       values",ylim=c(3.1,5.1),col="blue",lwd=2,cex.axis=1.5,cex.lab=1.5)
6   title(main=paste("European Central Bank yield curve observed at",time(ECBYieldCurve[n], sep=" ")))
7   grid();Sys.sleep(0.5)}

```

The next Figure 2.6 represents the output of the above script.

Figure 2.6: European Central Bank yield curves.*

Yield curve inversion

Increasing yield curves are typical of economic expansion phases. Decreasing yield curves can occur when central banks attempt to limit inflation by tightening interest rates, such as in the case of an economic recession. In this case, uncertainty triggers increased investment in long bonds whose rates tend to drop as a consequence, while reluctance to lend in the short term can lead to higher short rates.

*The animation works in Acrobat Reader on the entire pdf file.



Figure 2.7: August 2019 Federal Reserve yield curve inversion.*

The above Figure 2.7 illustrates a Federal Reserve (FED) yield curve inversions occurring in February and August 2019.

LIBOR (London Interbank Offered) Rates

Recall that the forward rate $f(t, T, S)$, $0 \leq t \leq T \leq S$, is defined using exponential compounding, from the relation

$$f(t, T, S) = -\frac{\log P(t, S) - \log P(t, T)}{S - T}. \quad (2.1.10)$$

In order to compute swaption prices one prefers to use forward rates as defined on the London Inter-Bank Offered Rates (LIBOR) market instead of the standard forward rates given by (2.1.10). Other types of LIBOR rates include EURIBOR (European Interbank Offered Rates), HIBOR (Hong Kong Interbank Offered Rates), SHIBOR (Shanghai Interbank Offered Rates), SIBOR (Singapore Interbank Offered Rates), TIBOR (Tokyo Interbank Offered Rates), etc. LIBOR rates are to be replaced by alternatives, including the Secured Overnight Financing Rate (SOFR), by the end of year 2021. In the sequel, the term LIBOR refers to the linear or simple compounding used in the next definition.

The forward LIBOR rate $L(t, T, S)$ for a loan on $[T, S]$ is defined using linear compounding, *i.e.* by replacing (2.1.10) with the relation

$$1 + (S - T)L(t, T, S) = \frac{P(t, T)}{P(t, S)}, \quad 0 \leq t \leq T,$$

which yields the following definition.

Definition 2.5 The forward LIBOR rate $L(t, T, S)$ at time t for a loan on $[T, S]$ is given by

$$L(t, T, S) = \frac{1}{S - T} \left(\frac{P(t, T)}{P(t, S)} - 1 \right), \quad 0 \leq t \leq T < S. \quad (2.1.11)$$

Note that (2.1.11) above yields the same formula for the (LIBOR) instantaneous forward rate

$$L(t, T) : = \lim_{S \searrow T} L(t, T, S)$$

*The animation works in Acrobat Reader on the entire pdf file.

$$\begin{aligned}
&= \lim_{S \searrow T} \frac{P(t, T) - P(t, S)}{(S - T)P(t, S)} \\
&= \lim_{\varepsilon \searrow 0} \frac{P(t, T) - P(t, T + \varepsilon)}{\varepsilon P(t, T + \varepsilon)} \\
&= \frac{1}{P(t, T)} \lim_{\varepsilon \searrow 0} \frac{P(t, T) - P(t, T + \varepsilon)}{\varepsilon} \\
&= -\frac{1}{P(t, T)} \frac{\partial P}{\partial T}(t, T) \\
&= -\frac{\partial}{\partial T} \log P(t, T) \\
&= f(t, T),
\end{aligned}$$

as in (2.1.4).

In addition, Relation (2.1.11) shows that the LIBOR rate can be viewed as a forward price $\widehat{X}_t = X_t / N_t$ with numéraire $N_t = (S - T)P(t, S)$ and $X_t = P(t, T) - P(t, S)$, according to Relation (3.2.4) of Chapter 3. As a consequence, from Proposition 3.4, the LIBOR rate $(L(t, T, S))_{t \in [T, S]}$ is a martingale under the forward measure $\widehat{\mathbb{P}}$ defined by

$$\frac{d\widehat{\mathbb{P}}}{d\mathbb{P}^*} = \frac{1}{P(0, S)} e^{-\int_0^S r_t dt}.$$

2.2 Interest Rate Swaps

The first interest rate swap occurred in 1981 between the World Bank, which was interested in borrowing German Marks and Swiss Francs, and IBM, which already had large amounts of those currencies but needed to borrow U.S. dollars.

The vanilla interest rate swap makes it possible to exchange a sequence of variable LIBOR rates $L(T, T_k, T_{k+1})$, $k = 1, 2, \dots, n - 1$, against a fixed rate κ over a succession of time intervals $[T_1, T_2], [T_2, T_3], \dots, [T_{n-1}, T_n]$ defining a *tenor structure*, see Section 4.1 for details.

Making the agreement fair results into an exchange of cashflows

$$\underbrace{(T_{k+1} - T_k)L(T, T_k, T_{k+1})}_{\text{floating leg}} - \underbrace{(T_{k+1} - T_k)\kappa}_{\text{fixed leg}},$$

at the dates T_2, T_3, \dots, T_n between the two parties, generating a cumulative discounted cash flow

$$\sum_{k=1}^{n-1} e^{-\int_T^{T_{k+1}} r_s ds} (T_{k+1} - T_k)(L(T, T_k, T_{k+1}) - \kappa),$$

at time $T = T_0$, in which we used simple (or linear) interest rate compounding. This corresponds to a *payer swap* in which the swap holder receives the *floating leg* and pays the *fixed leg* κ , whereas the holder of a *seller swap* receives the *fixed leg* κ and pays the *floating leg*.

The above cash flow is used to make the contract fair, and it can be priced at time T as

$$\mathbb{E}^* \left[\sum_{k=1}^{n-1} (T_{k+1} - T_k) e^{-\int_T^{T_{k+1}} r_s ds} (L(T, T_k, T_{k+1}) - \kappa) \mid \mathcal{F}_T \right]$$



$$\begin{aligned}
&= \sum_{k=1}^{n-1} (T_{k+1} - T_k) (L(T, T_k, T_{k+1}) - \kappa) \mathbf{E}^* \left[e^{-\int_T^{T_{k+1}} r_s ds} \mid \mathcal{F}_T \right] \\
&= \sum_{k=1}^{n-1} (T_{k+1} - T_k) P(T, T_{k+1}) (L(T, T_k, T_{k+1}) - \kappa).
\end{aligned} \tag{2.2.1}$$

The swap rate $S(T, T_1, T_n)$ is by definition the value of the rate κ that makes the contract fair by making the above cash flow $\mathcal{C}(T)$ vanish.

Definition 2.6 The LIBOR swap rate $S(T, T_1, T_n)$ is the value of the break-even rate κ that makes the contract fair by making the cash flow (2.2.1) vanish, i.e.

$$\sum_{k=1}^{n-1} (T_{k+1} - T_k) P(T, T_{k+1}) (L(T, T_k, T_{k+1}) - \kappa) = 0. \tag{2.2.2}$$

The next Proposition 2.7 makes use of the annuity numéraire

$$\begin{aligned}
P(T, T_1, T_n) &:= \mathbf{E}^* \left[\sum_{k=1}^{n-1} (T_{k+1} - T_k) e^{-\int_T^{T_{k+1}} r_s ds} \mid \mathcal{F}_T \right] \\
&= \sum_{k=1}^{n-1} (T_{k+1} - T_k) \mathbf{E}^* \left[e^{-\int_T^{T_{k+1}} r_s ds} \mid \mathcal{F}_T \right] \\
&= \sum_{k=1}^{n-1} (T_{k+1} - T_k) P(T, T_{k+1}), \quad 0 \leq T \leq T_1,
\end{aligned} \tag{2.2.3}$$

which represents the present value at time T of future \$1 receipts at times T_1, T_2, \dots, T_n , weighted by the lengths $T_{k+1} - T_k$ of the time intervals $(T_k, T_{k+1}]$, $k = 1, 2, \dots, n-1$.

The time intervals $(T_{k+1} - T_k)_{k=1,2,\dots,n-1}$ in the definition (2.2.3) of the annuity numéraire can be replaced by coupon payments $(c_{k+1})_{k=1,2,\dots,n-1}$ occurring at times $(T_{k+1})_{k=1,2,\dots,n-1}$, in which case the annuity numéraire becomes

$$\begin{aligned}
P(T, T_1, T_n) &:= \mathbf{E}^* \left[\sum_{k=1}^{n-1} c_{k+1} e^{-\int_T^{T_{k+1}} r_s ds} \mid \mathcal{F}_T \right] \\
&= \sum_{k=1}^{n-1} c_{k+1} \mathbf{E}^* \left[e^{-\int_T^{T_{k+1}} r_s ds} \mid \mathcal{F}_T \right] \\
&= \sum_{k=1}^{n-1} c_{k+1} P(T, T_{k+1}), \quad 0 \leq T \leq T_1,
\end{aligned} \tag{2.2.4}$$

which represents the value at time T of the future coupon payments discounted according to the bond prices $(P(T, T_{k+1}))_{k=1,2,\dots,n-1}$. This expression can also be used to define *amortizing swaps* in which the value of the notional decreases over time, or *accruing swaps* in which the value of the notional increases over time.

LIBOR swap rates

The LIBOR swap rate $S(t, T_1, T_n)$ is defined by solving Relation (2.2.2) for the forward rate $S(t, T_k, T_{k+1})$, i.e.

$$\sum_{k=1}^{n-1} (T_{k+1} - T_k) P(T, T_{k+1}) (L(T, T_k, T_{k+1}) - S(t, T_1, T_n)) = 0. \tag{2.2.5}$$

Proposition 2.7 The LIBOR swap rate $S(T, T_1, T_n)$ is given by

$$S(T, T_1, T_n) = \frac{1}{P(T, T_1, T_n)} \sum_{k=1}^{n-1} (T_{k+1} - T_k) P(T, T_{k+1}) L(T, T_k, T_{k+1}). \quad (2.2.6)$$

Proof. By definition, $S(T, T_1, T_n)$ is the (fixed) break-even rate over $[T_1, T_n]$ that will be agreed in exchange for the family of forward rates $L(T, T_k, T_{k+1})$, $k = 1, 2, \dots, n-1$, and it solves (2.2.5), i.e. we have

$$\begin{aligned} & \sum_{k=1}^{n-1} (T_{k+1} - T_k) P(T, T_{k+1}) L(T, T_k, T_{k+1}) - P(T, T_1, T_n) S(T, T_1, T_n) \\ &= \sum_{k=1}^{n-1} (T_{k+1} - T_k) P(T, T_{k+1}) L(T, T_k, T_{k+1}) \\ &\quad - S(T, T_1, T_n) \sum_{k=1}^{n-1} (T_{k+1} - T_k) P(T, T_{k+1}) \\ &= \sum_{k=1}^{n-1} (T_{k+1} - T_k) P(T, T_{k+1}) L(T, T_k, T_{k+1}) - S(T, T_1, T_n) P(T, T_1, T_n) \\ &= 0, \end{aligned}$$

which shows (2.2.6) by solving the above equation for $S(T, T_1, T_n)$. \square

The LIBOR swap rate $S(t, T_1, T_n)$ is defined by the same relation as (2.2.2), with the forward rate $L(t, T_k, T_{k+1})$ replaced with the LIBOR rate $L(t, T_k, T_{k+1})$. In this case, using the Definition 2.1.11 of LIBOR rates we obtain the next corollary.

Corollary 2.8 The LIBOR swap rate $S(t, T_1, T_n)$ is given by

$$S(t, T_1, T_n) = \frac{P(t, T_1) - P(t, T_n)}{P(t, T_1, T_n)}, \quad 0 \leq t \leq T_1. \quad (2.2.7)$$

Proof. By (2.2.6), (2.1.11) and a telescoping summation argument we have

$$\begin{aligned} S(t, T_1, T_n) &= \frac{1}{P(t, T_1, T_n)} \sum_{k=1}^{n-1} (T_{k+1} - T_k) P(t, T_{k+1}) L(t, T_k, T_{k+1}) \\ &= \frac{1}{P(t, T_1, T_n)} \sum_{k=1}^{n-1} P(t, T_{k+1}) \left(\frac{P(t, T_k)}{P(t, T_{k+1})} - 1 \right) \\ &= \frac{1}{P(t, T_1, T_n)} \sum_{k=1}^{n-1} (P(t, T_k) - P(t, T_{k+1})) \\ &= \frac{P(t, T_1) - P(t, T_n)}{P(t, T_1, T_n)}. \end{aligned} \quad (2.2.8)$$

\square

By (2.2.7), the bond prices $P(t, T_1)$ can be recovered from the values of the forward swap rates $S(t, T_1, T_n)$.



Clearly, a simple expression for the swap rate such as that of Corollary 2.8 cannot be obtained using the standard (*i.e.* non-LIBOR) rates defined in (2.1.10). Similarly, it will not be available for amortizing or accreting swaps because the telescoping summation argument does not apply to the expression (2.2.4) of the annuity numéraire.

When $n = 2$, the LIBOR swap rate $S(t, T_1, T_2)$ coincides with the LIBOR rate $L(t, T_1, T_2)$, as from (2.2.4) we have

$$\begin{aligned} S(t, T_1, T_2) &= \frac{P(t, T_1) - P(t, T_2)}{P(t, T_1, T_2)} \\ &= \frac{P(t, T_1) - P(t, T_2)}{(T_2 - T_1)P(t, T_2)} \\ &= L(t, T_1, T_2). \end{aligned} \quad (2.2.9)$$

Similarly to the case of LIBOR rates, Relation (2.2.7) shows that the LIBOR swap rate can be viewed as a forward price with (annuity) numéraire $N_t = P(t, T_1, T_n)$ and $X_t = P(t, T_1) - P(t, T_n)$. Consequently the LIBOR swap rate $(S(t, T_1, T_n))_{t \in [T, S]}$ is a martingale under the forward measure $\hat{\mathbb{P}}$ defined from (3.2.1) by

$$\frac{d\hat{\mathbb{P}}}{d\mathbb{P}^*} = \frac{P(T_1, T_1, T_n)}{P(0, T_1, T_n)} e^{-\int_0^{T_1} r_t dt}.$$

2.3 The HJM Model

In this section we turn to the modeling of instantaneous forward rate curves. From the beginning of this chapter we have started with the modeling of the short rate $(r_t)_{t \in \mathbb{R}_+}$, followed by its consequences on the pricing of bonds $P(t, T)$ and on the expressions of the forward rates $f(t, T, S)$ and $L(t, T, S)$.

In this section we choose a different starting point and consider the problem of directly modeling the instantaneous forward rate $f(t, T)$. The graph given in Figure 2.8 presents a possible random evolution of a forward interest rate curve using the Musiela convention, *i.e.* we will write

$$g(x) = f(t, t + x) = f(t, T), \quad (2.3.1)$$

under the substitution $x = T - t$, $x \geq 0$, and represent a sample of the instantaneous forward curve $x \mapsto f(t, t + x)$ for each $t \geq 0$.




Figure 2.8: Stochastic process of forward curves.

Definition 2.9 In the Heath-Jarrow-Morton (HJM) model, the instantaneous forward rate $f(t, T)$ is modeled under \mathbb{P}^* by a stochastic differential equation of the form

$$d_t f(t, T) = \alpha(t, T) dt + \sigma(t, T) dB_t, \quad 0 \leq t \leq T, \quad (2.3.2)$$

where $t \mapsto \alpha(t, T)$ and $t \mapsto \sigma(t, T)$, $0 \leq t \leq T$, are allowed to be random (adapted) processes.

In the above equation, the date T is fixed and the differential d_t is with respect to the time variable t .

Under basic Markovianity assumptions, a HJM model with deterministic coefficients $\alpha(t, T)$ and $\sigma(t, T)$ will yield a short rate process $(r_t)_{t \in \mathbb{R}_+}$ of the form

$$dr_t = (a(t) - b(t)r_t)dt + \sigma(t)dB_t,$$

see § 6.6 in [Privault, 2012](#), which is the [Hull and White, 1990](#) model, with the explicit solution

$$r_t = r_s e^{-\int_s^t b(\tau)d\tau} + \int_s^t e^{-\int_u^t b(\tau)d\tau} a(u)du + \int_s^t \sigma(u) e^{-\int_u^t b(\tau)d\tau} dB_u,$$

$0 \leq s \leq t$.

The HJM condition

How to “encode” absence of arbitrage in the defining HJM Equation (2.3.2) is an important question. Recall that under absence of arbitrage, the bond price $P(t, T)$ has been constructed as

$$P(t, T) = \mathbb{E}^* \left[\exp \left(- \int_t^T r_s ds \right) \middle| \mathcal{F}_t \right] = \exp \left(- \int_t^T f(t, s) ds \right), \quad (2.3.3)$$

cf. Proposition 2.3, hence the discounted bond price process is given by

$$t \mapsto \exp \left(- \int_0^t r_s ds \right) P(t, T) = \exp \left(- \int_0^t r_s ds - \int_t^T f(t, s) ds \right) \quad (2.3.4)$$

is a martingale under \mathbb{P}^* by Proposition 1.1 and Relation (2.1.5) in Proposition 2.3. This shows that \mathbb{P}^* is a risk-neutral probability measure, and by the first fundamental theorem of asset pricing we conclude that the market is without arbitrage opportunities.

Proposition 2.10 (HJM Condition [Heath, Jarrow, and Morton, 1992](#)). Under the condition

$$\alpha(t, T) = \sigma(t, T) \int_t^T \sigma(t, s) ds, \quad 0 \leq t \leq T, \quad (2.3.5)$$

which is known as the *HJM absence of arbitrage condition*, the discounted bond price process (2.3.4) is a martingale, and the probability measure \mathbb{P}^* is risk-neutral.

Proof. Using the process $(X_t)_{t \in [0, T]}$ defined as

$$X_t := \int_t^T f(t, s) ds = -\log P(t, T), \quad 0 \leq t \leq T,$$

such that $P(t, T) = e^{-X_t}$, we rewrite the spot forward rate, or yield

$$f(t, t, T) = \frac{1}{T-t} \int_t^T f(t, s) ds,$$



see (2.1.8), as

$$f(t, t, T) = \frac{1}{T-t} \int_t^T f(t, s) ds = \frac{X_t}{T-t}, \quad 0 \leq t \leq T,$$

where the dynamics of $t \mapsto f(t, s)$ is given by (2.3.2). We also use the extended Leibniz integral rule

$$d_t \int_t^T f(t, s) ds = -f(t, t) dt + \int_t^T d_t f(t, s) ds = -r_t dt + \int_t^T d_t f(t, s) ds,$$

see (2.1.6). This identity can be checked in the particular case where $f(t, s) = g(t)h(s)$ is a smooth function that satisfies the separation of variables property, as

$$\begin{aligned} d_t \left(\int_t^T g(t)h(s) ds \right) &= d_t \left(g(t) \int_t^T h(s) ds \right) \\ &= \int_t^T h(s) ds dg(t) + g(t) d_t \int_t^T h(s) ds \\ &= g'(t) \left(\int_t^T h(s) ds \right) dt - g(t)h(t) dt. \end{aligned}$$

We have

$$\begin{aligned} d_t X_t &= d_t \int_t^T f(t, s) ds \\ &= -f(t, t) dt + \int_t^T d_t f(t, s) ds \\ &= -f(t, t) dt + \int_t^T \alpha(t, s) ds dt + \int_t^T \sigma(t, s) ds dB_t \\ &= -r_t dt + \left(\int_t^T \alpha(t, s) ds \right) dt + \left(\int_t^T \sigma(t, s) ds \right) dB_t, \end{aligned}$$

hence

$$|d_t X_t|^2 = \left(\int_t^T \sigma(t, s) ds \right)^2 dt.$$

By Itô's calculus, we find

$$\begin{aligned} d_t P(t, T) &= d_t e^{-X_t} \\ &= -e^{-X_t} d_t X_t + \frac{1}{2} e^{-X_t} (d_t X_t)^2 \\ &= -e^{-X_t} d_t X_t + \frac{1}{2} e^{-X_t} \left(\int_t^T \sigma(t, s) ds \right)^2 dt \\ &= -e^{-X_t} \left(-r_t dt + \int_t^T \alpha(t, s) ds dt + \int_t^T \sigma(t, s) ds dB_t \right) \\ &\quad + \frac{1}{2} e^{-X_t} \left(\int_t^T \sigma(t, s) ds \right)^2 dt, \end{aligned}$$

and the discounted bond price satisfies

$$\begin{aligned} d_t \left(\exp \left(- \int_0^t r_s ds \right) P(t, T) \right) &= -r_t \exp \left(- \int_0^t r_s ds - X_t \right) dt + \exp \left(- \int_0^t r_s ds \right) d_t P(t, T) \\ &= -r_t \exp \left(- \int_0^t r_s ds - X_t \right) dt - \exp \left(- \int_0^t r_s ds - X_t \right) d_t X_t \\ &\quad + \frac{1}{2} \exp \left(- \int_0^t r_s ds - X_t \right) \left(\int_t^T \sigma(t, s) ds \right)^2 dt \end{aligned}$$

$$\begin{aligned}
&= -r_t \exp \left(- \int_0^t r_s ds - X_t \right) dt \\
&\quad - \exp \left(- \int_0^t r_s ds - X_t \right) \left(-r_t dt + \int_t^T \alpha(t,s) ds dt + \int_t^T \sigma(t,s) ds dB_t \right) \\
&\quad + \frac{1}{2} \exp \left(- \int_0^t r_s ds - X_t \right) \left(\int_t^T \sigma(t,s) ds \right)^2 dt \\
&= -\exp \left(- \int_0^t r_s ds - X_t \right) \int_t^T \sigma(t,s) ds dB_t \\
&\quad - \exp \left(- \int_0^t r_s ds - X_t \right) \left(\int_t^T \alpha(t,s) ds - \frac{1}{2} \left(\int_t^T \sigma(t,s) ds \right)^2 \right) dt.
\end{aligned}$$

Thus, the discounted bond price process

$$t \mapsto \exp \left(- \int_0^t r_s ds \right) P(t,T)$$

will be a martingale provided that

$$\int_t^T \alpha(t,s) ds - \frac{1}{2} \left(\int_t^T \sigma(t,s) ds \right)^2 = 0, \quad 0 \leq t \leq T. \quad (2.3.6)$$

Differentiating the above relation with respect to T yields

$$\alpha(t,T) = \sigma(t,T) \int_t^T \sigma(t,s) ds,$$

which is in fact equivalent to (2.3.6). \square

Forward Vasicek rates in the HJM model

The HJM coefficients in the Vasicek model are in fact deterministic, for example, taking $a = 0$, by (2.1.9) we have

$$d_t f(t,T) = \sigma^2 e^{-(T-t)b} \int_t^T e^{(t-s)b} ds dt + \sigma e^{-(T-t)b} dB_t,$$

i.e.

$$\alpha(t,T) = \sigma^2 e^{-(T-t)b} \int_t^T e^{(t-s)b} ds = \sigma^2 e^{-(T-t)b} \frac{1 - e^{-(T-t)b}}{b},$$

and $\sigma(t,T) = \sigma e^{-(T-t)b}$, and the HJM condition reads

$$\alpha(t,T) = \sigma^2 e^{-(T-t)b} \int_t^T e^{(t-s)b} ds = \sigma(t,T) \int_t^T \sigma(t,s) ds. \quad (2.3.7)$$

Random simulations of the Vasicek instantaneous forward rates are provided in Figures 2.9 and 2.10 using the Musiela convention (2.3.1).

Figure 2.9: Forward instantaneous curve $(t, x) \mapsto f(t, t + x)$ in the Vasicek model.*

Figure 2.10: Forward instantaneous curve $x \mapsto f(0, x)$ in the Vasicek model.[†]

For $x = 0$ the first “slice” of this surface is actually the short rate Vasicek process $r_t = f(t, t) = f(t, t + 0)$ which is represented in Figure 2.11 using another discretization.

[†]The animation works in Acrobat Reader on the entire pdf file.




Figure 2.11: Short-term interest rate curve $t \mapsto r_t$ in the Vasicek model.

2.4 Yield Curve Modeling

Nelson-Siegel parametrization of instantaneous forward rates

In the [Nelson and Siegel, 1987](#) parametrization the instantaneous forward rate curves are parametrized by 4 coefficients z_1, z_2, z_3, z_4 , as

$$g(x) = z_1 + (z_2 + z_3 x) e^{-x z_4}, \quad x \geq 0.$$

An example of a graph obtained by the Nelson-Siegel parametrization is given in Figure 2.12, for $z_1 = 1, z_2 = -10, z_3 = 100, z_4 = 10$.




Figure 2.12: Graph of $x \mapsto g(x)$ in the Nelson-Siegel model.

Svensson parametrization of instantaneous forward rates

The [Svensson, 1994](#) parametrization has the advantage to reproduce two humps instead of one, the location and height of which can be chosen via 6 parameters $z_1, z_2, z_3, z_4, z_5, z_6$ as

$$g(x) = z_1 + (z_2 + z_3 x) e^{-x z_4} + z_5 x e^{-x z_6}, \quad x \geq 0.$$

A typical graph of a Svensson parametrization is given in Figure 2.13, for $z_1 = 6.6, z_2 = -5, z_3 = -100, z_4 = 10, z_5 = -1/2, z_6 = 1$.




Figure 2.13: Graph of $x \mapsto g(x)$ in the Svensson model.

Figure 2.14 presents a fit of the market data of Figure 2.1 using a Svensson curve.




Figure 2.14: Fitting of a Svensson curve to market data.

The attached [IPython notebook](#) that may be run [here](#) to fit a Svensson curve to market data.

Vasicek parametrization

In the Vasicek model, the instantaneous forward rate process is given from (2.1.9) and (2.3.1) as

$$f(t, T) = \frac{a}{b} - \frac{\sigma^2}{2b^2} + \left(r_t - \frac{a}{b} + \frac{\sigma^2}{b^2} \right) e^{-bx} - \frac{\sigma^2}{2b^2} e^{-2bx}, \quad (2.4.1)$$

in the Musiela notation ($x = T - t$), and we have

$$\frac{\partial f}{\partial T}(t, T) = \left(a - br_t - \frac{\sigma^2}{b} (1 - e^{-(T-t)b}) \right) e^{-(T-t)b}.$$

We check that the derivative $\partial f / \partial T$ vanishes when $a - br_t + a - \sigma^2(1 - e^{-bx})/b = 0$, i.e.

$$e^{-bx} = 1 + \frac{b}{\sigma^2} (br_t - a),$$

which admits at most one solution, provided that $a > br_t$. As a consequence, the possible forward curves in the Vasicek model are limited to one change of “regime” per curve, as illustrated in Figure 2.15 for various values of r_t , and in Figure 2.16.




Figure 2.15: Graphs of forward rates with $b = 0.16$, $a/b = 0.04$, $r_0 = 2\%$, $\sigma = 4.5\%$.

The next Figure 2.16 is also using the parameters $b = 0.16$, $a/b = 0.04$, $r_0 = 2\%$, and $\sigma = 4.5\%$.




Figure 2.16: Forward instantaneous curve $(t, x) \mapsto f(t, t + x)$ in the Vasicek model.

One may think of constructing an instantaneous forward rate process taking values in the Svensson space, however this type of modeling is not consistent with absence of arbitrage, and it can be proved that the HJM curves cannot live in the Nelson-Siegel or Svensson spaces, see §3.5 of [Björk, 2004](#). In other words, it can be shown that the forward yield curves produced by the Vasicek model are included neither in the Nelson-Siegel space, nor in the Svensson space. In addition, the Vasicek yield curves do not appear to correctly model the market forward curves cf. also Figure 2.1 above.

Another way to deal with the curve fitting problem is to use deterministic shifts for the fitting of one forward curve, such as the initial curve at $t = 0$, cf. e.g. § 8.2 in [Privault, 2012](#).

Fitting the Nelson-Siegel and Svensson models to yield curve data

Recall that in the Nelson-Siegel parametrization the instantaneous forward rate curves are parametrized by four coefficients z_1, z_2, z_3, z_4 , as

$$f(t, t + x) = z_1 + (z_2 + z_3 x) e^{-x z_4}, \quad x \geq 0. \quad (2.4.2)$$

Taking $x = T - t$, the yield $f(t, t, T)$ is given as

$$f(t, t, T) = \frac{1}{T - t} \int_t^T f(t, s) ds$$



$$\begin{aligned}
&= \frac{1}{x} \int_0^x f(t, t+y) dy \\
&= z_1 + \frac{z_2}{x} \int_0^x e^{-yz_4} dy + \frac{z_3}{x} \int_0^x y e^{-yz_4} dy \\
&= z_1 + z_2 \frac{1 - e^{-xz_4}}{xz_4} + z_3 \frac{1 - e^{-xz_4} + xe^{-xz_4}}{xz_4}.
\end{aligned}$$

The yield $f(t, t+T)$ can be reparametrized as

$$f(t, t+x) = z_1 + (z_2 + z_3 x) e^{-xz_4} = \beta_0 + \beta_1 e^{-x/\lambda} + \frac{\beta_2}{\lambda} x e^{-x/\lambda}, \quad x \geq 0,$$

cf. [Charpentier, 2014](#), with $\beta_0 = z_1$, $\beta_1 = z_2$, $\beta_2 = z_3/z_4$, $\lambda = 1/z_4$.

```

1 require(YieldCurve);data(ECBYieldCurve)
2 mat.ECB<-c(3/12,0.5,1,2,3,4,5,6,7,8,9,10,11,12,13,14,15,16,17,18,19,20,21,22,23, 24,25,26,27,28,29,30)
3 first(ECBYieldCurve, '1 month');Nelson.Siegel(first(ECBYieldCurve, '1 month'), mat.ECB)

```

```

1 for (n in seq(from=70, to=290, by=10)) {
2   ECB.NS <- Nelson.Siegel(ECBYieldCurve[n], mat.ECB)
3   ECB.S <- Svensson(ECBYieldCurve[n], mat.ECB)
4   ECB.NS.yield.curve <- NSrates(ECB.NS, mat.ECB)
5   ECB.S.yield.curve <- Srates(ECB.S, mat.ECB, "Spot")
6   plot(mat.ECB, as.numeric(ECBYieldCurve[n]), type="o", lty=1, col=1, ylab="Interest rates",
7       xlab="Maturity in years", ylim=c(3.2,4.8))
8   lines(mat.ECB, as.numeric(ECB.NS.yield.curve), type="l", lty=3, col=2, lwd=2)
9   lines(mat.ECB, as.numeric(ECB.S.yield.curve), type="l", lty=2, col=6, lwd=2)
10  title(main=paste("ECB yield curve observed at", time(ECBYieldCurve[n], sep=" "), "vs fitted yield curve"))
11  legend('bottomright', legend=c("ECB data", "Nelson-Siegel", "Svensson"), col=c(1,2,6), lty=1, bg='gray90')
12  grid();Sys.sleep(2.5)}

```

Figure 2.17: ECB data vs fitted yield curve.*

2.5 Two-Factor Model

The correlation problem is another issue of concern when using the affine models considered so far, see [\(1.1.8\)](#) and [\(1.4.11\)](#). Let us compare three bond price simulations with maturity $T_1 = 10$,

*The animation works in Acrobat Reader on the entire pdf file.

$T_2 = 20$, and $T_3 = 30$ based on the same Brownian path, as given in Figure 2.18. Clearly, the bond prices

$$F(r_t, T_i) = P(t, T_i) = e^{A(t, T_i)r_t C(t, T_i)}, \quad 0 \leq t \leq T_i, \quad i = 1, 2,$$

with maturities T_1 and T_2 are linked by the relation

$$P(t, T_2) = P(t, T_1) \exp(A(t, T_2) - A(t, T_1) + r_t(C(t, T_2) - C(t, T_1))), \quad (2.5.1)$$

meaning that bond prices with different maturities could be deduced from each other, which is unrealistic.




Figure 2.18: Graph of $t \mapsto P(t, T_1), P(t, T_2), P(t, T_3)$.

In affine short rate models, by (2.5.1), $\log P(t, T_1)$ and $\log P(t, T_2)$ are linked by the affine relationship

$$\begin{aligned} \log P(t, T_2) &= \log P(t, T_1) + A(t, T_2) - A(t, T_1) + r_t(C(t, T_2) - C(t, T_1)) \\ &= \log P(t, T_1) + A(t, T_2) - A(t, T_1) + (C(t, T_2) - C(t, T_1)) \frac{\log P(t, T_1) - A(t, T_1)}{C(t, T_1)} \\ &= \left(1 + \frac{C(t, T_2) - C(t, T_1)}{A(t, T_1)}\right) \log P(t, T_1) + A(t, T_2) - A(t, T_1) \frac{C(t, T_2)}{C(t, T_1)} \end{aligned}$$

with constant coefficients, which yields the perfect correlation or anticorrelation

$$\text{Cor}(\log P(t, T_1), \log P(t, T_2)) = \pm 1,$$

depending on the sign of the coefficient $1 + (C(t, T_2) - C(t, T_1))/A(t, T_1)$, cf. § 8.3 in [Privault, 2012](#),

A solution to the correlation problem is to consider a two-factor model based on two control processes $(X_t)_{t \in \mathbb{R}_+}, (Y_t)_{t \in \mathbb{R}_+}$ which are solution of

$$\begin{cases} dX_t = \mu_1(t, X_t)dt + \sigma_1(t, X_t)dB_t^{(1)}, \\ dY_t = \mu_2(t, Y_t)dt + \sigma_2(t, Y_t)dB_t^{(2)}, \end{cases} \quad (2.5.2)$$

where $(B_t^{(1)})_{t \in \mathbb{R}_+}, (B_t^{(2)})_{t \in \mathbb{R}_+}$ are correlated Brownian motion, with

$$\text{Cov}(B_s^{(1)}, B_t^{(2)}) = \rho \min(s, t), \quad s, t \geq 0, \quad (2.5.3)$$



and

$$dB_t^{(1)} \cdot dB_t^{(2)} = \rho dt, \quad (2.5.4)$$

for some correlation parameter $\rho \in [-1, 1]$. In practice, $(B^{(1)})_{t \in \mathbb{R}_+}$ and $(B^{(2)})_{t \in \mathbb{R}_+}$ can be constructed from two independent Brownian motions $(W^{(1)})_{t \in \mathbb{R}_+}$ and $(W^{(2)})_{t \in \mathbb{R}_+}$, by letting

$$\begin{cases} B_t^{(1)} = W_t^{(1)}, \\ B_t^{(2)} = \rho W_t^{(1)} + \sqrt{1 - \rho^2} W_t^{(2)}, \end{cases} \quad t \geq 0,$$

and Relations (2.5.3) and (2.5.4) are easily satisfied from this construction.

In two-factor models one chooses to build the short-term interest rate r_t via

$$r_t := X_t + Y_t, \quad t \geq 0.$$

By the previous standard arbitrage arguments we define the price of a bond with maturity T as

$$\begin{aligned} P(t, T) : &= \mathbf{E}^* \left[\exp \left(- \int_t^T r_s ds \right) \middle| \mathcal{F}_t \right] \\ &= \mathbf{E}^* \left[\exp \left(- \int_t^T r_s ds \right) \middle| X_t, Y_t \right] \\ &= \mathbf{E}^* \left[\exp \left(- \int_t^T (X_s + Y_s) ds \right) \middle| X_t, Y_t \right] \\ &= F(t, X_t, Y_t), \end{aligned} \quad (2.5.5)$$

since the couple $(X_t, Y_t)_{t \in \mathbb{R}_+}$ is Markovian. Applying the Itô formula with two variables to

$$t \mapsto F(t, X_t, Y_t) = P(t, T) = \mathbf{E}^* \left[\exp \left(- \int_t^T r_s ds \right) \middle| \mathcal{F}_t \right],$$

and using the fact that the discounted process

$$t \mapsto e^{-\int_0^t r_s ds} P(t, T) = \mathbf{E}^* \left[\exp \left(- \int_0^T r_s ds \right) \middle| \mathcal{F}_t \right]$$

is an \mathcal{F}_t -martingale under \mathbb{P}^* , we can derive a PDE

$$\begin{aligned} &-(x+y)F(t, x, y) + \mu_1(t, x) \frac{\partial F}{\partial x}(t, x, y) + \mu_2(t, y) \frac{\partial F}{\partial y}(t, x, y) \\ &+ \frac{1}{2} \sigma_1^2(t, x) \frac{\partial^2 F}{\partial x^2}(t, x, y) + \frac{1}{2} \sigma_2^2(t, y) \frac{\partial^2 F}{\partial y^2}(t, x, y) \\ &+ \rho \sigma_1(t, x) \sigma_2(t, y) \frac{\partial^2 F}{\partial x \partial y}(t, x, y) + \frac{\partial F}{\partial t}(t, x, y) = 0, \end{aligned} \quad (2.5.6)$$

on \mathbb{R}^2 for the bond price $P(t, T)$. In the Vasicek model

$$\begin{cases} dX_t = -aX_t dt + \sigma dB_t^{(1)}, \\ dY_t = -bY_t dt + \eta dB_t^{(2)}, \end{cases}$$

this yields the solution $F(t, x, y)$ of (2.5.6) as

$$P(t, T) = F(t, X_t, Y_t) = F_1(t, X_t)F_2(t, Y_t) \exp(\rho U(t, T)), \quad (2.5.7)$$

where $F_1(t, X_t)$ and $F_2(t, Y_t)$ are the bond prices associated to X_t and Y_t in the Vasicek model, and

$$U(t, T) := \frac{\sigma\eta}{ab} \left(T - t + \frac{e^{-(T-t)a} - 1}{a} + \frac{e^{-(T-t)b} - 1}{b} - \frac{e^{-(a+b)(T-t)} - 1}{a+b} \right)$$

is a correlation term which vanishes when $(B_t^{(1)})_{t \in \mathbb{R}_+}$ and $(B_t^{(2)})_{t \in \mathbb{R}_+}$ are independent, i.e. when $\rho = 0$, cf. Ch. 4, Appendix A in [Brigo and Mercurio, 2006](#), § 8.4 of [Privault, 2012](#).

Partial differentiation of $\log P(t, T)$ with respect to T leads to the instantaneous forward rate

$$f(t, T) = f_1(t, T) + f_2(t, T) - \rho \frac{\sigma\eta}{ab} (1 - e^{-(T-t)a})(1 - e^{-(T-t)b}), \quad (2.5.8)$$

where $f_1(t, T), f_2(t, T)$ are the instantaneous forward rates corresponding to X_t and Y_t respectively, cf. § 8.4 of [Privault, 2012](#).

An example of a forward rate curve obtained in this way is given in Figure 2.19.




Figure 2.19: Graph of forward rates in a two-factor model.

Next, in Figure 2.20 we present a graph of the evolution of forward curves in a two-factor model.




Figure 2.20: Random evolution of instantaneous forward rates in a two-factor model.



2.6 The BGM Model

The models (HJM, affine, etc.) considered in the previous chapter suffer from various drawbacks such as nonpositivity of interest rates in Vasicek model, and lack of closed-form solutions in more complex models. The [Brace, Gatarek, and Musiela, 1997](#) (BGM) model has the advantage of yielding positive interest rates, and to permit to derive explicit formulas for the computation of prices for interest rate derivatives such as interest rate caps and swaptions on the LIBOR market.

In the BGM model we consider two bond prices $P(t, T_1), P(t, T_2)$ with maturities T_1, T_2 , and the forward measure

$$\frac{d\mathbb{P}_2}{d\mathbb{P}_2^*} = \frac{e^{-\int_0^{T_2} r_s ds}}{P(0, T_2)},$$

with numéraire $P(t, T_2)$, cf. (3.2.10). The forward LIBOR rate $L(t, T_1, T_2)$ is modeled as a driftless geometric Brownian motion under \mathbb{P}_2 , *i.e.*

$$\frac{dL(t, T_1, T_2)}{L(t, T_1, T_2)} = \gamma_1(t) dB_t, \quad (2.6.1)$$

$0 \leq t \leq T_1$, $i = 1, 2, \dots, n - 1$, for some deterministic volatility function of time $\gamma_1(t)$, with solution

$$L(u, T_1, T_2) = L(t, T_1, T_2) \exp \left(\int_t^u \gamma_1(s) dB_s - \frac{1}{2} \int_t^u |\gamma_1|^2(s) ds \right),$$

i.e. for $u = T_1$,

$$L(T_1, T_1, T_2) = L(t, T_1, T_2) \exp \left(\int_t^{T_1} \gamma_1(s) dB_s - \frac{1}{2} \int_t^{T_1} |\gamma_1|^2(s) ds \right).$$


Since $L(t, T_1, T_2)$ is a geometric Brownian motion under \mathbb{P}_2 , standard caplets can be priced at time $t \in [0, T_1]$ from the Black-Scholes formula.

In the next Table 2.1 we summarize some stochastic models used for interest rates.

	Model
Short rate r_t	Mean reverting SDEs
Instantaneous forward rate $f(t, s)$	HJM model
Forward rate $f(t, T, S)$	BGM model

Table 2.1: Stochastic interest rate models.

The following Graph 2.21 summarizes the notions introduced in this chapter.



¹Can be modeled by Vasicek and other short rate models

²Can be modeled from $dP(t, T)/P(t, T)$.

³Can be modeled in the BGM model

⁴Can be modeled in the HJM model

Figure 2.21: Roadmap of stochastic interest rate modeling.



Exercises

Exercise 2.1 We consider a bond with maturity T , priced $P(t, T) = \mathbb{E}^* \left[e^{-\int_t^T r_s ds} \mid \mathcal{F}_t \right]$ at time $t \in [0, T]$.

- Using the forward measure $\hat{\mathbb{P}}$ with numéraire $N_t = P(t, T)$, apply the change of numéraire formula (3.2.9) to compute the derivative $\frac{\partial P}{\partial T}(t, T)$.
- Using Relation (2.1.5), find an expression of the instantaneous forward rate $f(t, T)$ using the short rate r_T and the forward expectation $\hat{\mathbb{E}}$.
- Show that the instantaneous forward rate $(f(t, T))_{t \in [0, T]}$ is a martingale under the forward measure $\hat{\mathbb{P}}$.

Exercise 2.2 Consider a tenor structure $\{T_1, T_2\}$ and a bond with maturity T_2 and price given at time $t \in [0, T_2]$ by

$$P(t, T_2) = \exp \left(- \int_t^{T_2} f(t, s) ds \right), \quad t \in [0, T_2],$$

where the instantaneous yield curve $f(t, s)$ is parametrized as

$$f(t, s) = r_1 \mathbb{1}_{[0, T_1]}(s) + r_2 \mathbb{1}_{[T_1, T_2]}(s), \quad t \leq s \leq T_2.$$

Find a formula to estimate the values of r_1 and r_2 from the data of $P(0, T_2)$ and $P(T_1, T_2)$.

Same question when $f(t, s)$ is parametrized as

$$f(t, s) = r_1 s \mathbb{1}_{[0, T_1]}(s) + (r_1 T_1 + (s - T_1) r_2) \mathbb{1}_{[T_1, T_2]}(s), \quad t \leq s \leq T_2.$$

Exercise 2.3 Bridge model. Assume that the price $P(t, T)$ of a zero-coupon bond is modeled as

$$P(t, T) = e^{-\mu(T-t)+X_t^T}, \quad t \in [0, T],$$

where $\mu > 0$.

- Show that the terminal condition $P(T, T) = 1$ is satisfied.
- Compute the forward rate

$$f(t, T, S) = -\frac{1}{S-T} (\log P(t, S) - \log P(t, T)).$$

- Compute the instantaneous forward rate

$$f(t, T) = -\lim_{S \searrow T} \frac{1}{S-T} (\log P(t, S) - \log P(t, T)).$$

- Show that the limit $\lim_{T \searrow t} f(t, T)$ does not exist in $L^2(\Omega)$.

- Show that $P(t, T)$ satisfies the stochastic differential equation

$$\frac{dP(t, T)}{P(t, T)} = \sigma dB_t + \frac{\sigma^2}{2} dt - \frac{\log P(t, T)}{T-t} dt, \quad t \in [0, T].$$

- Rewrite the equation of Question (e)) as

$$\frac{dP(t, T)}{P(t, T)} = \sigma dB_t + r_t^T dt, \quad t \in [0, T],$$

where $(r_t^T)_{t \in [0, T]}$ is a process to be determined.

g) Show that we have the expression

$$P(t, T) = \mathbf{E}^* \left[e^{-\int_t^T r_s^T ds} \mid \mathcal{F}_t \right], \quad 0 \leq t \leq T.$$

h) Compute the conditional Radon-Nikodym density

$$\mathbf{E}^* \left[\frac{d\mathbb{P}_T}{d\mathbb{P}^*} \mid \mathcal{F}_t \right] = \frac{P(t, T)}{P(0, T)} e^{-\int_0^t r_s^T ds}$$

of the forward measure \mathbb{P}_T with respect to \mathbb{P}^* .

i) Show that the process

$$\hat{B}_t := B_t - \sigma t, \quad 0 \leq t \leq T,$$

is a standard Brownian motion under \mathbb{P}_T .

j) Compute the dynamics of X_t^S and $P(t, S)$ under \mathbb{P}_T .

Hint: Show that

$$-\mu(S-T) + \sigma(S-T) \int_0^t \frac{1}{S-s} dB_s = \frac{S-T}{S-t} \log P(t, S).$$

k) Compute the bond option price

$$\mathbf{E}^* \left[e^{-\int_t^T r_s^T ds} (P(T, S) - K)^+ \mid \mathcal{F}_t \right] = P(t, T) \mathbf{E}_T [(P(T, S) - K)^+ \mid \mathcal{F}_t],$$

$$0 \leq t < T < S.$$

Hint: Given X a Gaussian random variable with mean m and variance v^2 given \mathcal{F}_t , we have:

$$\begin{aligned} \mathbf{E}[(e^X - \kappa)^+ \mid \mathcal{F}_t] &= e^{m+v^2/2} \Phi \left(\frac{1}{v}(m+v^2 - \log \kappa) \right) \\ &\quad - \kappa \Phi \left(\frac{1}{v}(m - \log \kappa) \right). \end{aligned}$$

Exercise 2.4 Consider a short rate process $(r_t)_{t \in \mathbb{R}_+}$ of the form $r_t = h(t) + X_t$, where $h(t)$ is a deterministic function of time and $(X_t)_{\mathbb{R}_+}$ is a Vasicek process started at $X_0 = 0$.

- a) Compute the price $P(0, T)$ at time $t = 0$ of a bond with maturity T , using $h(t)$ and the function $A(T)$ defined in (1.4.17) for the pricing of Vasicek bonds.
- b) Show how the function $h(t)$ can be estimated from the market data of the initial instantaneous forward rate curve $f(0, t)$.

Exercise 2.5

- a) Given two LIBOR spot rates $L(t, t, T)$ and $L(t, t, S)$, compute the corresponding LIBOR forward rate $L(t, T, S)$.
- b) Assuming that $L(t, t, T) = 2\%$, $L(t, t, S) = 2.5\%$ and $t = 0$, $T = 1$, $S = 2T = 2$, would you buy a LIBOR forward contract over $[T, 2T]$ with rate $L(0, T, 2T)$ if $L(T, T, 2T)$ remained at $L(T, T, 2T) = L(0, 0, T) = 2\%$?

Exercise 2.6 (Exercise 1.4 continued).

- a) Compute the forward rate $f(t, T, S)$ in the Ho-Lee model (1.4.31) with constant deterministic volatility.

In the next questions we take $a = 0$.

- b) Compute the instantaneous forward rate $f(t, T)$ in this model.
- c) Derive the stochastic equation satisfied by the instantaneous forward rate $f(t, T)$.



- d) Check that the HJM absence of arbitrage condition is satisfied in this equation.

Exercise 2.7 Consider the two-factor Vasicek model

$$\begin{cases} dX_t = -bX_t dt + \sigma dB_t^{(1)}, \\ dY_t = -bY_t dt + \sigma dB_t^{(2)}, \end{cases}$$

where $(B_t^{(1)})_{t \in \mathbb{R}_+}$, $(B_t^{(2)})_{t \in \mathbb{R}_+}$ are correlated Brownian motion such that $dB_t^{(1)} \cdot dB_t^{(2)} = \rho dt$, for $\rho \in [-1, 1]$.

- a) Write down the expressions of the short rates X_t and Y_t .

Hint: They can be found in Section 1.1.

- b) Compute the variances $\text{Var}[X_t]$, $\text{Var}[Y_t]$, and the covariance $\text{Cov}(X_t, Y_t)$.

Hint: The expressions of $\text{Var}[X_t]$ and $\text{Var}[Y_t]$ can be found in Section 1.1.

- c) Compute the covariance $\text{Cov}(\log P(t, T_1), \log P(t, T_2))$ for the two-factor bond prices

$$P(t, T_1) = F_1(t, X_t, T_1) F_2(t, Y_t, T_1) e^{\rho U(t, T_1)}$$

and

$$P(t, T_2) = F_1(t, X_t, T_2) F_2(t, Y_t, T_2) e^{\rho U(t, T_2)},$$

where

$$\log F_1(t, x, T) = C_1^T + x A_1^T \quad \text{and} \quad \log F_2(t, x, T) = C_2^T + x A_2^T.$$

Hint: We have $\text{Cov}(X + Y, Z) = \text{Cov}(X, Z) + \text{Cov}(Y, Z)$ and $\text{Cov}(c, X) = 0$ when c is a constant.

Exercise 2.8 Stochastic string model (Santa-Clara and Sornette, 2001). Consider an instantaneous forward rate $f(t, x)$ solution of

$$d_t f(t, x) = \alpha x^2 dt + \sigma d_t B(t, x), \tag{2.6.2}$$

with a flat initial curve $f(0, x) = r$, where x represents the time to maturity, and $(B(t, x))_{(t, x) \in \mathbb{R}_+^2}$ is a standard *Brownian sheet* with covariance

$$\mathbb{E}[B(s, x)B(t, y)] = (\min(s, t))(\min(x, y)), \quad s, t, x, y \geq 0, \tag{2.6.3}$$

and initial conditions $B(t, 0) = B(0, x) = 0$ for all $t, x \geq 0$.

- a) Solve the equation (2.6.2) for $f(t, x)$.
b) Compute the short-term interest rate $r_t = f(t, 0)$.
c) Compute the value at time $t \in [0, T]$ of the bond price

$$P(t, T) = \exp\left(-\int_0^{T-t} f(t, x) dx\right)$$

with maturity T .

- d) Compute the variance $\mathbb{E}\left[\left(\int_0^{T-t} B(t, x) dx\right)^2\right]$ of the centered Gaussian random variable $\int_0^{T-t} B(t, x) dx$.
e) Compute the expected value $\mathbb{E}^*[P(t, T)]$.

- f) Find the value of α such that the discounted bond price

$$e^{-rt} P(t, T) = \exp\left(-rT - \frac{\alpha}{3}t(T-t)^3 - \sigma \int_0^{T-t} B(t,x)dx\right), \quad t \in [0, T].$$

satisfies $E^*[P(t, T)] = e^{-(T-t)r}$.

- g) Compute the bond option price $E^*\left[\exp\left(-\int_0^T r_s ds\right) (P(T, S) - K)^+\right]$ by the Black-Scholes formula, knowing that for any centered Gaussian random variable $X \simeq \mathcal{N}(0, v^2)$ with variance v^2 we have

$$\begin{aligned} E[(x e^{m+X} - K)^+] \\ = x e^{m+v^2/2} \Phi(m + \log(x/K))/v - K \Phi((m + \log(x/K))/v). \end{aligned}$$



3. Change of Numéraire and Forward Measures

Change of numéraire is a powerful technique for the pricing of options under random discount factors by the use of forward measures. It has applications to the pricing of interest rate derivatives and other types of options, including exchange options (Margrabe formula) and foreign exchange options (Garman-Kohlagen formula). The computation of self-financing hedging strategies by change of numéraire is treated in Section 3.5, and the change of numéraire technique will be applied to the pricing of interest rate derivatives such as bond options and swaptions in Chapter 4.

3.1	Notion of Numéraire	65
3.2	Change of Numéraire	68
3.3	Foreign Exchange	76
3.4	Pricing Exchange Options	83
3.5	Hedging by Change of Numéraire	85
	Exercises	88

3.1 Notion of Numéraire

A *numéraire* is any strictly positive $(\mathcal{F}_t)_{t \in \mathbb{R}_+}$ -adapted stochastic process $(N_t)_{t \in \mathbb{R}_+}$ that can be taken as a unit of reference when pricing an asset or a claim.

In general, the price S_t of an asset, when quoted in terms of the numéraire N_t , is given by

$$\widehat{S}_t := \frac{S_t}{N_t}, \quad t \geq 0.$$

Deterministic numéraires transformations are easy to handle, as change of numéraire by a constant factor is a formal algebraic transformation that does not involve any risk. This can be the case for

example when a currency is pegged to another currency,* for example the exchange rate of the French franc to the Euro was locked at €1 = FRF 6.55957 on 31 December 1998.

On the other hand, a random numéraire may involve new risks, and can allow for arbitrage opportunities.

Examples of numéraire processes $(N_t)_{t \in \mathbb{R}_+}$ include:

- *Money market account.*

Given $(r_t)_{t \in \mathbb{R}_+}$ a possibly random, time-dependent and $(\mathcal{F}_t)_{t \in \mathbb{R}_+}$ -adapted risk-free interest rate process, let[†]

$$N_t := \exp \left(\int_0^t r_s ds \right).$$

In this case,

$$\widehat{S}_t = \frac{S_t}{N_t} = e^{-\int_0^t r_s ds} S_t, \quad t \geq 0,$$

represents the discounted price of the asset at time 0.

- *Currency exchange rates*

In this case, $N_t := R_t$ denotes *e.g.* the EUR/SGD (EURSGD=X) exchange rate from a foreign currency (*e.g.* EUR) to a domestic currency (*e.g.* SGD), *i.e.* one unit of foreign currency (EUR) corresponds to R_t units of local currency (SGD). Let

$$\widehat{S}_t := \frac{S_t}{R_t}, \quad t \geq 0,$$

denote the price of a local (SG) asset quoted in units of the foreign currency (EUR). For example, if $R_t = 1.63$ and $S_t = \$1$, then

$$\widehat{S}_t = \frac{S_t}{R_t} = \frac{1}{1.63} \times \$1 \simeq €0.61,$$

and $1/R_t$ is the domestic SGD/EUR exchange rate. A question of interest is whether a local asset $\$S_t$, discounted according to a foreign risk-free rate r^f and priced in foreign currency as

$$e^{-r^f t} \frac{S_t}{R_t} = e^{-r^f t} \widehat{S}_t,$$

can be a martingale on the foreign market.

*Major currencies have started floating against each other since 1973, following the end of the system of fixed exchanged rates agreed upon at the Bretton Woods Conference, July 1-22, 1944.

[†]“Anyone who believes exponential growth can go on forever in a finite world is either a madman or an economist”, K.E. Boulding, 1973, page 248.



<p>My foreign currency account S_t grew by 5% this year.</p> <p>Q: Did I achieve a positive return?</p> <p>A:</p>	<p>My foreign currency account S_t grew by 5% this year.</p> <p>The foreign exchange rate dropped by 10%.</p> <p>Q: Did I achieve a positive return?</p> <p>A:</p>
--	---

(a) Scenario A.

(b) Scenario B.

Figure 3.1: Why change of numéraire?

- *Forward numéraire.*

The price $P(t, T)$ of a bond paying $P(T, T) = \$1$ at maturity T can be taken as numéraire. In this case, we have

$$N_t := P(t, T) = \mathbb{E}^* \left[e^{-\int_t^T r_s ds} \mid \mathcal{F}_t \right], \quad 0 \leq t \leq T.$$

Recall that

$$t \mapsto e^{-\int_0^t r_s ds} P(t, T) = \mathbb{E}^* \left[e^{-\int_0^T r_s ds} \mid \mathcal{F}_t \right], \quad 0 \leq t \leq T,$$

is an \mathcal{F}_t - martingale.

- *Annuity numéraire.*

Processes of the form

$$N_t := \sum_{k=1}^n (T_k - T_{k-1}) P(t, T_k), \quad 0 \leq t \leq T_1,$$

where $P(t, T_1), P(t, T_2), \dots, P(t, T_n)$ are bond prices with maturities $T_1 < T_2 < \dots < T_n$ arranged according to a *tenor structure*.

- *Combinations of the above:* for example a foreign money market account $e^{\int_0^t r_s^f ds} R_t$, expressed in local (or domestic) units of currency, where $(r_t^f)_{t \in \mathbb{R}_+}$ represents a short-term interest rate on the foreign market.

When the numéraire is a random process, the pricing of a claim whose value has been transformed under change of numéraire, *e.g.* under a change of currency, has to take into account the risks existing on the foreign market.

In particular, in order to perform a fair pricing, one has to determine a probability measure under which the transformed (or forward, or deflated) process $\widehat{S}_t = S_t / N_t$ will be a martingale, see Section 3.3 for details.

For example in case $N_t := e^{\int_0^t r_s ds}$ is the money market account, the risk-neutral probability measure \mathbb{P}^* is a measure under which the discounted price process

$$\widehat{S}_t = \frac{S_t}{N_t} = e^{-\int_0^t r_s ds} S_t, \quad t \geq 0,$$

is a martingale. In the next section, we will see that this property can be extended to any type of numéraire.

See Exercises 3.5 and 3.6 for other examples of numéraires.

3.2 Change of Numéraire

In this section we review the pricing of options by a change of measure associated to a numéraire N_t , cf. e.g. [Geman, El Karoui, and Rochet, 1995](#) and references therein.

Most of the results of this chapter rely on the following assumption, which expresses absence of arbitrage. In the foreign exchange setting where $N_t = R_t$, this condition states that the price of one unit of foreign currency is a martingale when quoted and discounted in the domestic currency.

Assumption 1. *The discounted numéraire*

$$t \mapsto M_t := e^{-\int_0^t r_s ds} N_t$$

is an \mathcal{F}_t -martingale under the risk-neutral probability measure \mathbb{P}^* .

Definition 3.1 Given $(N_t)_{t \in [0, T]}$ a numéraire process, the associated *forward measure* $\widehat{\mathbb{P}}$ is defined by the Radon-Nikodym density

$$\frac{d\widehat{\mathbb{P}}}{d\mathbb{P}^*} := \frac{M_T}{M_0} = e^{-\int_0^T r_s ds} \frac{N_T}{N_0}. \quad (3.2.1)$$

Recall that the above Relation (3.2.1) rewrites as

$$d\widehat{\mathbb{P}} = \frac{M_T}{M_0} d\mathbb{P}^* = e^{-\int_0^T r_s ds} \frac{N_T}{N_0} d\mathbb{P}^*,$$

which is equivalent to stating that

$$\begin{aligned} \widehat{\mathbb{E}}[F] &= \int_{\Omega} F(\omega) d\widehat{\mathbb{P}}(\omega) \\ &= \int_{\Omega} F(\omega) e^{-\int_0^T r_s ds} \frac{N_T}{N_0} d\mathbb{P}^*(\omega) \\ &= \mathbb{E}^* \left[F e^{-\int_0^T r_s ds} \frac{N_T}{N_0} \right], \end{aligned}$$

for all integrable \mathcal{F}_T -measurable random variables F . More generally, by (3.2.1) and the fact that the process

$$t \mapsto M_t := e^{-\int_0^t r_s ds} N_t$$

is an \mathcal{F}_t -martingale under \mathbb{P}^* under Assumption 1, we find that

$$\begin{aligned} \mathbb{E}^* \left[\frac{d\widehat{\mathbb{P}}}{d\mathbb{P}^*} \middle| \mathcal{F}_t \right] &= \mathbb{E}^* \left[\frac{N_T}{N_0} e^{-\int_0^T r_s ds} \middle| \mathcal{F}_t \right] \\ &= \mathbb{E}^* \left[\frac{M_T}{M_0} \middle| \mathcal{F}_t \right] \\ &= \frac{M_t}{M_0} \\ &= \frac{N_t}{N_0} e^{-\int_0^t r_s ds}, \quad 0 \leq t \leq T. \end{aligned} \quad (3.2.2)$$



In Proposition 3.5 we will show, as a consequence of next Lemma 3.2 below, that for any integrable random claim payoff C we have

$$\mathbf{E}^* \left[C e^{-\int_t^T r_s ds} N_T \mid \mathcal{F}_t \right] = N_t \widehat{\mathbf{E}}[C \mid \mathcal{F}_t], \quad 0 \leq t \leq T.$$

Similarly to the above, the Radon-Nikodym density $d\widehat{\mathbb{P}}_{|\mathcal{F}_t}/d\mathbb{P}_{|\mathcal{F}_t}^*$ of $\widehat{\mathbb{P}}_{|\mathcal{F}_t}$ with respect to $\mathbb{P}_{|\mathcal{F}_t}^*$ satisfies the relation

$$\begin{aligned} \widehat{\mathbf{E}}[F \mid \mathcal{F}_t] &= \int_{\Omega} F(\omega) d\widehat{\mathbb{P}}_{|\mathcal{F}_t}(\omega) \\ &= \int_{\Omega} F(\omega) e^{-\int_0^T r_s ds} \frac{d\widehat{\mathbb{P}}_{|\mathcal{F}_t}}{d\mathbb{P}_{|\mathcal{F}_t}^*} d\mathbb{P}_{|\mathcal{F}_t}^*(\omega) \\ &= \mathbf{E}^* \left[F \frac{d\widehat{\mathbb{P}}_{|\mathcal{F}_t}}{d\mathbb{P}_{|\mathcal{F}_t}^*} \mid \mathcal{F}_t \right], \end{aligned}$$

for all integrable \mathcal{F}_T -measurable random variables F , $0 \leq t \leq T$. Note that (3.2.2), which is \mathcal{F}_t -measurable, should not be confused with (3.2.3), which is \mathcal{F}_T -measurable.

Lemma 3.2 We have

$$\frac{d\widehat{\mathbb{P}}_{|\mathcal{F}_t}}{d\mathbb{P}_{|\mathcal{F}_t}^*} = \frac{M_T}{M_t} = e^{-\int_t^T r_s ds} \frac{N_T}{N_t}, \quad 0 \leq t \leq T. \quad (3.2.3)$$

Proof. The proof of (3.2.3) relies on an abstract version of the Bayes formula. For all bounded \mathcal{F}_t -measurable random variable G , by (3.2.2), the tower property of conditional expectations we have

$$\begin{aligned} \widehat{\mathbf{E}}[G\widehat{X}] &= \mathbf{E}^* \left[G\widehat{X} e^{-\int_0^T r_s ds} \frac{N_T}{N_0} \right] \\ &= \mathbf{E}^* \left[G \frac{N_t}{N_0} e^{-\int_0^t r_s ds} \mathbf{E}^* \left[\widehat{X} e^{-\int_t^T r_s ds} \frac{N_T}{N_t} \mid \mathcal{F}_t \right] \right] \\ &= \mathbf{E}^* \left[G \mathbf{E}^* \left[\frac{d\widehat{\mathbb{P}}}{d\mathbb{P}^*} \mid \mathcal{F}_t \right] \mathbf{E}^* \left[\widehat{X} e^{-\int_t^T r_s ds} \frac{N_T}{N_t} \mid \mathcal{F}_t \right] \right] \\ &= \mathbf{E}^* \left[G \frac{d\widehat{\mathbb{P}}}{d\mathbb{P}^*} \mathbf{E}^* \left[\widehat{X} e^{-\int_t^T r_s ds} \frac{N_T}{N_t} \mid \mathcal{F}_t \right] \right] \\ &= \widehat{\mathbf{E}} \left[G \mathbf{E}^* \left[\widehat{X} e^{-\int_t^T r_s ds} \frac{N_T}{N_t} \mid \mathcal{F}_t \right] \right], \end{aligned}$$

for all bounded random variables \widehat{X} , which shows that

$$\widehat{\mathbf{E}}[\widehat{X} \mid \mathcal{F}_t] = \mathbf{E}^* \left[\widehat{X} e^{-\int_t^T r_s ds} \frac{N_T}{N_t} \mid \mathcal{F}_t \right],$$

i.e. (3.2.3) holds. \square

We note that in case the numéraire $N_t = e^{\int_0^t r_s ds}$ is equal to the money market account we simply have $\widehat{\mathbb{P}} = \mathbb{P}^*$.

Definition 3.3 Given $(X_t)_{t \in \mathbb{R}_+}$ an asset price process, we define the process of forward (or

deflated) prices

$$\widehat{X}_t := \frac{X_t}{N_t}, \quad 0 \leq t \leq T. \quad (3.2.4)$$

The process $(\widehat{X}_t)_{t \in \mathbb{R}_+}$ represents the values at times t of X_t , expressed in units of the numéraire N_t . In the sequel, it will be useful to determine the dynamics of $(\widehat{X}_t)_{t \in \mathbb{R}_+}$ under the forward measure $\widehat{\mathbb{P}}$. The next proposition shows in particular that the process $(e^{\int_0^t r_s ds} / N_t)_{t \in \mathbb{R}_+}$ is an \mathcal{F}_t -martingale under $\widehat{\mathbb{P}}$.

Proposition 3.4 Let $(X_t)_{t \in \mathbb{R}_+}$ denote a continuous $(\mathcal{F}_t)_{t \in \mathbb{R}_+}$ -adapted asset price process such that

$$t \mapsto e^{-\int_0^t r_s ds} X_t, \quad t \geq 0,$$

is a martingale under \mathbb{P}^* . Then, under change of numéraire,

the deflated process $(\widehat{X}_t)_{t \in [0, T]} = (X_t / N_t)_{t \in [0, T]}$ of forward prices is an \mathcal{F}_t -martingale under $\widehat{\mathbb{P}}$,

provided that $(\widehat{X}_t)_{t \in [0, T]}$ is integrable under $\widehat{\mathbb{P}}$.

Proof. We show that

$$\widehat{\mathbb{E}} \left[\frac{X_t}{N_t} \mid \mathcal{F}_s \right] = \frac{X_s}{N_s}, \quad 0 \leq s \leq t. \quad (3.2.5)$$

Namely, for all bounded \mathcal{F}_s -measurable random variables G we note that using (3.2.2) under Assumption 1 we have

$$\begin{aligned} \widehat{\mathbb{E}} \left[G \frac{X_t}{N_t} \right] &= \mathbb{E}^* \left[G \frac{X_t}{N_t} \frac{d\widehat{\mathbb{P}}}{d\mathbb{P}^*} \right] \\ &= \mathbb{E}^* \left[\mathbb{E}^* \left[G \frac{X_t}{N_t} \frac{d\widehat{\mathbb{P}}}{d\mathbb{P}^*} \mid \mathcal{F}_t \right] \right] \\ &= \mathbb{E}^* \left[G \frac{X_t}{N_t} \mathbb{E}^* \left[\frac{d\widehat{\mathbb{P}}}{d\mathbb{P}^*} \mid \mathcal{F}_t \right] \right] \\ &= \mathbb{E}^* \left[G e^{-\int_0^t r_u du} \frac{X_t}{N_0} \right] \end{aligned} \quad (3.2.6)$$

$$= \mathbb{E}^* \left[G e^{-\int_0^s r_u du} \frac{X_s}{N_0} \right] \quad (3.2.7)$$

$$\begin{aligned} &= \mathbb{E}^* \left[G \frac{X_s}{N_s} \mathbb{E}^* \left[\frac{d\widehat{\mathbb{P}}}{d\mathbb{P}^*} \mid \mathcal{F}_s \right] \right] \\ &= \mathbb{E}^* \left[\mathbb{E}^* \left[G \frac{X_s}{N_s} \frac{d\widehat{\mathbb{P}}}{d\mathbb{P}^*} \mid \mathcal{F}_s \right] \right] \\ &= \mathbb{E}^* \left[G \frac{X_s}{N_s} \frac{d\widehat{\mathbb{P}}}{d\mathbb{P}^*} \right] \end{aligned}$$

$$= \widehat{\mathbb{E}} \left[G \frac{X_s}{N_s} \right], \quad 0 \leq s \leq t,$$

where from (3.2.6) to (3.2.7) we used the fact that

$$t \mapsto e^{-\int_0^t r_s ds} X_t$$

is an \mathcal{F}_t -martingale under \mathbb{P}^* . Finally, the identity

$$\widehat{\mathbb{E}}[G \widehat{X}_t] = \widehat{\mathbb{E}} \left[G \frac{X_t}{N_t} \right] = \widehat{\mathbb{E}} \left[G \frac{X_s}{N_s} \right] = \widehat{\mathbb{E}}[G \widehat{X}_s], \quad 0 \leq s \leq t,$$

for all bounded \mathcal{F}_s -measurable G , implies (3.2.5). \square

Pricing using change of numéraire

The change of numéraire technique is specially useful for pricing under random interest rates, in which case an expectation of the form

$$\mathbb{E}^* \left[e^{-\int_t^T r_s ds} C \mid \mathcal{F}_t \right]$$

becomes a *path integral*, see *e.g.* Dash, 2004 for a recent account of path integral methods in quantitative finance. The next proposition is the basic result of this section, it provides a way to price an option with arbitrary payoff C under a random discount factor $e^{-\int_t^T r_s ds}$ by use of the forward measure. It will be applied in Chapter 4 to the pricing of bond options and caplets, cf. Propositions 4.1, 4.3 and 4.5 below.

Proposition 3.5 An option with integrable claim payoff $C \in L^1(\Omega, \mathbb{P}^*, \mathcal{F}_T)$ is priced at time t as

$$\mathbb{E}^* \left[e^{-\int_t^T r_s ds} C \mid \mathcal{F}_t \right] = N_t \widehat{\mathbb{E}} \left[\frac{C}{N_T} \mid \mathcal{F}_t \right], \quad 0 \leq t \leq T, \quad (3.2.8)$$

provided that $C/N_T \in L^1(\Omega, \widehat{\mathbb{P}}, \mathcal{F}_T)$.

Proof. By Relation (3.2.3) in Lemma 3.2 we have

$$\begin{aligned} \mathbb{E}^* \left[e^{-\int_t^T r_s ds} C \mid \mathcal{F}_t \right] &= \mathbb{E}^* \left[\frac{d\widehat{\mathbb{P}}|_{\mathcal{F}_t}}{d\mathbb{P}^*|_{\mathcal{F}_t}} \frac{N_t}{N_T} C \mid \mathcal{F}_t \right] \\ &= N_t \mathbb{E}^* \left[\frac{d\widehat{\mathbb{P}}|_{\mathcal{F}_t}}{d\mathbb{P}^*|_{\mathcal{F}_t}} \frac{C}{N_T} \mid \mathcal{F}_t \right] \\ &= N_t \int_{\Omega} \frac{d\widehat{\mathbb{P}}|_{\mathcal{F}_t}}{d\mathbb{P}^*|_{\mathcal{F}_t}} \frac{C}{N_T} d\mathbb{P}^*|_{\mathcal{F}_t} \\ &= N_t \int_{\Omega} \frac{C}{N_T} d\widehat{\mathbb{P}}|_{\mathcal{F}_t} \\ &= N_t \widehat{\mathbb{E}} \left[\frac{C}{N_T} \mid \mathcal{F}_t \right], \quad 0 \leq t \leq T. \end{aligned}$$

Equivalently, we can write

$$\begin{aligned} N_t \widehat{\mathbb{E}} \left[\frac{C}{N_T} \mid \mathcal{F}_t \right] &= N_t \mathbb{E}^* \left[\frac{C}{N_T} \frac{d\widehat{\mathbb{P}}|_{\mathcal{F}_t}}{d\mathbb{P}^*|_{\mathcal{F}_t}} \mid \mathcal{F}_t \right] \\ &= \mathbb{E}^* \left[e^{-\int_t^T r_s ds} C \mid \mathcal{F}_t \right], \quad 0 \leq t \leq T. \end{aligned}$$

\square

Each application of the change of numéraire formula (3.2.8) will require to:

- pick a suitable numéraire $(N_t)_{t \in \mathbb{R}_+}$ satisfying Assumption 1,
 - make sure that the ratio C/N_T takes a sufficiently simple form,
 - use the Girsanov theorem in order to determine the dynamics of asset prices under the new probability measure $\widehat{\mathbb{P}}$,
- so as to compute the expectation under $\widehat{\mathbb{P}}$ on the right-hand side of (3.2.8).

Next, we consider further examples of numéraires and associated examples of option prices.

Examples:

- Money market account.*

Take $N_t := e^{\int_0^t r_s ds}$, where $(r_t)_{t \in \mathbb{R}_+}$ is a possibly random and time-dependent risk-free interest rate. In this case, Assumption 1 is clearly satisfied, we have $\widehat{\mathbb{P}} = \mathbb{P}^*$ and $d\mathbb{P}^*/d\widehat{\mathbb{P}}$, and (3.2.8) simply reads

$$\mathbb{E}^* \left[e^{-\int_t^T r_s ds} C \mid \mathcal{F}_t \right] = e^{\int_0^t r_s ds} \mathbb{E}^* \left[e^{-\int_0^T r_s ds} C \mid \mathcal{F}_t \right], \quad 0 \leq t \leq T,$$

which yields no particular information.

- Forward numéraire.*

Here, $N_t := P(t, T)$ is the price $P(t, T)$ of a bond maturing at time T , $0 \leq t \leq T$, and the discounted bond price process $\left(e^{-\int_0^t r_s ds} P(t, T) \right)_{t \in [0, T]}$ is an \mathcal{F}_t -martingale under \mathbb{P}^* , i.e. Assumption 1 is satisfied and $N_t = P(t, T)$ can be taken as numéraire. In this case, (3.2.8) shows that a random claim payoff C can be priced as

$$\mathbb{E}^* \left[e^{-\int_t^T r_s ds} C \mid \mathcal{F}_t \right] = P(t, T) \widehat{\mathbb{E}}[C \mid \mathcal{F}_t], \quad 0 \leq t \leq T, \quad (3.2.9)$$

since $N_T = P(T, T) = 1$, where the forward measure $\widehat{\mathbb{P}}$ satisfies

$$\frac{d\widehat{\mathbb{P}}}{d\mathbb{P}^*} = e^{-\int_0^T r_s ds} \frac{P(T, T)}{P(0, T)} = \frac{e^{-\int_0^T r_s ds}}{P(0, T)} \quad (3.2.10)$$

by (3.2.1).

- Annuity numéraires.*

We take

$$N_t := \sum_{k=1}^n (T_k - T_{k-1}) P(t, T_k)$$

where $P(t, T_1), \dots, P(t, T_n)$ are bond prices with maturities $T_1 < T_2 < \dots < T_n$. Here, (3.2.8) shows that a swaption on the cash flow $P(T, T_n) - P(T, T_1) - \kappa N_T$ can be priced as

$$\begin{aligned} \mathbb{E}^* \left[e^{-\int_t^T r_s ds} (P(T, T_n) - P(T, T_1) - \kappa N_T)^+ \mid \mathcal{F}_t \right] \\ = N_t \widehat{\mathbb{E}} \left[\left(\frac{P(T, T_n) - P(T, T_1)}{N_T} - \kappa \right)^+ \mid \mathcal{F}_t \right], \end{aligned}$$

$0 \leq t \leq T < T_1$, where $(P(T, T_n) - P(T, T_1))/N_T$ becomes a *swap rate*, cf. (2.2.7) in Proposition 2.8 and Section 4.5.



Girsanov theorem

We refer to *e.g.* Theorem III-35 page 132 of [Protter, 2004](#) for the following version of the Girsanov Theorem.

Theorem 3.6 Assume that $\widehat{\mathbb{P}}$ is equivalent^a to \mathbb{P}^* with Radon-Nikodym density $d\widehat{\mathbb{P}}/d\mathbb{P}^*$, and let $(W_t)_{t \in [0, T]}$ be a standard Brownian motion under \mathbb{P}^* . Then, letting

$$\Phi_t := \mathbb{E}^* \left[\frac{d\widehat{\mathbb{P}}}{d\mathbb{P}^*} \middle| \mathcal{F}_t \right], \quad 0 \leq t \leq T, \quad (3.2.11)$$

the process $(\widehat{W}_t)_{t \in [0, T]}$ defined by

$$d\widehat{W}_t := dW_t - \frac{1}{\Phi_t} d\Phi_t \bullet dW_t, \quad 0 \leq t \leq T, \quad (3.2.12)$$

is a standard Brownian motion under $\widehat{\mathbb{P}}$.

^aThis means that the Radon-Nikodym densities $d\widehat{\mathbb{P}}/d\mathbb{P}^*$ and $d\mathbb{P}^*/d\widehat{\mathbb{P}}$ exist and are strictly positive with \mathbb{P}^* and $\widehat{\mathbb{P}}$ -probability one, respectively.

In case the martingale $(\Phi_t)_{t \in [0, T]}$ takes the form

$$\Phi_t = \exp \left(- \int_0^t \psi_s dW_s - \frac{1}{2} \int_0^t |\psi_s|^2 ds \right), \quad 0 \leq t \leq T,$$

i.e.

$$d\Phi_t = -\psi_t \Phi_t dW_t, \quad 0 \leq t \leq T,$$

the Itô multiplication Table shows that Relation (3.2.12) reads

$$\begin{aligned} d\widehat{W}_t &= dW_t - \frac{1}{\Phi_t} d\Phi_t \bullet dW_t \\ &= dW_t - \frac{1}{\Phi_t} (-\psi_t \Phi_t dW_t) \bullet dW_t \\ &= dW_t + \psi_t dt, \quad 0 \leq t \leq T, \end{aligned}$$

and shows that the shifted process $(\widehat{W}_t)_{t \in [0, T]} = (W_t + \int_0^t \psi_s ds)_{t \in [0, T]}$ is a standard Brownian motion under $\widehat{\mathbb{P}}$, which is consistent with the Girsanov Theorem. The next result is another application of the Girsanov Theorem.

Proposition 3.7 The process $(\widehat{W}_t)_{t \in [0, T]}$ defined by

$$d\widehat{W}_t := dW_t - \frac{1}{N_t} dN_t \bullet dW_t, \quad 0 \leq t \leq T, \quad (3.2.13)$$

is a standard Brownian motion under $\widehat{\mathbb{P}}$.

Proof. Relation (3.2.2) shows that Φ_t defined in (3.2.11) satisfies

$$\begin{aligned} \Phi_t &= \mathbb{E}^* \left[\frac{d\widehat{\mathbb{P}}}{d\mathbb{P}^*} \middle| \mathcal{F}_t \right] \\ &= \mathbb{E}^* \left[\frac{N_T}{N_0} e^{-\int_0^T r_s ds} \middle| \mathcal{F}_t \right] \end{aligned}$$

$$= \frac{N_t}{N_0} e^{-\int_0^t r_s ds}, \quad 0 \leq t \leq T,$$

hence

$$\begin{aligned} d\Phi_t &= d\left(\frac{N_t}{N_0} e^{-\int_0^t r_s ds}\right) \\ &= -\Phi_t r_t dt + e^{-\int_0^t r_s ds} d\left(\frac{N_t}{N_0}\right) \\ &= -\Phi_t r_t dt + \frac{\Phi_t}{N_t} dN_t, \end{aligned}$$

which, by (3.2.12), yields

$$\begin{aligned} d\hat{W}_t &= dW_t - \frac{1}{\Phi_t} d\Phi_t \cdot dW_t \\ &= dW_t - \frac{1}{\Phi_t} \left(-\Phi_t r_t dt + \frac{\Phi_t}{N_t} dN_t \right) \cdot dW_t \\ &= dW_t - \frac{1}{N_t} dN_t \cdot dW_t, \end{aligned}$$

which is (3.2.13), from Relation (3.2.12) and the Itô multiplication Table. \square

The next Proposition 3.8 is consistent with the statement of Proposition 3.4, and in addition it specifies the dynamics of $(\hat{X}_t)_{t \in \mathbb{R}_+}$ under $\hat{\mathbb{P}}$ using the Girsanov Theorem 3.7. As a consequence, we have the next proposition, see Exercise 3.1 for another calculation based on geometric Brownian motion, and Exercise 3.10 for an extension to correlated Brownian motions.

Proposition 3.8 Assume that $(X_t)_{t \in \mathbb{R}_+}$ and $(N_t)_{t \in \mathbb{R}_+}$ satisfy the stochastic differential equations

$$dX_t = r_t X_t dt + \sigma_t^X X_t dW_t, \quad \text{and} \quad dN_t = r_t N_t dt + \sigma_t^N N_t dW_t, \quad (3.2.14)$$

where $(\sigma_t^X)_{t \in \mathbb{R}_+}$ and $(\sigma_t^N)_{t \in \mathbb{R}_+}$ are $(\mathcal{F}_t)_{t \in \mathbb{R}_+}$ -adapted volatility processes and $(W_t)_{t \in \mathbb{R}_+}$ is a standard Brownian motion under \mathbb{P}^* . Then we have

$$d\hat{X}_t = (\sigma_t^X - \sigma_t^N) \hat{X}_t d\hat{W}_t, \quad (3.2.15)$$

hence $(\hat{X}_t)_{t \in [0, T]}$ is given by the driftless geometric Brownian motion

$$\hat{X}_t = \hat{X}_0 \exp \left(\int_0^t (\sigma_s^X - \sigma_s^N) d\hat{W}_s - \frac{1}{2} \int_0^t (\sigma_s^X - \sigma_s^N)^2 ds \right), \quad 0 \leq t \leq T.$$

Proof. First we note that by (3.2.13) and (3.2.14),

$$d\hat{W}_t = dW_t - \frac{1}{N_t} dN_t \cdot dW_t = dW_t - \sigma_t^N dt, \quad t \geq 0,$$

is a standard Brownian motion under $\hat{\mathbb{P}}$. Next, by Itô's calculus and the Itô multiplication Table and (3.2.14) we have

$$\begin{aligned} d\left(\frac{1}{N_t}\right) &= -\frac{1}{N_t^2} dN_t + \frac{1}{N_t^3} dN_t \cdot dN_t \\ &= -\frac{1}{N_t^2} (r_t N_t dt + \sigma_t^N N_t dW_t) + \frac{|\sigma_t^N|^2}{N_t} dt \end{aligned}$$



$$\begin{aligned}
&= -\frac{1}{N_t^2} (r_t N_t dt + \sigma_t^N N_t (d\hat{W}_t + \sigma_t^N dt)) + \frac{|\sigma_t^N|^2}{N_t} dt \\
&= -\frac{1}{N_t} (r_t dt + \sigma_t^N d\hat{W}_t),
\end{aligned} \tag{3.2.16}$$

hence

$$\begin{aligned}
d\hat{X}_t &= d\left(\frac{X_t}{N_t}\right) \\
&= \frac{dX_t}{N_t} + X_t d\left(\frac{1}{N_t}\right) + dX_t \cdot d\left(\frac{1}{N_t}\right) \\
&= \frac{1}{N_t} (r_t X_t dt + \sigma_t^X X_t dW_t) - \frac{X_t}{N_t} (r_t dt + \sigma_t^N dW_t - |\sigma_t^N|^2 dt) \\
&\quad - \frac{1}{N_t} (r_t X_t dt + \sigma_t^X X_t dW_t) \cdot (r_t dt + \sigma_t^N dW_t - |\sigma_t^N|^2 dt) \\
&= \frac{1}{N_t} (r_t X_t dt + \sigma_t^X X_t dW_t) - \frac{X_t}{N_t} (r_t dt + \sigma_t^N dW_t) \\
&\quad + \frac{X_t}{N_t} |\sigma_t^N|^2 dt - \frac{X_t}{N_t} \sigma_t^X \sigma_t^N dt \\
&= \frac{X_t}{N_t} \sigma_t^X dW_t - \frac{X_t}{N_t} \sigma_t^N dW_t - \frac{X_t}{N_t} \sigma_t^X \sigma_t^N dt + X_t \frac{|\sigma_t^N|^2}{N_t} dt \\
&= \frac{X_t}{N_t} (\sigma_t^X dW_t - \sigma_t^N dW_t - \sigma_t^X \sigma_t^N dt + |\sigma_t^N|^2 dt) \\
&= \hat{X}_t (\sigma_t^X - \sigma_t^N) dW_t - \hat{X}_t (\sigma_t^X - \sigma_t^N) \sigma_t^N dt \\
&= \hat{X}_t (\sigma_t^X - \sigma_t^N) d\hat{W}_t,
\end{aligned}$$

since $d\hat{W}_t = dW_t - \sigma_t^N dt$, $0 \leq t \leq T$. □

We end this section with some comments on inverse changes of measure.

Inverse changes of measure

In the next proposition we compute the conditional inverse Radon-Nikodym density $d\mathbb{P}^*/d\hat{\mathbb{P}}$, see also (3.2.2).

Proposition 3.9 We have

$$\hat{\mathbb{E}} \left[\frac{d\mathbb{P}^*}{d\hat{\mathbb{P}}} \mid \mathcal{F}_t \right] = \frac{N_0}{N_t} \exp \left(\int_0^t r_s ds \right), \quad 0 \leq t \leq T, \tag{3.2.17}$$

and the process

$$t \mapsto \frac{M_0}{M_t} = \frac{N_0}{N_t} \exp \left(\int_0^t r_s ds \right), \quad 0 \leq t \leq T,$$

is an \mathcal{F}_t -martingale under $\hat{\mathbb{P}}$.

Proof. For all bounded and \mathcal{F}_t -measurable random variables F we have,

$$\begin{aligned}
\hat{\mathbb{E}} \left[F \frac{d\mathbb{P}^*}{d\hat{\mathbb{P}}} \right] &= \mathbb{E}^*[F] \\
&= \mathbb{E}^* \left[F \frac{N_t}{N_0} \right]
\end{aligned}$$

$$\begin{aligned}
&= \mathbf{E}^* \left[F \frac{N_T}{N_t} \exp \left(- \int_t^T r_s ds \right) \right] \\
&= \widehat{\mathbf{E}} \left[F \frac{N_0}{N_t} \exp \left(\int_0^t r_s ds \right) \right].
\end{aligned}$$

□

By (3.2.16) we also have

$$d \left(\frac{1}{N_t} \exp \left(\int_0^t r_s ds \right) \right) = - \frac{1}{N_t} \exp \left(\int_0^t r_s ds \right) \sigma_t^N d\widehat{W}_t,$$

which recovers the second part of Proposition 3.9, i.e. the martingale property of


$$t \mapsto \frac{M_0}{M_t} = \frac{1}{N_t} \exp \left(\int_0^t r_s ds \right)$$

under $\widehat{\mathbb{P}}$.

3.3 Foreign Exchange

Currency exchange is a typical application of change of numéraire, that illustrates the absence of arbitrage principle.

Let R_t denote the foreign exchange rate, i.e. R_t is the (possibly fractional) quantity of local currency that correspond to one unit of foreign currency, while $1/R_t$ represents the quantity of foreign currency that correspond to a unit of local currency.



Consider an investor that intends to exploit an “overseas investment opportunity” by

- at time 0, changing one unit of local currency into $1/R_0$ units of foreign currency,
- investing $1/R_0$ on the foreign market at the rate r^f , which will yield the amount e^{tr^f}/R_0 at time $t > 0$,
- changing back e^{tr^f}/R_0 into a quantity $e^{tr^f}R_t/R_0 = N_t/R_0$ of local currency.




Figure 3.2: Overseas investment opportunity.*

In other words, the foreign money market account e^{tr^f} is valued $e^{tr^f}R_t$ on the local (or domestic) market, and its discounted value on the local market is

$$e^{-tr^l + tr^f} R_t, \quad t > 0.$$

*For illustration purposes only. Not an advertisement.

The outcome of this investment will be obtained by a martingale comparison of $e^{tr^f}R_t/R_0$ to the amount e^{tr^l} that could have been obtained by investing on the local market.

Taking

$$N_t := e^{tr^f}R_t, \quad t \geq 0, \quad (3.3.1)$$

as *numéraire*, absence of arbitrage is expressed by Assumption 1, which states that the discounted numéraire process

$$t \mapsto e^{-rt}N_t = e^{-t(r^l - r^f)}R_t$$

is an \mathcal{F}_t -martingale under \mathbb{P}^* .

Next, we find a characterization of this arbitrage condition using the model parameters r, r^f, μ , by modeling the foreign exchange rates R_t according to a geometric Brownian motion (3.3.2).

Proposition 3.10 Assume that the foreign exchange rate R_t satisfies a stochastic differential equation of the form

$$dR_t = \mu R_t dt + \sigma R_t dW_t, \quad (3.3.2)$$

where $(W_t)_{t \in \mathbb{R}_+}$ is a standard Brownian motion under \mathbb{P}^* . Under the absence of arbitrage Assumption 1 for the numéraire (3.3.1), we have

$$\mu = r^l - r^f, \quad (3.3.3)$$

hence the exchange rate process satisfies

$$dR_t = (r^l - r^f)R_t dt + \sigma R_t dW_t. \quad (3.3.4)$$

under \mathbb{P}^* .

Proof. The equation (3.3.2) has solution

$$R_t = R_0 e^{\mu t + \sigma W_t - \sigma^2 t / 2}, \quad t \geq 0,$$

hence the discounted value of the foreign money market account e^{tr^f} on the local market is

$$e^{-tr^l} N_t = e^{-tr^l + tr^f} R_t = R_0 e^{(r^f - r^l + \mu)t + \sigma W_t - \sigma^2 t / 2}, \quad t \geq 0.$$

Under the absence of arbitrage Assumption 1, the process $e^{-(r^l - r^f)t} R_t = e^{-tr^l} N_t$ should be an \mathcal{F}_t -martingale under \mathbb{P}^* , and this holds provided that $r^f - r + \mu = 0$, which yields (3.3.3) and (3.3.4). \square

As a consequence of Proposition 3.10, under absence of arbitrage a local investor who buys a unit of foreign currency in the hope of a higher return $r^f >> r$ will have to face a lower (or even more negative) drift

$$\mu = r^l - r^f << 0$$

in his exchange rate R_t . The drift $\mu = r^l - r^f$ is also called the *cost of carrying* the foreign currency.

The local money market account $X_t := e^{t r^l}$ is valued $e^{t r^l} / R_t$ on the foreign market, and its discounted value at time $t \geq 0$ on the foreign market is

$$\begin{aligned}\frac{e^{(r^l - r^f)t}}{R_t} &= \frac{X_t}{N_t} \\ &= \frac{1}{R_0} e^{(r^l - r^f)t - \mu t - \sigma W_t + \sigma^2 t/2} \\ &= \frac{1}{R_0} e^{(r^l - r^f)t - \mu t - \sigma \hat{W}_t - \sigma^2 t/2},\end{aligned}\tag{3.3.5}$$

where

$$\begin{aligned}d\hat{W}_t &= dW_t - \frac{1}{N_t} dN_t \bullet dW_t \\ &= dW_t - \frac{1}{R_t} dR_t \bullet dW_t \\ &= dW_t - \sigma dt, \quad t \geq 0,\end{aligned}$$

is a standard Brownian motion under $\hat{\mathbb{P}}$ by (3.2.13). Under absence of arbitrage, the process $e^{-(r^l - r^f)t} R_t$ is an \mathcal{F}_t -martingale under \mathbb{P}^* and (3.3.5) is an \mathcal{F}_t -martingale under $\hat{\mathbb{P}}$ by Proposition 3.4, which recovers (3.3.3).

Proposition 3.11 Under the absence of arbitrage condition (3.3.3), the inverse exchange rate $1/R_t$ satisfies

$$d\left(\frac{1}{R_t}\right) = \frac{r^f - r^l}{R_t} dt - \frac{\sigma}{R_t} d\hat{W}_t,\tag{3.3.6}$$

under $\hat{\mathbb{P}}$, where $(R_t)_{t \in \mathbb{R}_+}$ is given by (3.3.4).

Proof. By (3.3.3), the exchange rate $1/R_t$ is written using Itô's calculus as

$$\begin{aligned}d\left(\frac{1}{R_t}\right) &= -\frac{1}{R_t^2} (\mu R_t dt + \sigma R_t dW_t) + \frac{1}{R_t^3} \sigma^2 R_t^2 dt \\ &= -\frac{\mu - \sigma^2}{R_t} dt - \frac{\sigma}{R_t} dW_t \\ &= -\frac{\mu}{R_t} dt - \frac{\sigma}{R_t} d\hat{W}_t \\ &= \frac{r^f - r^l}{R_t} dt - \frac{\sigma}{R_t} d\hat{W}_t,\end{aligned}$$

where $(\hat{W}_t)_{t \in \mathbb{R}_+}$ is a standard Brownian motion under $\hat{\mathbb{P}}$. \square

Consequently, under absence of arbitrage, a foreign investor who buys a unit of the local currency in the hope of a higher return $r >> r^f$ will have to face a lower (or even more negative) drift $-\mu = r^f - r$ in his exchange rate $1/R_t$ as written in (3.3.6) under $\hat{\mathbb{P}}$.

Foreign exchange options

We now price a foreign exchange call option with payoff $(R_T - \kappa)^+$ under \mathbb{P}^* on the exchange rate R_T by the Black-Scholes formula as in the next proposition, also known as the [Garman and Kohlhagen, 1983](#) formula. The foreign exchange call option is designed for a local buyer of foreign currency.



Proposition 3.12 (Garman and Kohlhagen, 1983 formula for call options). Consider the exchange rate process $(R_t)_{t \in \mathbb{R}_+}$ given by (3.3.4). The price of the foreign exchange call option on R_T with maturity T and strike price $\kappa > 0$ is given in local currency units as

$$e^{-(T-t)r^l} \mathbf{E}^* [(R_T - \kappa)^+ | \mathcal{F}_t] = e^{-(T-t)r^f} R_t \Phi_+(t, R_t) - \kappa e^{-(T-t)r^l} \Phi_-(t, R_t), \quad (3.3.7)$$

$0 \leq t \leq T$, where

$$\Phi_+(t, x) = \Phi \left(\frac{\log(x/\kappa) + (T-t)(r^l - r^f + \sigma^2/2)}{\sigma\sqrt{T-t}} \right),$$

and

$$\Phi_-(t, x) = \Phi \left(\frac{\log(x/\kappa) + (T-t)(r^l - r^f - \sigma^2/2)}{\sigma\sqrt{T-t}} \right).$$

Proof. As a consequence of (3.3.4), we find the numéraire dynamics

$$\begin{aligned} dN_t &= d(e^{tr^f} R_t) \\ &= r^f e^{tr^f} R_t dt + e^{tr^f} dR_t \\ &= r e^{tr^f} R_t dt + \sigma e^{tr^f} R_t dW_t \\ &= r N_t dt + \sigma N_t dW_t. \end{aligned}$$

Hence, a standard application of the Black-Scholes formula yields

$$\begin{aligned} e^{-(T-t)r^l} \mathbf{E}^* [(R_T - \kappa)^+ | \mathcal{F}_t] &= e^{-(T-t)r^l} \mathbf{E}^* [(e^{-Tr^f} N_T - \kappa)^+ | \mathcal{F}_t] \\ &= e^{-(T-t)r^l} e^{-Tr^f} \mathbf{E}^* [(N_T - \kappa e^{Tr^f})^+ | \mathcal{F}_t] \\ &= e^{-Tr^f} \left(N_t \Phi \left(\frac{\log(N_t e^{-Tr^f}/\kappa) + (r + \sigma^2/2)(T-t)}{\sigma\sqrt{T-t}} \right) \right. \\ &\quad \left. - \kappa e^{Tr^f - (T-t)r^l} \Phi \left(\frac{\log(N_t e^{-Tr^f}/\kappa) + (r^l - \sigma^2/2)(T-t)}{\sigma\sqrt{T-t}} \right) \right) \\ &= e^{-Tr^f} \left(N_t \Phi \left(\frac{\log(R_t/\kappa) + (T-t)(r^l - r^f + \sigma^2/2)}{\sigma\sqrt{T-t}} \right) \right. \\ &\quad \left. - \kappa e^{Tr^f - (T-t)r^l} \Phi \left(\frac{\log(R_t/\kappa) + (T-t)(r^l - r^f - \sigma^2/2)}{\sigma\sqrt{T-t}} \right) \right) \\ &= e^{-(T-t)r^f} R_t \Phi_+(t, R_t) - \kappa e^{-(T-t)r^l} \Phi_-(t, R_t). \end{aligned}$$

A similar conclusion can be reached by directly applying (3.3.4). \square

Similarly, from (3.3.6) rewritten as

$$d \left(\frac{e^{tr^l}}{R_t} \right) = r^f \frac{e^{tr^l}}{R_t} dt - \sigma \frac{e^{tr^l}}{R_t} d\hat{W}_t,$$

a foreign exchange put option with payoff $(1/\kappa - 1/R_T)^+$ can be priced under $\hat{\mathbb{P}}$ in a Black-Scholes model by taking e^{tr^l}/R_t as underlying asset price, r^f as risk-free interest rate, and $-\sigma$

as volatility parameter. The foreign exchange put option is designed for the foreign seller of local currency, see for example the **buy back guarantee*** which is a typical example of a foreign exchange put option.

Proposition 3.13 (Garman and Kohlhagen, 1983 formula for put options). Consider the exchange rate process $(R_t)_{t \in \mathbb{R}_+}$ given by (3.3.4). The price of the foreign exchange call option on R_T with maturity T and strike price $1/\kappa > 0$ is given in foreign currency units as

$$\begin{aligned} & e^{-(T-t)r^f} \widehat{\mathbf{E}} \left[\left(\frac{1}{\kappa} - \frac{1}{R_T} \right)^+ \middle| \mathcal{F}_t \right] \\ &= \frac{e^{-(T-t)r^f}}{\kappa} \Phi_- \left(t, \frac{1}{R_t} \right) - \frac{e^{-(T-t)r^f}}{R_t} \Phi_+ \left(t, \frac{1}{R_t} \right), \end{aligned} \quad (3.3.8)$$

$0 \leq t \leq T$, where

$$\Phi_+(t, x) := \Phi \left(-\frac{\log(\kappa x) + (T-t)(r^f - r^l + \sigma^2/2)}{\sigma \sqrt{T-t}} \right),$$

and

$$\Phi_-(t, x) := \Phi \left(-\frac{\log(\kappa x) + (T-t)(r^f - r^l - \sigma^2/2)}{\sigma \sqrt{T-t}} \right).$$

Proof. The Black-Scholes formula yields

$$\begin{aligned} & e^{-(T-t)r^f} \widehat{\mathbf{E}} \left[\left(\frac{1}{\kappa} - \frac{1}{R_T} \right)^+ \middle| \mathcal{F}_t \right] = e^{-(T-t)r^f} e^{-Tr^l} \widehat{\mathbf{E}} \left[\left(\frac{e^{Tr^l}}{\kappa} - \frac{e^{Tr^l}}{R_T} \right)^+ \middle| \mathcal{F}_t \right] \\ &= \frac{1}{\kappa} e^{-(T-t)r^f} \Phi_- \left(t, \frac{1}{R_t} \right) - \frac{e^{-(T-t)r^f}}{R_t} \Phi_+ \left(t, \frac{1}{R_t} \right), \end{aligned}$$

which is the symmetric of (3.3.7) by exchanging R_t with $1/R_t$, and r with r^f . \square

Call/put duality for foreign exchange options

Let $N_t = e^{t'r^f} R_t$, where R_t is an exchange rate with respect to a foreign currency and r_f is the foreign market interest rate.

Proposition 3.14 The foreign exchange call and put options on the local and foreign markets are linked by the call/put duality relation

$$e^{-(T-t)r^f} \mathbf{E}^* [(R_T - \kappa)^+ \mid \mathcal{F}_t] = \kappa R_t e^{-(T-t)r^f} \widehat{\mathbf{E}} \left[\left(\frac{1}{\kappa} - \frac{1}{R_T} \right)^+ \middle| \mathcal{F}_t \right], \quad (3.3.9)$$

between a put option with strike price $1/\kappa$ and a (possibly fractional) quantity $1/(\kappa R_t)$ of call option(s) with strike price κ .

*Right-click to open or save the attachment.



Proof. By application of change of numéraire from Proposition 3.5 and (3.2.8) we have

$$\widehat{\mathbf{E}} \left[\frac{1}{e^{Tr^f} R_T} (R_T - \kappa)^+ \mid \mathcal{F}_t \right] = \frac{1}{N_t} e^{-(T-t)r^l} \mathbf{E}^* [(R_T - \kappa)^+ \mid \mathcal{F}_t],$$

hence

$$\begin{aligned} e^{-(T-t)r^f} \widehat{\mathbf{E}} \left[\left(\frac{1}{\kappa} - \frac{1}{R_T} \right)^+ \mid \mathcal{F}_t \right] &= e^{-(T-t)r^f} \widehat{\mathbf{E}} \left[\frac{1}{\kappa R_T} (R_T - \kappa)^+ \mid \mathcal{F}_t \right] \\ &= \frac{1}{\kappa} e^{tr^f} \widehat{\mathbf{E}} \left[\frac{1}{e^{Tr^f} R_T} (R_T - \kappa)^+ \mid \mathcal{F}_t \right] \\ &= \frac{1}{\kappa N_t} e^{tr^f - (T-t)r^l} \mathbf{E}^* [(R_T - \kappa)^+ \mid \mathcal{F}_t] \\ &= \frac{1}{\kappa R_t} e^{-(T-t)r^l} \mathbf{E}^* [(R_T - \kappa)^+ \mid \mathcal{F}_t]. \end{aligned}$$

□

In the Black-Scholes case, the duality (3.3.9) can be directly checked by verifying that (3.3.8) coincides with

$$\begin{aligned} &\frac{1}{\kappa R_t} e^{-(T-t)r^l} \mathbf{E}^* [(R_T - \kappa)^+ \mid \mathcal{F}_t] \\ &= \frac{1}{\kappa R_t} e^{-(T-t)r^l} e^{-Tr^f} \mathbf{E}^* [(e^{Tr^f} R_T - \kappa e^{Tr^f})^+ \mid \mathcal{F}_t] \\ &= \frac{1}{\kappa R_t} e^{-(T-t)r^l} e^{-Tr^f} \mathbf{E}^* [(N_T - \kappa e^{Tr^f})^+ \mid \mathcal{F}_t] \\ &= \frac{1}{\kappa R_t} (e^{-(T-t)r^f} R_t \Phi_+^c(t, R_t) - \kappa e^{-(T-t)r^l} \Phi_-^c(t, R_t)) \\ &= \frac{1}{\kappa} e^{-(T-t)r^f} \Phi_+^c(t, R_t) - \frac{e^{-(T-t)r^l}}{R_t} \Phi_-^c(t, R_t) \\ &= \frac{1}{\kappa} e^{-(T-t)r^f} \Phi_- \left(t, \frac{1}{R_t} \right) - \frac{e^{-(T-t)r^l}}{R_t} \Phi_+ \left(t, \frac{1}{R_t} \right), \end{aligned}$$

where

$$\Phi_+^c(t, x) := \Phi \left(\frac{\log(x/\kappa) + (T-t)(r^l - r^f + \sigma^2/2)}{\sigma \sqrt{T-t}} \right),$$

and

$$\Phi_-^c(t, x) := \Phi \left(\frac{\log(x/\kappa) + (T-t)(r^l - r^f - \sigma^2/2)}{\sigma \sqrt{T-t}} \right).$$

	Local market	Foreign market
Measure	\mathbb{P}^*	$\hat{\mathbb{P}}$
Discount factor	$t \mapsto e^{-r^l t}$	$t \mapsto e^{-r^f t}$
Martingale	$t \mapsto e^{-t r^l} N_t = e^{-t(r^l - r^f)} R_t$	$t \mapsto \frac{X_t}{N_t} = \frac{e^{(r^l - r^f)t}}{R_t}$
Option	$e^{-(T-t)r^l} \mathbb{E}^* [(R_T - \kappa)^+ \mathcal{F}_t]$	$e^{-(T-t)r^f} \hat{\mathbb{E}} \left[\left(\frac{1}{\kappa} - \frac{1}{R_T} \right)^+ \middle \mathcal{F}_t \right]$
Application	Local purchase of foreign currency	Foreign selling of local currency

Table 3.1: Local vs foreign exchange options.

Example - Buy back guarantee

The put option priced

$$\begin{aligned} & e^{-(T-t)r^f} \hat{\mathbb{E}} \left[\left(\frac{1}{\kappa} - \frac{1}{R_T} \right)^+ \middle| \mathcal{F}_t \right] \\ &= \frac{1}{\kappa} e^{-(T-t)r^f} \Phi_- \left(t, \frac{1}{R_t} \right) - \frac{e^{-(T-t)r^l}}{R_t} \Phi_+ \left(t, \frac{1}{R_t} \right) \end{aligned}$$

on the foreign market corresponds to a **buy back guarantee*** in currency exchange. In the case of an option “at the money” with $\kappa = R_t$ and $r = r^f \simeq 0$, we find

$$\begin{aligned} \hat{\mathbb{E}} \left[\left(\frac{1}{R_t} - \frac{1}{R_T} \right)^+ \middle| \mathcal{F}_t \right] &= \frac{1}{R_t} \times \left(\Phi \left(\frac{\sigma \sqrt{T-t}}{2} \right) - \Phi \left(-\frac{\sigma \sqrt{T-t}}{2} \right) \right) \\ &= \frac{1}{R_t} \times \left(2\Phi \left(\frac{\sigma \sqrt{T-t}}{2} \right) - 1 \right). \end{aligned}$$

For example, let R_t denote the USD/EUR (USDEUR=X) exchange rate from a foreign currency (USD) to a local currency (EUR), i.e. one unit of the foreign currency (USD) corresponds to $R_t = 1/1.23$ units of local currency (EUR). Taking $T-t = 30$ days and $\sigma = 10\%$, we find that the foreign currency put option allowing the foreign sale of one EURO back into USDs is priced at the money in USD as

$$\begin{aligned} \hat{\mathbb{E}} \left[\left(\frac{1}{R_t} - \frac{1}{R_T} \right)^+ \middle| \mathcal{F}_t \right] &= 1.23(2\Phi(0.05 \times \sqrt{31/365}) - 1) \\ &= 1.23(2 \times 0.505813 - 1) \\ &= \$0.01429998 \end{aligned}$$

*Right-click to open or save the attachment.



per USD, or €0.011626 per exchanged unit of EURO. Based on a displayed option price of €4.5 and in order to make the contract fair, this would translate into an average amount of $4.5/0.011626 \simeq €387$ exchanged at the counter by customers subscribing to the buy back guarantee.

3.4 Pricing Exchange Options

Based on Proposition 3.4, we model the process \widehat{X}_t of forward prices as a continuous martingale under $\widehat{\mathbb{P}}$, written as

$$d\widehat{X}_t = \widehat{\sigma}_t d\widehat{W}_t, \quad t \geq 0, \quad (3.4.1)$$

where $(\widehat{W}_t)_{t \in \mathbb{R}_+}$ is a standard Brownian motion under $\widehat{\mathbb{P}}$ and $(\widehat{\sigma}_t)_{t \in \mathbb{R}_+}$ is an $(\mathcal{F}_t)_{t \in \mathbb{R}_+}$ -adapted stochastic volatility process. More precisely, we assume that $(\widehat{X}_t)_{t \in \mathbb{R}_+}$ has the dynamics

$$d\widehat{X}_t = \widehat{\sigma}_t(\widehat{X}_t) d\widehat{W}_t, \quad (3.4.2)$$

where $x \mapsto \widehat{\sigma}_t(x)$ is a local volatility function which is Lipschitz in x , uniformly in $t \geq 0$. The Markov property of the diffusion process $(\widehat{X}_t)_{t \in \mathbb{R}_+}$, cf. Theorem V-6-32 of Protter, 2004, shows that when \widehat{g} is a deterministic payoff function, the conditional expectation $\widehat{\mathbb{E}}[\widehat{g}(\widehat{X}_T) | \mathcal{F}_t]$ can be written using a (measurable) function $\widehat{C}(t, x)$ of t and \widehat{X}_t , as

$$\widehat{\mathbb{E}}[\widehat{g}(\widehat{X}_T) | \mathcal{F}_t] = \widehat{C}(t, \widehat{X}_t), \quad 0 \leq t \leq T.$$

Consequently, a vanilla option with claim payoff $C := N_T \widehat{g}(\widehat{X}_T)$ can be priced from Proposition 3.5 as

$$\begin{aligned} \mathbf{E}^* \left[e^{-\int_t^T r_s ds} N_T \widehat{g}(\widehat{X}_T) \mid \mathcal{F}_t \right] &= N_t \widehat{\mathbb{E}}[\widehat{g}(\widehat{X}_T) | \mathcal{F}_t] \\ &= N_t \widehat{C}(t, \widehat{X}_t), \quad 0 \leq t \leq T. \end{aligned} \quad (3.4.3)$$

In the next Proposition 3.15 we state the Margrabe, 1978 formula for the pricing of exchange options by the zero interest rate Black-Scholes formula. It will be applied in particular in Proposition 4.3 below for the pricing of bond options. Here, $(N_t)_{t \in \mathbb{R}_+}$ denotes any numéraire process satisfying Assumption 1.

Proposition 3.15 (Margrabe, 1978 formula). Assume that $\widehat{\sigma}_t(\widehat{X}_t) = \widehat{\sigma}(t)\widehat{X}_t$, i.e. the martingale $(\widehat{X}_t)_{t \in [0, T]}$ is a (driftless) geometric Brownian motion under $\widehat{\mathbb{P}}$ with deterministic volatility $(\widehat{\sigma}(t))_{t \in [0, T]}$. Then we have

$$\mathbf{E}^* \left[e^{-\int_t^T r_s ds} (X_T - \kappa N_T)^+ \mid \mathcal{F}_t \right] = X_t \Phi_+^0(t, \widehat{X}_t) - \kappa N_t \Phi_-^0(t, \widehat{X}_t), \quad (3.4.4)$$

$t \in [0, T]$, where

$$\Phi_+^0(t, x) = \Phi \left(\frac{\log(x/\kappa)}{v(t, T)} + \frac{v(t, T)}{2} \right), \quad \Phi_-^0(t, x) = \Phi \left(\frac{\log(x/\kappa)}{v(t, T)} - \frac{v(t, T)}{2} \right), \quad (3.4.5)$$

and $v^2(t, T) = \int_t^T \widehat{\sigma}^2(s) ds$.

Proof. Taking $g(x) := (x - \kappa)^+$ in (3.4.3), the call option with payoff

$$\begin{aligned}(X_T - \kappa N_T)^+ &= N_T (\widehat{X}_T - \kappa)^+ \\ &= N_T \left(\widehat{X}_t \exp \left(\int_t^T \widehat{\sigma}(s) d\widehat{W}_s - \frac{1}{2} \int_t^T |\widehat{\sigma}(s)|^2 ds \right) - \kappa \right)^+, \end{aligned}$$

and floating strike price κN_T is priced by (3.4.3) as

$$\begin{aligned}\mathbf{E}^* \left[e^{-\int_t^T r_s ds} (X_T - \kappa N_T)^+ \mid \mathcal{F}_t \right] &= \mathbf{E}^* \left[e^{-\int_t^T r_s ds} N_T (\widehat{X}_T - \kappa)^+ \mid \mathcal{F}_t \right] \\ &= N_t \widehat{\mathbf{E}}[(\widehat{X}_T - \kappa)^+ \mid \mathcal{F}_t] \\ &= N_t \widehat{C}(t, \widehat{X}_t), \end{aligned}$$

where the function $\widehat{C}(t, \widehat{X}_t)$ is given by the Black-Scholes formula

$$\widehat{C}(t, x) = x \Phi_+^0(t, x) - \kappa \Phi_-^0(t, x),$$

with zero interest rate, since $(\widehat{X}_t)_{t \in [0, T]}$ is a driftless geometric Brownian motion which is an \mathcal{F}_t -martingale under $\widehat{\mathbb{P}}$, and \widehat{X}_T is a lognormal random variable with variance coefficient $v^2(t, T) = \int_t^T \widehat{\sigma}^2(s) ds$. Hence we have

$$\begin{aligned}\mathbf{E}^* \left[e^{-\int_t^T r_s ds} (X_T - \kappa N_T)^+ \mid \mathcal{F}_t \right] &= N_t \widehat{C}(t, \widehat{X}_t) \\ &= N_t \widehat{X}_t \Phi_+^0(t, \widehat{X}_t) - \kappa N_t \Phi_-^0(t, \widehat{X}_t), \end{aligned}$$

$t \geq 0$. □

In particular, from Proposition 3.8 and (3.2.15), we can take $\widehat{\sigma}(t) = \sigma_t^X - \sigma_t^N$ when $(\sigma_t^X)_{t \in \mathbb{R}_+}$ and $(\sigma_t^N)_{t \in \mathbb{R}_+}$ are deterministic.

Examples:

- a) When the short rate process $(r(t))_{t \in [0, T]}$ is a *deterministic* function of time and $N_t = e^{\int_t^T r(s) ds}$, $0 \leq t \leq T$, we have $\widehat{\mathbb{P}} = \mathbb{P}^*$ and Proposition 3.15 yields the [Merton, 1973](#) “zero interest rate” version of the Black-Scholes formula

$$\begin{aligned}e^{-\int_t^T r(s) ds} \mathbf{E}^* [(X_T - \kappa)^+ \mid \mathcal{F}_t] \\ = X_t \Phi_+^0 \left(t, e^{\int_t^T r(s) ds} X_t \right) - \kappa e^{-\int_t^T r(s) ds} \Phi_-^0 \left(t, e^{\int_t^T r(s) ds} X_t \right), \end{aligned}$$

where Φ_+^0 and Φ_-^0 are defined in (3.4.5) and $(X_t)_{t \in \mathbb{R}_+}$ satisfies the equation

$$\frac{dX_t}{X_t} = r(t) dt + \widehat{\sigma}(t) dW_t, \quad i.e. \quad \frac{d\widehat{X}_t}{\widehat{X}_t} = \widehat{\sigma}(t) dW_t, \quad 0 \leq t \leq T.$$

- b) In the case of pricing under a *forward numéraire*, i.e. when $N_t = P(t, T)$, $t \in [0, T]$, we get

$$\mathbf{E}^* \left[e^{-\int_t^T r_s ds} (X_T - \kappa)^+ \mid \mathcal{F}_t \right] = X_t \Phi_+^0(t, \widehat{X}_t) - \kappa P(t, T) \Phi_-^0(t, \widehat{X}_t),$$

$0 \leq t \leq T$, since $N_T = P(T, T) = 1$. In particular, when $X_t = P(t, S)$ the above formula allows us to price a bond call option on $P(T, S)$ as

$$\mathbf{E}^* \left[e^{-\int_t^T r_s ds} (P(T, S) - \kappa)^+ \mid \mathcal{F}_t \right] = P(t, S) \Phi_+^0(t, \widehat{X}_t) - \kappa P(t, T) \Phi_-^0(t, \widehat{X}_t),$$

$0 \leq t \leq T$, provided that the martingale $\widehat{X}_t = P(t, S) / P(t, T)$ under $\widehat{\mathbb{P}}$ is given by a geometric Brownian motion, cf. Section 4.2.



3.5 Hedging by Change of Numéraire

We use the stochastic integral representation of the forward claim payoffs and change of numéraire in order to compute self-financing portfolio strategies. Our hedging portfolios will be built on the assets (X_t, N_t) , not on X_t and the money market account $B_t = e^{\int_0^t r_s ds}$, extending the classical hedging portfolios that are available from the Black-Scholes formula, using a technique from [Jamshidian, 1996](#), cf. also [Privault and Teng, 2012](#).

Consider a claim with random payoff C , typically an interest rate derivative, cf. Chapter 4. Assume that the forward claim payoff $C/N_T \in L^2(\Omega)$ has the stochastic integral representation

$$\widehat{C} := \frac{C}{N_T} = \widehat{\mathbf{E}} \left[\frac{C}{N_T} \right] + \int_0^T \widehat{\phi}_t d\widehat{X}_t, \quad (3.5.1)$$

where $(\widehat{X}_t)_{t \in [0, T]}$ is given by (3.4.1) and $(\widehat{\phi}_t)_{t \in [0, T]}$ is a square-integrable adapted process under $\widehat{\mathbb{P}}$, from which it follows that the forward claim price

$$\widehat{V}_t := \frac{V_t}{N_t} = \frac{1}{N_t} \mathbf{E}^* \left[e^{-\int_t^T r_s ds} C \mid \mathcal{F}_t \right] = \widehat{\mathbf{E}} \left[\frac{C}{N_T} \mid \mathcal{F}_t \right], \quad 0 \leq t \leq T,$$

is an \mathcal{F}_t -martingale under $\widehat{\mathbb{P}}$, that can be decomposed as

$$\widehat{V}_t = \widehat{\mathbf{E}}[\widehat{C} \mid \mathcal{F}_t] = \widehat{\mathbf{E}} \left[\frac{C}{N_T} \right] + \int_0^t \widehat{\phi}_s d\widehat{X}_s, \quad 0 \leq t \leq T. \quad (3.5.2)$$

The next Proposition 3.16 extends the argument of [Jamshidian, 1996](#) to the general framework of pricing using change of numéraire. Note that this result differs from the standard formula that uses the money market account $B_t = e^{\int_0^t r_s ds}$ for hedging instead of N_t , cf. e.g. [Geman, El Karoui, and Rochet, 1995](#) pages 453-454.

Proposition 3.16 Letting $\widehat{\eta}_t := \widehat{V}_t - \widehat{X}_t \widehat{\phi}_t$, with $\widehat{\phi}_t$ defined in (3.5.2), $0 \leq t \leq T$, the portfolio allocation

$$(\widehat{\phi}_t, \widehat{\eta}_t)_{t \in [0, T]}$$

with value

$$V_t = \widehat{\phi}_t X_t + \widehat{\eta}_t N_t, \quad 0 \leq t \leq T,$$

is self-financing in the sense that

$$dV_t = \widehat{\phi}_t dX_t + \widehat{\eta}_t dN_t,$$

and it hedges the claim payoff C , i.e.

$$V_t = \widehat{\phi}_t X_t + \widehat{\eta}_t N_t = \mathbf{E}^* \left[e^{-\int_t^T r_s ds} C \mid \mathcal{F}_t \right], \quad 0 \leq t \leq T. \quad (3.5.3)$$

Proof. In order to check that the portfolio allocation $(\widehat{\phi}_t, \widehat{\eta}_t)_{t \in [0, T]}$ hedges the claim payoff C it suffices to check that (3.5.3) holds since by (3.2.8) the price V_t at time $t \in [0, T]$ of the hedging portfolio satisfies

$$V_t = N_t \widehat{V}_t = \mathbf{E}^* \left[e^{-\int_t^T r_s ds} C \mid \mathcal{F}_t \right], \quad 0 \leq t \leq T.$$

Next, we show that the portfolio allocation $(\widehat{\phi}_t, \widehat{\eta}_t)_{t \in [0, T]}$ is self-financing. By *numéraire invariance*, cf. e.g. page 184 of [Protter, 2001](#), we have, using the relation $d\widehat{V}_t = \widehat{\phi}_t d\widehat{X}_t$ from (3.5.2),

$$dV_t = d(N_t \widehat{V}_t)$$

$$\begin{aligned}
&= \widehat{V}_t dN_t + N_t d\widehat{V}_t + dN_t \cdot d\widehat{V}_t \\
&= \widehat{V}_t dN_t + N_t \widehat{\phi}_t d\widehat{X}_t + \widehat{\phi}_t dN_t \cdot d\widehat{X}_t \\
&= \widehat{\phi}_t \widehat{X}_t dN_t + N_t \widehat{\phi}_t d\widehat{X}_t + \widehat{\phi}_t dN_t \cdot d\widehat{X}_t + (\widehat{V}_t - \widehat{\phi}_t \widehat{X}_t) dN_t \\
&= \widehat{\phi}_t d(N_t \widehat{X}_t) + \widehat{\eta}_t dN_t \\
&= \widehat{\phi}_t dX_t + \widehat{\eta}_t dN_t.
\end{aligned}$$

□

We now consider an application to the forward Delta hedging of European-type options with payoff $C = N_T \widehat{g}(\widehat{X}_T)$ where $\widehat{g} : \mathbb{R} \rightarrow \mathbb{R}$ and $(\widehat{X}_t)_{t \in \mathbb{R}_+}$ has the Markov property as in (3.4.2), where $\widehat{\sigma} : \mathbb{R}_+ \times \mathbb{R}$ is a *deterministic* function. Assuming that the function $\widehat{C}(t, x)$ defined by

$$\widehat{V}_t := \widehat{\mathbf{E}}[\widehat{g}(\widehat{X}_T) | \mathcal{F}_t] = \widehat{C}(t, \widehat{X}_t)$$

is \mathcal{C}^2 on \mathbb{R}_+ , we have the following corollary of Proposition 3.16, which extends the Black-Scholes Delta hedging technique to the general change of numéraire setup.

Corollary 3.17 Letting $\widehat{\eta}_t = \widehat{C}(t, \widehat{X}_t) - \widehat{X}_t \frac{\partial \widehat{C}}{\partial x}(t, \widehat{X}_t)$, $0 \leq t \leq T$, the portfolio allocation

$$\left(\frac{\partial \widehat{C}}{\partial x}(t, \widehat{X}_t), \widehat{\eta}_t \right)_{t \in [0, T]}$$

with value

$$V_t = \widehat{\eta}_t N_t + X_t \frac{\partial \widehat{C}}{\partial x}(t, \widehat{X}_t), \quad t \geq 0,$$

is self-financing and hedges the claim payoff $C = N_T \widehat{g}(\widehat{X}_T)$.

Proof. This result follows directly from Proposition 3.16 by noting that by Itô's formula, and the martingale property of \widehat{V}_t under $\widehat{\mathbb{P}}$ the stochastic integral representation (3.5.2) is given by

$$\begin{aligned}
\widehat{V}_T &= \widehat{C} \\
&= \widehat{g}(\widehat{X}_T) \\
&= \widehat{C}(0, \widehat{X}_0) + \int_0^T \frac{\partial \widehat{C}}{\partial t}(t, \widehat{X}_t) dt + \int_0^T \frac{\partial \widehat{C}}{\partial x}(t, \widehat{X}_t) d\widehat{X}_t \\
&\quad + \frac{1}{2} \int_0^T \frac{\partial^2 \widehat{C}}{\partial x^2}(t, \widehat{X}_t) |\widehat{\sigma}_t|^2 dt \\
&= \widehat{C}(0, \widehat{X}_0) + \int_0^T \frac{\partial \widehat{C}}{\partial x}(t, \widehat{X}_t) d\widehat{X}_t \\
&= \widehat{\mathbf{E}} \left[\frac{C}{N_T} \right] + \int_0^T \widehat{\phi}_t d\widehat{X}_t, \quad 0 \leq t \leq T,
\end{aligned}$$

hence

$$\widehat{\phi}_t = \frac{\partial \widehat{C}}{\partial x}(t, \widehat{X}_t), \quad 0 \leq t \leq T.$$

□

In the case of an exchange option with payoff function

$$C = (X_T - \kappa N_T)^+ = N_T (\widehat{X}_T - \kappa)^+$$



on the geometric Brownian motion $(\widehat{X}_t)_{t \in [0, T]}$ under $\widehat{\mathbb{P}}$ with

$$\widehat{\sigma}_t(\widehat{X}_t) = \widehat{\sigma}(t)\widehat{X}_t, \quad (3.5.4)$$

where $(\widehat{\sigma}(t))_{t \in [0, T]}$ is a deterministic volatility function of time, we have the following corollary on the hedging of exchange options based on the Margrabe, 1978 formula (3.4.4).

Corollary 3.18 The decomposition

$$\mathbb{E}^* \left[e^{-\int_t^T r_s ds} (X_T - \kappa N_T)^+ \mid \mathcal{F}_t \right] = X_t \Phi_+^0(t, \widehat{X}_t) - \kappa N_t \Phi_-^0(t, \widehat{X}_t)$$

yields a self-financing portfolio allocation $(\Phi_+^0(t, \widehat{X}_t), -\kappa \Phi_-^0(t, \widehat{X}_t))_{t \in [0, T]}$ in the assets (X_t, N_t) , that hedges the claim payoff $C = (X_T - \kappa N_T)^+$.

Proof. We apply Corollary 3.17 and the relation

$$\frac{\partial \widehat{C}}{\partial x}(t, x) = \Phi_+^0(t, x), \quad x \in \mathbb{R},$$

for the function $\widehat{C}(t, x) = x \Phi_+^0(t, x) - \kappa \Phi_-^0(t, x)$. \square

Note that the Delta hedging method requires the computation of the function $\widehat{C}(t, x)$ and that of the associated finite differences, and may not apply to path-dependent claims.

Examples:

- a) When the short rate process $(r(t))_{t \in [0, T]}$ is a *deterministic* function of time and $N_t = e^{\int_t^T r(s) ds}$, Corollary 3.18 yields the usual Black-Scholes hedging strategy

$$\begin{aligned} & (\Phi_+(t, \widehat{X}_t), -\kappa e^{\int_0^T r(s) ds} \Phi_-(t, X_t))_{t \in [0, T]} \\ &= (\Phi_+^0(t, e^{\int_t^T r(s) ds} \widehat{X}_t), -\kappa e^{\int_0^T r(s) ds} \Phi_-^0(t, e^{\int_t^T r(s) ds} X_t))_{t \in [0, T]}, \end{aligned}$$

in the assets $(X_t, e^{\int_0^t r(s) ds})$, that hedges the claim payoff $C = (X_T - \kappa)^+$, with

$$\Phi_+(t, x) := \Phi \left(\frac{\log(x/\kappa) + \int_t^T r(s) ds + (T-t)\sigma^2/2}{\sigma\sqrt{T-t}} \right),$$

and

$$\Phi_-(t, x) := \Phi \left(\frac{\log(x/\kappa) + \int_t^T r(s) ds - (T-t)\sigma^2/2}{\sigma\sqrt{T-t}} \right).$$

- b) In case $N_t = P(t, T)$ and $X_t = P(t, S)$, $0 \leq t \leq T < S$, Corollary 3.18 shows that when $(\widehat{X}_t)_{t \in [0, T]}$ is modeled as the geometric Brownian motion (3.5.4) under $\widehat{\mathbb{P}}$, the bond call option with payoff $(P(T, S) - \kappa)^+$ can be hedged as

$$\mathbb{E}^* \left[e^{-\int_t^T r_s ds} (P(T, S) - \kappa)^+ \mid \mathcal{F}_t \right] = P(t, S) \Phi_+(t, \widehat{X}_t) - \kappa P(t, T) \Phi_-(t, \widehat{X}_t)$$

by the self-financing portfolio allocation

$$(\Phi_+(t, \widehat{X}_t), -\kappa \Phi_-(t, \widehat{X}_t))_{t \in [0, T]}$$

in the assets $(P(t, S), P(t, T))$, i.e. one needs to hold the quantity $\Phi_+(t, \widehat{X}_t)$ of the bond maturing at time S , and to short a quantity $\kappa \Phi_-(t, \widehat{X}_t)$ of the bond maturing at time T .

Exercises

Exercise 3.1 Let $(B_t)_{t \in \mathbb{R}_+}$ be a standard Brownian motion started at 0 under the risk-neutral probability measure \mathbb{P}^* . Consider a numéraire $(N_t)_{t \in \mathbb{R}_+}$ given by

$$N_t := N_0 e^{\eta B_t - \eta^2 t / 2}, \quad t \geq 0,$$

and a risky asset $(X_t)_{t \in \mathbb{R}_+}$ given by

$$X_t := X_0 e^{\sigma B_t - \sigma^2 t / 2}, \quad t \geq 0,$$

in a market with risk-free interest rate $r = 0$. Let $\widehat{\mathbb{P}}$ denote the forward measure relative to the numéraire $(N_t)_{t \in \mathbb{R}_+}$, under which the process $\widehat{X}_t := X_t / N_t$ of forward prices is known to be a martingale.

- a) Using the Itô formula, compute

$$\begin{aligned} d\widehat{X}_t &= d\left(\frac{X_t}{N_t}\right) \\ &= \frac{X_0}{N_0} d(e^{(\sigma-\eta)B_t - (\sigma^2 - \eta^2)t/2}). \end{aligned}$$

- b) Explain why the exchange option price $\mathbb{E}^*[(X_T - \lambda N_T)^+]$ at time 0 has the Black-Scholes form

$$\begin{aligned} \mathbb{E}^*[(X_T - \lambda N_T)^+] & \quad (3.5.5) \\ &= X_0 \Phi\left(\frac{\log(\widehat{X}_0/\lambda)}{\widehat{\sigma}\sqrt{T}} + \frac{\widehat{\sigma}\sqrt{T}}{2}\right) - \lambda N_0 \Phi\left(\frac{\log(\widehat{X}_0/\lambda)}{\widehat{\sigma}\sqrt{T}} - \frac{\widehat{\sigma}\sqrt{T}}{2}\right). \end{aligned}$$

Hints:

- (i) Use the change of numéraire identity

$$\mathbb{E}^*[(X_T - \lambda N_T)^+] = N_0 \widehat{\mathbb{E}}[(\widehat{X}_T - \lambda)^+].$$

- (ii) The forward price \widehat{X}_t is a martingale under the forward measure $\widehat{\mathbb{P}}$ relative to the numéraire $(N_t)_{t \in \mathbb{R}_+}$.

- c) Give the value of $\widehat{\sigma}$ in terms of σ and η .

Exercise 3.2 Let $(B_t^{(1)})_{t \in \mathbb{R}_+}$ and $(B_t^{(2)})_{t \in \mathbb{R}_+}$ be correlated standard Brownian motions started at 0 under the risk-neutral probability measure \mathbb{P}^* , with correlation $\text{Corr}(B_s^{(1)}, B_t^{(2)}) = \rho \min(s, t)$, i.e. $dB_t^{(1)} \cdot dB_t^{(2)} = \rho dt$. Consider two asset prices $(S_t^{(1)})_{t \in \mathbb{R}_+}$ and $(S_t^{(2)})_{t \in \mathbb{R}_+}$ given by the geometric Brownian motions

$$S_t^{(1)} := S_0^{(1)} e^{rt + \sigma B_t^{(1)} - \sigma^2 t / 2}, \quad \text{and} \quad S_t^{(2)} := S_0^{(2)} e^{rt + \eta B_t^{(2)} - \eta^2 t / 2}, \quad t \geq 0.$$

Let $\widehat{\mathbb{P}}_2$ denote the forward measure with numéraire $(N_t)_{t \in \mathbb{R}_+} := (S_t^{(2)})_{t \in \mathbb{R}_+}$ and Radon-Nikodym density

$$\frac{d\widehat{\mathbb{P}}_2}{d\mathbb{P}^*} = e^{-rT} \frac{S_T^{(2)}}{S_0^{(2)}} = e^{\eta B_T^{(2)} - \eta^2 T / 2}.$$

- a) Using the Girsanov Theorem 3.7, determine the shifts $(\widehat{B}_t^{(1)})_{t \in \mathbb{R}_+}$ and $(\widehat{B}_t^{(2)})_{t \in \mathbb{R}_+}$ of $(B_t^{(1)})_{t \in \mathbb{R}_+}$ and $(B_t^{(2)})_{t \in \mathbb{R}_+}$ which are standard Brownian motions under $\widehat{\mathbb{P}}_2$.



b) Using the Itô formula, compute

$$\begin{aligned} d\hat{S}_t^{(1)} &= d\left(\frac{S_t^{(1)}}{S_t^{(2)}}\right) \\ &= \frac{S_0^{(1)}}{S_0^{(2)}} d\left(e^{\sigma B_t^{(1)} - \eta B_t^{(2)} - (\sigma^2 - \eta^2)t/2}\right), \end{aligned}$$

and write the answer in terms of the martingales $d\hat{B}_t^{(1)}$ and $d\hat{B}_t^{(2)}$.

c) Using change of numéraire, explain why the exchange option price

$$e^{-rT} \mathbb{E}^* [(S_T^{(1)} - \lambda S_T^{(2)})^+]$$

at time 0 has the Black-Scholes form

$$\begin{aligned} e^{-rT} \mathbb{E}^* [(S_T^{(1)} - \lambda S_T^{(2)})^+] &= S_0^{(1)} \Phi\left(\frac{\log(\hat{S}_0^{(1)}/\lambda)}{\hat{\sigma}\sqrt{T}} + \frac{\hat{\sigma}\sqrt{T}}{2}\right) \\ &\quad - \lambda S_0^{(2)} \Phi\left(\frac{\log(\hat{S}_0^{(1)}/\lambda)}{\hat{\sigma}\sqrt{T}} - \frac{\hat{\sigma}\sqrt{T}}{2}\right), \end{aligned}$$

where the value of $\hat{\sigma}$ can be expressed in terms of σ and η .

Exercise 3.3 Consider two zero-coupon bond prices of the form $P(t, T) = F(t, r_t)$ and $P(t, S) = G(t, r_t)$, where $(r_t)_{t \in \mathbb{R}_+}$ is a short-term interest rate process. Taking $N_t := P(t, T)$ as a numéraire defining the forward measure $\hat{\mathbb{P}}$, compute the dynamics of $(P(t, S))_{t \in [0, T]}$ under $\hat{\mathbb{P}}$ using a standard Brownian motion $(\hat{W}_t)_{t \in [0, T]}$ under $\hat{\mathbb{P}}$.

Exercise 3.4 Forward contracts. Using a change of numéraire argument for the numéraire $N_t := P(t, T)$, $t \in [0, T]$, compute the price at time $t \in [0, T]$ of a forward (or future) contract with payoff $P(T, S) - K$ in a bond market with short-term interest rate $(r_t)_{t \in \mathbb{R}_+}$. How would you hedge this forward contract?

Exercise 3.5 (Question 2.7 page 17 of [Downes, Joshi, and Denson, 2008](#)). Consider a price process $(S_t)_{t \in \mathbb{R}_+}$ given by $dS_t = rS_t dt + \sigma S_t dB_t$ under the risk-neutral probability measure \mathbb{P}^* , where $r \in \mathbb{R}$ and $\sigma > 0$, and the option with payoff

$$S_T (S_T - K)^+ = \max(S_T (S_T - K), 0)$$

at maturity T .

a) Show that the option payoff can be rewritten as

$$(S_T (S_T - K))^+ = N_T (S_T - K)^+$$

for a suitable choice of numéraire process $(N_t)_{t \in [0, T]}$.

- b) Rewrite the option price $e^{-(T-t)r} \mathbb{E}^* [(S_T (S_T - K))^+ | \mathcal{F}_t]$ using a forward measure $\hat{\mathbb{P}}$ and a change of numéraire argument.
- c) Find the dynamics of $(S_t)_{t \in \mathbb{R}_+}$ under the forward measure $\hat{\mathbb{P}}$.
- d) Price the option with payoff

$$S_T (S_T - K)^+ = \max(S_T (S_T - K), 0)$$

at time $t \in [0, T]$ using the Black-Scholes formula.

Exercise 3.6 Consider the risk asset with dynamics $dS_t = rS_t dt + \sigma S_t dW_t$ with constant $r \in \mathbb{R}$ and $\sigma > 0$ under the risk-neutral measure \mathbb{P}^* . The power call option with payoff $(S_T^n - K^n)^+$, $n \geq 1$, is priced at time $t \in [0, T]$ as

$$\begin{aligned} & e^{-(T-t)r} \mathbb{E}^* [(S_T^n - K^n)^+ | \mathcal{F}_t] \\ &= e^{-(T-t)r} \mathbb{E}^* [S_T^n \mathbb{1}_{\{S_T \geq K\}} | \mathcal{F}_t] - K^n e^{-(T-t)r} \mathbb{E}^* [\mathbb{1}_{\{S_T \geq K\}} | \mathcal{F}_t] \end{aligned}$$

under the risk-neutral measure \mathbb{P}^* .

- a) Write down the density $\frac{d\hat{\mathbb{P}}}{d\mathbb{P}^*}$ using the numéraire process

$$N_t := S_t^n e^{-(n-1)n\sigma^2 t/2 - (n-1)rt}, \quad t \in [0, T].$$

- b) Construct a standard Brownian motion \hat{W}_t under $\hat{\mathbb{P}}$.
 c) Compute the term $e^{-(T-t)r} \mathbb{E}^* [S_T^n \mathbb{1}_{\{S_T \geq K\}} | \mathcal{F}_t]$ using change of numéraire.

Hint: We have

$$\begin{aligned} \mathbb{P}^*(S_T \geq K | \mathcal{F}_t) &= \mathbb{E}^* [\mathbb{1}_{\{S_T \geq K\}} | \mathcal{F}_t] \\ &= \Phi \left(\frac{\log(S_t/K) + (r^l - \sigma^2/2)(T-t)}{\sigma\sqrt{T-t}} \right). \end{aligned}$$

Exercise 3.7 Bond options. Consider two bonds with maturities T and S , with prices $P(t, T)$ and $P(t, S)$ given by

$$\frac{dP(t, T)}{P(t, T)} = r_t dt + \zeta_t^T dW_t,$$

and

$$\frac{dP(t, S)}{P(t, S)} = r_t dt + \zeta_t^S dW_t,$$

where $(\zeta^T(s))_{s \in [0, T]}$ and $(\zeta^S(s))_{s \in [0, S]}$ are deterministic volatility functions of time.

- a) Show, using Itô's formula, that

$$d \left(\frac{P(t, S)}{P(t, T)} \right) = \frac{P(t, S)}{P(t, T)} (\zeta^S(t) - \zeta^T(t)) d\hat{W}_t,$$

where $(\hat{W}_t)_{t \in \mathbb{R}_+}$ is a standard Brownian motion under $\hat{\mathbb{P}}$.

- b) Show that

$$P(T, S) = \frac{P(t, S)}{P(t, T)} \exp \left(\int_t^T (\zeta^S(s) - \zeta^T(s)) d\hat{W}_s - \frac{1}{2} \int_t^T |\zeta^S(s) - \zeta^T(s)|^2 ds \right).$$

Let $\hat{\mathbb{P}}$ denote the forward measure associated to the numéraire

$$N_t := P(t, T), \quad 0 \leq t \leq T.$$

- c) Show that for all $S, T > 0$ the price at time t

$$\mathbb{E}^* \left[e^{-\int_t^T r_s ds} (P(T, S) - \kappa)^+ | \mathcal{F}_t \right]$$

of a bond call option on $P(T, S)$ with payoff $(P(T, S) - \kappa)^+$ is equal to

$$\mathbb{E}^* \left[e^{-\int_t^T r_s ds} (P(T, S) - \kappa)^+ | \mathcal{F}_t \right] \tag{3.5.6}$$

$$= P(t, S) \Phi \left(\frac{v}{2} + \frac{1}{v} \log \frac{P(t, S)}{\kappa P(t, T)} \right) - \kappa P(t, T) \Phi \left(-\frac{v}{2} + \frac{1}{v} \log \frac{P(t, S)}{\kappa P(t, T)} \right),$$

where

$$v^2 = \int_t^T |\zeta^S(s) - \zeta^T(s)|^2 ds.$$

- d) Compute the self-financing hedging strategy that hedges the bond option using a portfolio based on the assets $P(t, T)$ and $P(t, S)$.

Exercise 3.8 Consider two risky assets S_1 and S_2 modeled by the geometric Brownian motions

$$S_1(t) = e^{\sigma_1 W_t + \mu t} \quad \text{and} \quad S_2(t) = e^{\sigma_2 W_t + \mu t}, \quad t \geq 0, \quad (3.5.7)$$

where $(W_t)_{t \in \mathbb{R}_+}$ is a standard Brownian motion under \mathbb{P} .

- a) Find a condition on r, μ and σ_2 so that the discounted price process $e^{-rt} S_2(t)$ is a martingale under \mathbb{P} .
b) Assume that $r - \mu = \sigma_2^2/2$, and let

$$X_t = e^{(\sigma_2^2 - \sigma_1^2)t/2} S_1(t), \quad t \geq 0.$$

Show that the discounted process $e^{-rt} X_t$ is a martingale under \mathbb{P} .

- c) Taking $N_t = S_2(t)$ as numéraire, show that the forward process $\hat{X}(t) = X_t / N_t$ is a martingale under the forward measure $\hat{\mathbb{P}}$ defined by the Radon-Nikodym density

$$\frac{d\hat{\mathbb{P}}}{d\mathbb{P}} = e^{-rT} \frac{N_T}{N_0}.$$

Recall that

$$\hat{W}_t := W_t - \sigma_2 t$$

is a standard Brownian motion under $\hat{\mathbb{P}}$.

- d) Using the relation

$$e^{-rT} \mathbb{E}^*[(S_1(T) - S_2(T))^+] = N_0 \hat{\mathbb{E}} \left[\frac{(S_1(T) - S_2(T))^+}{N_T} \right],$$

compute the price

$$e^{-rT} \mathbb{E}^*[(S_1(T) - S_2(T))^+]$$

of the exchange option on the assets S_1 and S_2 .

Exercise 3.9 Compute the price $e^{-(T-t)r} \mathbb{E}^* [\mathbb{1}_{\{R_T \geq \kappa\}} | \mathcal{F}_t]$ at time $t \in [0, T]$ of a cash-or-nothing “binary” foreign exchange call option with maturity T and strike price κ on the foreign exchange rate process $(R_t)_{t \in \mathbb{R}_+}$ given by

$$dR_t = (r^I - r^F) R_t dt + \sigma R_t dW_t,$$

where $(W_t)_{t \in \mathbb{R}_+}$ is a standard Brownian motion under \mathbb{P}^* .

Hint: We have the relation

$$\mathbb{P}^*(x e^{X+\mu} \geq \kappa) = \Phi \left(\frac{\mu - \log(\kappa/x)}{\sqrt{\text{Var}[X]}} \right)$$

for $X \sim \mathcal{N}(0, \text{Var}[X])$ a centered Gaussian random variable.

Exercise 3.10 Extension of Proposition 3.8 to correlated Brownian motions. Assume that $(S_t)_{t \in \mathbb{R}_+}$ and $(N_t)_{t \in \mathbb{R}_+}$ satisfy the stochastic differential equations

$$dS_t = r_t S_t dt + \sigma_t^S S_t dW_t^S, \quad \text{and} \quad dN_t = \eta_t N_t dt + \sigma_t^N N_t dW_t^N,$$

where $(W_t^S)_{t \in \mathbb{R}_+}$ and $(W_t^N)_{t \in \mathbb{R}_+}$ have the correlation

$$dW_t^S \cdot dW_t^N = \rho dt,$$

where $\rho \in [-1, 1]$.

- a) Show that $(W_t^N)_{t \in \mathbb{R}_+}$ can be written as

$$W_t^N = \rho W_t^S + \sqrt{1 - \rho^2} W_t, \quad t \geq 0,$$

where $(W_t)_{t \in \mathbb{R}_+}$ is a standard Brownian motion under \mathbb{P}^* , independent of $(W_t^S)_{t \in \mathbb{R}_+}$.

- b) Letting $X_t = S_t/N_t$, show that dX_t can be written as

$$dX_t = (r_t - \eta_t + (\sigma_t^N)^2 - \rho \sigma_t^N \sigma_t^S) X_t dt + \widehat{\sigma}_t X_t dW_t^X,$$

where $(W_t^X)_{t \in \mathbb{R}_+}$ is a standard Brownian motion under \mathbb{P}^* and $\widehat{\sigma}_t$ is to be computed.

Exercise 3.11 Quanto options (Exercise 9.5 in Shreve, 2004). Consider an asset priced S_t at time t , with

$$dS_t = rS_t dt + \sigma^S S_t dW_t^S,$$

and an exchange rate $(R_t)_{t \in \mathbb{R}_+}$ given by

$$dR_t = (r - r^f) R_t dt + \sigma^R R_t dW_t^R,$$

from (3.3.3) in Proposition 3.10, where $(W_t^R)_{t \in \mathbb{R}_+}$ is written as

$$W_t^R = \rho W_t^S + \sqrt{1 - \rho^2} W_t, \quad t \geq 0,$$

where $(W_t)_{t \in \mathbb{R}_+}$ is a standard Brownian motion under \mathbb{P}^* , independent of $(W_t^S)_{t \in \mathbb{R}_+}$, i.e., we have

$$dW_t^R \cdot dW_t^S = \rho dt,$$

where ρ is a correlation coefficient.

- a) Let

$$a = r^I - r^f + \rho \sigma^R \sigma^S - (\sigma^R)^2$$

and $X_t = e^{at} S_t / R_t$, $t \geq 0$, and show by Exercise 3.10 that dX_t can be written as

$$dX_t = rX_t dt + \widehat{\sigma} X_t dW_t^X,$$

where $(W_t^X)_{t \in \mathbb{R}_+}$ is a standard Brownian motion under \mathbb{P}^* and $\widehat{\sigma}$ is to be determined.

- b) Compute the price

$$e^{-(T-t)r} \mathbb{E}^* \left[\left(\frac{S_T}{R_T} - \kappa \right)^+ \middle| \mathcal{F}_t \right]$$

of the quanto option at time $t \in [0, T]$.



4. Pricing of Interest Rate Derivatives

Interest rate derivatives are option contracts whose payoffs can be based on fixed-income securities such as bonds, or on cash flows exchanged in *e.g.* interest rate swaps. In this chapter we consider the pricing and hedging of interest rate and fixed income derivatives such as bond options, caplets, caps and swaptions, using the change of numéraire technique and forward measures.

4.1	Forward Measures and Tenor Structure	93
4.2	Bond Options	97
4.3	Caplet Pricing	98
4.4	Forward Swap Measures	102
4.5	Swaption Pricing	103
	Exercises	111

4.1 Forward Measures and Tenor Structure

The maturity dates are arranged according to a discrete *tenor structure*

$$\{0 = T_0 < T_1 < T_2 < \dots < T_n\}.$$

A sample of forward interest rate curve data is given in Table 4.1, which contains the values of $(T_1, T_2, \dots, T_{23})$ and of $\{f(t, t + T_i, t + T_i + \delta)\}_{i=1,2,\dots,23}$, with $t = 07/05/2003$ and $\delta = \text{six months}$.

Maturity	2D	1W	1M	2M	3M	1Y	2Y	3Y	4Y	5Y	6Y	7Y
Rate (%)	2.55	2.53	2.56	2.52	2.48	2.34	2.49	2.79	3.07	3.31	3.52	3.71
Maturity	8Y	9Y	10Y	11Y	12Y	13Y	14Y	15Y	20Y	25Y	30Y	
Rate (%)	3.88	4.02	4.14	4.23	4.33	4.40	4.47	4.54	4.74	4.83	4.86	

Table 4.1: Forward rates arranged according to a tenor structure.

Recall that by definition of $P(t, T_i)$ and absence of arbitrage the discounted bond price process

$$t \mapsto e^{-\int_0^t r_s ds} P(t, T_i), \quad 0 \leq t \leq T_i,$$

is an \mathcal{F}_t -martingale under the probability measure $\mathbb{P}^* = \mathbb{P}$, hence it satisfies the Assumption 1 on page 68 for $i = 1, 2, \dots, n$. As a consequence the bond price process can be taken as a numéraire

$$N_t^{(i)} := P(t, T_i), \quad 0 \leq t \leq T_i,$$

in the definition

$$\frac{d\hat{\mathbb{P}}_i}{d\mathbb{P}^*} = \frac{1}{P(0, T_i)} e^{-\int_0^{T_i} r_s ds} \quad (4.1.1)$$

of the *forward measure* $\hat{\mathbb{P}}_i$. The following proposition will allow us to price contingent claims using the forward measure $\hat{\mathbb{P}}_i$, it is a direct consequence of Proposition 3.5, noting that here we have $P(T_i, T_i) = 1$.

Proposition 4.1 For all sufficiently integrable random variables F we have

$$\mathbb{E}^* \left[F e^{-\int_t^{T_i} r_s ds} \mid \mathcal{F}_t \right] = P(t, T_i) \hat{\mathbb{E}}_i [F \mid \mathcal{F}_t], \quad 0 \leq t \leq T_i, \quad i = 1, 2, \dots, n. \quad (4.1.2)$$

Recall that by Proposition 3.4, the deflated process

$$t \mapsto \frac{P(t, T_j)}{P(t, T_i)}, \quad 0 \leq t \leq \min(T_i, T_j),$$

is an \mathcal{F}_t -martingale under $\hat{\mathbb{P}}_i$ for all $T_i, T_j \geq 0$.

In the sequel we assume as in (1.4.9) that the dynamics of the bond price $P(t, T_i)$ is given by

$$\frac{dP(t, T_i)}{P(t, T_i)} = r_t dt + \zeta_i(t) dW_t, \quad (4.1.3)$$

for $i = 1, 2, \dots, n$, where $(W_t)_{t \in \mathbb{R}_+}$ is a standard Brownian motion under \mathbb{P}^* and $(r_t)_{t \in \mathbb{R}_+}$ and $(\zeta_i(t))_{t \in \mathbb{R}_+}$ are adapted processes with respect to the filtration $(\mathcal{F}_t)_{t \in \mathbb{R}_+}$ generated by $(W_t)_{t \in \mathbb{R}_+}$, i.e.

$$P(t, T_i) = P(0, T_i) \exp \left(\int_0^t r_s ds + \int_0^t \zeta_i(s) dW_s - \frac{1}{2} \int_0^t |\zeta_i(s)|^2 ds \right),$$

$0 \leq t \leq T_i, i = 1, 2, \dots, n$.

Forward Brownian motions

Proposition 4.2 For all $i = 1, 2, \dots, n$, the process

$$\widehat{W}_t^i := W_t - \int_0^t \zeta_i(s) ds, \quad 0 \leq t \leq T_i, \quad (4.1.4)$$

is a standard Brownian motion under the forward measure $\widehat{\mathbb{P}}_i$.

Proof. The Girsanov Proposition 3.7 applied to the numéraire

$$N_t^{(i)} := P(t, T_i), \quad 0 \leq t \leq T_i,$$

as in (3.2.13), shows that

$$\begin{aligned} d\widehat{W}_t^i &:= dW_t - \frac{1}{N_t^{(i)}} dN_t^{(i)} \bullet dW_t \\ &= dW_t - \frac{1}{P(t, T_i)} dP(t, T_i) \bullet dW_t \\ &= dW_t - \frac{1}{P(t, T_i)} (P(t, T_i) r_t dt + \zeta_i(t) P(t, T_i) dW_t) \bullet dW_t \\ &= dW_t - \zeta_i(t) dt, \end{aligned}$$

is a standard Brownian motion under the forward measure $\widehat{\mathbb{P}}_i$ for all $i = 1, 2, \dots, n$. \square

We have

$$d\widehat{W}_t^i = dW_t - \zeta_i(t) dt, \quad i = 1, 2, \dots, n, \quad (4.1.5)$$

and

$$d\widehat{W}_t^j = dW_t - \zeta_j(t) dt = d\widehat{W}_t^i + (\zeta_i(t) - \zeta_j(t)) dt, \quad i, j = 1, 2, \dots, n,$$

which shows that $(\widehat{W}_t^j)_{t \in \mathbb{R}_+}$ has drift $(\zeta_i(t) - \zeta_j(t))_{t \in \mathbb{R}_+}$ under $\widehat{\mathbb{P}}_i$.

Bond price dynamics under the forward measure

In order to apply Proposition 4.1 and to compute the price

$$\mathbb{E}^* \left[e^{-\int_t^{T_i} r_s ds} C \mid \mathcal{F}_t \right] = P(t, T_i) \widehat{\mathbb{E}}_i [C \mid \mathcal{F}_t],$$

of a random claim payoff C , it can be useful to determine the dynamics of the underlying variables r_t , $f(t, T, S)$, and $P(t, T)$ via their stochastic differential equations written under the forward measure $\widehat{\mathbb{P}}_i$.

As a consequence of Proposition 4.2 and (4.1.3), the dynamics of $t \mapsto P(t, T_j)$ under $\widehat{\mathbb{P}}_i$ is given by

$$\frac{dP(t, T_j)}{P(t, T_j)} = r_t dt + \zeta_i(t) \zeta_j(t) dt + \zeta_j(t) d\widehat{W}_t^i, \quad i, j = 1, 2, \dots, n, \quad (4.1.6)$$

where $(\widehat{W}_t^i)_{t \in \mathbb{R}_+}$ is a standard Brownian motion under $\widehat{\mathbb{P}}_i$, and we have

$$\begin{aligned} P(t, T_j) &= P(0, T_j) \exp \left(\int_0^t r_s ds + \int_0^t \zeta_j(s) dW_s - \frac{1}{2} \int_0^t |\zeta_j(s)|^2 ds \right) && [\text{under } \mathbb{P}^*] \\ &= P(0, T_j) \exp \left(\int_0^t r_s ds + \int_0^t \zeta_j(s) d\widehat{W}_s^i + \frac{1}{2} \int_0^t |\zeta_j(s)|^2 ds \right) && [\text{under } \widehat{\mathbb{P}}_j] \end{aligned}$$

$$\begin{aligned}
&= P(0, T_j) \exp \left(\int_0^t r_s ds + \int_0^t \zeta_j(s) d\hat{W}_s^i + \int_0^t \zeta_j(s) \zeta_i(s) ds - \frac{1}{2} \int_0^t |\zeta_j(s)|^2 ds \right) \text{ [under } \hat{\mathbb{P}}_i] \\
&= P(0, T_j) \exp \left(\int_0^t r_s ds + \int_0^t \zeta_j(s) d\hat{W}_s^i - \frac{1}{2} \int_0^t |\zeta_j(s) - \zeta_i(s)|^2 ds + \frac{1}{2} \int_0^t |\zeta_i(s)|^2 ds \right),
\end{aligned}$$

$t \in [0, T_j]$, $i, j = 1, 2, \dots, n$. Consequently, the forward price $P(t, T_j) / P(t, T_i)$ can be written as

$$\begin{aligned}
&\frac{P(t, T_j)}{P(t, T_i)} \\
&= \frac{P(0, T_j)}{P(0, T_i)} \exp \left(\int_0^t (\zeta_j(s) - \zeta_i(s)) d\hat{W}_s^j + \frac{1}{2} \int_0^t |\zeta_j(s) - \zeta_i(s)|^2 ds \right) \text{ [under } \hat{\mathbb{P}}_j] \\
&= \frac{P(0, T_j)}{P(0, T_i)} \exp \left(\int_0^t (\zeta_j(s) - \zeta_i(s)) d\hat{W}_s^i - \frac{1}{2} \int_0^t |\zeta_i(s) - \zeta_j(s)|^2 ds \right), \text{ [under } \hat{\mathbb{P}}_i]
\end{aligned} \tag{4.1.7}$$

$t \in [0, \min(T_i, T_j)]$, $i, j = 1, 2, \dots, n$, which also follows from Proposition 3.8.

Short rate dynamics under the forward measure

In case the short rate process $(r_t)_{t \in \mathbb{R}_+}$ is given as the (Markovian) solution to the stochastic differential equation

$$dr_t = \mu(t, r_t) dt + \sigma(t, r_t) dW_t,$$

by (4.1.5) its dynamics will be given under $\hat{\mathbb{P}}_i$ by

$$\begin{aligned}
dr_t &= \mu(t, r_t) dt + \sigma(t, r_t) (\zeta_i(t) dt + d\hat{W}_t^i) \\
&= \mu(t, r_t) dt + \sigma(t, r_t) \zeta_i(t) dt + \sigma(t, r_t) d\hat{W}_t^i.
\end{aligned} \tag{4.1.8}$$

In the case of the Vašíček, 1977 model, by (1.4.10) we have

$$dr_t = (a - br_t) dt + \sigma dW_t,$$

and

$$\zeta_i(t) = -\frac{\sigma}{b} (1 - e^{-b(T_i-t)}), \quad 0 \leq t \leq T_i,$$

hence from (4.1.8) we have

$$d\hat{W}_t^i = dW_t - \zeta_i(t) dt = dW_t + \frac{\sigma}{b} (1 - e^{-b(T_i-t)}) dt,$$

and

$$dr_t = (a - br_t) dt - \frac{\sigma^2}{b} (1 - e^{-b(T_i-t)})^2 dt + \sigma d\hat{W}_t^i \tag{4.1.9}$$

and we obtain

$$\frac{dP(t, T_i)}{P(t, T_i)} = r_t dt + \frac{\sigma^2}{b^2} (1 - e^{-b(T_i-t)})^2 dt - \frac{\sigma}{b} (1 - e^{-b(T_i-t)}) d\hat{W}_t^i,$$

from (1.4.10).



4.2 Bond Options

The next proposition can be obtained as an application of the Margrabe formula (3.4.4) of Proposition 3.15 by taking $X_t = P(t, T_j)$, $N_t^{(i)} = P(t, T_i)$, and $\widehat{X}_t = X_t / N_t^{(i)} = P(t, T_j) / P(t, T_i)$. In the Vasicek model, this formula has been first obtained in [Jamshidian, 1989](#).

We work with a standard Brownian motion $(W_t)_{t \in \mathbb{R}_+}$ under \mathbb{P}^* , generating the filtration $(\mathcal{F}_t)_{t \in \mathbb{R}_+}$, and an $(\mathcal{F}_t)_{t \in \mathbb{R}_+}$ -adapted short rate process $(r_t)_{t \in \mathbb{R}_+}$.

Proposition 4.3 Let $0 \leq T_i \leq T_j$ and assume as in (1.4.9) that the dynamics of the bond prices $P(t, T_i)$, $P(t, T_j)$ under \mathbb{P}^* are given by

$$\frac{dP(t, T_i)}{P(t, T_i)} = r_i dt + \zeta_i(t) dW_t, \quad \frac{dP(t, T_j)}{P(t, T_j)} = r_j dt + \zeta_j(t) dW_t,$$

where $(\zeta_i(t))_{t \in \mathbb{R}_+}$ and $(\zeta_j(t))_{t \in \mathbb{R}_+}$ are *deterministic* volatility functions. Then the price of a bond call option on $P(T_i, T_j)$ with payoff

$$C := (P(T_i, T_j) - \kappa)^+$$

can be written as

$$\begin{aligned} & \mathbb{E}^* \left[e^{-\int_t^{T_i} r_s ds} (P(T_i, T_j) - \kappa)^+ \mid \mathcal{F}_t \right] \\ &= P(t, T_j) \Phi \left(\frac{v(t, T_i)}{2} + \frac{1}{v(t, T_i)} \log \frac{P(t, T_j)}{\kappa P(t, T_i)} \right) \\ & \quad - \kappa P(t, T_i) \Phi \left(-\frac{v(t, T_i)}{2} + \frac{1}{v(t, T_i)} \log \frac{P(t, T_j)}{\kappa P(t, T_i)} \right), \end{aligned} \tag{4.2.1}$$

where $v^2(t, T_i) := \int_t^{T_i} |\zeta_i(s) - \zeta_j(s)|^2 ds$ and

$$\Phi(x) := \frac{1}{\sqrt{2\pi}} \int_{-\infty}^x e^{-y^2/2} dy, \quad x \in \mathbb{R},$$

is the Gaussian cumulative distribution function.

Proof. First, we note that using $N_t^{(i)} := P(t, T_i)$ as a numéraire the price of a bond call option on $P(T_i, T_j)$ with payoff $F = (P(T_i, T_j) - \kappa)^+$ can be written from Proposition 3.5 using the forward measure $\widehat{\mathbb{P}}_i$, or directly by (3.2.9), as

$$\mathbb{E}^* \left[e^{-\int_t^{T_i} r_s ds} (P(T_i, T_j) - \kappa)^+ \mid \mathcal{F}_t \right] = P(t, T_i) \widehat{\mathbb{E}}_i [(P(T_i, T_j) - \kappa)^+ \mid \mathcal{F}_t]. \tag{4.2.2}$$

Next, by (4.1.7) or by solving (3.2.15) in Proposition 3.8 we can write $P(T_i, T_j)$ as the geometric Brownian motion

$$\begin{aligned} P(T_i, T_j) &= \frac{P(T_i, T_j)}{P(T_i, T_i)} \\ &= \frac{P(t, T_j)}{P(t, T_i)} \exp \left(\int_t^{T_i} (\zeta_j(s) - \zeta_i(s)) d\widehat{W}_s^i - \frac{1}{2} \int_t^{T_i} |\zeta_i(s) - \zeta_j(s)|^2 ds \right), \end{aligned}$$

under the forward measure $\widehat{\mathbb{P}}_i$, and rewrite (4.2.2) as

$$\begin{aligned} & \mathbb{E}^* \left[e^{-\int_t^{T_i} r_s ds} (P(T_i, T_j) - \kappa)^+ \mid \mathcal{F}_t \right] \\ &= P(t, T_i) \widehat{\mathbb{E}}_i \left[\left(\frac{P(t, T_j)}{P(t, T_i)} e^{\int_t^{T_i} (\zeta_j(s) - \zeta_i(s)) d\widehat{W}_s^i - \frac{1}{2} \int_t^{T_i} |\zeta_i(s) - \zeta_j(s)|^2 ds} - \kappa \right)^+ \mid \mathcal{F}_t \right] \\ &= \widehat{\mathbb{E}}_i \left[\left(P(t, T_j) e^{\int_t^{T_i} (\zeta_j(s) - \zeta_i(s)) d\widehat{W}_s^i - \frac{1}{2} \int_t^{T_i} |\zeta_i(s) - \zeta_j(s)|^2 ds} - \kappa P(t, T_i) \right)^+ \mid \mathcal{F}_t \right]. \end{aligned}$$

Since $(\zeta_i(s))_{s \in [0, T_i]}$ and $(\zeta_j(s))_{s \in [0, T_j]}$ in (4.1.3) are deterministic volatility functions, $P(T_i, T_j)$ is a lognormal random variable given \mathcal{F}_t under $\widehat{\mathbb{P}}_i$ and we price the bond option by the zero-rate Black-Scholes formula

$$Bl(P(t, T_j), \kappa P(t, T_i), v(t, T_i) / \sqrt{T_i - t}, 0, T_i - t)$$

with underlying asset price $P(t, T_j)$, strike level $\kappa P(t, T_i)$, volatility parameter

$$\frac{v(t, T_i)}{\sqrt{T_i - t}} = \sqrt{\frac{\int_t^{T_i} |\zeta_i(s) - \zeta_j(s)|^2 ds}{T_i - t}},$$

time to maturity $T_i - t$, and zero interest rate, which yields (4.2.1). \square

Note that from Corollary 3.17 the decomposition (4.2.1) gives the self-financing portfolio in the assets $P(t, T_i)$ and $P(t, T_j)$ for the claim with payoff $(P(T_i, T_j) - \kappa)^+$.

In the Vasicek case the above bond option price could also be computed from the joint distribution of $(r_T, \int_t^T r_s ds)$, which is Gaussian, or from the dynamics (4.1.6)-(4.1.9) of $P(t, T)$ and r_t under $\widehat{\mathbb{P}}_i$, cf. § 7.3 of Privault, 2012, and Kim, 2002 for the CIR and other short rate models with correlated Brownian motions.

4.3 Caplet Pricing

An interest rate caplet is an option contract that offers protection against the fluctuations of a variable (or floating) rate with respect to a fixed rate κ . The payoff of a caplet on the yield (or spot forward rate) $L(T_i, T_i, T_{i+1})$ with strike level κ can be written as

$$(L(T_i, T_i, T_{i+1}) - \kappa)^+,$$

and priced at time $t \in [0, T_i]$ from Proposition 3.5 using the forward measure $\widehat{\mathbb{P}}_{i+1}$ as

$$\begin{aligned} & \mathbb{E}^* \left[e^{-\int_t^{T_{i+1}} r_s ds} (L(T_i, T_i, T_{i+1}) - \kappa)^+ \mid \mathcal{F}_t \right] \\ &= P(t, T_{i+1}) \widehat{\mathbb{E}}_{i+1} [(L(T_i, T_i, T_{i+1}) - \kappa)^+ \mid \mathcal{F}_t], \end{aligned} \tag{4.3.1}$$

by taking $N_t^{(i+1)} = P(t, T_{i+1})$ as a numéraire.

Proposition 4.4 The LIBOR rate

$$L(t, T_i, T_{i+1}) := \frac{1}{T_{i+1} - T_i} \left(\frac{P(t, T_i)}{P(t, T_{i+1})} - 1 \right), \quad 0 \leq t \leq T_i < T_{i+1},$$

is a martingale under the forward measure $\widehat{\mathbb{P}}_{i+1}$ defined in (4.1.1).



Proof. The LIBOR rate $L(t, T_i, T_{i+1})$ is a deflated process according to the forward numéraire process $(P(t, T_{i+1}))_{t \in [0, T_{i+1}]}$. Therefore, by Proposition 3.4 it is a martingale under $\widehat{\mathbb{P}}_{i+1}$. \square

The caplet on $L(T_i, T_i, T_{i+1})$ can be priced at time $t \in [0, T_i]$ as

$$\begin{aligned} & \mathbf{E}^* \left[e^{-\int_t^{T_{i+1}} r_s ds} (L(T_i, T_i, T_{i+1}) - \kappa)^+ \middle| \mathcal{F}_t \right] \\ &= \mathbf{E}^* \left[e^{-\int_t^{T_{i+1}} r_s ds} \left(\frac{1}{T_{i+1} - T_i} \left(\frac{P(t, T_i)}{P(t, T_{i+1})} - 1 \right) - \kappa \right)^+ \middle| \mathcal{F}_t \right], \end{aligned} \quad (4.3.2)$$

where the discount factor is counted from the settlement date T_{i+1} . The next pricing formula (4.3.4) allows us to price and hedge a caplet using a portfolio based on the bonds $P(t, T_i)$ and $P(t, T_{i+1})$, cf. (4.3.8) below, when $L(t, T_i, T_{i+1})$ is modeled in the BGM model of Section 2.6.

Proposition 4.5 (Black caplet formula). Assume that $L(t, T_i, T_{i+1})$ is modeled in the BGM model as

$$\frac{dL(t, T_i, T_{i+1})}{L(t, T_i, T_{i+1})} = \gamma_i(t) d\widehat{W}_t^{i+1}, \quad (4.3.3)$$

$0 \leq t \leq T_i$, $i = 1, 2, \dots, n-1$, where $\gamma_i(t)$ is a deterministic volatility function of time $t \in [0, T_i]$, $i = 1, 2, \dots, n-1$. The caplet on $L(T_i, T_i, T_{i+1})$ with strike level κ is priced at time $t \in [0, T_i]$ as

$$\begin{aligned} & (T_{i+1} - T_i) \mathbf{E}^* \left[e^{-\int_t^{T_{i+1}} r_s ds} (L(T_i, T_i, T_{i+1}) - \kappa)^+ \middle| \mathcal{F}_t \right] \\ &= (P(t, T_i) - P(t, T_{i+1})) \Phi(d_+(t, T_i)) - \kappa (T_{i+1} - T_i) P(t, T_{i+1}) \Phi(d_-(t, T_i)), \end{aligned} \quad (4.3.4)$$

$0 \leq t \leq T_i$, where

$$d_+(t, T_i) = \frac{\log(L(t, T_i, T_{i+1}) / \kappa) + (T_i - t) \sigma_i^2(t, T_i) / 2}{\sigma_i(t, T_i) \sqrt{T_i - t}}, \quad (4.3.5)$$

and

$$d_-(t, T_i) = \frac{\log(L(t, T_i, T_{i+1}) / \kappa) - (T_i - t) \sigma_i^2(t, T_i) / 2}{\sigma_i(t, T_i) \sqrt{T_i - t}}, \quad (4.3.6)$$

and

$$|\sigma_i(t, T_i)|^2 = \frac{1}{T_i - t} \int_t^{T_i} |\gamma_i(s)|^2 ds. \quad (4.3.7)$$

Proof. Taking $P(t, T_{i+1})$ as a numéraire, the forward price

$$\widehat{X}_t := \frac{P(t, T_i)}{P(t, T_{i+1})} = 1 + (T_{i+1} - T_i) L(T_i, T_i, T_{i+1})$$

and the forward LIBOR rate process $(L(t, T_i, T_{i+1}))_{t \in [0, T_i]}$ are martingales under $\widehat{\mathbb{P}}_{i+1}$ by Proposition 4.4, $i = 1, 2, \dots, n-1$. More precisely, by (4.3.3) we have

$$L(T_i, T_i, T_{i+1}) = L(t, T_i, T_{i+1}) \exp \left(\int_t^{T_i} \gamma_i(s) d\widehat{W}_s^{i+1} - \frac{1}{2} \int_t^{T_i} |\gamma_i(s)|^2 ds \right),$$

$0 \leq t \leq T_i$, i.e. $t \mapsto L(t, T_i, T_{i+1})$ is a geometric Brownian motion with time-dependent volatility $\gamma_i(t)$ under $\widehat{\mathbb{P}}_{i+1}$. Hence by (4.3.1), since $N_{T_{i+1}}^{(i+1)} = 1$, we have

$$\begin{aligned} & \mathbf{E}^* \left[e^{-\int_t^{T_{i+1}} r_s ds} (L(T_i, T_i, T_{i+1}) - \kappa)^+ \mid \mathcal{F}_t \right] \\ &= P(t, T_{i+1}) \widehat{\mathbf{E}}_{i+1} [(L(T_i, T_i, T_{i+1}) - \kappa)^+ \mid \mathcal{F}_t] \\ &= P(t, T_{i+1}) (L(t, T_i, T_{i+1}) \Phi(d_+(t, T_i)) - \kappa \Phi(d_-(t, T_i))) \\ &= P(t, T_{i+1}) \text{Bl}(L(t, T_i, T_{i+1}), \kappa, \sigma_i(t, T_i), 0, T_i - t), \end{aligned}$$

$t \in [0, T_i]$, where

$$\text{Bl}(x, \kappa, \sigma, 0, \tau) = x \Phi(d_+(t, T_i)) - \kappa \Phi(d_-(t, T_i))$$

is the zero-interest rate Black-Scholes function, with

$$|\sigma_i(t, T_i)|^2 = \frac{1}{T_i - t} \int_t^{T_i} |\gamma_i(s)|^2 ds.$$

Therefore, we obtain

$$\begin{aligned} & (T_{i+1} - T_i) \mathbf{E}^* \left[e^{-\int_t^{T_{i+1}} r_s ds} (L(T_i, T_i, T_{i+1}) - \kappa)^+ \mid \mathcal{F}_t \right] \\ &= P(t, T_{i+1}) L(t, T_i, T_{i+1}) \Phi(d_+(t, T_i)) - \kappa P(t, T_{i+1}) \Phi(d_-(t, T_i)) \\ &= P(t, T_{i+1}) \left(\frac{P(t, T_i)}{P(t, T_{i+1})} - 1 \right) \Phi(d_+(t, T_i)) - \kappa (T_{i+1} - T_i) P(t, T_{i+1}) \Phi(d_-(t, T_i)), \end{aligned}$$

which yields (4.3.4). \square

In addition, by Corollary 3.17 we obtain the self-financing portfolio strategy

$$(\Phi(d_+(t, T_i)), -\Phi(d_+(t, T_i)) - \kappa (T_{i+1} - T_i) \Phi(d_-(t, T_i))) \quad (4.3.8)$$

in the bonds $(P(t, T_i), P(t, T_{i+1}))$ with maturities T_i and T_{i+1} , cf. Corollary 3.18 and Privault and Teng, 2012.

The formula (4.3.4) can be applied to options on underlying futures or forward contracts on commodities whose prices are modeled according to (4.3.3), as in the next corollary.

Corollary 4.6 (Black, 1976 formula). Let $L(t, T_i, T_{i+1})$ be modeled as in (4.3.3) and let the bond price $P(t, T_{i+1})$ be given as $P(t, T_{i+1}) = e^{-(T_{i+1}-t)r}$. Then, (4.3.4) becomes

$$e^{-(T_{i+1}-t)r} L(t, T_i, T_{i+1}) \Phi(d_+(t, T_i)) - \kappa e^{-(T_{i+1}-t)r} \Phi(d_-(t, T_i)),$$

$$0 \leq t \leq T_i.$$

Floorlet pricing

The floorlet on $L(T_i, T_i, T_{i+1})$ with strike level κ is a contract with payoff $(\kappa - L(T_i, T_i, T_{i+1}))^+$. Floorlets are analog to put options and can be similarly priced by the call/put parity in the Black-Scholes formula.

Proposition 4.7 Assume that $L(t, T_i, T_{i+1})$ is modeled in the BGM model as in (4.3.3). The floorlet on $L(T_i, T_i, T_{i+1})$ with strike level κ is priced at time $t \in [0, T_i]$ as



$$(T_{i+1} - T_i) \mathbf{E}^* \left[e^{-\int_t^{T_{i+1}} r_s ds} (\kappa - L(T_i, T_i, T_{i+1}))^+ \mid \mathcal{F}_t \right] \\ = \kappa (T_{i+1} - T_i) P(t, T_{i+1}) \Phi(-d_-(T_i - t)) - (P(t, T_i) - P(t, T_{i+1})) \Phi(-d_+(T_i - t)), \quad (4.3.9)$$

$0 \leq t \leq T_i$, where $d_+(t, T_i)$, $d_-(t, T_i)$ and $|\sigma_i(t, T_i)|^2$ are defined in (4.3.5)-(4.3.7).

Proof. We have

$$(T_{i+1} - T_i) \mathbf{E}^* \left[e^{-\int_t^{T_{i+1}} r_s ds} (\kappa - L(T_i, T_i, T_{i+1}))^+ \mid \mathcal{F}_t \right] \\ = (T_{i+1} - T_i) P(t, T_{i+1}) \widehat{\mathbf{E}}_{i+1} [(\kappa - L(T_i, T_i, T_{i+1}))^+ \mid \mathcal{F}_t] \\ = (T_{i+1} - T_i) P(t, T_{i+1}) (\kappa \Phi(-d_-(T_i - t)) - (T_{i+1} - T_i) L(t, T_i, T_{i+1}) \Phi(-d_+(T_i - t))) \\ = (T_{i+1} - T_i) P(t, T_{i+1}) \kappa \Phi(-d_-(T_i - t)) - (P(t, T_i) - P(t, T_{i+1})) \Phi(-d_+(T_i - t)),$$

$0 \leq t \leq T_i$. \square

Cap pricing

More generally, one can consider interest rate caps that are relative to a given tenor structure $\{T_1, T_2, \dots, T_n\}$, with discounted payoff

$$\sum_{k=1}^{n-1} (T_{k+1} - T_k) e^{-\int_t^{T_{k+1}} r_s ds} (L(T_k, T_k, T_{k+1}) - \kappa)^+.$$

Pricing formulas for interest rate caps are easily deduced from analog formulas for caplets, since the payoff of a cap can be decomposed into a sum of caplet payoffs. Thus, the cap price at time $t \in [0, T_1]$ is given by

$$\mathbf{E}^* \left[\sum_{k=1}^{n-1} (T_{k+1} - T_k) e^{-\int_t^{T_{k+1}} r_s ds} (L(T_k, T_k, T_{k+1}) - \kappa)^+ \mid \mathcal{F}_t \right] \\ = \sum_{k=1}^{n-1} (T_{k+1} - T_k) \mathbf{E}^* \left[e^{-\int_t^{T_{k+1}} r_s ds} (L(T_k, T_k, T_{k+1}) - \kappa)^+ \mid \mathcal{F}_t \right] \\ = \sum_{k=1}^{n-1} (T_{k+1} - T_k) P(t, T_{k+1}) \widehat{\mathbf{E}}_{k+1} [(L(T_k, T_k, T_{k+1}) - \kappa)^+ \mid \mathcal{F}_t]. \quad (4.3.10)$$

In the BGM model (4.3.3) the interest rate cap with payoff

$$\sum_{k=1}^{n-1} (T_{k+1} - T_k) (L(T_k, T_k, T_{k+1}) - \kappa)^+$$

can be priced at time $t \in [0, T_1]$ by the Black formula

$$\sum_{k=1}^{n-1} (T_{k+1} - T_k) P(t, T_{k+1}) \text{Bl}(L(t, T_k, T_{k+1}), \kappa, \sigma_k(t, T_k), 0, T_k - t),$$

where

$$|\sigma_k(t, T_k)|^2 = \frac{1}{T_k - t} \int_t^{T_k} |\gamma_k|^2(s) ds.$$

4.4 Forward Swap Measures

In this section we introduce the forward swap (or annuity) measures, or annuity measures, to be used for the pricing of swaptions, and we study their properties. We start with the definition of the *annuity numéraire*

$$N_t^{(i,j)} := P(t, T_i, T_j) = \sum_{k=i}^{j-1} (T_{k+1} - T_k) P(t, T_{k+1}), \quad 0 \leq t \leq T_i, \quad (4.4.1)$$

with in particular, when $j = i + 1$,

$$P(t, T_i, T_{i+1}) = (T_{i+1} - T_i) P(t, T_{i+1}), \quad 0 \leq t \leq T_i.$$

$1 \leq i < n$. The annuity numéraire can be also used to price a *bond ladder*. It satisfies the following martingale property, which can be proved by linearity and the fact that $t \mapsto e^{-\int_0^t r_s ds} P(t, T_k)$ is a martingale for all $k = 1, 2, \dots, n$, under Assumption 1.

R The discounted annuity numéraire

$$t \mapsto e^{-\int_0^t r_s ds} P(t, T_i, T_j) = e^{-\int_0^t r_s ds} \sum_{k=i}^{j-1} (T_{k+1} - T_k) P(t, T_{k+1}), \quad 0 \leq t \leq T_i,$$

is a martingale under \mathbb{P}^* .

The forward swap measure $\widehat{\mathbb{P}}_{i,j}$ is defined, according to Definition 3.1, by

$$\frac{d\widehat{\mathbb{P}}_{i,j}}{d\mathbb{P}^*} := e^{-\int_0^{T_i} r_s ds} \frac{N_{T_i}^{(i,j)}}{N_0^{(i,j)}} = e^{-\int_0^{T_i} r_s ds} \frac{P(T_i, T_i, T_j)}{P(0, T_i, T_j)}, \quad (4.4.2)$$

$1 \leq i < j \leq n$.

R We have

$$\begin{aligned} \mathbb{E}^* \left[\frac{d\widehat{\mathbb{P}}_{i,j}}{d\mathbb{P}^*} \middle| \mathcal{F}_t \right] &= \frac{1}{P(0, T_i, T_j)} \mathbb{E}^* \left[e^{-\int_0^{T_i} r_s ds} P(T_i, T_i, T_j) \middle| \mathcal{F}_t \right] \\ &= \frac{1}{P(0, T_i, T_j)} \mathbb{E}^* \left[e^{-\int_0^{T_i} r_s ds} \sum_{k=i}^{j-1} (T_{k+1} - T_k) P(T_i, T_{k+1}) \middle| \mathcal{F}_t \right] \\ &= \frac{1}{P(0, T_i, T_j)} \sum_{k=i}^{j-1} (T_{k+1} - T_k) \mathbb{E}^* \left[e^{-\int_0^{T_i} r_s ds} P(T_i, T_{k+1}) \middle| \mathcal{F}_t \right] \\ &= \frac{1}{P(0, T_i, T_j)} e^{-\int_0^{T_i} r_s ds} \sum_{k=i}^{j-1} (T_{k+1} - T_k) P(t, T_{k+1}) \\ &= e^{-\int_0^t r_s ds} \frac{P(t, T_i, T_j)}{P(0, T_i, T_j)}, \end{aligned}$$

$0 \leq t \leq T_i$, by Remark 4.4, and

$$\frac{d\widehat{\mathbb{P}}_{i,j}|_{\mathcal{F}_t}}{d\mathbb{P}^*|_{\mathcal{F}_t}} = e^{-\int_t^{T_i} r_s ds} \frac{P(T_i, T_i, T_j)}{P(t, T_i, T_j)}, \quad 0 \leq t \leq T_{i+1}, \quad (4.4.3)$$

by Relation (3.2.3) in Lemma 3.2.

Proposition 4.8 The LIBOR swap rate

$$S(t, T_i, T_j) = \frac{P(t, T_i) - P(t, T_j)}{P(t, T_i, T_j)} = v_i^{i,j}(t) - v_j^{i,j}(t), \quad 0 \leq t \leq T_i,$$

see Corollary 2.8, is a martingale under the forward swap measure $\widehat{\mathbb{P}}_{i,j}$.

Proof. We use the fact that the deflated process

$$t \mapsto v_k^{i,j}(t) := \frac{P(t, T_k)}{P(t, T_i, T_j)}, \quad i, j, k = 1, 2, \dots, n,$$

is an \mathcal{F}_t -martingale under $\widehat{\mathbb{P}}_{i,j}$ by Proposition 3.4. \square

The following pricing formula is then stated for a given integrable claim with payoff of the form $P(T_i, T_i, T_j)F$, using the forward swap measure $\widehat{\mathbb{P}}_{i,j}$:

$$\begin{aligned} \mathbf{E}^* \left[e^{-\int_t^{T_i} r_s ds} P(T_i, T_i, T_j) F \mid \mathcal{F}_t \right] &= P(t, T_i, T_j) \mathbf{E}^* \left[F \frac{d\widehat{\mathbb{P}}_{i,j}|_{\mathcal{F}_t}}{d\mathbb{P}^*|_{\mathcal{F}_t}} \mid \mathcal{F}_t \right] \\ &= P(t, T_i, T_j) \widehat{\mathbf{E}}_{i,j}[F \mid \mathcal{F}_t], \end{aligned} \quad (4.4.4)$$

after applying (4.4.2) and (4.4.3) on the last line, or Proposition 3.5.

4.5 Swaption Pricing

A payer (or call) swaption gives the option, but not the obligation, to enter an interest rate swap as payer of a fixed rate κ and as receiver of floating LIBOR rates $L(T_i, T_k, T_{k+1})$ at time T_{k+1} , $k = i, \dots, j-1$, and has the payoff

$$\begin{aligned} &\left(\sum_{k=i}^{j-1} (T_{k+1} - T_k) \mathbf{E}^* \left[e^{-\int_{T_i}^{T_{k+1}} r_s ds} \mid \mathcal{F}_{T_i} \right] (L(T_i, T_k, T_{k+1}) - \kappa) \right)^+ \\ &= \left(\sum_{k=i}^{j-1} (T_{k+1} - T_k) P(T_i, T_{k+1}) (L(T_i, T_k, T_{k+1}) - \kappa) \right)^+ \end{aligned} \quad (4.5.1)$$

at time T_i . This swaption can be priced at time $t \in [0, T_i]$ under the risk-neutral probability measure \mathbb{P}^* as

$$\mathbf{E}^* \left[e^{-\int_t^{T_i} r_s ds} \left(\sum_{k=i}^{j-1} (T_{k+1} - T_k) P(T_i, T_{k+1}) (L(T_i, T_k, T_{k+1}) - \kappa) \right)^+ \mid \mathcal{F}_t \right], \quad (4.5.2)$$

$t \in [0, T_i]$. When $j = i + 1$, the swaption price (4.5.2) coincides with the price at time t of a caplet on $[T_i, T_{i+1}]$ up to a factor $\delta_i := T_{i+1} - T_i$, since

$$\begin{aligned} &\mathbf{E}^* \left[e^{-\int_t^{T_i} r_s ds} ((T_{i+1} - T_i) P(T_i, T_{i+1}) (L(T_i, T_i, T_{i+1}) - \kappa))^+ \mid \mathcal{F}_t \right] \\ &= (T_{i+1} - T_i) \mathbf{E}^* \left[e^{-\int_t^{T_i} r_s ds} P(T_i, T_{i+1}) (L(T_i, T_i, T_{i+1}) - \kappa)^+ \mid \mathcal{F}_t \right] \\ &= (T_{i+1} - T_i) \mathbf{E}^* \left[e^{-\int_t^{T_i} r_s ds} \mathbf{E}^* \left[e^{-\int_{T_i}^{T_{i+1}} r_s ds} \mid \mathcal{F}_{T_i} \right] (L(T_i, T_i, T_{i+1}) - \kappa)^+ \mid \mathcal{F}_t \right] \\ &= (T_{i+1} - T_i) \mathbf{E}^* \left[\mathbf{E}^* \left[e^{-\int_t^{T_i} r_s ds} e^{-\int_{T_i}^{T_{i+1}} r_s ds} (L(T_i, T_i, T_{i+1}) - \kappa)^+ \mid \mathcal{F}_{T_i} \right] \mid \mathcal{F}_t \right] \end{aligned}$$

$$= (T_{i+1} - T_i) \mathbf{E}^* \left[e^{-\int_t^{T_{i+1}} r_s ds} (L(T_i, T_i, T_{i+1}) - \kappa)^+ \middle| \mathcal{F}_t \right], \quad (4.5.3)$$

$0 \leq t \leq T_i$, which coincides with the caplet price (4.3.1) up to the factor $T_{i+1} - T_i$. Unlike in the case of interest rate caps, the sum in (4.5.2) can not be taken out of the positive part. Nevertheless, the price of the swaption can be bounded as in the next proposition.

Proposition 4.9 The payer swaption price (4.5.2) can be upper bounded by the interest rate cap price (4.3.10) as

$$\begin{aligned} & \mathbf{E}^* \left[e^{-\int_t^{T_i} r_s ds} \left(\sum_{k=i}^{j-1} (T_{k+1} - T_k) P(T_i, T_{k+1}) (L(T_i, T_k, T_{k+1}) - \kappa) \right)^+ \middle| \mathcal{F}_t \right] \\ & \leq \sum_{k=i}^{j-1} (T_{k+1} - T_k) \mathbf{E}^* \left[e^{-\int_t^{T_{k+1}} r_s ds} (L(T_i, T_k, T_{k+1}) - \kappa)^+ \middle| \mathcal{F}_t \right], \end{aligned}$$

$$0 \leq t \leq T_i.$$

Proof. Due to the inequality

$$(x_1 + x_2 + \cdots + x_m)^+ \leq x_1^+ + x_2^+ + \cdots + x_m^+, \quad x_1, x_2, \dots, x_m \in \mathbb{R},$$

we have

$$\begin{aligned} & \mathbf{E}^* \left[e^{-\int_t^{T_i} r_s ds} \left(\sum_{k=i}^{j-1} (T_{k+1} - T_k) P(T_i, T_{k+1}) (L(T_i, T_k, T_{k+1}) - \kappa) \right)^+ \middle| \mathcal{F}_t \right] \\ & \leq \mathbf{E}^* \left[e^{-\int_t^{T_i} r_s ds} \sum_{k=i}^{j-1} (T_{k+1} - T_k) P(T_i, T_{k+1}) (L(T_i, T_k, T_{k+1}) - \kappa)^+ \middle| \mathcal{F}_t \right] \\ & = \sum_{k=i}^{j-1} (T_{k+1} - T_k) \mathbf{E}^* \left[e^{-\int_t^{T_i} r_s ds} P(T_i, T_{k+1}) (L(T_i, T_k, T_{k+1}) - \kappa)^+ \middle| \mathcal{F}_t \right] \\ & = \sum_{k=i}^{j-1} (T_{k+1} - T_k) \mathbf{E}^* \left[e^{-\int_t^{T_i} r_s ds} \mathbf{E}^* \left[e^{-\int_t^{T_{k+1}} r_s ds} \middle| \mathcal{F}_{T_i} \right] (L(T_i, T_k, T_{k+1}) - \kappa)^+ \middle| \mathcal{F}_t \right] \\ & = \sum_{k=i}^{j-1} (T_{k+1} - T_k) \mathbf{E}^* \left[\mathbf{E}^* \left[e^{-\int_t^{T_{k+1}} r_s ds} (L(T_i, T_k, T_{k+1}) - \kappa)^+ \middle| \mathcal{F}_{T_i} \right] \middle| \mathcal{F}_t \right] \\ & = \sum_{k=i}^{j-1} (T_{k+1} - T_k) \mathbf{E}^* \left[e^{-\int_t^{T_{k+1}} r_s ds} (L(T_i, T_k, T_{k+1}) - \kappa)^+ \middle| \mathcal{F}_t \right] \\ & = \mathbf{E}^* \left[\sum_{k=i}^{j-1} (T_{k+1} - T_k) e^{-\int_t^{T_{k+1}} r_s ds} (L(T_i, T_k, T_{k+1}) - \kappa)^+ \middle| \mathcal{F}_t \right], \end{aligned}$$

$$0 \leq t \leq T_i.$$

□

The payoff of the payer swaption can be rewritten as in the following lemma which is a direct consequence of the definition of the swap rate $S(T_i, T_i, T_j)$, see Proposition 2.7 and Corollary 2.8.

Lemma 4.10 The payer swaption payoff (4.5.1) at time T_i with swap rate $\kappa = S(t, T_j, T_j)$ can be rewritten as

$$\left(\sum_{k=i}^{j-1} (T_{k+1} - T_k) P(T_i, T_{k+1}) (L(T_i, T_k, T_{k+1}) - \kappa) \right)^+$$



$$= (P(T_i, T_i) - P(T_i, T_j) - \kappa P(T_i, T_i, T_j))^+ \quad (4.5.4)$$

$$= P(T_i, T_i, T_j) (S(T_i, T_i, T_j) - \kappa)^+. \quad (4.5.5)$$

Proof. The relation

$$\sum_{k=i}^{j-1} (T_{k+1} - T_k) P(t, T_{k+1}) (L(t, T_k, T_{k+1}) - S(t, T_i, T_j)) = 0$$

that defines the forward swap rate $S(t, T_i, T_j)$ shows that

$$\begin{aligned} & \sum_{k=i}^{j-1} (T_{k+1} - T_k) P(t, T_{k+1}) L(t, T_k, T_{k+1}) \\ &= S(t, T_i, T_j) \sum_{k=i}^{j-1} (T_{k+1} - T_k) P(t, T_{k+1}) \\ &= P(t, T_i, T_j) S(t, T_i, T_j) \\ &= P(t, T_i) - P(t, T_j) \end{aligned}$$

as in the proof of Corollary 2.8, hence by the definition (4.4.1) of $P(t, T_i, T_j)$ we have

$$\begin{aligned} & \sum_{k=i}^{j-1} (T_{k+1} - T_k) P(t, T_{k+1}) (L(t, T_k, T_{k+1}) - \kappa) \\ &= P(t, T_i) - P(t, T_j) - \kappa P(t, T_i, T_j) \\ &= P(t, T_i, T_j) (S(t, T_i, T_j) - \kappa), \end{aligned}$$

and for $t = T_i$ we get

$$\begin{aligned} & \left(\sum_{k=i}^{j-1} (T_{k+1} - T_k) P(T_i, T_{k+1}) (L(T_i, T_k, T_{k+1}) - \kappa) \right)^+ \\ &= P(T_i, T_i, T_j) (S(T_i, T_i, T_j) - \kappa)^+. \end{aligned}$$

□

The next proposition simply states that a payer swaption on the LIBOR rate can be priced as a European call option on the swap rate $S(T_i, T_i, T_j)$ under the forward swap measure $\widehat{\mathbb{P}}_{i,j}$.

Proposition 4.11 The price (4.5.2) of the payer swaption with payoff

$$\left(\sum_{k=i}^{j-1} (T_{k+1} - T_k) P(T_i, T_{k+1}) (L(T_i, T_k, T_{k+1}) - \kappa) \right)^+ \quad (4.5.6)$$

on the LIBOR market can be written under the forward swap measure $\widehat{\mathbb{P}}_{i,j}$ as the European call price

$$P(t, T_i, T_j) \widehat{\mathbb{E}}_{i,j} [(S(T_i, T_i, T_j) - \kappa)^+ | \mathcal{F}_t], \quad 0 \leq t \leq T_i,$$

on the swap rate $S(T_i, T_i, T_j)$.

Proof. As a consequence of (4.4.4) and Lemma 4.10, we find

$$\mathbb{E}^* \left[e^{-\int_t^{T_i} r_s ds} \left(\sum_{k=i}^{j-1} (T_{k+1} - T_k) P(T_i, T_{k+1}) (L(T_i, T_k, T_{k+1}) - \kappa) \right)^+ \middle| \mathcal{F}_t \right]$$

$$= \mathbf{E}^* \left[e^{-\int_t^{T_i} r_s ds} (P(T_i, T_i) - P(T_i, T_j) - \kappa P(T_i, T_i, T_j))^+ \middle| \mathcal{F}_t \right] \quad (4.5.7)$$

$$= \mathbf{E}^* \left[e^{-\int_t^{T_i} r_s ds} P(T_i, T_i, T_j) (S(T_i, T_i, T_j) - \kappa)^+ \middle| \mathcal{F}_t \right]$$

$$= P(t, T_i, T_j) \mathbf{E}^* \left[\frac{d\hat{\mathbb{P}}_{i,j|\mathcal{F}_t}}{d\mathbb{P}_{|\mathcal{F}_t}^*} (S(T_i, T_i, T_j) - \kappa)^+ \middle| \mathcal{F}_t \right]$$

$$= P(t, T_i, T_j) \hat{\mathbf{E}}_{i,j} [(S(T_i, T_i, T_j) - \kappa)^+ \mid \mathcal{F}_t]. \quad (4.5.8)$$

□

In the next Proposition 4.12 we price the payer swaption with payoff (4.5.6) or equivalently (4.5.5), by modeling the swap rate $(S(t, T_i, T_j))_{0 \leq t \leq T_i}$ using standard Brownian motion $(\hat{W}_t^{i,j})_{0 \leq t \leq T_i}$ under the swap forward measure $\hat{\mathbb{P}}_{i,j}$. See Exercise 4 for swaption pricing without the Black-Scholes formula.

Proposition 4.12 (Black swaption formula). Assume that the LIBOR swap rate (2.2.7) is modeled as a geometric Brownian motion under $\hat{\mathbb{P}}_{i,j}$, i.e.

$$dS(t, T_i, T_j) = S(t, T_i, T_j) \hat{\sigma}_{i,j}(t) d\hat{W}_t^{i,j}, \quad (4.5.9)$$

where $(\hat{\sigma}_{i,j}(t))_{t \in \mathbb{R}_+}$ is a deterministic volatility function of time. Then the payer swaption with payoff

$$(P(T, T_i) - P(T, T_j) - \kappa P(T_i, T_i, T_j))^+ = P(T_i, T_i, T_j) (S(T_i, T_i, T_j) - \kappa)^+$$

can be priced using the Black-Scholes call formula as

$$\begin{aligned} & \mathbf{E}^* \left[e^{-\int_t^{T_i} r_s ds} P(T_i, T_i, T_j) (S(T_i, T_i, T_j) - \kappa)^+ \middle| \mathcal{F}_t \right] \\ &= (P(t, T_i) - P(t, T_j)) \Phi(d_+(T_i - t)) \\ &\quad - \kappa \Phi(d_-(T_i - t)) \sum_{k=i}^{j-1} (T_{k+1} - T_k) P(t, T_{k+1}), \end{aligned}$$

where

$$d_+(T_i - t) = \frac{\log(S(t, T_i, T_j) / \kappa) + \sigma_{i,j}^2(t, T_i)(T_i - t)/2}{\sigma_{i,j}(t, T_i)\sqrt{T_i - t}}, \quad (4.5.10)$$

and

$$d_-(T_i - t) = \frac{\log(S(t, T_i, T_j) / \kappa) - \sigma_{i,j}^2(t, T_i)(T_i - t)/2}{\sigma_{i,j}(t, T_i)\sqrt{T_i - t}}, \quad (4.5.11)$$

and

$$|\sigma_{i,j}(t, T_i)|^2 = \frac{1}{T_i - t} \int_t^{T_i} |\hat{\sigma}_{i,j}(s)|^2 ds, \quad 0 \leq t \leq T. \quad (4.5.12)$$

Proof. Since $S(t, T_i, T_j)$ is a geometric Brownian motion with volatility function $(\hat{\sigma}(t))_{t \in \mathbb{R}_+}$ under $\hat{\mathbb{P}}_{i,j}$, by (4.5.4)-(4.5.5) in Lemma 4.10 or (4.5.7)-(4.5.8) we have

$$\mathbf{E}^* \left[e^{-\int_t^{T_i} r_s ds} P(T_i, T_i, T_j) (S(T_i, T_i, T_j) - \kappa)^+ \middle| \mathcal{F}_t \right]$$

$$= \mathbf{E}^* \left[e^{-\int_t^T r_s ds} (P(T, T_i) - P(T, T_j) - \kappa P(T_i, T_i, T_j))^+ \middle| \mathcal{F}_t \right]$$



$$\begin{aligned}
&= P(t, T_i, T_j) \widehat{\mathbf{E}}_{i,j} [(S(T_i, T_i, T_j) - \kappa)^+ | \mathcal{F}_t] \\
&= P(t, T_i, T_j) \text{Bl}(S(t, T_i, T_j), \kappa, \sigma_{i,j}(t, T_i), 0, T_i - t) \\
&= P(t, T_i, T_j) (S(t, T_i, T_j) \Phi_+(t, S(t, T_i, T_j)) - \kappa \Phi_-(t, S(t, T_i, T_j))) \\
&= (P(t, T_i) - P(t, T_j)) \Phi_+(t, S(t, T_i, T_j)) - \kappa P(t, T_i, T_j) \Phi_-(t, S(t, T_i, T_j)) \\
&= (P(t, T_i) - P(t, T_j)) \Phi_+(t, S(t, T_i, T_j)) \\
&\quad - \kappa \Phi_-(t, S(t, T_i, T_j)) \sum_{k=i}^{j-1} (T_{k+1} - T_k) P(t, T_{k+1}).
\end{aligned}$$

□

In addition, the hedging strategy

$$\begin{aligned}
&(\Phi_+(t, S(t, T_i, T_j)), -\kappa \Phi_-(t, S(t, T_i, T_j)) (T_{i+1} - T_i), \dots \\
&\dots, -\kappa \Phi_-(t, S(t, T_i, T_j)) (T_{j-1} - T_{j-2}), -\Phi_+(t, S(t, T_i, T_j)))
\end{aligned}$$

based on the assets $(P(t, T_i), \dots, P(t, T_j))$ is self-financing by Corollary 3.18, see also [Privault and Teng, 2012](#). Similarly to the above, a receiver (or put) swaption gives the option, but not the obligation, to enter an interest rate swap as receiver of a fixed rate κ and as payer of floating LIBOR rates $L(T_i, T_k, T_{k+1})$ at times T_{i+1}, \dots, T_j , and can be priced as in the next proposition.

Proposition 4.13 Assume that the LIBOR swap rate (2.2.7) is modeled as the geometric Brownian motion (4.5.9) under the forward swap measure $\widehat{P}_{i,j}$. Then the receiver swaption with payoff

$$(\kappa P(T_i, T_i, T_j) - (P(T, T_i) - P(T, T_j)))^+ = P(T_i, T_i, T_j) (\kappa - S(T_i, T_i, T_j))^+$$

can be priced using the Black-Scholes put formula as

$$\begin{aligned}
&\mathbb{E}^* \left[e^{-\int_t^{T_i} r_s ds} P(T_i, T_i, T_j) (\kappa - S(T_i, T_i, T_j))^+ \middle| \mathcal{F}_t \right] \\
&= \kappa \Phi(-d_-(T_i - t)) \sum_{k=i}^{j-1} (T_{k+1} - T_k) P(t, T_{k+1}) \\
&\quad - (P(t, T_i) - P(t, T_j)) \Phi(-d_+(T_i - t)),
\end{aligned}$$

where $d_+(T_i - t)$, and $d_-(T_i - t)$ and $|\sigma_{i,j}(t, T_i)|^2$ are defined in (4.5.10)-(4.5.12).

Swaption prices can also be computed by an approximation formula, from the exact dynamics of the swap rate $S(t, T_i, T_j)$ under the forward swap measure $\widehat{P}_{i,j}$, based on the bond price dynamics of the form (4.1.3), cf. [Schoenmakers, 2005](#), page 17.

Swaption volatilities can be estimated from swaption prices as implied volatilities from the Black pricing formula:




Figure 4.1: Implied swaption volatilities.

Implied swaption volatilities can then be used to calibrate the BGM model, cf. [Schoenmakers, 2005](#), [Privault and Wei, 2009](#), § 11.4 of [Privault, 2012](#).

Bermudan swaption pricing in Quantlib

The Bermudan swaption on the tenor structure $\{T_i, \dots, T_j\}$ is priced as the supremum

$$\begin{aligned} & \sup_{l \in \{i, \dots, j-1\}} \mathbb{E}^* \left[e^{-\int_t^{T_l} r_s ds} \left(\sum_{k=l}^{j-1} \delta_k P(T_l, T_{k+1}) (L(T_l, T_k, T_{k+1}) - \kappa) \right)^+ \middle| \mathcal{F}_t \right] \\ &= \sup_{l \in \{i, \dots, j-1\}} \mathbb{E}^* \left[e^{-\int_t^{T_l} r_s ds} (P(T_l, T_l) - P(T_l, T_j) - \kappa P(T_l, T_l, T_j))^+ \middle| \mathcal{F}_t \right] \\ &= \sup_{l \in \{i, \dots, j-1\}} \mathbb{E}^* \left[e^{-\int_t^{T_l} r_s ds} P(T_l, T_l, T_j) (S(T_l, T_l, T_j) - \kappa)^+ \middle| \mathcal{F}_t \right], \end{aligned}$$

where the supremum is over all stopping times taking values in $\{T_i, \dots, T_j\}$.

Bermudan swaptions can be priced using this [Rcode*](#) in (R)quantlib, with the following output:

Summary of pricing results for Bermudan Swaption

Price (in bp) of Bermudan swaption is 24.92137

Strike is NULL (ATM strike is 0.05)

Model used is: Hull-White using analytic formulas

Calibrated model parameters are:

a = 0.04641

sigma = 0.005869

This modified [code†](#) can be used in particular the pricing of ordinary swaptions, with the output:

Summary of pricing results for Bermudan Swaption

Price (in bp) of Bermudan swaption is 22.45436

Strike is NULL (ATM strike is 0.05)

Model used is: Hull-White using analytic formulas

*Click to open or download.

†Click to open or download.



Calibrated model parameters are:

a = 0.07107

sigma = 0.006018

Table 4.2 summarizes some possible uses of change of numéraire in option pricing.

Application	Asset price	Numéraire process	Option payoff	Forward measure \hat{P}	Dedicated process	Option price	Change of numéraire formula
Risk-neutral pricing	S_t	$N_t = e^{\int_0^t r_s ds}$	C	$\frac{d\hat{P}}{dP^*} = 1$	$\bar{S}_t = e^{-\int_0^t r_s ds} S_t$	$\mathbf{E}^* \left[e^{-\int_t^T r_s ds} C \mid \mathcal{F}_t \right]$	$e^{\int_0^T r_s ds} \mathbf{E}^* \left[e^{-\int_0^T r_s ds} C \mid \mathcal{F}_t \right]$
Exchange option	S_t	N_t	$(S_T - \kappa N_T)^+$	$\frac{d\hat{P}}{dP^*} = e^{-\int_0^T r_s ds} \frac{N_T}{N_0}$	$\hat{X}_t = \frac{S_t}{N_t}$	$\mathbf{E}^* \left[e^{-\int_t^T r_s ds} (S_T - \kappa N_T)^+ \mid \mathcal{F}_t \right]$	$N_t \hat{\mathbb{E}}_t \left[(\hat{X}_T - \kappa)^+ \mid \mathcal{F}_t \right]$
Exotic	$S_t = S_0 e^{rt + \sigma W_t - \sigma^2 t/2}$	$N_t = S_t$	$S_t (S_T - K)^+$	$\frac{d\hat{P}}{dP^*} = e^{-(T-t)r} \frac{S_T}{S_0}$	$\hat{X}_t = 1$	$\mathbf{E}^* \left[e^{-(T-t)r} (S_T - K)^+ \mid \mathcal{F}_t \right]$	$S_t \hat{\mathbb{E}}_t \left[(S_T - K)^+ \mid \mathcal{F}_t \right]$
Foreign exchange	$e^{rt} R_t$	$N_t = e^{rt} R_t$	$(R_T - \kappa)^+$	$\frac{d\hat{P}}{dP^*} = e^{(r'-r)T} \frac{R_T}{R_0}$	$\hat{X}_t = 1$	$e^{rT} \mathbf{E}^* \left[e^{-(T-t)r} (R_T - \kappa) \mid \mathcal{F}_t \right]$	$e^{rt} R_t \hat{\mathbb{E}}_t \left[\left(1 - \frac{\kappa}{R_t} \right)^+ \mid \mathcal{F}_t \right]$
Bond option	$P(t, S)$	$N_t = P(t, T)$	$(P(T, S) - K)^+$	$\frac{d\hat{P}_T}{dP^*} = \frac{e^{-\int_0^T r_s ds}}{P(0, T)}$	$\hat{X}_t = \frac{P(t, S)}{P(t, T)}$	$\mathbf{E}^* \left[e^{-\int_t^T r_s ds} (P(T, S) - K)^+ \mid \mathcal{F}_t \right]$	$P(t, T) \hat{\mathbb{E}}_t \left[(P(T, S) - K)^+ \mid \mathcal{F}_t \right]$
Caplets and caps	$P(t, T)$	$N_t = P(t, S)$	$(S - T)(L(T, T, S) - \kappa)^+$	$\frac{d\hat{P}_S}{dP^*} = \frac{e^{-\int_0^S r_s ds}}{P(0, S)}$	$L(t, T, S) = \frac{1}{S - T} \left(\frac{P(t, T)}{P(t, S)} - 1 \right)$	$(S - T) \mathbf{E}^* \left[e^{-\int_t^S r_s ds} (L(T, T, S) - \kappa)^+ \mid \mathcal{F}_t \right]$	$(S - T) \hat{\mathbb{E}}_S \left[(L(T, T, S) - \kappa)^+ \mid \mathcal{F}_t \right]$
Swaption	$P(t, T_1), P(t, T_n)$	$N_t = P(t, T_1, T_n)$	$(P(T_1, T_1) - P(T_1, T_n) - \kappa P(T_1, T_1, T_n))^+$	$\frac{d\hat{P}_{T_n}}{dP^*} = e^{-\int_0^{T_n} r_s ds} \frac{P(T_1, T_1, T_n)}{P(0, T_1, T_n)}$	$S(t, T_1, T_n) = \frac{P(t, T_1) - P(t, T_n)}{P(t, T_1, T_n)}$	$\mathbf{E}^* \left[e^{-\int_t^{T_n} r_s ds} (P(T_1, T_1) - P(T_1, T_n) - \kappa P(T_1, T_1, T_n))^+ \mid \mathcal{F}_t \right]$	$P(t, T_1, T_n) \hat{\mathbb{E}}_{t,n} \left[(S(T_1, T_1, T_n) - \kappa)^+ \mid \mathcal{F}_t \right]$
Black-Scholes	$S_t = S_0 e^{rt + \sigma W_t - \sigma^2 t/2}$	$N_t = S_t^{\alpha} e^{-(\alpha-1)\sigma^2 t/2 - (\alpha-1)r t}$	$(S_T^{\alpha} - K^{\alpha})^+$	$\frac{d\hat{P}}{dP^*} = e^{\alpha W_T - (\alpha-1)\sigma^2 T/2 + (\alpha-1)r T}$	$\hat{X}_t = S_T^{\alpha-1} e^{(\alpha-1)\sigma^2 T/2 + (\alpha-1)r T}$	$e^{-(T-t)r} \mathbf{E}^* \left[S_T^{\alpha} \mathbb{1}_{\{S_T > K\}} \mid \mathcal{F}_t \right] - K^{\alpha} e^{-(T-t)r} \mathbf{E}^* \left[\mathbb{1}_{\{S_T \geq K\}} \mid \mathcal{F}_t \right]$	$e^{(\alpha-1)\sigma^2 T/2 + (\alpha-1)r T} N \mathbb{E}^* \left[\mathbb{1}_{\{S_T > K\}} \mid \mathcal{F}_t \right] - K^{\alpha} e^{-(T-t)r} \mathbf{E}^* \left[\mathbb{1}_{\{S_T \geq K\}} \mid \mathcal{F}_t \right]$

Table 4.2: A list of numéraire processes and their applications.

Exercises

Exercise 4.1 Consider a floorlet on a three-month LIBOR rate in nine month's time, with a notional principal amount of \$10,000 per interest rate percentage point. The term structure is flat at 3.95% per year with discrete compounding, the volatility of the forward LIBOR rate in nine months is 10%, and the floor rate is 4.5%.

- What are the key assumptions on the LIBOR rate in nine month in order to apply Black's formula to price this floorlet?
- Compute the price of this floorlet using Black's formula as an application of Proposition 4.7 and (4.3.9), using the functions $\Phi(d_+)$ and $\Phi(d_-)$.

Exercise 4.2 Consider a payer swaption giving its holder the right, but not the obligation, to enter into a 3-year annual pay swap in four years, where a fixed rate of 5% will be paid and the LIBOR rate will be received. Assume that the yield curve is flat at 5% with continuous annual compounding and the volatility of the swap rate is 20%. The notional principal is \$100,000 per interest rate percentage point.

- What are the key assumptions in order to apply Black's formula to value this swaption?
- Compute the price of this swaption using Black's formula as an application of Proposition 4.12.

Exercise 4.3 Consider a *receiver* swaption giving its holder the right, but not the obligation, to enter into a 2-year annual pay swap in three years, where a fixed rate of 5% will be received and the LIBOR rate will be paid. Assume that the yield curve is flat at 2% with continuous annual compounding and the volatility of the swap rate is 10%. The notional principal is \$10,000 per percentage point, and the swaption price is quoted in basis points. Write down the expression of the price of this swaption using Black's formula.

Exercise 4.4 Consider two bonds with maturities T_1 and T_2 , $T_1 < T_2$, which follow the stochastic differential equations

$$dP(t, T_1) = r_t P(t, T_1) dt + \zeta_1(t) P(t, T_1) dW_t$$

and

$$dP(t, T_2) = r_t P(t, T_2) dt + \zeta_2(t) P(t, T_2) dW_t.$$

- Using Itô calculus, show that the forward process $P(t, T_2)/P(t, T_1)$ is a driftless geometric Brownian motion driven by $d\hat{W}_t := dW_t - \zeta_1(t)dt$ under the T_1 -forward measure $\hat{\mathbb{P}}$.
- Compute the price $\mathbb{E}^* \left[e^{-\int_t^{T_1} r_s ds} (K - P(T_1, T_2))^+ \mid \mathcal{F}_t \right]$ of a bond put option at time $t \in [0, T_1]$ using change of numéraire and the Black-Scholes formula.

Hint: Given X a Gaussian random variable with mean m and variance v^2 given \mathcal{F}_t , we have:

$$\begin{aligned} \mathbb{E}[(\kappa - e^X)^+ \mid \mathcal{F}_t] &= \kappa \Phi \left(-\frac{1}{v}(m - \log \kappa) \right) \\ &\quad - e^{m+v^2/2} \Phi \left(-\frac{1}{v}(m + v^2 - \log \kappa) \right). \end{aligned} \tag{4.5.13}$$

Exercise 4.5 Given two bonds with maturities T, S and prices $P(t, T), P(t, S)$, consider the LIBOR rate

$$L(t, T, S) := \frac{P(t, T) - P(t, S)}{(S - T)P(t, S)}$$

at time $t \in [0, T]$, modeled as

$$dL(t, T, S) = \mu_t L(t, T, S) dt + \sigma L(t, T, S) dW_t, \quad 0 \leq t \leq T, \quad (4.5.14)$$

where $(W_t)_{t \in [0, T]}$ is a standard Brownian motion under the risk-neutral probability measure \mathbb{P}^* , $\sigma > 0$ is a constant, and $(\mu_t)_{t \in [0, T]}$ is an adapted process. Let

$$F(t) := \mathbb{E}^* \left[e^{-\int_t^S r_s ds} (\kappa - L(T, T, S))^+ \mid \mathcal{F}_t \right]$$

denote the price at time t of a floorlet option with strike level κ , maturity T , and payment date S .

- a) Rewrite the value of $F(t)$ using the forward measure $\widehat{\mathbb{P}}_S$ with maturity S .
- b) What is the dynamics of $L(t, T, S)$ under the forward measure $\widehat{\mathbb{P}}_S$?
- c) Write down the value of $F(t)$ using the Black-Scholes formula.

Hint: Given X a centered Gaussian random variable with variance v^2 , we have

$$\mathbb{E}^*[(\kappa - e^{m+X})^+] = \kappa \Phi(-(m - \log \kappa)/v) - e^{m+v^2/2} \Phi(-v - (m - \log \kappa)/v),$$

where Φ denotes the Gaussian cumulative distribution function.

Exercise 4.6 Jamshidian's trick (Jamshidian, 1989). Consider a family $(P(t, T_k))_{k=i, \dots, j}$ of bond prices defined from a short rate process $(r_t)_{t \in \mathbb{R}_+}$. We assume that the bond prices are functions $P(T_i, T_{k+1}) = F_{k+1}(T_i, r_{T_i})$ of r_{T_i} that are *increasing* in the variable r_{T_i} , for all $k = i, i+1, \dots, j-1$.

- a) Compute the price $P(t, T_i, T_j)$ of the annuity numeraire paying coupons c_{i+1}, \dots, c_j at times T_{i+1}, \dots, T_j in terms of the bond prices

$$P(t, T_{i+1}), \dots, P(t, T_j).$$

- b) Show that the payoff

$$(P(T_i, T_i) - P(T_i, T_j) - \kappa P(T_i, T_i, T_j))^+$$

of a European swaption can be rewritten as

$$\left(1 - \kappa \sum_{k=i}^{j-1} \tilde{c}_{k+1} P(T_i, T_{k+1}) \right)^+,$$

by writing \tilde{c}_k in terms of c_k , $k = i+1, \dots, j$.

- c) Assuming that the bond prices are functions $P(T_i, T_{k+1}) = F_k(T_i, r_{T_i})$ of r_{T_i} that are *increasing* in the variable r_{T_i} , for all $k = i, \dots, j-1$, show, choosing γ_k such that

$$\kappa \sum_{k=i}^{j-1} c_{k+1} F_{k+1}(T_i, \gamma_k) = 1,$$

that the European swaption with payoff

$$(P(T_i, T_i) - P(T_i, T_j) - \kappa P(T_i, T_i, T_j))^+ = \left(1 - \kappa \sum_{k=i}^{j-1} c_{k+1} P(T_i, T_{k+1}) \right)^+,$$

where c_j contains the final coupon payment, can be priced as a weighted sum of bond put options under the forward measure $\widehat{\mathbb{P}}_i$ with numéraire $N_t^{(i)} := P(t, T_i)$.



Exercise 4.7 Path freezing. Consider n bonds with prices $(P(t, T_i))_{i=1,\dots,n}$ and the bond option with payoff

$$\left(\sum_{i=2}^n c_i P(T_0, T_i) - \kappa P(T_0, T_1) \right)^+ = P(T_0, T_1) (X_{T_0} - \kappa)^+,$$

where $N_t := P(t, T_1)$ is taken as numeraire and

$$X_t := \frac{1}{P(t, T_1)} \sum_{i=2}^n c_i P(t, T_i) = \sum_{i=2}^n c_i \widehat{P}(t, T_i), \quad 0 \leq t \leq T_1.$$

with $\widehat{P}(t, T_i) := P(t, T_i) / P(t, T_1)$, $i = 2, 3, \dots, n$.

- a) Assuming that the deflated bond price $(\widehat{P}(t, T_i))_{t \in [0, T_i]}$ has the (martingale) dynamics $d\widehat{P}(t, T_i) = \sigma_i(t) \widehat{P}(t, T_i) d\widehat{W}_t$ under the forward measure $\widehat{\mathbb{P}}_1$, where $(\sigma_i(t))_{t \in \mathbb{R}_+}$ is a deterministic function, write down the dynamics of X_t as $dX_t = \sigma_t X_t d\widehat{W}_t$, where σ_t is to be computed explicitly.
- b) Approximating $(\widehat{P}(t, T_i))_{t \in [0, T_i]}$ by $\widehat{P}(0, T_i)$ and $(P(t, T_2, T_n))_{t \in [0, T_2]}$ by $P(0, T_2, T_n)$, find a deterministic approximation $\widehat{\sigma}(t)$ of σ_t , and deduce an expression of the option price

$$\mathbb{E}^* \left[e^{-\int_0^{T_1} r_s ds} \left(\sum_{i=2}^n c_i P(T_0, T_i) - \kappa P(T_0, T_1) \right)^+ \right] = P(0, T_1) \widehat{\mathbb{E}}[(X_{T_0} - \kappa)^+]$$

using the Black-Scholes formula.

Hint: Given X a centered Gaussian random variable with variance v^2 , we have:

$$\mathbb{E}[(x e^{X-v^2/2} - \kappa)^+] = x \Phi(v/2 + (\log(x/\kappa))/v) - \kappa \Phi(-v/2 + (\log(x/\kappa))/v).$$

Exercise 4.8 (Exercise 1.4 continued). We work in the short rate model

$$dr_t = \sigma dB_t,$$

where $(B_t)_{t \in \mathbb{R}_+}$ is a standard Brownian motion under \mathbb{P}^* , and $\widehat{\mathbb{P}}_2$ is the forward measure defined by

$$\frac{d\widehat{\mathbb{P}}_2}{d\mathbb{P}^*} = \frac{1}{P(0, T_2)} e^{-\int_0^{T_2} r_s ds}.$$

- a) State the expressions of $\zeta_1(t)$ and $\zeta_2(t)$ in

$$\frac{dP(t, T_i)}{P(t, T_i)} = r_t dt + \zeta_i(t) dB_t, \quad i = 1, 2,$$

and the dynamics of the $P(t, T_1)/P(t, T_2)$ under $\widehat{\mathbb{P}}_2$, where $P(t, T_1)$ and $P(t, T_2)$ are bond prices with maturities T_1 and T_2 .

Hint: Use Exercise 1.4 and the relation (1.4.9).

- b) State the expression of the forward rate $f(t, T_1, T_2)$.
- c) Compute the dynamics of $f(t, T_1, T_2)$ under the forward measure $\widehat{\mathbb{P}}_2$ with

$$\frac{d\widehat{\mathbb{P}}_2}{d\mathbb{P}^*} = \frac{1}{P(0, T_2)} e^{-\int_0^{T_2} r_s ds}.$$

d) Compute the price

$$(T_2 - T_1) \mathbb{E}^* \left[e^{-\int_t^{T_2} r_s ds} (f(T_1, T_1, T_2) - \kappa)^+ \mid \mathcal{F}_t \right]$$

of an interest rate cap at time $t \in [0, T_1]$, using the expectation under the forward measure $\widehat{\mathbb{P}}_2$.

e) Compute the dynamics of the swap rate process

$$S(t, T_1, T_2) = \frac{P(t, T_1) - P(t, T_2)}{(T_2 - T_1)P(t, T_2)}, \quad t \in [0, T_1],$$

under $\widehat{\mathbb{P}}_2$.

f) Using (4.5.3), compute the swaption price

$$(T_2 - T_1) \mathbb{E}^* \left[e^{-\int_t^{T_1} r_s ds} P(T_1, T_2) (S(T_1, T_1, T_2) - \kappa)^+ \mid \mathcal{F}_t \right]$$

on the swap rate $S(T_1, T_1, T_2)$ using the expectation under the forward swap measure $\widehat{\mathbb{P}}_{1,2}$.

Exercise 4.9 Consider three zero-coupon bonds $P(t, T_1)$, $P(t, T_2)$ and $P(t, T_3)$ with maturities $T_1 = \delta$, $T_2 = 2\delta$ and $T_3 = 3\delta$ respectively, and the forward LIBOR $L(t, T_1, T_2)$ and $L(t, T_2, T_3)$ defined by

$$L(t, T_i, T_{i+1}) = \frac{1}{\delta} \left(\frac{P(t, T_i)}{P(t, T_{i+1})} - 1 \right), \quad i = 1, 2.$$

Assume that $L(t, T_1, T_2)$ and $L(t, T_2, T_3)$ are modeled in the BGM model by

$$\frac{dL(t, T_1, T_2)}{L(t, T_1, T_2)} = e^{-at} d\widehat{W}_t^{(2)}, \quad 0 \leq t \leq T_1, \tag{4.5.15}$$

and $L(t, T_2, T_3) = b$, $0 \leq t \leq T_2$, for some constants $a, b > 0$, where $\widehat{W}_t^{(2)}$ is a standard Brownian motion under the forward measure $\widehat{\mathbb{P}}_2$ defined by

$$\frac{d\widehat{\mathbb{P}}_2}{d\mathbb{P}^*} = \frac{e^{-\int_0^{T_2} r_s ds}}{P(0, T_2)}.$$

- a) Compute $L(t, T_1, T_2)$, $0 \leq t \leq T_2$ by solving Equation (4.5.15).
- b) Show that the price at time t of the caplet with strike level κ can be written as

$$\mathbb{E}^* \left[e^{-\int_t^{T_2} r_s ds} (L(T_1, T_1, T_2) - \kappa)^+ \mid \mathcal{F}_t \right] = P(t, T_2) \widehat{\mathbb{E}}_2 \left[(L(T_1, T_1, T_2) - \kappa)^+ \mid \mathcal{F}_t \right],$$

where $\widehat{\mathbb{E}}_2$ denotes the expectation under the forward measure $\widehat{\mathbb{P}}_2$.

- c) Using the hint below, compute the price at time t of the caplet with strike level κ on $L(T_1, T_1, T_2)$.
- d) Compute

$$\frac{P(t, T_1)}{P(t, T_1, T_3)}, \quad 0 \leq t \leq T_1, \quad \text{and} \quad \frac{P(t, T_3)}{P(t, T_1, T_3)}, \quad 0 \leq t \leq T_2,$$

in terms of b and $L(t, T_1, T_2)$, where $P(t, T_1, T_3)$ is the annuity numéraire

$$P(t, T_1, T_3) = \delta P(t, T_2) + \delta P(t, T_3), \quad 0 \leq t \leq T_2.$$



- e) Compute the dynamics of the swap rate

$$t \mapsto S(t, T_1, T_3) = \frac{P(t, T_1) - P(t, T_3)}{P(t, T_1, T_3)}, \quad 0 \leq t \leq T_1,$$

i.e. show that we have

$$dS(t, T_1, T_3) = \sigma_{1,3}(t) S(t, T_1, T_3) d\hat{W}_t^{(2)},$$

where $\sigma_{1,3}(t)$ is a stochastic process to be determined.

- f) Using the Black-Scholes formula, compute an approximation of the swaption price

$$\begin{aligned} & \mathbf{E}^* \left[e^{-\int_t^{T_1} r_s ds} P(T_1, T_1, T_3) (S(T_1, T_1, T_3) - \kappa)^+ \mid \mathcal{F}_t \right] \\ &= P(t, T_1, T_3) \widehat{\mathbf{E}}_2 [(S(T_1, T_1, T_3) - \kappa)^+ \mid \mathcal{F}_t], \end{aligned}$$

at time $t \in [0, T_1]$. You will need to approximate $\sigma_{1,3}(s)$, $s \geq t$, by “freezing” all random terms at time t .

Hint: Given X a centered Gaussian random variable with variance v^2 , we have

$$\mathbf{E}^* [(e^{m+X} - \kappa)^+] = e^{m+v^2/2} \Phi(v + (m - \log \kappa)/v) - \kappa \Phi((m - \log \kappa)/v),$$

where Φ denotes the Gaussian cumulative distribution function.

Exercise 4.10 Bond option hedging. Consider a portfolio allocation $(\xi_t^T, \xi_t^S)_{t \in [0, T]}$ made of two bonds with maturities T, S , and value

$$V_t = \xi_t^T P(t, T) + \xi_t^S P(t, S), \quad 0 \leq t \leq T,$$

at time t . We assume that the portfolio is self-financing, i.e.

$$dV_t = \xi_t^T dP(t, T) + \xi_t^S dP(t, S), \quad 0 \leq t \leq T, \tag{4.5.16}$$

and that it *hedges* the claim payoff $(P(T, S) - \kappa)^+$, so that

$$\begin{aligned} V_t &= \mathbf{E}^* \left[e^{-\int_t^T r_s ds} (P(T, S) - \kappa)^+ \mid \mathcal{F}_t \right] \\ &= P(t, T) \mathbf{E}_T [(P(T, S) - \kappa)^+ \mid \mathcal{F}_t], \quad 0 \leq t \leq T. \end{aligned}$$

- a) Show that we have

$$\begin{aligned} & \mathbf{E}^* \left[e^{-\int_t^T r_s ds} (P(T, S) - K)^+ \mid \mathcal{F}_t \right] \\ &= P(0, T) \mathbf{E}_T [(P(T, S) - K)^+] + \int_0^t \xi_s^T dP(s, T) + \int_0^t \xi_s^S dP(s, S). \end{aligned}$$

- b) Show that under the self-financing condition (4.5.16), the deflated portfolio value $\tilde{V}_t = e^{-\int_0^t r_s ds} V_t$ satisfies

$$d\tilde{V}_t = \xi_t^T d\tilde{P}(t, T) + \xi_t^S d\tilde{P}(t, S),$$

where

$$\tilde{P}(t, T) := e^{-\int_0^t r_s ds} P(t, T), \quad t \in [0, T],$$

and

$$\tilde{P}(t, S) := e^{-\int_0^t r_s ds} P(t, S), \quad t \in [0, S],$$

denote the discounted bond prices.

- c) From now on we work in the framework of Proposition 4.3, and we let the function $C(x, v)$ be defined by

$$C(X_t, v(t, T)) := \mathbf{E}_T [(P(T, S) - K)^+ | \mathcal{F}_t],$$

where X_t is the forward price $X_t := P(t, S) / P(t, T)$, $t \in [0, T]$, and

$$\nu^2(t, T) := \int_t^T |\sigma_s^S - \sigma_s^T|^2 ds.$$

Show that

$$\begin{aligned} \mathbf{E}_T [(P(T, S) - K)^+ | \mathcal{F}_t] &= \mathbf{E}_T [(P(T, S) - K)^+] \\ &\quad + \int_0^t \frac{\partial C}{\partial x}(X_u, v(u, T)) dX_u, \quad t \geq 0. \end{aligned}$$

Hint: Use the martingale property and the Itô formula.

- d) Show that the deflated portfolio value $\widehat{V}_t = V_t / P(t, T)$ satisfies

$$\begin{aligned} d\widehat{V}_t &= \frac{\partial C}{\partial x}(X_t, v(t, T)) dX_t \\ &= \frac{P(t, S)}{P(t, T)} \frac{\partial C}{\partial x}(X_t, v(t, T)) (\sigma_t^S - \sigma_t^T) d\widehat{B}_t^T. \end{aligned}$$

- e) Show that

$$dV_t = P(t, S) \frac{\partial C}{\partial x}(X_t, v(t, T)) (\sigma_t^S - \sigma_t^T) dB_t + \widehat{V}_t dP(t, T).$$

- f) Show that

$$d\widetilde{V}_t = \widetilde{P}(t, S) \frac{\partial C}{\partial x}(X_t, v(t, T)) (\sigma_t^S - \sigma_t^T) dB_t + \widehat{V}_t d\widetilde{P}(t, T).$$

- g) Compute the hedging strategy $(\xi_t^T, \xi_t^S)_{t \in [0, T]}$ of the bond option.

- h) Show that

$$\frac{\partial C}{\partial x}(x, v) = \Phi \left(\frac{\log(x/K) + \tau v^2 / 2}{\sqrt{\tau} v} \right),$$

and compute the hedging strategy $(\xi_t^T, \xi_t^S)_{t \in [0, T]}$ in terms of the normal cumulative distribution function Φ .

Exercise 4.11 Consider a LIBOR rate $L(t, T, S)$, $t \in [0, T]$, modeled as $dL(t, T, S) = \mu_t L(t, T, S) dt + \sigma(t) L(t, T, S) dW_t$, $0 \leq t \leq T$, where $(W_t)_{t \in [0, T]}$ is a standard Brownian motion under the risk-neutral probability measure \mathbb{P}^* , $(\mu_t)_{t \in [0, T]}$ is an adapted process, and $\sigma(t) > 0$ is a deterministic volatility function of time t .

- a) What is the dynamics of $L(t, T, S)$ under the forward measure $\widehat{\mathbb{P}}$ with numéraire $N_t := P(t, S)$?
b) Rewrite the price

$$\mathbf{E}^* \left[e^{-\int_t^S r_s ds} \phi(L(T, T, S)) | \mathcal{F}_t \right] \tag{4.5.17}$$

at time $t \in [0, T]$ of an option with payoff function ϕ using the forward measure $\widehat{\mathbb{P}}$.

- c) Write down the above option price (4.5.17) using an integral.

Exercise 4.12 Given n bonds with maturities T_1, T_2, \dots, T_n , consider the annuity numéraire

$$P(t, T_i, T_j) = \sum_{k=i}^{j-1} (T_{k+1} - T_k) P(t, T_{k+1})$$

and the swap rate

$$S(t, T_i, T_j) = \frac{P(t, T_i) - P(t, T_j)}{P(t, T_i, T_j)}$$



at time $t \in [0, T_i]$, modeled as

$$dS(t, T_i, T_j) = \mu_t S(t, T_i, T_j) dt + \sigma S(t, T_i, T_j) dW_t, \quad 0 \leq t \leq T_i, \quad (4.5.18)$$

where $(W_t)_{t \in [0, T_i]}$ is a standard Brownian motion under the risk-neutral probability measure \mathbb{P}^* , $(\mu_t)_{t \in [0, T]}$ is an adapted process and $\sigma > 0$ is a constant. Let

$$\mathbb{E}^* \left[e^{-\int_t^{T_i} r_s ds} P(T_i, T_i, T_j) \phi(S(T_i, T_i, T_j)) \mid \mathcal{F}_t \right] \quad (4.5.19)$$

at time $t \in [0, T_i]$ of an option with payoff function ϕ .

- a) Rewrite the option price (4.5.19) at time $t \in [0, T_i]$ using the forward swap measure $\widehat{\mathbb{P}}_{i,j}$ defined from the annuity numéraire $P(t, T_i, T_j)$.
- b) What is the dynamics of $S(t, T_i, T_j)$ under the forward swap measure $\widehat{\mathbb{P}}_{i,j}$?
- c) Write down the above option price (4.5.17) using a Gaussian integral.
- d) Apply the above to the computation at time $t \in [0, T_i]$ of the put swaption price

$$\mathbb{E}^* \left[e^{-\int_t^{T_i} r_s ds} P(T_i, T_i, T_j) (\kappa - S(T_i, T_i, T_j))^+ \mid \mathcal{F}_t \right]$$

with strike level κ , using the Black-Scholes formula.

Hint: Given X a centered Gaussian random variable with variance v^2 , we have

$$\mathbb{E}[(\kappa - e^{m+X})^+] = \kappa \Phi(-(m - \log \kappa)/v) - e^{m+v^2/2} \Phi(-v - (m - \log \kappa)/v),$$

where Φ denotes the Gaussian cumulative distribution function.

Exercise 4.13 Consider a bond market with two bonds with maturities T_1, T_2 , whose prices $P(t, T_1), P(t, T_2)$ at time t are given by

$$\frac{dP(t, T_1)}{P(t, T_1)} = r_t dt + \zeta_1(t) dB_t, \quad \frac{dP(t, T_2)}{P(t, T_2)} = r_t dt + \zeta_2(t) dB_t,$$

where $(r_t)_{t \in \mathbb{R}_+}$ is a short-term interest rate process, $(B_t)_{t \in \mathbb{R}_+}$ is a standard Brownian motion generating a filtration $(\mathcal{F}_t)_{t \in \mathbb{R}_+}$, and $\zeta_1(t), \zeta_2(t)$ are volatility processes. The LIBOR rate $L(t, T_1, T_2)$ is defined by

$$L(t, T_1, T_2) = \frac{P(t, T_1) - P(t, T_2)}{P(t, T_2)}.$$

Recall that a caplet on the LIBOR market can be priced at time $t \in [0, T_1]$ as

$$\begin{aligned} & \mathbb{E} \left[e^{-\int_t^{T_2} r_s ds} (L(T_1, T_1, T_2) - \kappa)^+ \mid \mathcal{F}_t \right] \\ &= P(t, T_2) \widehat{\mathbb{E}} \left[(L(T_1, T_1, T_2) - \kappa)^+ \mid \mathcal{F}_t \right], \end{aligned} \quad (4.5.20)$$

under the forward measure $\widehat{\mathbb{P}}$ defined by

$$\frac{d\widehat{\mathbb{P}}}{d\mathbb{P}^*} = e^{-\int_0^{T_1} r_s ds} \frac{P(T_1, T_2)}{P(0, T_2)},$$

under which

$$\widehat{B}_t := B_t - \int_0^t \zeta_2(s) ds, \quad t \in \mathbb{R}_+, \quad (4.5.21)$$

is a standard Brownian motion.

In the sequel we let $L_t = L(t, T_1, T_2)$ for simplicity of notation.

- a) Using Itô calculus, show that the LIBOR rate satisfies

$$dL_t = L_t \sigma(t) d\hat{B}_t, \quad 0 \leq t \leq T_1, \quad (4.5.22)$$

where the LIBOR rate volatility is given by

$$\sigma(t) = \frac{P(t, T_1)(\zeta_1(t) - \zeta_2(t))}{P(t, T_1) - P(t, T_2)}.$$

- b) Solve the equation (4.5.22) on the interval $[t, T_1]$, and compute L_{T_1} from the initial condition L_t .
c) Assuming that $\sigma(t)$ in (4.5.22) is a deterministic volatility function of time t , show that the price

$$P(t, T_2) \widehat{\mathbb{E}}[(L_{T_1} - \kappa)^+ | \mathcal{F}_t]$$

of the caplet can be written as $P(t, T_2)C(L_t, v(t, T_1))$, where $v^2(t, T_1) = \int_t^{T_1} |\sigma(s)|^2 ds$, and $C(L_t, v(t, T_1))$ is a function of L_t and $v(t, T_1)$.

- d) Consider a portfolio allocation $(\xi_t^{(1)}, \xi_t^{(2)})_{t \in [0, T_1]}$ made of bonds with maturities T_1, T_2 and value

$$V_t = \xi_t^{(1)} P(t, T_1) + \xi_t^{(2)} P(t, T_2),$$

at time $t \in [0, T_1]$. We assume that the portfolio is self-financing, i.e.

$$dV_t = \xi_t^{(1)} dP(t, T_1) + \xi_t^{(2)} dP(t, T_2), \quad 0 \leq t \leq T_1, \quad (4.5.23)$$

and that it *hedges* the claim payoff $(L_{T_1} - \kappa)^+$, so that

$$\begin{aligned} V_t &= \mathbb{E} \left[e^{-\int_t^{T_1} r_s ds} (P(T_1, T_2)(L_{T_1} - \kappa))^+ \middle| \mathcal{F}_t \right] \\ &= P(t, T_2) \widehat{\mathbb{E}}[(L_{T_1} - \kappa)^+ | \mathcal{F}_t], \end{aligned}$$

$0 \leq t \leq T_1$. Show that we have

$$\begin{aligned} &\mathbb{E} \left[e^{-\int_t^{T_1} r_s ds} (P(T_1, T_2)(L_{T_1} - \kappa))^+ \middle| \mathcal{F}_t \right] \\ &= P(0, T_2) \widehat{\mathbb{E}}[(L_{T_1} - \kappa)^+] + \int_0^t \xi_s^{(1)} dP(s, T_1) + \int_0^t \xi_s^{(2)} dP(s, T_1), \end{aligned}$$

$0 \leq t \leq T_1$.

- e) Show that under the self-financing condition (4.5.23), the discounted portfolio value $\tilde{V}_t = e^{-\int_0^t r_s ds} V_t$ satisfies

$$d\tilde{V}_t = \xi_t^{(1)} d\tilde{P}(t, T_1) + \xi_t^{(2)} d\tilde{P}(t, T_2),$$

where $\tilde{P}(t, T_1) := e^{-\int_0^t r_s ds} P(t, T_1)$ and $\tilde{P}(t, T_2) := e^{-\int_0^t r_s ds} P(t, T_2)$ denote the discounted bond prices.

- f) Show that

$$\widehat{\mathbb{E}}[(L_{T_1} - \kappa)^+ | \mathcal{F}_t] = \widehat{\mathbb{E}}[(L_{T_1} - \kappa)^+] + \int_0^t \frac{\partial C}{\partial x}(L_u, v(u, T_1)) dL_u,$$

and that the deflated portfolio value $\widehat{V}_t = V_t / P(t, T_2)$ satisfies

$$d\widehat{V}_t = \frac{\partial C}{\partial x}(L_t, v(t, T_1)) dL_t = \sigma(t) L_t \frac{\partial C}{\partial x}(L_t, v(t, T_1)) d\hat{B}_t.$$

Hint: use the martingale property and the Itô formula.

- g) Show that

$$dV_t = (P(t, T_1) - P(t, T_2)) \frac{\partial C}{\partial x}(L_t, v(t, T_1)) \sigma(t) dB_t + \widehat{V}_t dP(t, T_2).$$



h) Show that

$$\begin{aligned} d\tilde{V}_t &= \frac{\partial C}{\partial x}(L_t, v(t, T_1))d(\tilde{P}(t, T_1) - \tilde{P}(t, T_2)) \\ &\quad + \left(\hat{V}_t - L_t \frac{\partial C}{\partial x}(L_t, v(t, T_1)) \right) d\tilde{P}(t, T_2), \end{aligned}$$

and deduce the values of the hedging portfolio allocation $(\xi_t^{(1)}, \xi_t^{(2)})_{t \in \mathbb{R}_+}$.

Exercise 4.14 Consider a bond market with tenor structure $\{T_i, \dots, T_j\}$ and $j-i+1$ bonds with maturities T_i, \dots, T_j , whose prices $P(t, T_i), \dots, P(t, T_j)$ at time t are given by

$$\frac{dP(t, T_k)}{P(t, T_k)} = r_t dt + \zeta_k(t) dB_t, \quad k = i, \dots, j,$$

where $(r_t)_{t \in \mathbb{R}_+}$ is a short-term interest rate process and $(B_t)_{t \in \mathbb{R}_+}$ denotes a standard Brownian motion generating a filtration $(\mathcal{F}_t)_{t \in \mathbb{R}_+}$, and $\zeta_i(t), \dots, \zeta_j(t)$ are volatility processes.

The swap rate $S(t, T_i, T_j)$ is defined by

$$S(t, T_i, T_j) = \frac{P(t, T_i) - P(t, T_j)}{P(t, T_i, T_j)},$$

where

$$P(t, T_i, T_j) = \sum_{k=i}^{j-1} (T_{k+1} - T_k) P(t, T_{k+1})$$

is the annuity numéraire. Recall that a swaption on the LIBOR market can be priced at time $t \in [0, T_i]$ as

$$\begin{aligned} \mathbf{E}^* \left[e^{-\int_t^{T_i} r_s ds} \left(\sum_{k=i}^{j-1} (T_{k+1} - T_k) P(T_i, T_{k+1}) (S(T_i, T_k, T_{k+1}) - \kappa) \right)^+ \middle| \mathcal{F}_t \right] \\ = P(t, T_i, T_j) \mathbf{E}_{i,j} [(S(T_i, T_i, T_j) - \kappa)^+ \mid \mathcal{F}_t], \end{aligned} \quad (4.5.24)$$

under the forward swap measure $\widehat{\mathbb{P}}_{i,j}$ defined by

$$\frac{d\widehat{\mathbb{P}}_{i,j}}{d\mathbb{P}^*} = e^{-\int_0^{T_i} r_s ds} \frac{P(T_i, T_i, T_j)}{P(0, T_i, T_j)}, \quad 1 \leq i < j \leq n,$$

under which

$$\widehat{B}_t^{i,j} := B_t - \sum_{k=i}^{j-1} (T_{k+1} - T_k) \frac{P(t, T_{k+1})}{P(t, T_i, T_j)} \zeta_{k+1}(t) dt \quad (4.5.25)$$

is a standard Brownian motion. Recall that the swap rate can be modeled as

$$dS(t, T_i, T_j) = S(t, T_i, T_j) \sigma_{i,j}(t) d\widehat{B}_t^{i,j}, \quad 0 \leq t \leq T_i, \quad (4.5.26)$$

where the swap rate volatilities are given by

$$\begin{aligned} \sigma_{i,j}(t) &= \sum_{l=i}^{j-1} (T_{l+1} - T_l) \frac{P(t, T_{l+1})}{P(t, T_i, T_j)} (\zeta_i(t) - \zeta_{l+1}(t)) \\ &\quad + \frac{P(t, T_j)}{P(t, T_i) - P(t, T_j)} (\zeta_i(t) - \zeta_j(t)) \end{aligned} \quad (4.5.27)$$

$1 \leq i, j \leq n$, cf. e.g. Proposition 10.8 in [Privault, 2012](#). In the sequel we denote $S_t = S(t, T_i, T_j)$ for simplicity of notation.

- a) Solve the equation (4.5.26) on the interval $[t, T_i]$, and compute $S(T_i, T_i, T_j)$ from the initial condition $S(t, T_i, T_j)$.
- b) Assuming that $\sigma_{i,j}(t)$ is a deterministic volatility function of time t for $1 \leq i, j \leq n$, show that the price (4.5.8) of the swaption can be written as

$$P(t, T_i, T_j)C(S_t, v(t, T_i)),$$

where

$$v^2(t, T_i) = \int_t^{T_i} |\sigma_{i,j}(s)|^2 ds,$$

and $C(x, v)$ is a function to be specified using the Black-Scholes formula $B(x, K, \sigma, r, \tau)$, with the relation

$$\mathbf{E}[(x e^{m+X} - K)^+] = \Phi(v + (m + \log(x/K))/v) - K\Phi((m + \log(x/K))/v),$$

where X is a centered Gaussian random variable with variance v^2 .

- c) Consider a portfolio allocation $(\xi_t^{(i)}, \dots, \xi_t^{(j)})_{t \in [0, T_i]}$ made of bonds with maturities T_i, \dots, T_j and value

$$V_t = \sum_{k=i}^j \xi_t^{(k)} P(t, T_k),$$

at time $t \in [0, T_i]$. We assume that the portfolio is self-financing, *i.e.*

$$dV_t = \sum_{k=i}^j \xi_t^{(k)} dP(t, T_k), \quad 0 \leq t \leq T_i, \quad (4.5.28)$$

and that it *hedges* the claim payoff $(S(T_i, T_i, T_j) - \kappa)^+$, so that

$$\begin{aligned} V_t &= \mathbf{E}^* \left[e^{-\int_t^{T_i} r_s ds} \left(\sum_{k=i}^{j-1} (T_{k+1} - T_k) P(T_i, T_{k+1}) (L(T_i, T_k, T_{k+1}) - \kappa) \right)^+ \middle| \mathcal{F}_t \right] \\ &= P(t, T_i, T_j) \mathbf{E}_{i,j} [(S(T_i, T_i, T_j) - \kappa)^+ | \mathcal{F}_t], \end{aligned}$$

$0 \leq t \leq T_i$. Show that

$$\begin{aligned} \mathbf{E}^* \left[e^{-\int_t^{T_i} r_s ds} \left(\sum_{k=i}^{j-1} (T_{k+1} - T_k) P(T_i, T_{k+1}) (L(T_i, T_k, T_{k+1}) - \kappa) \right)^+ \middle| \mathcal{F}_t \right] \\ = P(0, T_i, T_j) \mathbf{E}_{i,j} [(S(T_i, T_i, T_j) - \kappa)^+] + \sum_{k=i}^j \int_0^t \xi_s^{(k)} dP(s, T_i), \end{aligned}$$

$0 \leq t \leq T_i$.

- d) Show that under the self-financing condition (4.5.28), the discounted portfolio value $\tilde{V}_t = e^{-\int_0^t r_s ds} V_t$ satisfies

$$d\tilde{V}_t = \sum_{k=i}^j \xi_t^{(k)} d\tilde{P}(t, T_k),$$

where $\tilde{P}(t, T_k) = e^{-\int_0^t r_s ds} P(t, T_k)$, $k = i, i+1, \dots, j$, denote the discounted bond prices.

- e) Show that

$$\begin{aligned} \mathbf{E}_{i,j} [(S(T_i, T_i, T_j) - \kappa)^+ | \mathcal{F}_t] \\ = \mathbf{E}_{i,j} [(S(T_i, T_i, T_j) - \kappa)^+] + \int_0^t \frac{\partial C}{\partial x} (S_u, v(u, T_i)) dS_u. \end{aligned}$$

Hint: use the martingale property and the Itô formula.

- f) Show that the deflated portfolio value $\hat{V}_t = V_t / P(t, T_i, T_j)$ satisfies

$$d\hat{V}_t = \frac{\partial C}{\partial x} (S_t, v(t, T_i)) dS_t = S_t \frac{\partial C}{\partial x} (S_t, v(t, T_i)) \sigma_t^{i,j} d\hat{B}_t^{i,j}.$$



g) Show that

$$dV_t = (P(t, T_i) - P(t, T_j)) \frac{\partial C}{\partial x}(S_t, v(t, T_i)) \sigma_t^{i,j} dB_t + \hat{V}_t dP(t, T_i, T_j).$$

h) Show that

$$\begin{aligned} dV_t &= S_t \zeta_i(t) \frac{\partial C}{\partial x}(S_t, v(t, T_i)) \sum_{k=i}^{j-1} (T_{k+1} - T_k) P(t, T_{k+1}) dB_t \\ &\quad + (\hat{V}_t - S_t \frac{\partial C}{\partial x}(S_t, v(t, T_i))) \sum_{k=i}^{j-1} (T_{k+1} - T_k) P(t, T_{k+1}) \zeta_{k+1}(t) dB_t \\ &\quad + \frac{\partial C}{\partial x}(S_t, v(t, T_i)) P(t, T_j) (\zeta_i(t) - \zeta_j(t)) dB_t. \end{aligned}$$

i) Show that

$$\begin{aligned} d\tilde{V}_t &= \frac{\partial C}{\partial x}(S_t, v(t, T_i)) d(\tilde{P}(t, T_i) - \tilde{P}(t, T_j)) \\ &\quad + (\hat{V}_t - S_t \frac{\partial C}{\partial x}(S_t, v(t, T_i))) d\tilde{P}(t, T_i, T_j). \end{aligned}$$

j) Show that

$$\frac{\partial C}{\partial x}(x, v(t, T_i)) = \Phi \left(\frac{\log(x/K)}{v(t, T_i)} + \frac{v(t, T_i)}{2} \right).$$

k) Show that we have

$$\begin{aligned} d\tilde{V}_t &= \Phi \left(\frac{\log(S_t/K)}{v(t, T_i)} + \frac{v(t, T_i)}{2} \right) d(\tilde{P}(t, T_i) - \tilde{P}(t, T_j)) \\ &\quad - \kappa \Phi \left(\frac{\log(S_t/K)}{v(t, T_i)} - \frac{v(t, T_i)}{2} \right) d\tilde{P}(t, T_i, T_j). \end{aligned}$$

l) Show that the hedging strategy is given by

$$\begin{aligned} \xi_t^{(i)} &= \Phi \left(\frac{\log(S_t/K)}{v(t, T_i)} + \frac{v(t, T_i)}{2} \right), \\ \xi_t^{(j)} &= -\Phi \left(\frac{\log(S_t/K)}{v(t, T_i)} + \frac{v(t, T_i)}{2} \right) - \kappa(T_j - T_{j-1}) \Phi \left(\frac{\log(S_t/K)}{v(t, T_i)} - \frac{v(t, T_i)}{2} \right), \end{aligned}$$

and

$$\xi_t^{(k)} = -\kappa(T_{k+1} - T_k) \Phi \left(\frac{\log(S_t/K)}{v(t, T_i)} - \frac{v(t, T_i)}{2} \right), \quad i \leq k \leq j-2.$$

5. Reduced-Form Approach to Credit Risk

The reduced-form approach to credit risk modeling focuses on modeling default probabilities as stochastic processes, in contrast to the structural approach in which bankruptcy is modeled from the firm's asset value. The modeling of default risk using failure rate processes and exogeneous random variables results into the use of enlarged filtration that can incorporate information on default events.

5.1	Survival Probabilities	123
5.2	Stochastic Default	125
5.3	Defaultable Bonds	128
	Exercises	131

5.1 Survival Probabilities

Given $t > 0$, let $\mathbb{P}(\tau > t)$ denote the probability that a random system with lifetime τ survives at least t years. Assuming that survival probabilities $\mathbb{P}(\tau > t)$ are strictly positive for all $t > 0$, we can compute the conditional probability for that system to survive up to time T , given that it was still functioning at time $t \in [0, T]$, as

$$\mathbb{P}(\tau > T \mid \tau > t) = \frac{\mathbb{P}(\tau > T \text{ and } \tau > t)}{\mathbb{P}(\tau > t)} = \frac{\mathbb{P}(\tau > T)}{\mathbb{P}(\tau > t)}, \quad 0 \leq t \leq T,$$

with

$$\begin{aligned}\mathbb{P}(\tau \leq T \mid \tau > t) &= 1 - \mathbb{P}(\tau > T \mid \tau > t) \\ &= \frac{\mathbb{P}(\tau > t) - \mathbb{P}(\tau > T)}{\mathbb{P}(\tau > t)}\end{aligned}$$

$$\begin{aligned}
&= \frac{\mathbb{P}(\tau \leq T) - \mathbb{P}(\tau \leq t)}{\mathbb{P}(\tau > t)} \\
&= \frac{\mathbb{P}(t < \tau \leq T)}{\mathbb{P}(\tau > t)}, \quad 0 \leq t \leq T.
\end{aligned} \tag{5.1.1}$$

Such survival probabilities are typically found in life (or mortality) tables:

Age t	$\mathbb{P}(\tau \leq t+1 \tau > t)$
20	0.0894%
30	0.1008%
40	0.2038%
50	0.4458%
60	0.9827%

Table 5.1: Mortality table.

The corresponding conditional survival probability distribution can be computed as follows:

$$\begin{aligned}
\mathbb{P}(\tau \in dx | \tau > t) &= \mathbb{P}(x < \tau \leq x + dx | \tau > t) \\
&= \mathbb{P}(\tau \leq x + dx | \tau > t) - \mathbb{P}(\tau \leq x | \tau > t) \\
&= \frac{\mathbb{P}(\tau \leq x + dx) - \mathbb{P}(\tau \leq x)}{\mathbb{P}(\tau > t)} \\
&= \frac{1}{\mathbb{P}(\tau > t)} d\mathbb{P}(\tau \leq x) \\
&= -\frac{1}{\mathbb{P}(\tau > t)} d\mathbb{P}(\tau > x), \quad x > t.
\end{aligned}$$

Proposition 5.1 The *failure rate* function, defined as

$$\lambda(t) := \frac{\mathbb{P}(\tau \leq t + dt | \tau > t)}{dt},$$

satisfies

$$\mathbb{P}(\tau > t) = \exp\left(-\int_0^t \lambda(u) du\right), \quad t \geq 0. \tag{5.1.2}$$

Proof. By (5.1.1), we have

$$\begin{aligned}
\lambda(t) &:= \frac{\mathbb{P}(\tau \leq t + dt | \tau > t)}{dt} \\
&= \frac{1}{\mathbb{P}(\tau > t)} \frac{\mathbb{P}(t < \tau \leq t + dt)}{dt} \\
&= \frac{1}{\mathbb{P}(\tau > t)} \frac{\mathbb{P}(\tau > t) - \mathbb{P}(\tau > t + dt)}{dt} \\
&= -\frac{d}{dt} \log \mathbb{P}(\tau > t) \\
&= -\frac{1}{\mathbb{P}(\tau > t)} \frac{d}{dt} \mathbb{P}(\tau > t), \quad t > 0,
\end{aligned}$$

and the differential equation

$$\frac{d}{dt} \mathbb{P}(\tau > t) = -\lambda(t) \mathbb{P}(\tau > t),$$



which can be solved as in (5.1.2) under the initial condition $\mathbb{P}(\tau > 0) = 1$. \square

Proposition 5.1 allows us to rewrite the (conditional) survival probability as

$$\mathbb{P}(\tau > T \mid \tau > t) = \frac{\mathbb{P}(\tau > T)}{\mathbb{P}(\tau > t)} = \exp\left(-\int_t^T \lambda(u)du\right), \quad 0 \leq t \leq T,$$

with

$$\mathbb{P}(\tau > t+h \mid \tau > t) = e^{-\lambda(t)h} \simeq 1 - \lambda(t)h, \quad [h \searrow 0],$$

and

$$\mathbb{P}(\tau \leq t+h \mid \tau > t) = 1 - e^{-\lambda(t)h} \simeq \lambda(t)h, \quad [h \searrow 0],$$

as h tends to 0. When the failure rate $\lambda(t) = \lambda > 0$ is a constant function of time, Relation (5.1.2) shows that

$$\mathbb{P}(\tau > T) = e^{-\lambda T}, \quad T \geq 0,$$

i.e. τ has the exponential distribution with parameter λ . Note that given $(\tau_n)_{n \geq 1}$ a sequence of i.i.d. exponentially distributed random variables, letting

$$T_n = \tau_1 + \tau_2 + \cdots + \tau_n, \quad n \geq 1,$$

defines the sequence of jump times of a standard Poisson process with intensity $\lambda > 0$.

5.2 Stochastic Default

When the random time τ is a *stopping time* with respect to $(\mathcal{F}_t)_{t \in \mathbb{R}_+}$ we have

$$\{\tau > t\} \in \mathcal{F}_t, \quad t \geq 0,$$

i.e. the knowledge of whether default or bankruptcy has already occurred at time t is contained in \mathcal{F}_t , $t \in \mathbb{R}_+$, cf. e.g. Section 9.3 of [Privault, 2014](#). As a consequence, we can write

$$\mathbb{P}(\tau > t \mid \mathcal{F}_t) = \mathbf{E}[\mathbb{1}_{\{\tau > t\}} \mid \mathcal{F}_t] = \mathbb{1}_{\{\tau > t\}}, \quad t \in \mathbb{R}_+.$$

In the sequel we will not assume that τ is an \mathcal{F}_t -stopping time, and by analogy with (5.1.2) we will write $\mathbb{P}(\tau > t \mid \mathcal{F}_t)$ as

$$\mathbb{P}(\tau > t \mid \mathcal{F}_t) = \exp\left(-\int_0^t \lambda_u du\right), \quad t \geq 0, \tag{5.2.1}$$

where the failure rate function $(\lambda_t)_{t \in \mathbb{R}_+}$ is modeled as a random process adapted to a filtration $(\mathcal{F}_t)_{t \in \mathbb{R}_+}$.

The process $(\lambda_t)_{t \in \mathbb{R}_+}$ can also be chosen among the classical mean-reverting diffusion processes, including jump-diffusion processes. In [Lando, 1998](#), the process $(\lambda_t)_{t \in \mathbb{R}_+}$ is constructed as $\lambda_t := h(X_t)$, $t \in \mathbb{R}_+$, where h is a nonnegative function and $(X_t)_{t \in \mathbb{R}_+}$ is a stochastic process generating the filtration $(\mathcal{F}_t)_{t \in \mathbb{R}_+}$.

The default time τ is then *defined* as

$$\tau := \inf \left\{ t \in \mathbb{R}_+ : \int_0^t h(X_u) du \geq L \right\},$$

where L is an exponentially distributed random variable with parameter $\mu > 0$ and distribution function $\mathbb{P}(L > x) = e^{-\mu x}$, $x \geq 0$, independent of $(\mathcal{F}_t)_{t \in \mathbb{R}_+}$. In this case, as τ is not an $(\mathcal{F}_t)_{t \in \mathbb{R}_+}$ -stopping time, we have

$$\begin{aligned}\mathbb{P}(\tau > t \mid \mathcal{F}_t) &= \mathbb{P}\left(\int_0^t h(X_u)du < L \mid \mathcal{F}_t\right) \\ &= \exp\left(-\mu \int_0^t h(X_u)du\right) \\ &= \exp\left(-\mu \int_0^t \lambda_u du\right), \quad t \geq 0.\end{aligned}$$

Definition 5.2 Let $(\mathcal{G}_t)_{t \in \mathbb{R}_+}$ be the filtration defined by $\mathcal{G}_\infty := \mathcal{F}_\infty \vee \sigma(\tau)$ and

$$\mathcal{G}_t := \{B \in \mathcal{G}_\infty : \exists A \in \mathcal{F}_t \text{ such that } A \cap \{\tau > t\} = B \cap \{\tau > t\}\}, \quad (5.2.2)$$

with $\mathcal{F}_t \subset \mathcal{G}_t$, $t \in \mathbb{R}_+$.

In other words, \mathcal{G}_t contains insider information on whether default at time τ has occurred or not before time t , and τ is a $(\mathcal{G}_t)_{t \in \mathbb{R}_+}$ -stopping time. Note that this information on τ may not be available to a generic user who has only access to the smaller filtration $(\mathcal{F}_t)_{t \in \mathbb{R}_+}$. The next key Lemma 5.3, see Lando, 1998, Guo, Jarrow, and Menn, 2007, allows us to price a contingent claim given the information in the larger filtration $(\mathcal{G}_t)_{t \in \mathbb{R}_+}$, by only using information in $(\mathcal{F}_t)_{t \in \mathbb{R}_+}$ and factoring in the default rate factor $\exp\left(-\int_t^T \lambda_u du\right)$.

Lemma 5.3 (Guo, Jarrow, and Menn, 2007, Theorem 1) For any \mathcal{F}_T -measurable integrable random variable F , we have

$$\mathbb{E}[F \mathbb{1}_{\{\tau > T\}} \mid \mathcal{G}_t] = \mathbb{1}_{\{\tau > t\}} \mathbb{E}\left[F \exp\left(-\int_t^T \lambda_u du\right) \mid \mathcal{F}_t\right].$$

Proof. By (5.2.1) we have

$$\frac{\mathbb{P}(\tau > T \mid \mathcal{F}_T)}{\mathbb{P}(\tau > t \mid \mathcal{F}_t)} = \frac{e^{-\int_0^T \lambda_u du}}{e^{-\int_0^t \lambda_u du}} = \exp\left(-\int_t^T \lambda_u du\right),$$

hence, since F is \mathcal{F}_T -measurable,

$$\begin{aligned}\mathbb{1}_{\{\tau > t\}} \mathbb{E}\left[F \exp\left(-\int_t^T \lambda_u du\right) \mid \mathcal{F}_t\right] &= \mathbb{1}_{\{\tau > t\}} \mathbb{E}\left[F \frac{\mathbb{P}(\tau > T \mid \mathcal{F}_T)}{\mathbb{P}(\tau > t \mid \mathcal{F}_t)} \mid \mathcal{F}_t\right] \\ &= \frac{\mathbb{1}_{\{\tau > t\}}}{\mathbb{P}(\tau > t \mid \mathcal{F}_t)} \mathbb{E}\left[F \mathbb{E}[\mathbb{1}_{\{\tau > T\}} \mid \mathcal{F}_T] \mid \mathcal{F}_t\right] \\ &= \frac{\mathbb{1}_{\{\tau > t\}}}{\mathbb{P}(\tau > t \mid \mathcal{F}_t)} \mathbb{E}\left[\mathbb{E}[F \mathbb{1}_{\{\tau > T\}} \mid \mathcal{F}_T] \mid \mathcal{F}_t\right] \\ &= \mathbb{1}_{\{\tau > t\}} \frac{\mathbb{E}[F \mathbb{1}_{\{\tau > T\}} \mid \mathcal{F}_t]}{\mathbb{P}(\tau > t \mid \mathcal{F}_t)} \\ &= \mathbb{1}_{\{\tau > t\}} \mathbb{E}[F \mathbb{1}_{\{\tau > T\}} \mid \mathcal{G}_t] \\ &= \mathbb{E}[F \mathbb{1}_{\{\tau > T\}} \mid \mathcal{G}_t], \quad 0 \leq t \leq T.\end{aligned}$$

In the last step of the above argument we used the key relation

$$\mathbb{1}_{\{\tau > t\}} \mathbb{E}\left[F \mathbb{1}_{\{\tau > T\}} \mid \mathcal{G}_t\right] = \frac{\mathbb{1}_{\{\tau > t\}}}{\mathbb{P}(\tau > t \mid \mathcal{F}_t)} \mathbb{E}\left[F \mathbb{1}_{\{\tau > T\}} \mid \mathcal{F}_t\right],$$



cf. Relation (75.2) in § XX-75 page 186 of [Dellacherie, Maisonneuve, and Meyer, 1992](#), Theorem VI-3-14 page 371 of [Protter, 2004](#), and Lemma 3.1 of [Elliott, Jeanblanc, and Yor, 2000](#), under the conditional probability measure $\mathbb{P}_{|\mathcal{F}_t}$, $0 \leq t \leq T$. Indeed, according to (5.2.2), for any $B \in \mathcal{G}_t$ we have, for some event $A \in \mathcal{F}_t$,

$$\begin{aligned} \mathbf{E} [\mathbb{1}_B \mathbb{1}_{\{\tau > t\}} F \mathbb{1}_{\{\tau > T\}}] &= \mathbf{E} [\mathbb{1}_{B \cap \{\tau > t\}} F \mathbb{1}_{\{\tau > T\}}] \\ &= \mathbf{E} [\mathbb{1}_{A \cap \{\tau > t\}} F \mathbb{1}_{\{\tau > T\}}] \\ &= \mathbf{E} [\mathbb{1}_A \mathbb{1}_{\{\tau > t\}} F \mathbb{1}_{\{\tau > T\}}] \\ &= \mathbf{E} \left[\mathbb{1}_A \mathbb{1}_{\{\tau > t\}} \frac{\mathbf{E}[\mathbb{1}_{\{\tau > t\}} | \mathcal{F}_t]}{\mathbb{P}(\tau > t | \mathcal{F}_t)} F \mathbb{1}_{\{\tau > T\}} \right] \\ &= \mathbf{E} \left[\frac{\mathbf{E}[\mathbb{1}_A \mathbb{1}_{\{\tau > t\}} | \mathcal{F}_t]}{\mathbb{P}(\tau > t | \mathcal{F}_t)} F \mathbb{1}_{\{\tau > T\}} \right] \\ &= \mathbf{E} \left[\frac{\mathbf{E}[\mathbb{1}_A \mathbb{1}_{\{\tau > t\}} | \mathcal{F}_t]}{\mathbb{P}(\tau > t | \mathcal{F}_t)} \mathbf{E}[F \mathbb{1}_{\{\tau > T\}} | \mathcal{F}_t] \right] \\ &= \mathbf{E} \left[\frac{\mathbb{1}_A \mathbb{1}_{\{\tau > t\}}}{\mathbb{P}(\tau > t | \mathcal{F}_t)} \mathbf{E}[F \mathbb{1}_{\{\tau > T\}} | \mathcal{F}_t] \right] \\ &= \mathbf{E} \left[\frac{\mathbf{E}[\mathbb{1}_A \mathbb{1}_{\{\tau > t\}} | \mathcal{F}_t]}{\mathbb{P}(\tau > t | \mathcal{F}_t)} \mathbf{E}[F \mathbb{1}_{\{\tau > T\}} | \mathcal{F}_t] \right] \\ &= \mathbf{E} \left[\frac{\mathbb{1}_A \mathbb{1}_{\{\tau > t\}}}{\mathbb{P}(\tau > t | \mathcal{F}_t)} \mathbf{E}[F \mathbb{1}_{\{\tau > T\}} | \mathcal{F}_t] \right] \\ &= \mathbf{E} \left[\frac{\mathbb{1}_B \mathbb{1}_{\{\tau > t\}}}{\mathbb{P}(\tau > t | \mathcal{F}_t)} \mathbf{E}[F \mathbb{1}_{\{\tau > T\}} | \mathcal{F}_t] \right], \end{aligned}$$

hence by a standard characterization of conditional expectations, we have

$$\mathbf{E} [\mathbb{1}_{\{\tau > t\}} F \mathbb{1}_{\{\tau > T\}} | \mathcal{G}_t] = \frac{\mathbb{1}_{\{\tau > t\}}}{\mathbb{P}(\tau > t | \mathcal{F}_t)} \mathbf{E}[F \mathbb{1}_{\{\tau > T\}} | \mathcal{F}_t]$$

□

Taking $F = 1$ in Lemma 5.3 allows one to write the survival probability up to time T , given the information known up to time t , as

$$\begin{aligned} \mathbb{P}(\tau > T | \mathcal{G}_t) &= \mathbf{E} [\mathbb{1}_{\{\tau > T\}} | \mathcal{G}_t] \\ &= \mathbb{1}_{\{\tau > t\}} \mathbf{E} \left[\exp \left(- \int_t^T \lambda_u du \right) \middle| \mathcal{F}_t \right], \quad 0 \leq t \leq T. \end{aligned} \tag{5.2.3}$$

In particular, applying Lemma 5.3 for $t = T$ and $F = 1$ shows that

$$\mathbf{E} [\mathbb{1}_{\{\tau > t\}} | \mathcal{G}_t] = \mathbb{1}_{\{\tau > t\}},$$

which shows that $\{\tau > t\} \in \mathcal{G}_t$ for all $t > 0$, and recovers the fact that τ is a $(\mathcal{G}_t)_{t \in \mathbb{R}_+}$ -stopping time, while in general, τ is not $(\mathcal{F}_t)_{t \in \mathbb{R}_+}$ -stopping time.

The computation of $\mathbb{P}(\tau > T | \mathcal{G}_t)$ according to (5.2.3) is then similar to that of a bond price, by considering the failure rate $\lambda(t)$ as a “virtual” short-term interest rate. In particular the failure rate $\lambda(t, T)$ can be modeled in the HJM framework, cf. e.g. Chapter 11.4 of [Privault, 2014](#), and

$$\mathbb{P}(\tau > T | \mathcal{G}_t) = \mathbf{E} \left[\exp \left(- \int_t^T \lambda(t, u) du \right) \middle| \mathcal{F}_t \right]$$

can then be computed by applying HJM bond pricing techniques.

The computation of expectations given \mathcal{G}_t as in Lemma 5.3 can be useful for pricing under insider trading, in which the insider has access to the augmented filtration \mathcal{G}_t while the ordinary trader has only access to \mathcal{F}_t , therefore generating two different prices $\mathbb{E}^*[F | \mathcal{F}_t]$ and $\mathbb{E}^*[F | \mathcal{G}_t]$ for the same claim payoff F under the same risk-neutral probability measure \mathbb{P}^* . This leads to the issue of computing the dynamics of the underlying asset price by decomposing it using a \mathcal{F}_t -martingale *vs* a \mathcal{G}_t -martingale instead of using different forward measures as in *e.g.* § 12.1 of Privault, 2014. This can be obtained by the technique of enlargement of filtration, see Jeulin, 1980, Jacod, 1985, Yor, 1985, Elliott and Jeanblanc, 1999.

5.3 Defaultable Bonds

Bond pricing models are generally based on the terminal condition $P(T, T) = \$1$ according to which the bond payoff at maturity is always equal to \$1, and default does not occurs. In this chapter we allow for the possibility of default at a random time τ , in which case the terminal payoff of a bond is allowed to vanish at maturity.

The price $P_d(t, T)$ at time t of a default bond with maturity T , (random) default time τ and (possibly random) recovery rate $\xi \in [0, 1]$ is given by

$$\begin{aligned} P_d(t, T) &= \mathbb{E}^* \left[\mathbb{1}_{\{\tau > T\}} \exp \left(- \int_t^T r_u du \right) \middle| \mathcal{G}_t \right] \\ &\quad + \mathbb{E}^* \left[\xi \mathbb{1}_{\{\tau \leq T\}} \exp \left(- \int_t^T r_u du \right) \middle| \mathcal{G}_t \right], \quad 0 \leq t \leq T. \end{aligned}$$

Proposition 5.4 The default bond with maturity T and default time τ can be priced at time $t \in [0, T]$ as

$$\begin{aligned} P_d(t, T) &= \mathbb{1}_{\{\tau > t\}} \mathbb{E}^* \left[\exp \left(- \int_t^T (r_u + \lambda_u) du \right) \middle| \mathcal{F}_t \right] \\ &\quad + \mathbb{E}^* \left[\xi \mathbb{1}_{\{\tau \leq T\}} \exp \left(- \int_t^T r_u du \right) \middle| \mathcal{G}_t \right], \quad 0 \leq t \leq T. \end{aligned}$$

Proof. We take $F = \exp \left(- \int_t^T r_u du \right)$ in Lemma 5.3, which shows that

$$\mathbb{E}^* \left[\mathbb{1}_{\{\tau > t\}} \exp \left(- \int_t^T r_u du \right) \middle| \mathcal{G}_t \right] = \mathbb{1}_{\{\tau > t\}} \mathbb{E}^* \left[\exp \left(- \int_t^T (r_u + \lambda_u) du \right) \middle| \mathcal{F}_t \right],$$

cf. *e.g.* Lando, 1998, Duffie and Singleton, 2003, Guo, Jarrow, and Menn, 2007. \square

In the case of complete default (zero-recovery) we have $\xi = 0$ and

$$P_d(t, T) = \mathbb{1}_{\{\tau > t\}} \mathbb{E}^* \left[\exp \left(- \int_t^T (r_s + \lambda_s) ds \right) \middle| \mathcal{F}_t \right], \quad 0 \leq t \leq T. \quad (5.3.1)$$

From the above expression (5.3.1) we note that the effect of the presence of a default time τ is to decrease the bond price, which can be viewed as an increase of the short rate by the amount λ_u . In



a simple setting where the interest rate $r > 0$ and failure rate $\lambda > 0$ are constant, the default bond price becomes

$$P_d(t, T) = \mathbb{1}_{\{\tau>t\}} e^{-(r+\lambda)(T-t)}, \quad 0 \leq t \leq T.$$

In this case, the failure rate λ can be estimated at time $t \in [0, T]$ from a default bond price $P_d(t, T)$ and a non-default bond price $P(t, T) = e^{-(T-t)r}$ as

$$\lambda = \frac{1}{T-t} \log \frac{P(t, T)}{P_d(t, T)}.$$

Finally, from *e.g.* Proposition 12.1 of [Privault, 2014](#) the bond price (5.3.1) can also be expressed under the forward measure $\hat{\mathbb{P}}$ with maturity T , as

$$\begin{aligned} P_d(t, T) &= \mathbb{1}_{\{\tau>t\}} \mathbb{E}^* \left[\exp \left(- \int_t^T (r_s + \lambda_s) ds \right) \mid \mathcal{F}_t \right] \\ &= \mathbb{1}_{\{\tau>t\}} \mathbb{E}^* \left[\exp \left(- \int_t^T r_s ds \right) \mid \mathcal{F}_t \right] \hat{\mathbb{E}} \left[\exp \left(- \int_t^T \lambda_s ds \right) \mid \mathcal{F}_t \right] \\ &= \mathbb{1}_{\{\tau>t\}} N_t \hat{\mathbb{P}}(\tau > T \mid \mathcal{G}_t), \end{aligned}$$

where $(N_t)_{t \in \mathbb{R}_+}$ is the numéraire process

$$N_t := P(t, T) = \mathbb{E}^* \left[\exp \left(- \int_t^T r_s ds \right) \mid \mathcal{F}_t \right], \quad 0 \leq t \leq T,$$

and by (5.2.3),

$$\hat{\mathbb{P}}(\tau > T \mid \mathcal{G}_t) = \mathbb{1}_{\{\tau>t\}} \hat{\mathbb{E}} \left[\exp \left(- \int_t^T \lambda_s ds \right) \mid \mathcal{F}_t \right]$$

is the survival probability under the forward measure $\hat{\mathbb{P}}$ defined as

$$\frac{d\hat{\mathbb{P}}}{d\mathbb{P}} := \frac{N_T}{N_0} e^{-\int_0^T r_t dt},$$

see [Chen and Huang, 2001](#), [Chen, Cheng, et al., 2008](#),

Estimating the default rates

Recall that the price of a default bond with maturity T , (random) default time τ and (possibly random) recovery rate $\xi \in [0, 1]$ is given by

$$\begin{aligned} P_d(t, T) &= \mathbb{1}_{\{\tau>t\}} \mathbb{E}^* \left[\exp \left(- \int_t^T (r_u + \lambda_u) du \right) \mid \mathcal{F}_t \right] \\ &\quad + \mathbb{E}^* \left[\xi \mathbb{1}_{\{\tau \leq T\}} \exp \left(- \int_t^T r_u du \right) \mid \mathcal{G}_t \right], \quad 0 \leq t \leq T, \end{aligned}$$

where ξ denotes the recovery rate. We consider a simplified deterministic step function model with zero recovery rate and tenor structure

$$\{t = T_0 < T_1 < \dots < T_n = T\},$$

where

$$r(t) = \sum_{l=0}^{n-1} r_l \mathbb{1}_{(T_l, T_{l+1}]}(t) \quad \text{and} \quad \lambda(t) = \sum_{l=0}^{n-1} \lambda_l \mathbb{1}_{(T_l, T_{l+1}]}(t), \quad t \in \mathbb{R}_+. \quad (5.3.2)$$

i) Estimating the default rates from default bond prices.

We have

$$\begin{aligned} P_d(t, T_k) &= \mathbb{1}_{\{\tau>t\}} \exp \left(- \int_t^{T_k} (r(u) + \lambda(u)) du \right) \\ &= \mathbb{1}_{\{\tau>t\}} \exp \left(- \sum_{l=0}^{k-1} (r_l + \lambda_l)(T_{l+1} - T_l) \right), \end{aligned}$$

$k = 1, 2, \dots, n$, from which we can infer

$$\lambda_k = -r_k + \frac{1}{T_{k+1} - T_k} \log \frac{P_d(t, T_k)}{P_d(t, T_{k+1})} > 0, \quad k = 0, 1, \dots, n-1.$$

ii) Estimating (implied) default probabilities $\mathbb{P}^*(\tau < T | \mathcal{G}_t)$ from default rates.

Based on the expression

$$\begin{aligned} \mathbb{P}^*(\tau > T | \mathcal{G}_t) &= \mathbb{E}^* [\mathbb{1}_{\{\tau>T\}} | \mathcal{G}_t] \\ &= \mathbb{1}_{\{\tau>t\}} \mathbb{E}^* \left[\exp \left(- \int_t^T \lambda_u du \right) \mid \mathcal{F}_t \right], \quad 0 \leq t \leq T, \end{aligned} \tag{5.3.3}$$

of the survival probability up to time T , and given the information known up to time t , in terms of the hazard rate process $(\lambda_u)_{u \in \mathbb{R}_+}$ adapted to a filtration $(\mathcal{F}_t)_{t \in \mathbb{R}_+}$, we find

$$\begin{aligned} \mathbb{P}(\tau > T | \mathcal{G}_{T_k}) &= \mathbb{1}_{\{\tau>T_k\}} \exp \left(- \int_{T_k}^T \lambda_u du \right) \\ &= \mathbb{1}_{\{\tau>t\}} \exp \left(- \sum_{l=k}^{n-1} \lambda_l (T_{l+1} - T_l) \right), \quad k = 0, 1, \dots, n-1, \end{aligned}$$

where

$$\mathcal{G}_t = \mathcal{F}_t \vee \sigma(\{\tau \leq u\} : 0 \leq u \leq t), \quad t \in \mathbb{R}_+,$$

i.e. \mathcal{G}_t contains the additional information on whether default at time τ has occurred or not before time t .

In Table 5.2, bond ratings are determined according to hazard (or failure) rate thresholds.

Bond Credit Ratings	Moody's		S & P	
	Municipal	Corporate	Municipal	Corporate
Aaa/AAAs	0.00	0.52	0.00	0.60
Aa/AA	0.06	0.52	0.00	1.50
A/A	0.03	1.29	0.23	2.91
Baa/BBB	0.13	4.64	0.32	10.29
Ba/BB	2.65	19.12	1.74	29.93
B/B	11.86	43.34	8.48	53.72
Caa-C/CCC-C	16.58	69.18	44.81	69.19
Investment Grade	0.07	2.09	0.20	4.14
Non-Invest. Grade	4.29	31.37	7.37	42.35
All	0.10	9.70	0.29	12.98

Table 5.2: Cumulative historic default rates (in percentage).*

*Source: Moody's, S&P.

Exercises

Exercise 5.1 Consider a standard zero-coupon bond with constant yield $r > 0$ and a defaultable (risky) bond with constant yield r_d and default probability $\alpha \in (0, 1)$. Find a relation between r, r_d, α and the bond maturity T .

Exercise 5.2 A standard zero-coupon bond with constant yield $r > 0$ and maturity T is priced $P(t, T) = e^{-(T-t)r}$ at time $t \in [0, T]$. Assume that the company can get bankrupt at a random time $t + \tau$, and default on its final \$1 payment if $\tau < T - t$.

- a) Explain why the defaultable bond price $P_d(t, T)$ can be expressed as

$$P_d(t, T) = e^{-(T-t)r} \mathbf{E}^* [\mathbb{1}_{\{\tau > T-t\}}]. \quad (5.3.4)$$

- b) Assuming that the default time τ is exponentially distributed with parameter $\lambda > 0$, compute the default bond price $P_d(t, T)$ using (5.3.4).
- c) Find a formula that can estimate the parameter λ from the risk-free rate r and the market data $P_M(t, T)$ of the defaultable bond price at time $t \in [0, T]$.

Exercise 5.3 Consider a (random) default time τ with cumulative distribution function

$$\mathbb{P}(\tau > t \mid \mathcal{F}_t) = \exp\left(-\int_0^t \lambda_u du\right), \quad t \in \mathbb{R}_+,$$

where λ_t is a (random) default rate process which is adapted to the filtration $(\mathcal{F}_t)_{t \in \mathbb{R}_+}$. Recall that the probability of survival up to time T , given the information known up to time t , is given by

$$\mathbb{P}(\tau > T \mid \mathcal{G}_t) = \mathbb{1}_{\{\tau > t\}} \mathbf{E}^* \left[\exp\left(-\int_t^T \lambda_u du\right) \mid \mathcal{F}_t \right],$$

where $\mathcal{G}_t = \mathcal{F}_t \vee \sigma(\{\tau < u\} : 0 \leq u \leq t), t \in \mathbb{R}_+$, is the filtration defined by adding the default time information to the history $(\mathcal{F}_t)_{t \in \mathbb{R}_+}$. In this framework, the price $P(t, T)$ of defaultable bond with maturity T , short-term interest rate r_t and (random) default time τ is given by

$$\begin{aligned} P(t, T) &= \mathbf{E}^* \left[\mathbb{1}_{\{\tau > T\}} \exp\left(-\int_t^T r_u du\right) \mid \mathcal{G}_t \right] \\ &= \mathbb{1}_{\{\tau > t\}} \mathbf{E}^* \left[\exp\left(-\int_t^T (r_u + \lambda_u) du\right) \mid \mathcal{F}_t \right]. \end{aligned} \quad (5.3.5)$$

In the sequel we assume that the processes $(r_t)_{t \in \mathbb{R}_+}$ and $(\lambda_t)_{t \in \mathbb{R}_+}$ are modeled according to the Vasicek processes

$$\begin{cases} dr_t = -ar_t dt + \sigma dB_t^{(1)}, \\ d\lambda_t = -b\lambda_t dt + \eta dB_t^{(2)}, \end{cases}$$

where $(B_t^{(1)})_{t \in \mathbb{R}_+}$ and $(B_t^{(2)})_{t \in \mathbb{R}_+}$ are standard $(\mathcal{F}_t)_{t \in \mathbb{R}_+}$ -Brownian motions with correlation $\rho \in [-1, 1]$, and $dB_t^{(1)} \cdot dB_t^{(2)} = \rho dt$.

- a) Give a justification for the fact that

$$\mathbf{E}^* \left[\exp\left(-\int_t^T (r_u + \lambda_u) du\right) \mid \mathcal{F}_t \right]$$

can be written as a function $F(t, r_t, \lambda_t)$ of t, r_t and $\lambda_t, t \in [0, T]$.

b) Show that

$$t \mapsto \exp\left(-\int_0^t (r_s + \lambda_s) ds\right) \mathbb{E}^*\left[\exp\left(-\int_t^T (r_u + \lambda_u) du\right) \mid \mathcal{F}_t\right]$$

is an \mathcal{F}_t -martingale under \mathbb{P} .

- c) Use the Itô formula with two variables to derive a PDE on \mathbb{R}^2 for the function $F(t, x, y)$.
d) Taking $r_0 := 0$, show that we have

$$\int_t^T r_s ds = C(a, t, T) r_t + \sigma \int_t^T C(a, s, T) dB_s^{(1)},$$

and

$$\int_t^T \lambda_s ds = C(b, t, T) \lambda_t + \eta \int_t^T C(b, s, T) dB_s^{(2)},$$

where

$$C(a, t, T) = -\frac{1}{a}(\mathbb{e}^{-(T-t)a} - 1).$$

- e) Show that the random variable

$$\int_t^T r_s ds + \int_t^T \lambda_s ds$$

is has a Gaussian distribution, and compute its conditional mean

$$\mathbb{E}^*\left[\int_t^T r_s ds + \int_t^T \lambda_s ds \mid \mathcal{F}_t\right]$$

and variance

$$\text{Var}\left[\int_t^T r_s ds + \int_t^T \lambda_s ds \mid \mathcal{F}_t\right],$$

conditionally to \mathcal{F}_t .

- f) Compute $P(t, T)$ from its expression (5.3.5) as a conditional expectation.
g) Show that the solution $F(t, x, y)$ to the 2-dimensional PDE of Question (c)) is

$$\begin{aligned} F(t, x, y) &= \exp(-C(a, t, T)x - C(b, t, T)y) \\ &\quad \times \exp\left(\frac{\sigma^2}{2} \int_t^T C^2(a, s, T) ds + \frac{\eta^2}{2} \int_t^T C^2(b, s, T) ds\right) \\ &\quad \times \exp\left(\rho \sigma \eta \int_t^T C(a, s, T) C(b, s, T) ds\right). \end{aligned}$$

- h) Show that the defaultable bond price $P(t, T)$ can also be written as

$$P(t, T) = \mathbb{e}^{U(t, T)} \mathbb{P}(\tau > T \mid \mathcal{G}_t) \mathbb{E}^*\left[\exp\left(-\int_t^T r_s ds\right) \mid \mathcal{F}_t\right],$$

where

$$U(t, T) = \rho \frac{\sigma \eta}{ab} (T - t - C(a, t, T) - C(b, t, T) + C(a + b, t, T)).$$

- i) By partial differentiation of $\log P(t, T)$ with respect to T , compute the corresponding instantaneous short rate

$$f(t, T) = -\frac{\partial}{\partial T} \log P(t, T).$$

- j) Show that $\mathbb{P}(\tau > T \mid \mathcal{G}_t)$ can be written using an HJM type default rate as

$$\mathbb{P}(\tau > T \mid \mathcal{G}_t) = \mathbb{1}_{\{\tau > t\}} \exp\left(-\int_t^T f_2(t, u) du\right),$$

where

$$f_2(t, u) = \lambda_t \mathbb{e}^{-(u-t)b} - \frac{\eta^2}{2} C^2(b, t, u).$$



- k) Show how the result of Question (h)) can be simplified when $(B_t^{(1)})_{t \in \mathbb{R}_+}$ and $(B_t^{(2)})_{t \in \mathbb{R}_+}$ are independent.

Exercise Solutions

Chapter 1

Exercise 1.1

a) We have

$$\begin{aligned}
 dr_t &= r_0 d e^{-bt} + \frac{a}{b} d(1 - e^{-bt}) + \sigma d \left(e^{-bt} \int_0^t e^{bs} dB_s \right) \\
 &= -br_0 e^{-bt} dt + a e^{-bt} dt + \sigma e^{-bt} d \int_0^t e^{bs} dB_s + \sigma \int_0^t e^{bs} dB_s d e^{-bt} \\
 &= -br_0 e^{-bt} dt + a e^{-bt} dt + \sigma e^{-bt} e^{bt} dB_t - \sigma b \int_0^t e^{bs} dB_s e^{-bt} dt \\
 &= -br_0 e^{-bt} dt + a e^{-bt} dt + \sigma dB_t - \sigma b \int_0^t e^{bs} dB_s e^{-bt} dt \\
 &= -br_0 e^{-bt} dt + a e^{-bt} dt + \sigma dB_t - b \left(r_t - r_0 e^{-bt} - \frac{a}{b} (1 - e^{-bt}) \right) dt \\
 &= (a - br_t) dt + \sigma dB_t,
 \end{aligned}$$

which shows that r_t solves (1.4.29).

b) We note that

$$\begin{aligned}
 r_t &= r_0 e^{-bt} + \frac{a}{b} (1 - e^{-bt}) + \sigma \int_0^t e^{-(t-u)b} dB_u \\
 &= r_0 e^{-bs} e^{-(t-s)b} + \frac{a}{b} e^{-(t-s)b} (1 - e^{-bs}) \\
 &\quad + \frac{a}{b} (1 - e^{(t-s)b}) + \sigma e^{-(t-s)b} \int_0^s e^{-(s-u)b} dB_u + \sigma \int_s^t e^{-(t-u)b} dB_u \\
 &= r_s e^{-(t-s)b} + \frac{a}{b} (1 - e^{(t-s)b}) + \sigma \int_s^t e^{-(t-u)b} dB_u, \quad 0 \leq s \leq t.
 \end{aligned}$$

Hence, assuming that r_s has the $\mathcal{N}(a/b, \sigma^2/(2b))$ distribution, the distribution of r_t is Gaussian with mean

$$\begin{aligned}
 \mathbf{E}[r_t] &= e^{-(t-s)b} \mathbf{E}[r_s] + \frac{a}{b} (1 - e^{(t-s)b}) \\
 &= \frac{a}{b} e^{-(t-s)b} + \frac{a}{b} (1 - e^{(t-s)b}) \\
 &= \frac{a}{b},
 \end{aligned}$$

and variance

$$\begin{aligned}
 \text{Var}[r_t] &= \text{Var} \left[r_s e^{-(t-s)b} + \frac{a}{b} (1 - e^{(t-s)b}) + \sigma \int_s^t e^{-(t-u)b} dB_u \right] \\
 &= \text{Var} \left[r_s e^{-(t-s)b} + \sigma \int_s^t e^{-(t-u)b} dB_u \right] \\
 &= \text{Var} [r_s e^{-(t-s)b}] + \text{Var} \left[\sigma \int_s^t e^{-(t-u)b} dB_u \right] \\
 &= e^{-2(t-s)b} \text{Var}[r_s] + \sigma^2 \text{Var} \left[\int_s^t e^{-(t-u)b} dB_u \right] \\
 &= e^{-2(t-s)b} \frac{\sigma^2}{2b} + \sigma^2 \int_s^t e^{-2(t-u)b} du \\
 &= e^{-2(t-s)b} \frac{\sigma^2}{2b} + \sigma^2 \int_0^{t-s} e^{-2bu} du \\
 &= \frac{\sigma^2}{2b}, \quad t \geq 0.
 \end{aligned}$$

Exercise 1.2

- i) The zero-coupon bond price $P(t, T)$ in the Vasicek model is given by

$$\log P(t, T) = A(T-t) + r_t C(T-t), \quad 0 \leq t \leq T,$$

where

$$C(T-t) := -\frac{1}{b}(1 - e^{-(T-t)b}),$$

and

$$\begin{aligned}
 A(T-t) &:= \frac{4ab - 3\sigma^2}{4b^3} + \frac{\sigma^2 - 2ab}{2b^2}(T-t) \\
 &\quad + \frac{\sigma^2 - ab}{b^3} e^{-(T-t)b} - \frac{\sigma^2}{4b^3} e^{-2(T-t)b} \quad 0 \leq t \leq T.
 \end{aligned} \tag{A.1}$$

Since $\lim_{T \rightarrow \infty} C(T-t)/(T-t) = 0$ and

$$\lim_{T \rightarrow \infty} A(T-t)/(T-t) = (\sigma^2 - 2ab)/(2b^2),$$

we find

$$r_\infty = -\lim_{T \rightarrow \infty} \frac{\log P(t, T)}{T-t} = -\frac{\sigma^2 - 2ab}{2b^2} = \frac{a}{b} - \frac{\sigma^2}{2b^2} = \lim_{t \rightarrow \infty} (\mathbb{E}[r_t] - \text{Var}[r_t]).$$

- ii) We have

$$\begin{aligned}
 \log \frac{P(t, T)}{P(0, T)} &= \log P(t, T) - \log P(0, T) \\
 &= A(T-t) - A(0) + r_t (C(T-t) - C(0)) \\
 &= -t \frac{\sigma^2 - 2ab}{2b^2} + \frac{\sigma^2 - ab}{b^3} e^{-bt} (e^{bt} - 1) - \frac{\sigma^2}{4b^3} e^{-2bt} (e^{2bt} - 1),
 \end{aligned}$$

hence

$$\lim_{T \rightarrow \infty} \log \frac{P(t, T)}{P(0, T)} = -t \frac{\sigma^2 - 2ab}{2b^2} = \left(\frac{a}{b} - \frac{\sigma^2}{2b^2} \right) t = r_\infty t,$$

and*

$$\lim_{T \rightarrow \infty} \log \frac{P(t, T)}{P(0, T)} = e^{t(a/b - \sigma^2/(2b^2))} = e^{r_\infty t},$$

*The log function is continuous on $(0, \infty)$.



which shows that the yield of the long-bond return is the asymptotic bond yield r_∞ .

Remark: By (1.4.15) of the course notes, the Vasicek bond price $P(t, T)$ can be rewritten in terms of the asymptotic bond yield r_∞ as

$$P(t, T) = e^{-(T-t)r_\infty + (r_t - r_\infty)C(T-t) - \sigma^2 C^2(T-t)/(4b)}, \quad 0 \leq t \leq T,$$

see e.g. Relation (3.12) in [Brody, Hughston, and Meier, 2018](#).

Exercise 1.3 An estimator of σ can be obtained from the orthogonality relation

$$\begin{aligned} \sum_{l=0}^{n-1} ((\tilde{r}_{t_{l+1}} - a\Delta t - (1 - b\Delta t)\tilde{r}_{t_l})^2 - \sigma^2 \Delta t (\tilde{r}_{t_l})^{2\gamma}) &= \sigma^2 \sum_{l=0}^{n-1} (\tilde{r}_{t_l})^{2\gamma} ((Z_l)^2 - \Delta t) \\ &\simeq 0, \end{aligned}$$

which is due to the independence of t_{t_l} and Z_l , $l = 0, \dots, n-1$, and yields

$$\hat{\sigma}^2 = \frac{\sum_{l=0}^{n-1} (\tilde{r}_{t_{l+1}} - a\Delta t - (1 - b\Delta t)\tilde{r}_{t_l})^2}{\Delta t \sum_{l=0}^{n-1} (\tilde{r}_{t_l})^{2\gamma}}.$$

Regarding the estimation of γ , we can combine the above relation with the second orthogonality relation

$$\begin{aligned} \sum_{l=0}^{n-1} ((\tilde{r}_{t_{l+1}} - a\Delta t - (1 - b\Delta t)\tilde{r}_{t_l})^2 - \sigma^2 \Delta t (\tilde{r}_{t_l})^{2\gamma}) \tilde{r}_{t_l} \\ = \sigma^2 \sum_{l=0}^{n-1} (\tilde{r}_{t_l})^{2\gamma+1} ((Z_l)^2 - \Delta t) \\ \simeq 0, \end{aligned}$$

cf. § 2.2 of [Faff and Gray, 2006](#). One may also attempt to minimize the residual

$$\sum_{l=0}^{n-1} ((\tilde{r}_{t_{l+1}} - a\Delta t - (1 - b\Delta t)\tilde{r}_{t_l})^2 - \sigma^2 \Delta t (\tilde{r}_{t_l})^{2\gamma})^2$$

by equating the following derivatives to zero, as

$$\begin{aligned} \frac{\partial}{\partial \sigma} \sum_{l=0}^{n-1} ((\tilde{r}_{t_{l+1}} - a\Delta t - (1 - b\Delta t)\tilde{r}_{t_l})^2 - \sigma^2 \Delta t (\tilde{r}_{t_l})^{2\gamma})^2 \\ = -4\sigma \sum_{l=0}^{n-1} (\tilde{r}_{t_l})^{2\gamma} ((\tilde{r}_{t_{l+1}} - a\Delta t - (1 - b\Delta t)\tilde{r}_{t_l})^2 - \sigma^2 \Delta t (\tilde{r}_{t_l})^{2\gamma}) \\ = 0, \end{aligned}$$

hence

$$\sum_{l=0}^{n-1} (\tilde{r}_{t_l})^{2\gamma} (\tilde{r}_{t_{l+1}} - a\Delta t - (1 - b\Delta t)\tilde{r}_{t_l})^2 - \sigma^2 \Delta t \sum_{l=0}^{n-1} (\tilde{r}_{t_l})^{4\gamma} = 0,$$

which yields

$$\hat{\sigma}^2 = \frac{\sum_{l=0}^{n-1} (\tilde{r}_{t_l})^{2\gamma} (\tilde{r}_{t_{l+1}} - a\Delta t - (1 - b\Delta t)\tilde{r}_{t_l})^2}{\Delta t \sum_{l=0}^{n-1} (\tilde{r}_{t_l})^{4\gamma}}. \quad (\text{A.2})$$

We also have

$$\begin{aligned} & \frac{\partial}{\partial \gamma} \sum_{l=0}^{n-1} \left((\tilde{r}_{t_{l+1}} - a\Delta t - (1 - b\Delta t)\tilde{r}_{t_l})^2 - \sigma^2 \Delta t (\tilde{r}_{t_l})^{2\gamma} \right)^2 \\ &= -4\sigma^2 \Delta t \sum_{l=0}^{n-1} (\tilde{r}_{t_l})^{2\gamma} ((\tilde{r}_{t_{l+1}} - a\Delta t - (1 - b\Delta t)\tilde{r}_{t_l})^2 - \sigma^2 \Delta t (\tilde{r}_{t_l})^{2\gamma}) \log \tilde{r}_{t_l} \\ &= 0, \end{aligned}$$

which yields

$$\hat{\sigma}^2 = \frac{\sum_{l=0}^{n-1} (\tilde{r}_{t_l})^{2\gamma} (\tilde{r}_{t_{l+1}} - a\Delta t - (1 - b\Delta t)\tilde{r}_{t_l})^2 \log \tilde{r}_{t_l}}{\Delta t \sum_{l=0}^{n-1} (\tilde{r}_{t_l})^{4\gamma} \log \tilde{r}_{t_l}}, \quad (\text{A.3})$$

and shows that γ can be estimated by matching Relations (A.2) and (A.3), i.e.

$$\begin{aligned} & \frac{\sum_{l=0}^{n-1} (\tilde{r}_{t_l})^{2\gamma} (\tilde{r}_{t_{l+1}} - a\Delta t - (1 - b\Delta t)\tilde{r}_{t_l})^2}{\sum_{l=0}^{n-1} (\tilde{r}_{t_l})^{4\gamma}} \\ &= \frac{\sum_{l=0}^{n-1} (\tilde{r}_{t_l})^{2\gamma} (\tilde{r}_{t_{l+1}} - a\Delta t - (1 - b\Delta t)\tilde{r}_{t_l})^2 \log \tilde{r}_{t_l}}{\sum_{l=0}^{n-1} (\tilde{r}_{t_l})^{4\gamma} \log \tilde{r}_{t_l}}. \end{aligned}$$

Remarks.

- i) Estimators of a and b can be obtained by minimizing the residual

$$\sum_{l=0}^{n-1} (\tilde{r}_{t_{l+1}} - a\Delta t - (1 - b\Delta t)\tilde{r}_{t_l})^2$$

as in the [Vašíček, 1977](#) model, i.e. from the equations

$$\sum_{l=0}^{n-1} (\tilde{r}_{t_{l+1}} - a\Delta t - (1 - b\Delta t)\tilde{r}_{t_l}) = 0$$

and

$$\sum_{l=0}^{n-1} (\tilde{r}_{t_{l+1}} - a\Delta t - (1 - b\Delta t)\tilde{r}_{t_l}) \tilde{r}_{t_l} = 0.$$

- ii) Instead of using the (generalised) method of moments, parameter estimation for stochastic differential equations can be achieved by maximum likelihood estimation, see e.g. [Lindström, 2007](#) and references therein.

Exercise 1.4

- a) We have $r_t = r_0 + at + \sigma B_t$, and

$$F(t, r_t) = F(t, r_0 + at + \sigma B_t),$$



hence by Proposition 1.2 the PDE satisfied by $F(t,x)$ is

$$-xF(t,x) + \frac{\partial F}{\partial t}(t,x) + a\frac{\partial F}{\partial x}(t,x) + \frac{1}{2}\sigma^2\frac{\partial^2 F}{\partial x^2}(t,x) = 0, \quad (\text{A.4})$$

with terminal condition $F(T,x) = 1$.

b) We have $r_t = r_0 + at + \sigma B_t$ and

$$\begin{aligned} F(t,r_t) &= \mathbf{E}^* \left[\exp \left(- \int_t^T r_s ds \right) \middle| \mathcal{F}_t \right] \\ &= \mathbf{E}^* \left[\exp \left(-r_0(T-t) - a \int_t^T s ds - \sigma \int_t^T B_s ds \right) \middle| \mathcal{F}_t \right] \\ &= \mathbf{E}^* \left[e^{-r_0(T-t)-a(T^2-t^2)/2} \exp \left(-(T-t)B_t - \sigma \int_t^T (T-s) dB_s \right) \middle| \mathcal{F}_t \right] \\ &= e^{-r_0(T-t)-a(T^2-t^2)/2-(T-t)\sigma B_t} \mathbf{E}^* \left[\exp \left(-\sigma \int_t^T (T-s) dB_s \right) \middle| \mathcal{F}_t \right] \\ &= e^{-r_0(T-t)-a(T-t)(T+t)/2-(T-t)\sigma B_t} \mathbf{E}^* \left[\exp \left(-\sigma \int_t^T (T-s) dB_s \right) \right] \\ &= \exp \left(-(T-t)r_t - a(T-t)^2/2 + \frac{\sigma^2}{2} \int_t^T (T-s)^2 ds \right) \\ &= \exp(-(T-t)r_t - a(T-t)^2/2 + (T-t)^3 \sigma^2/6), \end{aligned}$$

hence $F(t,x) = \exp(-(T-t)x - a(T-t)^2/2 + (T-t)^3 \sigma^2/6)$.

Note that the PDE (A.4) can also be solved by looking for a solution of the form $F(t,x) = e^{A(T-t)+xC(T-t)}$, in which case one would find $A(s) = -as^2/2 + \sigma^2 s^3/6$ and $C(s) = -s$.

c) We check that the function $F(t,x)$ of Question (b)) satisfies the PDE (A.4) of Question (a)), since $F(T,x) = 1$ and

$$\begin{aligned} -xF(t,x) + \left(x + a(T-t) - \frac{\sigma^2}{2}(T-t)^2 \right) F(t,x) - a(T-t)F(t,x) \\ + \frac{\sigma^2}{2}(T-t)^2 F(t,x) = 0. \end{aligned}$$

Exercise 1.5 We check from (1.4.34) and the differentiation rule $d \int_0^t f(u) du = f(t)dt$ that

$$\begin{aligned} dr_t &= \alpha\beta d \left(S_t \int_0^t \frac{1}{S_u} du \right) + r_0 dt S_t \\ &= \alpha\beta S_t d \int_0^t \frac{1}{S_u} du + \alpha\beta \int_0^t \frac{1}{S_u} du dS_t + r_0 dt S_t \\ &= \alpha\beta \frac{S_t}{S_t} dt + \alpha\beta \int_0^t \frac{S_t}{S_u} du \frac{dS_t}{S_t} + r_0 dt S_t \\ &= \alpha\beta dt + (r_t - r_0 S_t) \frac{dS_t}{S_t} + r_0 dt S_t \\ &= \alpha\beta dt + r_t \frac{dS_t}{S_t} \\ &= \alpha\beta dt + r_t(-\beta dt + \sigma dB_t) \\ &= \beta(\alpha - r_t)dt + \sigma dB_t, \quad t \geq 0. \end{aligned}$$

Exercise 1.6

a) Applying the Itô formula

$$df(r_t) = f'(r_t)dr_t + \frac{1}{2}f''(r_t)(dr_t)^2$$

to the function $f(x) = x^{2-\gamma}$ with

$$f'(x) = (2-\gamma)x^{1-\gamma} \text{ and } f''(x) = (2-\gamma)(1-\gamma)x^{-\gamma},$$

we have

$$\begin{aligned} dR_t &= dr_t^{2-\gamma} \\ &= df(r_t) \\ &= f'(r_t)dr_t + \frac{1}{2}f''(r_t)(dr_t)^2 \\ &= f'(r_t)((\beta r_t^{\gamma-1} + \alpha r_t)dt + \sigma r_t^{\gamma/2}dB_t)dr_t + \\ &\quad \frac{1}{2}f''(r_t)((\beta r_t^{\gamma-1} + \alpha r_t)dt + \sigma r_t^{\gamma/2}dB_t)^2 \\ &= f'(r_t)((\beta r_t^{\gamma-1} + \alpha r_t)dt + \sigma r_t^{\gamma/2}dB_t)dr_t + \frac{\sigma^2}{2}f''(r_t)r_t^\gamma dt \\ &= (2-\gamma)r_t^{1-\gamma}((\beta r_t^{\gamma-1} + \alpha r_t)dt + \sigma r_t^{\gamma/2}dB_t) + \frac{\sigma^2}{2}(2-\gamma)(1-\gamma)r_t^\gamma r_t^{-\gamma}dt \\ &= (2-\gamma)(\beta + \alpha r_t^{2-\gamma})dt + \frac{\sigma^2}{2}(2-\gamma)(1-\gamma)dt + \sigma(2-\gamma)r_t^{1-\gamma/2}dB_t \\ &= (2-\gamma)\left(\beta + \frac{\sigma^2}{2}(1-\gamma) + \alpha R_t\right)dt + (2-\gamma)\sigma\sqrt{R_t}dB_t. \end{aligned}$$

We conclude that the process $R_t = r_t^{2-\gamma}$ follows the CIR equation

$$dR_t = b(a - R_t)dt + \eta\sqrt{R_t}dB_t$$

with initial condition $R_0 = r_0^{2-\gamma}$ and coefficients

$$b = (2-\gamma)\alpha, \quad a = \frac{1}{\alpha}\left(\beta + (1-\gamma)\frac{\sigma^2}{2}\right), \quad \text{and} \quad \eta = (2-\gamma)\sigma.$$

b) By Itô's formula and the relation $P(t, T) = F(t, r_t)$, $t \in [0, T]$, we have

$$\begin{aligned} d\left(e^{-\int_0^t r_s ds} P(t, T)\right) &= -r_t e^{-\int_0^t r_s ds} P(t, T)dt + e^{-\int_0^t r_s ds} dP(t, T) \\ &= -r_t e^{-\int_0^t r_s ds} F(t, r_t)dt + e^{-\int_0^t r_s ds} dF(t, r_t) \\ &= -r_t e^{-\int_0^t r_s ds} F(t, r_t)dt + e^{-\int_0^t r_s ds} \frac{\partial F}{\partial t}(t, r_t)dt + e^{-\int_0^t r_s ds} \frac{\partial F}{\partial x}(t, r_t)dr_t \\ &\quad + e^{-\int_0^t r_s ds} \frac{1}{2} \frac{\partial^2 F}{\partial x^2}(t, r_t)(dr_t)^2 \\ &= -r_t e^{-\int_0^t r_s ds} F(t, r_t)dt + e^{-\int_0^t r_s ds} \frac{\partial F}{\partial x}(t, r_t)((\beta r_t^{-(1-\gamma)} + \alpha r_t)dt + \sigma r_t^{\gamma/2}dB_t) \\ &\quad + e^{-\int_0^t r_s ds} \left(\frac{\partial F}{\partial t}(t, r_t) + \frac{1}{2}\sigma^2 r_t^\gamma \frac{\partial^2 F}{\partial x^2}(t, r_t)\right)dt \\ &= e^{-\int_0^t r_s ds} \sigma r_t^{\gamma/2} \frac{\partial F}{\partial x}(t, r_t)dB_t \\ &\quad + e^{-\int_0^t r_s ds} \left(-r_t F(t, r_t) + (\beta r_t^{-(1-\gamma)} + \alpha r_t) \frac{\partial F}{\partial x}(t, r_t)\right) \end{aligned}$$



$$+ \frac{1}{2} \sigma^2 r_t^\gamma \frac{\partial^2 F}{\partial x^2}(t, r_t) + \frac{\partial F}{\partial t}(t, r_t) \Big) dt. \quad (\text{A.5})$$

Given that $t \mapsto e^{-\int_0^t r_s ds} P(t, T)$ is a martingale, the above expression (A.5) should only contain terms in dB_t and all terms in dt should vanish inside (A.5). This leads to the identities

$$\left\{ \begin{array}{l} r_t F(t, r_t) = (\beta r_t^{-(1-\gamma)} + \alpha r_t) \frac{\partial F}{\partial x}(t, r_t) + \frac{1}{2} \sigma^2 r_t^\gamma \frac{\partial^2 F}{\partial x^2}(t, r_t) + \frac{\partial F}{\partial t}(t, r_t) \\ d \left(e^{-\int_0^t r_s ds} P(t, T) \right) = \sigma e^{-\int_0^t r_s ds} r_t^{\gamma/2} \frac{\partial F}{\partial x}(t, r_t) dB_t, \end{array} \right.$$

and to the PDE

$$xF(t, x) = \frac{\partial F}{\partial t}(t, x) + (\beta x^{-(1-\gamma)} + \alpha x) \frac{\partial F}{\partial x}(t, x) + \frac{\sigma^2}{2} x^\gamma \frac{\partial^2 F}{\partial x^2}(t, x).$$

Exercise 1.7

a) The discounted bond prices process $e^{-\int_0^t r_s ds} F(t, r_t)$ is a martingale, and we have

$$\begin{aligned} d \left(e^{-\int_0^t r_s ds} F(t, r_t) \right) &= e^{-\int_0^t r_s ds} \left(-r_t F(t, r_t) dt + \frac{\partial F}{\partial t}(t, r_t) dt + \frac{\partial F}{\partial x}(t, r_t) dr_t + \frac{\sigma^2}{2} \frac{\partial^2 F}{\partial x^2}(t, r_t) (dr_t)^2 \right) \\ &= e^{-\int_0^t r_s ds} \left(-r_t F(t, r_t) dt + \frac{\partial F}{\partial t}(t, r_t) dt + \frac{\partial F}{\partial x}(t, r_t) (-ar_t dt + \sigma \sqrt{r_t} dB_t) \right) \\ &\quad + r_t e^{-\int_0^t r_s ds} \frac{\sigma^2}{2} \frac{\partial^2 F}{\partial x^2}(t, r_t) dt, \end{aligned}$$

hence $F(t, x)$ satisfies the *affine PDE*

$$-xF(t, x) + \frac{\partial F}{\partial t}(t, x) - ax \frac{\partial F}{\partial x}(t, x) + x \frac{\sigma^2}{2} \frac{\partial^2 F}{\partial x^2}(t, x) = 0. \quad (\text{A.7})$$

b) Plugging $F(t, x) = e^{A(T-t)+xC(T-t)}$ into the PDE (A.7) shows that

$$\begin{aligned} e^{A(T-t)+xC(T-t)} \left(-x - A'(T-t) - xC'(T-t) - axC(T-t) + \frac{\sigma^2 x}{2} C^2(T-t) \right) \\ = 0. \end{aligned}$$

Taking successively $x = 0$ and $x = 1$ in the above relation then yields the two equations

$$\left\{ \begin{array}{l} A'(T-t) = 0, \\ -1 - C'(T-t) - aC(T-t) + \frac{\sigma^2}{2} C^2(T-t) = 0. \end{array} \right.$$

Remark: The initial condition $A(0) = 0$ shows that $A(s) = 1$, and it can be shown from the condition $C(0) = 0$ that

$$C(T-t) = \frac{2(1 - e^{\gamma(T-t)})}{2\gamma + (a + \gamma)(e^{\gamma(T-t)} - 1)}, \quad t \in [0, T],$$

with $\gamma = \sqrt{a^2 + 2\sigma^2}$, see e.g. Eq. (3.25) page 66 of Brigo and Mercurio, 2006.

Exercise 1.8

- a) The payoff of the convertible bond is given by $\text{Max}(\alpha S_\tau, P(\tau, T))$.
b) We have

$$\begin{aligned}\text{Max}(\alpha S_\tau, P(\tau, T)) &= P(\tau, T) \mathbb{1}_{\{\alpha S_\tau \leq P(\tau, T)\}} + \alpha S_\tau \mathbb{1}_{\{\alpha S_\tau > P(\tau, T)\}} \\ &= P(\tau, T) + (\alpha S_\tau - P(\tau, T)) \mathbb{1}_{\{\alpha S_\tau > P(\tau, T)\}} \\ &= P(\tau, T) + (\alpha S_\tau - P(\tau, T))^+ \\ &= P(\tau, T) + \alpha(S_\tau - P(\tau, T)/\alpha)^+, \end{aligned}$$

where the latter European call option payoff has the strike price $K := P(\tau, T)/\alpha$.

- c) From the Markov property applied at time $t \in [0, \tau]$, we will write the corporate bond price as a function $C(t, S_t, r_t)$ of the underlying asset price and interest rate, hence we have

$$C(t, S_t, r_t) = \mathbf{E}^* \left[e^{-\int_t^\tau r_s ds} \text{Max}(\alpha S_\tau, P(\tau, T)) \middle| \mathcal{F}_t \right].$$

The martingale property follows from the equalities

$$\begin{aligned}e^{-\int_0^t r_s ds} C(t, S_t, r_t) &= e^{-\int_0^t r_s ds} \mathbf{E}^* \left[e^{-\int_t^\tau r_s ds} \text{Max}(\alpha S_\tau, P(\tau, T)) \middle| \mathcal{F}_t \right] \\ &= \mathbf{E}^* \left[e^{-\int_0^\tau r_s ds} \text{Max}(\alpha S_\tau, P(\tau, T)) \middle| \mathcal{F}_t \right].\end{aligned}$$

- d) We have

$$\begin{aligned}d \left(e^{-\int_0^t r_s ds} C(t, S_t, r_t) \right) &= -r_t e^{-\int_0^t r_s ds} C(t, S_t, r_t) dt + e^{-\int_0^t r_s ds} \frac{\partial C}{\partial t}(t, S_t, r_t) dt \\ &\quad + e^{-\int_0^t r_s ds} \frac{\partial C}{\partial x}(t, S_t, r_t) (r_t S_t dt + \sigma S_t dB_t^{(1)}) \\ &\quad + e^{-\int_0^t r_s ds} \frac{\partial C}{\partial y}(t, S_t, r_t) (\gamma(t, r_t) dt + \eta(t, S_t) dB_t^{(2)}) \\ &\quad + e^{-\int_0^t r_s ds} \frac{\sigma^2}{2} S_t^2 \frac{\partial^2 C}{\partial x^2}(t, S_t, r_t) dt + e^{-\int_0^t r_s ds} \eta^2(t, r_t) \frac{1}{2} \frac{\partial^2 C}{\partial y^2}(t, S_t, r_t) dt \\ &\quad + \rho \sigma S_t \eta(t, r_t) e^{-\int_0^t r_s ds} \frac{\partial^2 C}{\partial x \partial y}(t, S_t, r_t) dt.\end{aligned}\tag{A.8}$$

The martingale property of $(e^{-\int_0^t r_s ds} C(t, S_t, r_t))_{t \in \mathbb{R}_+}$ shows that the sum of terms in factor of dt vanishes in the above Relation (A.8), which yields the PDE

$$\begin{aligned}0 &= -y C(t, x, y) + \frac{\partial C}{\partial t}(t, x, y) dt + ry \frac{\partial C}{\partial x}(t, x, y) + \gamma(t, y) \frac{\partial C}{\partial y}(t, x, y) \\ &\quad + \frac{\sigma^2}{2} x^2 \frac{\partial^2 C}{\partial x^2}(t, x, y) + \eta^2(t, y) \frac{1}{2} \frac{\partial^2 C}{\partial y^2}(t, x, y) + \rho \sigma x \eta(t, y) \frac{\partial^2 C}{\partial x \partial y}(t, x, y),\end{aligned}$$

with the terminal condition

$$C(\tau, x, y) = \text{Max}(\alpha x, F(\tau, y)), \quad \text{where } F(\tau, r_\tau) = P(\tau, T)$$

is the bond pricing function.

- e) The convertible bond can be priced as

$$\begin{aligned}\mathbf{E}^* \left[e^{-\int_t^\tau r_s ds} \text{Max}(\alpha S_\tau, P(\tau, T)) \middle| \mathcal{F}_t \right] &= \mathbf{E}^* \left[e^{-\int_t^\tau r_s ds} P(\tau, T) \middle| \mathcal{F}_t \right] + \alpha \mathbf{E}^* \left[e^{-\int_t^\tau r_s ds} (S_\tau - P(\tau, T)/\alpha)^+ \middle| \mathcal{F}_t \right] \\ &= P(t, T) + \alpha P(t, T) \widehat{\mathbf{E}}[(S_\tau/P(\tau, T) - 1/\alpha)^+ \middle| \mathcal{F}_t].\end{aligned}\tag{A.9}$$



f) By Proposition 3.8 we find

$$dZ_t = (\sigma - \sigma_B(t))Z_t d\hat{W}_t,$$

where $(\hat{W}_t)_{t \in \mathbb{R}_+}$ is a standard Brownian motion under the forward measure \hat{P} .

g) By modeling $(Z_t)_{t \in \mathbb{R}_+}$ as the geometric Brownian motion

$$dZ_t = \sigma(t)Z_t d\hat{W}_t,$$

(A.9) shows that the convertible bond is priced as

$$P(t, T) + \alpha S_t \Phi(d_+) - P(t, T) \Phi(d_-),$$

where

$$d_+ = \frac{1}{v(t, T)} \left(\log \frac{S_t}{P(t, T)} + \frac{v^2(t, \tau)}{2} \right),$$

$$d_- = \frac{1}{v(t, T)} \left(\log \frac{S_t}{P(t, T)} - \frac{v^2(t, \tau)}{2} \right),$$

$$\text{and } v^2(t, T) = \int_t^T \sigma^2(s, T) ds, 0 < t < T.$$

Exercise 1.9 We have

$$\begin{aligned} \frac{\partial}{\partial r} P_c(0, n) &= \frac{\partial}{\partial r} \left(\frac{1}{(1+r)^n} + \frac{c}{r} \left(1 - \frac{1}{(1+r)^n} \right) \right) \\ &= -\frac{n}{(1+r)^{n+1}} - \frac{c}{r^2} \left(1 - \frac{1}{(1+r)^n} \right) + \frac{nc}{r(1+r)^{n+1}}, \end{aligned}$$

hence

$$\begin{aligned} D_c(0, n) &= -\frac{1+r}{P_c(0, n)} \frac{\partial}{\partial r} P_c(0, n) \\ &= -\frac{n}{(1+r)^n} - \frac{(1+r)c}{r^2} \left(1 - \frac{1}{(1+r)^n} \right) + \frac{nc}{r(1+r)^n} \\ &\quad \frac{1}{(1+r)^n} + \frac{c}{r} \left(1 - \frac{1}{(1+r)^n} \right) \\ &= -\frac{-nr - \frac{1+r}{r} (c((1+r)^n - 1)) + nc}{r + c((1+r)^n - 1)} \\ &= -\frac{1+r - nr - \frac{1+r}{r} (r + c((1+r)^n - 1)) + nc}{r + c((1+r)^n - 1)} \\ &= \frac{1+r}{r} - \frac{1+r+n(c-r)}{r+c((1+r)^n-1)} \\ &= \frac{(1-c/r)n}{1+c((1+r)^n-1)/r} + \frac{1+r}{r} \left(\frac{c((1+r)^n-1)}{r+c((1+r)^n-1)} \right), \end{aligned}$$

with $D_0(0, n) = n$. We note that

$$\lim_{n \rightarrow \infty} D_c(0, n) = \lim_{n \rightarrow \infty} \left(\frac{1+r}{r} - \frac{1+r+n(c-r)}{r+c((1+r)^n-1)} \right) = 1 + \frac{1}{r}.$$

When n becomes large, the duration (or relative sensitivity) of the bond price converges to $1 + 1/r$ whenever the (nonnegative) coupon amount c is nonzero, otherwise the bond duration of $P_c(0, n)$

is n . In particular, the presence of a nonzero coupon makes the duration (or relative sensitivity) of the bond price bounded as n increases, whereas the duration n of $P_0(0,n)$ goes to infinity as n increases.

As a consequence, the presence of the coupon tends to put an upper limit the risk and sensitivity of bond prices with respect to the market interest rate r as n becomes large, which can be used for bond *immunization*. Note that the duration $D_c(0,n)$ can also be written as the relative average

$$D_c(0,n) = \frac{1}{P_c(0,n)} \left(\frac{n}{(1+r)^n} + c \sum_{k=1}^n \frac{k}{(1+r)^k} \right)$$

of zero coupon bond durations, weighted by their respective zero-coupon prices.

Exercise 1.10

a) We have

$$P(t,T) = P(s,T) \exp \left(\int_s^t r_u du + \int_s^t \sigma_u^T dB_u - \frac{1}{2} \int_s^t |\sigma_u^T|^2 du \right),$$

$$0 \leq s \leq t \leq T.$$

b) We have

$$d \left(e^{-\int_0^t r_s ds} P(t,T) \right) = e^{-\int_0^t r_s ds} \sigma_t^T P(t,T) dB_t,$$

which gives a martingale after integration, from the properties of the Itô integral.

c) By the martingale property of the previous question we have

$$\begin{aligned} \mathbf{E} \left[e^{-\int_0^T r_s ds} \middle| \mathcal{F}_t \right] &= \mathbf{E} \left[P(T,T) e^{-\int_0^T r_s ds} \middle| \mathcal{F}_t \right] \\ &= P(t,T) e^{-\int_0^t r_s ds}, \quad 0 \leq t \leq T. \end{aligned}$$

d) By the previous question we have

$$\begin{aligned} P(t,T) &= e^{\int_0^t r_s ds} \mathbf{E} \left[e^{-\int_0^T r_s ds} \middle| \mathcal{F}_t \right] \\ &= \mathbf{E} \left[e^{\int_0^t r_s ds} e^{-\int_0^T r_s ds} \middle| \mathcal{F}_t \right] \\ &= \mathbf{E} \left[e^{-\int_t^T r_s ds} \middle| \mathcal{F}_t \right], \quad 0 \leq t \leq T, \end{aligned}$$

since $e^{-\int_0^t r_s ds}$ is an \mathcal{F}_t -measurable random variable.

e) We have

$$\begin{aligned} \frac{P(t,S)}{P(t,T)} &= \frac{P(s,S)}{P(s,T)} \exp \left(\int_s^t (\sigma_u^S - \sigma_u^T) dB_u - \frac{1}{2} \int_s^t (|\sigma_u^S|^2 - |\sigma_u^T|^2) du \right) \\ &= \frac{P(s,S)}{P(s,T)} \exp \left(\int_s^t (\sigma_u^S - \sigma_u^T) dB_u^T - \frac{1}{2} \int_s^t (\sigma_u^S - \sigma_u^T)^2 du \right), \end{aligned}$$

$0 \leq t \leq T$, hence letting $s = t$ and $t = T$ in the above expression, we find

$$P(T,S) = \frac{P(t,S)}{P(t,T)} \exp \left(\int_t^T (\sigma_s^S - \sigma_s^T) dB_s^T - \frac{1}{2} \int_t^T (\sigma_s^S - \sigma_s^T)^2 ds \right).$$

f) We have

$$\begin{aligned} P(t,T) \widehat{\mathbf{E}}[(P(T,S) - \kappa)^+] &= P(t,T) \widehat{\mathbf{E}} \left[\left(\frac{P(t,S)}{P(t,T)} e^{\int_t^T (\sigma_s^S - \sigma_s^T) dB_s^T - \int_t^T (\sigma_s^S - \sigma_s^T)^2 ds / 2} - \kappa \right)^+ \right] \\ &= P(t,T) \mathbf{E}^* [(e^X - \kappa)^+ | \mathcal{F}_t] \\ &= P(t,T) e^{m(t) + v^2(t)/2} \Phi \left(\frac{v(t)}{2} + \frac{1}{v(t)} (m(t) + v^2(t)/2 - \log \kappa) \right) \end{aligned}$$



$$-\kappa P(t, T) \Phi \left(-\frac{v(t)}{2} + \frac{1}{v(t)} (m(t) + v^2(t)/2 - \log \kappa) \right),$$

where

$$m(t) := \log \frac{P(t, S)}{P(t, T)} - \frac{1}{2} \int_t^T (\sigma_s^S - \sigma_s^T)^2 ds,$$

and X is a centered Gaussian random variable with variance

$$v^2(t) := \int_t^T (\sigma_s^S - \sigma_s^T)^2 ds,$$

given \mathcal{F}_t . This yields

$$\begin{aligned} & P(t, T) \widehat{\mathbb{E}}[(P(T, S) - \kappa)^+] \\ &= P(t, S) \Phi \left(\frac{v(t)}{2} + \frac{1}{v(t)} \log \frac{P(t, S)}{\kappa P(t, T)} \right) - \kappa P(t, T) \Phi \left(-\frac{v(t)}{2} + \frac{1}{v(t)} \log \frac{P(t, S)}{\kappa P(t, T)} \right). \end{aligned}$$

Exercise 1.11 From Proposition 1.2, the bond pricing PDE is

$$\begin{cases} \frac{\partial F}{\partial t}(t, x) = xF(t, x) - (\alpha - \beta x) \frac{\partial F}{\partial x}(t, x) - \frac{1}{2} \sigma^2 x^2 \frac{\partial^2 F}{\partial x^2}(t, x) \\ F(T, x) = 1. \end{cases}$$

We search for a solution of the form

$$F(t, x) = e^{A(T-t) - xB(T-t)},$$

with $A(0) = B(0) = 0$, which implies

$$\begin{cases} A'(s) = 0 \\ B'(s) + \beta B(s) + \frac{1}{2} \sigma^2 B^2(s) = 1, \end{cases}$$

hence in particular $A(s) = 0$, $s \in \mathbb{R}$, and $B(s)$ solves a Riccati equation, whose solution can be checked to be

$$B(s) = \frac{2(e^{\gamma s} - 1)}{2\gamma + (\beta + \gamma)(e^{\gamma s} - 1)},$$

with $\gamma = \sqrt{\beta^2 + 2\sigma^2}$.

Exercise 1.12

a) We have

$$\begin{cases} y_{0,1} = -\frac{1}{T_1} \log P(0, T_1) = 9.53\%, \\ y_{0,2} = -\frac{1}{T_2} \log P(0, T_2) = 9.1\%, \\ y_{1,2} = -\frac{1}{T_2 - T_1} \log \frac{P(0, T_2)}{P(T_1, T_2)} = 8.6\%, \end{cases}$$

with $T_1 = 1$ and $T_2 = 2$.

b) We have

$$P_c(1, 2) = (\$1 + \$0.1) \times P_0(1, 2) = (\$1 + \$0.1) \times e^{-(T_2 - T_1)y_{1,2}} = \$1.00914,$$

and

$$\begin{aligned} P_c(0, 2) &= (\$1 + \$0.1) \times P_0(0, 2) + \$0.1 \times P_0(0, 1) \\ &= (\$1 + \$0.1) \times e^{-(T_2 - T_1)y_{0,2}} + \$0.1 \times e^{-(T_2 - T_1)y_{0,1}} \\ &= \$1.00831. \end{aligned}$$

Exercise 1.13

- a) The discretization $r_{t_{k+1}} := r_{t_k} + (a - br_{t_k})\Delta t \pm \sigma\sqrt{\Delta t}$, does not lead to a binomial tree as r_{t_2} could be obtained in four different ways from r_{t_0} as

$$r_{t_2} = r_{t_1}(1 - b\Delta t) + a\Delta t \pm \sigma\sqrt{\Delta t}$$

$$= \begin{cases} (r_{t_0}(1 - b\Delta t) + a\Delta t + \sigma\sqrt{\Delta t})(1 - b\Delta t) + a\Delta t + \sigma\sqrt{\Delta t} \\ (r_{t_0}(1 - b\Delta t) + a\Delta t + \sigma\sqrt{\Delta t})(1 - b\Delta t) + a\Delta t - \sigma\sqrt{\Delta t} \\ (r_{t_0}(1 - b\Delta t) + a\Delta t - \sigma\sqrt{\Delta t})(1 - b\Delta t) + a\Delta t + \sigma\sqrt{\Delta t} \\ (r_{t_0}(1 - b\Delta t) + a\Delta t - \sigma\sqrt{\Delta t})(1 - b\Delta t) + a\Delta t - \sigma\sqrt{\Delta t}. \end{cases}$$

- b) By the Girsanov Theorem, the process $(r_t/\sigma)_{t \in [0, T]}$ with

$$\frac{dr_t}{\sigma} = \frac{a - br_t}{\sigma} dt + dB_t$$

is a standard Brownian motion under the probability measure \mathbb{Q} with Radon-Nikodym density

$$\begin{aligned} \frac{d\mathbb{Q}}{d\mathbb{P}} &= \exp\left(-\frac{1}{\sigma} \int_0^T (a - br_t) dB_t - \frac{1}{2\sigma^2} \int_0^T (a - br_t)^2 dt\right) \\ &\simeq \exp\left(-\frac{1}{\sigma^2} \int_0^T (a - br_t)(dr_t - (a - br_t)dt) - \frac{1}{2\sigma^2} \int_0^T (a - br_t)^2 dt\right) \\ &= \exp\left(-\frac{1}{\sigma^2} \int_0^T (a - br_t) dr_t + \frac{1}{2\sigma^2} \int_0^T (a - br_t)^2 dt\right). \end{aligned}$$

In other words, if we generate $(r_t/\sigma)_{t \in [0, T]}$ and the increments $\sigma^{-1}dr_t \simeq \pm\sqrt{\Delta t}$ as a standard Brownian motion under \mathbb{Q} , then, under the probability measure \mathbb{P} such that

$$\frac{d\mathbb{P}}{d\mathbb{Q}} = \exp\left(\frac{1}{\sigma^2} \int_0^T (a - br_t) dr_t - \frac{1}{2\sigma^2} \int_0^T (a - br_t)^2 dt\right),$$

the process

$$dB_t = \frac{dr_t}{\sigma} - \frac{a - br_t}{\sigma} dt$$

will be a standard Brownian motion under \mathbb{P} , and the samples

$$dr_t = (a - br_t)dt + \sigma dB_t$$

of $(r_t)_{t \in [0, T]}$ will be distributed as a Vasicek process under \mathbb{P} .

- c) Approximating the standard Brownian increment $\sigma^{-1}dr_t$ under \mathbb{Q} by $\pm\sqrt{\Delta t}$, we have

$$\begin{aligned} 2^{T/\Delta T} \prod_{0 < t < T} \left(\frac{1}{2} \pm \frac{a - br_t}{2\sigma} \sqrt{\Delta t} \right) &= \prod_{0 < t < T} \left(1 \pm \frac{a - br_t}{\sigma} \sqrt{\Delta t} \right) \\ &= \exp\left(\log \prod_{0 < t < T} \left(1 \pm \frac{a - br_t}{\sigma} \sqrt{\Delta t} \right)\right) \\ &= \exp\left(\sum_{0 < t < T} \log \left(1 \pm \frac{a - br_t}{\sigma} \sqrt{\Delta t} \right)\right) \\ &\simeq \exp\left(\sum_{0 < t < T} \frac{a - br_t}{\sigma} (\pm\sqrt{\Delta t}) - \frac{1}{2} \sum_{0 < t < T} \frac{(a - br_t)^2}{\sigma^2} \Delta t\right) \\ &\simeq \exp\left(\frac{1}{\sigma^2} \int_0^T (a - br_t) dr_t - \frac{1}{2\sigma^2} \int_0^T (a - br_t)^2 dt\right) \end{aligned}$$



$$= \frac{dP}{dQ}.$$

d) We check that

$$\begin{aligned}\mathbb{E}[\Delta r_{t_1} | r_{t_0}] &= (a - br_{t_0})\Delta t \\ &= p(r_{t_0})\sigma\sqrt{\Delta t} - (1 - p(r_{t_0}))\sigma\sqrt{\Delta t} \\ &= \sigma p(r_{t_0})\sqrt{\Delta t} - \sigma q(r_{t_0})\sqrt{\Delta t},\end{aligned}$$

with

$$p(r_{t_0}) = \frac{1}{2} + \frac{a - br_{t_0}}{2\sigma}\sqrt{\Delta t} \quad \text{and} \quad q(r_{t_0}) = \frac{1}{2} - \frac{a - br_{t_0}}{2\sigma}\sqrt{\Delta t}.$$

Similarly, we have

$$\begin{aligned}\mathbb{E}[\Delta r_{t_2} | r_{t_1}] &= (a - br_{t_1})\Delta t \\ &= p(r_{t_1})\sigma\sqrt{\Delta t} - (1 - p(r_{t_1}))\sigma\sqrt{\Delta t} \\ &= \sigma p(r_{t_1})\sqrt{\Delta t} - \sigma q(r_{t_1})\sqrt{\Delta t},\end{aligned}$$

with

$$p(r_{t_1}) = \frac{1}{2} + \frac{a - br_{t_1}}{2\sigma}\sqrt{\Delta t}, \quad q(r_{t_1}) = \frac{1}{2} - \frac{a - br_{t_1}}{2\sigma}\sqrt{\Delta t}.$$

Exercise 1.14

a) We have

$$P(1,2) = \mathbb{E}^* \left[\frac{100}{1+r_1} \right] = \frac{100}{2(1+r_1^u)} + \frac{100}{2(1+r_1^d)}.$$

b) We have

$$P(0,2) = \frac{100}{2(1+r_0)(1+r_1^u)} + \frac{100}{2(1+r_0)(1+r_1^d)}.$$

c) We have $P(0,1) = 91.74 = 100/(1+r_0)$, hence

$$r_0 = \frac{100 - P(0,1)}{P(0,1)} = 100/91.74 - 1 = 0.090037061 \simeq 9\%.$$

d) We have

$$83.40 = P(0,2) = \frac{P(0,1)}{2(1+r_1^u)} + \frac{P(0,1)}{2(1+r_1^d)}$$

and $r_1^u/r_1^d = e^{2\sigma\sqrt{\Delta t}}$, hence

$$83.40 = P(0,2) = \frac{P(0,1)}{2(1+r_1^d e^{2\sigma\sqrt{\Delta t}})} + \frac{P(0,1)}{2(1+r_1^d)}$$

or

$$e^{2\sigma\sqrt{\Delta t}}(r_1^d)^2 + 2r_1^d e^{\sigma\sqrt{\Delta t}} \cosh(\sigma\sqrt{\Delta t}) \left(1 - \frac{P(0,1)}{2P(0,2)} \right) + 1 - \frac{P(0,1)}{P(0,2)} = 0,$$

and

$$\begin{aligned}r_1^d &= e^{-\sigma\sqrt{\Delta t}} \left(\cosh(\sigma\sqrt{\Delta t}) \left(\frac{P(0,1)}{2P(0,2)} - 1 \right) \right. \\ &\quad \left. \pm \sqrt{\left(\frac{P(0,1)}{2P(0,2)} - 1 \right)^2 \cosh^2(\sigma\sqrt{\Delta t}) + \frac{P(0,1)}{P(0,2)} - 1} \right) \\ &= 0.078684844 \simeq 7.87\%,\end{aligned}$$

and

$$r_1^u = r_1^d e^{2\sigma\sqrt{\Delta t}}$$

$$\begin{aligned}
&= e^{\sigma\sqrt{\Delta t}} \left(\cosh(\sigma\sqrt{\Delta t}) \left(\frac{P(0,1)}{2P(0,2)} - 1 \right) \right. \\
&\quad \left. \pm \sqrt{\left(\frac{P(0,1)}{2P(0,2)} - 1 \right)^2 \cosh^2(\sigma\sqrt{\Delta t}) + \frac{P(0,1)}{P(0,2)} - 1} \right) \\
&= 0.122174525 \simeq 12.22\%.
\end{aligned}$$

We also find

$$\mu = \frac{1}{\Delta t} \left(\sigma\sqrt{\Delta t} + \log \frac{r_1^d}{r_0} \right) = \frac{1}{\Delta t} \left(-\sigma\sqrt{\Delta t} + \log \frac{r_1^u}{r_0} \right) = 0.085229181 \simeq 8.52\%.$$

Exercise 1.15

- a) When $n = 1$ the relation (1.4.38) shows that $\tilde{f}(t, t, T_1) = f(t, t, T_1)$ with $F(t, x) = c_1 e^{-(T_1-1)x}$ and $P(t, T_1) = c_1 e^{f(t, t, T_1)}$, hence

$$D(t, T_1) := -\frac{1}{P(t, T_1)} \frac{\partial F}{\partial x}(t, f(t, t, T_1)) = T_1 - t, \quad 0 \leq t \leq T_1.$$

- b) In general, we have

$$\begin{aligned}
D(t, T_n) &= -\frac{1}{P(t, T_n)} \frac{\partial F}{\partial x}(t, \tilde{f}(t, t, T_n)) \\
&= \frac{1}{P(t, T_n)} \sum_{k=1}^n (T_k - t) c_k e^{-(T_k-t)\tilde{f}(t,t,T_n)} \\
&= \sum_{k=1}^n (T_k - t) w_k,
\end{aligned}$$

where

$$\begin{aligned}
w_k &:= \frac{c_k}{P(t, T_n)} e^{-(T_k-t)\tilde{f}(t,t,T_n)} \\
&= \frac{c_k e^{-(T_k-t)\tilde{f}(t,t,T_n)}}{\sum_{l=1}^n c_l e^{-(T_l-t)f(t,t,T_l)}} \\
&= \frac{c_k e^{-(T_k-t)\tilde{f}(t,t,T_n)}}{\sum_{l=1}^n c_l e^{-(T_l-t)\tilde{f}(t,t,T_n)}}, \quad k = 1, 2, \dots, n,
\end{aligned}$$

and the weights w_1, w_2, \dots, w_n satisfy

$$\sum_{k=1}^n w_k = 1.$$

- c) We have

$$\begin{aligned}
C(t, T_n) &= \frac{1}{P(t, T_n)} \frac{\partial^2 F}{\partial x^2}(t, \tilde{f}(t, t, T_n)) \\
&= \sum_{k=1}^n (T_k - t)^2 w_k \\
&= \sum_{k=1}^n (T_k - t - D(t, T_n))^2 w_k + 2D(t, T_n) \sum_{k=1}^n (T_k - t) w_k - (D(t, T_n))^2 \\
&= (D(t, T_n))^2 + \sum_{k=1}^n (T_k - t - D(t, T_n))^2 w_k
\end{aligned}$$



$$= (D(t, T_n))^2 + (S(t, T_n))^2,$$

where

$$(S(t, T_n))^2 := \sum_{k=1}^n (T_k - t - D(t, T_n))^2 w_k.$$

d) We have

$$\begin{aligned} D(t, T_n) &= \frac{1}{P(t, T_n)} \sum_{k=1}^n c_k B(T_k - t) e^{A(T_k - t) + B(T_k - t)f_\alpha(t, t, T_n)} \\ &= \frac{1}{e^{A(T_n - t) + B(T_n - t)f_\alpha(t, t, T_n)}} \sum_{k=1}^n c_k B(T_k - t) e^{A(T_k - t) + B(T_k - t)f_\alpha(t, t, T_n)} \\ &= \sum_{k=1}^n c_k B(T_k - t) e^{A(T_k - t) - A(T_n - t) + (B(T_k - t) - B(T_n - t))f_\alpha(t, t, T_n)}. \end{aligned}$$

e) We have

$$\begin{aligned} D(t, T_n) &= \frac{1}{b} \sum_{k=1}^n (1 - e^{-(T_k - t)b}) c_k e^{A(T_k - t) - A(T_n - t) + (e^{-(T_n - t)b} - e^{-(T_k - t)b})f_\alpha(t, t, T_n)/b} \\ &= \frac{1}{b} \sum_{k=1}^n (1 - e^{-(T_k - t)b}) c_k e^{A(T_k - t) - A(T_n - t)} (P(t, t + \alpha(T_n - t)))^{\frac{(e^{-(T_n - t)b} - e^{-(T_k - t)b})}{(T_n - t)\alpha b}}. \end{aligned}$$

Chapter 2

Exercise 2.1

a) By partial differentiation with respect to T under the expectation $\widehat{\mathbb{E}}$, we have

$$\begin{aligned} \frac{\partial P}{\partial T}(t, T) &= \frac{\partial}{\partial T} \mathbb{E}^* \left[e^{-\int_t^T r_s ds} \mid \mathcal{F}_t \right] \\ &= \mathbb{E}^* \left[-r_T e^{-\int_t^T r_s ds} \mid \mathcal{F}_t \right] \\ &= -P(t, T) \widehat{\mathbb{E}}[r_T \mid \mathcal{F}_t]. \end{aligned}$$

b) As a consequence of Question (a)), we find

$$f(t, T) = -\frac{1}{P(t, T)} \frac{\partial P}{\partial T}(t, T) = \widehat{\mathbb{E}}[r_T \mid \mathcal{F}_t], \quad 0 \leq t \leq T, \quad (\text{A.10})$$

see Relation (22) page 10 of Mamon, 2004.

c) The martingale property of $(f(t, T))_{t \in [0, T]}$ is a under the forward measure $\widehat{\mathbb{E}}$ follows from Relation (A.10) and the from the tower property of conditional expectations.

Remark. In the Vasicek model, by (1.1.2) we have

$$\begin{aligned} r_T &= r_0 e^{-bT} + \frac{a}{b} (1 - e^{-bT}) + \sigma \int_0^T e^{-(T-s)b} dW_s \\ &= r_0 e^{-bT} + \frac{a}{b} (1 - e^{-bT}) + \sigma \int_0^T e^{-(T-s)b} d\widehat{W}_s \\ &\quad - \frac{\sigma^2}{b} \int_0^T e^{-(T-s)b} (1 - e^{-(T-s)b}) ds \\ &= r_0 e^{-bT} + \frac{a}{b} (1 - e^{-bT}) + \sigma \int_0^T e^{-(T-s)b} d\widehat{W}_s \\ &\quad - \frac{\sigma^2}{b} \int_0^T e^{-(T-s)b} ds + \frac{\sigma^2}{b} \int_0^T e^{-2(T-s)b} ds, \end{aligned}$$

hence

$$\widehat{\mathbb{E}}[r_T \mid \mathcal{F}_t] = r_0 e^{-bT} + \frac{a}{b} (1 - e^{-bT}) + \sigma \int_0^t e^{-(T-s)b} d\widehat{W}_s$$

$$\begin{aligned}
& -\frac{\sigma^2}{b} \int_0^T e^{-(T-s)b} ds + \frac{\sigma^2}{b} \int_0^T e^{-2(T-s)b} ds \\
= & r_0 e^{-bT} + \frac{a}{b} (1 - e^{-bT}) + \sigma \int_0^t e^{-(T-s)b} dW_s \\
& + \frac{\sigma^2}{b} \int_0^t e^{-(T-s)b} (1 - e^{-(T-s)b}) ds \\
& - \frac{\sigma^2}{b} \int_0^T e^{-(T-s)b} ds + \frac{\sigma^2}{b} \int_0^T e^{-2(T-s)b} ds \\
= & r_0 e^{-bT} + \frac{a}{b} (1 - e^{-bT}) + e^{-(T-t)b} \left(r_t - r_0 e^{-bt} - \frac{a}{b} (1 - e^{-bt}) \right) \\
& + \frac{\sigma^2}{b} \int_0^t e^{-(T-s)b} ds - \frac{\sigma^2}{b} \int_0^t e^{-2(T-s)b} ds \\
& - \frac{\sigma^2}{b} \int_0^T e^{-(T-s)b} ds + \frac{\sigma^2}{b} \int_0^T e^{-2(T-s)b} ds \\
= & \frac{a}{b} + e^{-(T-t)b} \left(r_t - \frac{a}{b} \right) - \frac{\sigma^2}{b} \int_t^T e^{-(T-s)b} ds + \frac{\sigma^2}{b} \int_t^T e^{-2(T-s)b} ds \\
= & \frac{a}{b} + e^{-(T-t)b} \left(r_t - \frac{a}{b} \right) - \frac{\sigma^2}{b} \int_0^{T-t} e^{-bs} ds + \frac{\sigma^2}{b} \int_0^{T-t} e^{-2bs} ds \\
= & \frac{a}{b} + e^{-(T-t)b} \left(r_t - \frac{a}{b} \right) - \frac{\sigma^2}{b^2} (1 - e^{-(T-t)b}) + \frac{\sigma^2}{b^2} (1 - e^{-2(T-t)b}) \\
= & \frac{a}{b} - \frac{\sigma^2}{2b^2} + e^{-(T-t)b} \left(r_t - \frac{a}{b} + \frac{\sigma^2}{b^2} \right) - \frac{\sigma^2}{b^2} e^{-2(T-t)b},
\end{aligned}$$

which recovers (2.4.1).

Exercise 2.2 We have

$$P(0, T_2) = \exp \left(- \int_0^{T_2} f(t, s) ds \right) = e^{-r_1 T_1 - r_2 (T_2 - T_1)}, \quad t \in [0, T_2],$$

and

$$P(T_1, T_2) = \exp \left(- \int_{T_1}^{T_2} f(t, s) ds \right) = e^{-r_2 (T_2 - T_1)}, \quad t \in [0, T_2],$$

from which we deduce

$$r_2 = -\frac{1}{T_2 - T_1} \log P(T_1, T_2),$$

and

$$\begin{aligned}
r_1 &= -r_2 \frac{T_2 - T_1}{T_1} - \frac{1}{T_1} \log P(0, T_2) \\
&= \frac{1}{T_1} \log P(T_1, T_2) - \frac{1}{T_1} \log P(0, T_2) \\
&= -\frac{1}{T_1} \log \frac{P(0, T_2)}{P(T_1, T_2)}.
\end{aligned}$$

Exercise 2.3

a) We check that $P(T, T) = e^{X_T^T} = 1$.

b) We have

$$f(t, T, S) = -\frac{1}{S - T} (X_t^S - X_t^T - \mu(S - T))$$



$$\begin{aligned}
&= \mu - \sigma \frac{1}{S-T} \left((S-t) \int_0^t \frac{1}{S-s} dB_s - (T-t) \int_0^t \frac{1}{T-s} dB_s \right) \\
&= \mu - \sigma \frac{1}{S-T} \int_0^t \left(\frac{S-t}{S-s} - \frac{T-t}{T-s} \right) dB_s \\
&= \mu - \sigma \frac{1}{S-T} \int_0^t \frac{(T-s)(S-t) - (T-t)(S-s)}{(S-s)(T-s)} dB_s \\
&= \mu + \frac{\sigma}{S-T} \int_0^t \frac{(s-t)(S-T)}{(S-s)(T-s)} dB_s.
\end{aligned}$$

c) We have

$$f(t, T) = \mu - \sigma \int_0^t \frac{t-s}{(T-s)^2} dB_s.$$

d) We note that

$$\lim_{T \searrow t} f(t, T) = \mu - \sigma \int_0^t \frac{1}{t-s} dB_s$$

does not exist in $L^2(\Omega)$.

e) By Itô's calculus we have

$$\begin{aligned}
\frac{dP(t, T)}{P(t, T)} &= \sigma dB_t + \frac{1}{2} \sigma^2 dt + \mu dt - \frac{X_t^T}{T-t} dt \\
&= \sigma dB_t + \frac{1}{2} \sigma^2 dt - \frac{\log P(t, T)}{T-t} dt, \quad t \in [0, T].
\end{aligned}$$

f) Letting

$$\begin{aligned}
r_t^T &:= \mu + \frac{1}{2} \sigma^2 - \frac{X_t^T}{T-t} \\
&= \mu + \frac{1}{2} \sigma^2 - \sigma \int_0^t \frac{dB_s}{T-s},
\end{aligned}$$

by Question (e)) we find that

$$\frac{dP(t, T)}{P(t, T)} = \sigma dB_t + r_t^T dt, \quad 0 \leq t \leq T.$$

g) The equation of Question (f)) can be solved as

$$P(t, T) = P(0, T) \exp \left(\sigma B_t - \frac{\sigma^2 t}{2} + \int_0^t r_s^T ds \right), \quad 0 \leq t \leq T,$$

hence the process

$$P(t, T) \exp \left(- \int_0^t r_s^T ds \right) = P(0, T) \exp \left(\sigma B_t - \frac{\sigma^2 t}{2} \right), \quad 0 \leq t \leq T,$$

is a martingale under \mathbb{P}^* , with the relation

$$\begin{aligned}
P(t, T) \exp \left(- \int_0^t r_s^T ds \right) &= \mathbb{E}^* \left[P(T, T) \exp \left(- \int_0^T r_s^T ds \right) \middle| \mathcal{F}_t \right] \\
&= \mathbb{E}^* \left[\exp \left(- \int_0^T r_s^T ds \right) \middle| \mathcal{F}_t \right], \quad 0 \leq t \leq T,
\end{aligned}$$

showing that

$$\begin{aligned}
P(t, T) &= \exp \left(\int_0^t r_s^T ds \right) \mathbb{E}^* \left[\exp \left(- \int_0^T r_s^T ds \right) \middle| \mathcal{F}_t \right] \\
&= \mathbb{E}^* \left[\exp \left(\int_0^t r_s^T ds \right) \exp \left(- \int_0^T r_s^T ds \right) \middle| \mathcal{F}_t \right] \\
&= \mathbb{E}^* \left[\exp \left(- \int_t^T r_s^T ds \right) \middle| \mathcal{F}_t \right], \quad 0 \leq t \leq T.
\end{aligned}$$

h) By Question (g)) we have

$$\mathbb{E} \left[\frac{d\mathbb{P}_T}{d\mathbb{P}} \mid \mathcal{F}_t \right] = \frac{P(t, T)}{P(0, T)} e^{-\int_0^t r_s^T ds} = e^{\sigma B_t - \sigma^2 t / 2}, \quad 0 \leq t \leq T.$$

i) By the Girsanov Theorem, the process $\hat{B}_t := B_t - \sigma t$ is a standard Brownian motion under \mathbb{P}_T .

j) We have

$$\begin{aligned} \log P(T, S) &= -\mu(S-T) + \sigma \int_0^T \frac{S-T}{S-s} dB_s \\ &= -\mu(S-T) + \sigma \int_0^t \frac{S-T}{S-s} dB_s + \sigma \int_t^T \frac{S-T}{S-s} dB_s \\ &= \frac{S-T}{S-t} \log P(t, S) + \sigma \int_t^T \frac{S-T}{S-s} dB_s \\ &= \frac{S-T}{S-t} \log P(t, S) + \sigma \int_t^T \frac{S-T}{S-s} d\hat{B}_s + \sigma^2 \int_t^T \frac{S-T}{S-s} ds \\ &= \frac{S-T}{S-t} \log P(t, S) + \sigma \int_t^T \frac{S-T}{S-s} d\hat{B}_s + (S-T)\sigma^2 \log \frac{S-t}{S-T}, \end{aligned}$$

$$0 < T < S.$$

k) We have

$$\begin{aligned} &P(t, T) \mathbb{E}_T [(P(T, S) - K)^+ \mid \mathcal{F}_t] \\ &= P(t, T) \mathbb{E} [(e^{X_t} - K)^+ \mid \mathcal{F}_t] \\ &= P(t, T) e^{m_t + v_t^2 / 2} \Phi \left(\frac{v_t}{2} + \frac{1}{v_t} (m_t + v_t^2 / 2 - \log \kappa) \right) \\ &\quad - \kappa P(t, T) \Phi \left(-\frac{v_t}{2} + \frac{1}{v_t} (m_t + v_t^2 / 2 - \log \kappa) \right) \\ &= P(t, T) e^{m_t + v_t^2 / 2} \Phi \left(v_t + \frac{1}{v_t} (m_t - \log \kappa) \right) - \kappa P(t, T) \Phi \left(\frac{1}{v_t} (m_t - \log \kappa) \right), \end{aligned}$$

with

$$m_t = \frac{S-T}{S-t} \log P(t, S) + (S-T)\sigma^2 \log \frac{S-t}{S-T}$$

and

$$\begin{aligned} v_t^2 &= \sigma^2 \int_t^T \frac{(S-T)^2}{(S-s)^2} ds \\ &= (S-T)^2 \sigma^2 \left(\frac{1}{S-T} - \frac{1}{S-t} \right) \\ &= (S-T) \sigma^2 \frac{(T-t)}{(S-t)}, \end{aligned}$$

hence

$$\begin{aligned} &P(t, T) \mathbb{E}_T [(P(T, S) - K)^+ \mid \mathcal{F}_t] \\ &= P(t, T) (P(t, S))^{(S-T)(S-t)} \left(\frac{S-t}{S-T} \right)^{(S-T)\sigma^2} e^{v_t^2 / 2} \\ &\quad \times \Phi \left(v_t + \frac{1}{v_t} \log \left(\frac{(P(t, S))^{(S-T)(S-t)}}{\kappa} \left(\frac{S-t}{S-T} \right)^{(S-T)\sigma^2} \right) \right) \\ &\quad - \kappa P(t, T) \Phi \left(\frac{1}{v_t} \log \left(\frac{(P(t, S))^{(S-T)(S-t)}}{\kappa} \left(\frac{S-t}{S-T} \right)^{(S-T)\sigma^2} \right) \right). \end{aligned}$$

Exercise 2.4



a) In the Vašíček, 1977 model, we have

$$\begin{aligned} P(t, T) &= \mathbb{E} \left[\exp \left(- \int_t^T r_s ds \right) \right] \\ &= \mathbb{E} \left[\exp \left(- \int_t^T h(s) ds - \int_t^T X_s ds \right) \right] \\ &= \exp \left(- \int_t^T h(s) ds \right) \mathbb{E} \left[\exp \left(- \int_t^T X_s ds \right) \right] \\ &= \exp \left(- \int_t^T h(s) ds + A(T-t) + X_t C(T-t) \right), \quad 0 \leq t \leq T, \end{aligned}$$

hence, since $X_0 = 0$ we find $P(0, T) = \exp \left(- \int_0^T h(s) ds + A(T) \right)$.

b) By the identification

$$\begin{aligned} P(t, T) &= \exp \left(- \int_t^T h(s) ds + A(T-t) + X_t C(T-t) \right) \\ &= \exp \left(- \int_t^T f(t, s) ds \right), \end{aligned}$$

we find

$$\int_t^T h(s) ds = \int_t^T f(t, s) ds + A(T-t) + X_t C(T-t),$$

and by differentiation with respect to T this yields

$$h(T) = f(t, T) + A'(T-t) + X_t C'(T-t), \quad 0 \leq t \leq T,$$

where

$$A(T-t) = \frac{4ab - 3\sigma^2}{4b^3} + \frac{\sigma^2 - 2ab}{2b^2}(T-t) + \frac{\sigma^2 - ab}{b^3} e^{-b(T-t)} - \frac{\sigma^2}{4b^3} e^{-2b(T-t)}.$$

Given an initial market data curve $f^M(0, T)$, the matching $f^M(0, T) = f(0, T)$ can be achieved at time $t = 0$ by letting

$$h(T) := f^M(0, T) + A'(T) = f^M(0, T) + \frac{\sigma^2 - 2ab}{2b^2} - \frac{\sigma^2 - ab}{b^2} e^{-bT} + \frac{\sigma^2}{2b^2} e^{-2bT},$$

$T > 0$. Note however that in general, at time $t \in (0, T]$ we will have

$$h(T) = f(t, T) + A'(T-t) + X_t C'(T-t) = f^M(0, T) + A'(T),$$

and the relation

$$f(t, T) = f^M(0, T) + A'(T) - A'(T-t) - X_t C'(T-t), \quad t \in [0, T],$$

will allow us to match market data at time $t = 0$ only, i.e. for the initial curve. In any case, model calibration is to be done at time $t = 0$.

Exercise 2.5 From the definition

$$L(t, t, T) = \frac{1}{T-t} \left(\frac{1}{P(t, T)} - 1 \right),$$

we have

$$P(t, T) = \frac{1}{1 + (T-t)L(t, t, T)},$$

and similarly

$$P(t, S) = \frac{1}{1 + (S-t)L(t, t, S)}.$$

Hence we get

$$\begin{aligned} L(t, T, S) &= \frac{1}{S-T} \left(\frac{P(t, T)}{P(t, S)} - 1 \right) \\ &= \frac{1}{S-T} \left(\frac{1 + (S-t)L(t, t, S)}{1 + (T-t)L(t, t, T)} - 1 \right) \\ &= \frac{1}{S-T} \left(\frac{(S-t)L(t, t, S) - (T-t)L(t, t, T)}{1 + (T-t)L(t, t, T)} \right). \end{aligned}$$

When $T = \text{one year}$ and $L(0, 0, T) = 2\%$, $L(0, 0, 2T) = 2.5\%$ we find

$$L(t, T, S) = \frac{1}{T} \left(\frac{2TL(0, 0, 2T) - TL(0, 0, T)}{1 + TL(0, 0, T)} \right) = \frac{2 \times 0.025 - 0.02}{1 + 0.02} = 2.94\%,$$

so we would not prefer a spot rate at $L(T, T, 2T) = 2\%$ over a forward contract with rate $L(0, T, 2T) = 2.94\%$.

Exercise 2.6 (Exercise 1.4 continued).

a) By Definition 2.1, we have

$$\begin{aligned} f(t, T, S) &= \frac{1}{S-T} (\log P(t, T) - \log P(t, S)) \\ &= \frac{1}{S-T} \left(\left(-(T-t)r_t + \frac{\sigma^2}{6}(T-t)^3 \right) - \left(-(S-t)r_t + \frac{\sigma^2}{6}(S-t)^3 \right) \right) \\ &= r_t + \frac{1}{S-T} \frac{\sigma^2}{6} ((T-t)^3 - (S-t)^3). \end{aligned}$$

b) We have

$$f(t, T) = -\frac{\partial}{\partial T} \log P(t, T) = r_t - \frac{\sigma^2}{2}(T-t)^2.$$

c) We have

$$d_t f(t, T) = (T-t)\sigma^2 dt + \sigma dB_t.$$

d) The HJM condition (2.3.5) is satisfied since the drift of $d_t f(t, T)$ equals $\sigma \int_t^T \sigma ds$.

Exercise 2.7

a) We have

$$X_t = X_0 e^{-bt} + \sigma \int_0^t e^{-(t-s)b} dB_s^{(1)}$$

and

$$Y_t = Y_0 e^{-bt} + \sigma \int_0^t e^{-(t-s)b} dB_s^{(2)}, \quad t \in \mathbb{R}_+,$$

see (1.1.2).

b) We have

$$\text{Var}[X_t] = \text{Var}[Y_t] = \frac{\sigma^2}{2b} (1 - e^{-2bt}), \quad t \in \mathbb{R}_+,$$

see page 2, and therefore

$$\begin{aligned} \text{Cov}(X_t, Y_t) &= \text{Cov} \left(X_0 e^{-bt} + \sigma \int_0^t e^{-(t-s)b} dB_s^{(1)}, Y_0 e^{-bt} + \sigma \int_0^t e^{-(t-s)b} dB_s^{(2)} \right) \\ &= \sigma^2 \text{Cov} \left(\int_0^t e^{-(t-s)b} dB_s^{(1)}, \int_0^t e^{-(t-s)b} dB_s^{(2)} \right) \end{aligned}$$



$$\begin{aligned}
&= \sigma^2 \mathbb{E} \left[\int_0^t e^{-(t-s)b} dB_s^{(1)} \int_0^t e^{-(t-s)b} dB_s^{(2)} \right] \\
&= \rho \sigma^2 \int_0^t e^{-2(t-s)b} ds \\
&= \rho \frac{\sigma^2}{2b} (1 - e^{-2bt}), \quad t \in \mathbb{R}_+.
\end{aligned}$$

c) We have

$$\begin{aligned}
&\text{Cov}(\log P(t, T_1), \log P(t, T_2)) \\
&= \text{Cov}(\log(F_1(t, X_t, T_1) F_2(t, Y_t, T_2) e^{\rho U(t, T_1)}), \log(F_1(t, X_t, T_1) F_2(t, Y_t, T_2) e^{\rho U(t, T_2)})) \\
&= \text{Cov}(C_1^{T_1} + X_t A_1^{T_1} + C_2^{T_1} + Y_t A_2^{T_1}, C_1^{T_2} + X_t A_1^{T_2} + C_2^{T_2} + Y_t A_2^{T_2}) \\
&= \text{Cov}(X_t A_1^{T_1} + Y_t A_2^{T_1}, X_t A_1^{T_2} + Y_t A_2^{T_2}) \\
&= A_1^{T_1} A_1^{T_2} \text{Var}[X_t] + A_2^{T_1} A_2^{T_2} \text{Var}[Y_t] + (A_1^{T_1} A_2^{T_2} + A_1^{T_2} A_2^{T_1}) \text{Cov}(X_t, Y_t) \\
&= (A_1^{T_1} A_1^{T_2} + A_2^{T_1} A_2^{T_2} + \rho(A_1^{T_1} A_2^{T_2} + A_1^{T_2} A_2^{T_1})) \text{Var}[X_t].
\end{aligned}$$

When $\text{Cov}(X_t, Y_t) = \rho \text{Var}[X_t] = \rho \text{Var}[Y_t]$, we find the correlation

$$\begin{aligned}
\text{Cov}(\log P(t, T_1), \log P(t, T_2)) &= \frac{\text{Cov}(\log P(t, T_1), \log P(t, T_2))}{\sqrt{\text{Var}[\log P(t, T_1)] \text{Var}[\log P(t, T_2)]}} \\
&= \frac{A_1^{T_1} A_1^{T_2} + A_2^{T_1} A_2^{T_2} + \rho(A_1^{T_1} A_2^{T_2} + A_1^{T_2} A_2^{T_1})}{\sqrt{(A_1^{T_1})^2 + (A_2^{T_1})^2 + \rho(A_1^{T_1} A_2^{T_1} + A_1^{T_2} A_2^{T_1})}} \\
&\quad \times \frac{1}{\sqrt{(A_1^{T_2})^2 + (A_2^{T_2})^2 + \rho(A_1^{T_2} A_2^{T_2} + A_1^{T_1} A_2^{T_1})}}.
\end{aligned}$$

When $\rho = 1$, we find

$$\begin{aligned}
&\text{Cov}(\log P(t, T_1), \log P(t, T_2)) \\
&= \frac{A_1^{T_1} A_1^{T_2} + A_2^{T_1} A_2^{T_2} + A_1^{T_1} A_2^{T_2} + A_1^{T_2} A_2^{T_1}}{\sqrt{(A_1^{T_1})^2 + (A_2^{T_1})^2 + A_1^{T_1} A_2^{T_1} + A_1^{T_2} A_2^{T_1}} \sqrt{(A_1^{T_2})^2 + (A_2^{T_2})^2 + A_1^{T_2} A_2^{T_2} + A_1^{T_1} A_2^{T_2}}} \\
&= \frac{(A_1^{T_1} + A_2^{T_1})(A_1^{T_2} + A_2^{T_2})}{|A_1^{T_1} + A_2^{T_1}| |A_1^{T_2} + A_2^{T_2}|} \\
&= \pm 1.
\end{aligned}$$

For example, if $A_1^{T_1} = 4$, $A_1^{T_2} = 1$, $A_2^{T_1} = 1$ and $A_2^{T_2} = 4$, we find

$$\text{Cov}(\log P(t, T_1), \log P(t, T_2)) = \frac{8 + 17\rho}{17 + 8\rho}.$$




Figure S.1: Log bond prices correlation graph in the two-factor model.

Exercise 2.8

a) We have

$$f(t, x) = f(0, x) + \alpha \int_0^t s^2 ds + \sigma \int_0^t d_s B(s, x) = r + \alpha t x^2 + \sigma B(t, x).$$

b) We have

$$r_t = f(t, 0) = r + B(t, 0) = r.$$

c) We have

$$\begin{aligned} P(t, T) &= \exp\left(-\int_t^T f(t, s) ds\right) \\ &= \exp\left(-(T-t)r - \alpha t \int_0^{T-t} s^2 ds - \sigma \int_0^{T-t} B(t, x) dx\right) \\ &= \exp\left(-(T-t)r - \frac{\alpha}{3}t(T-t)^3 - \sigma \int_0^{T-t} B(t, x) dx\right), \quad t \in [0, T]. \end{aligned}$$

d) Using (2.6.3), we have

$$\begin{aligned} \mathbb{E}\left[\left(\int_0^{T-t} B(t, x) dx\right)^2\right] &= \int_0^{T-t} \int_0^{T-t} \mathbb{E}[B(t, x)B(t, y)] dx dy \\ &= t \int_0^{T-t} \int_0^{T-t} \min(x, y) dx dy \\ &= 2t \int_0^{T-t} \int_0^y x dx dy = \frac{1}{3}t(T-t)^3. \end{aligned}$$

e) By Question (d)) we have

$$\begin{aligned} \mathbb{E}[P(t, T)] &= \mathbb{E}\left[\exp\left(-(T-t)r - \frac{\alpha}{3}t(T-t)^3 - \sigma \int_0^{T-t} B(t, x) dx\right)\right] \\ &= \exp\left(-(T-t)r - \frac{\alpha}{3}t(T-t)^3\right) \mathbb{E}\left[\exp\left(-\sigma \int_0^{T-t} B(t, x) dx\right)\right] \\ &= \exp\left(-(T-t)r - \frac{\alpha}{3}t(T-t)^3\right) \exp\left(\frac{\sigma^2}{2} \text{Var}\left[\int_0^{T-t} B(t, x) dx\right]\right) \\ &= \exp\left(-(T-t)r - \frac{\alpha}{3}t(T-t)^3 + \frac{\sigma^2}{6}t(T-t)^3\right), \quad 0 \leq t \leq T. \end{aligned}$$

f) By Question (e)) we check that the required relation is satisfied if

$$-\frac{\alpha}{3}t(T-t)^3 + \frac{\sigma^2}{6}t(T-t)^3 = 0,$$

i.e. $\alpha = \sigma^2/2$.

Remark: In order to derive an analog of the HJM absence of arbitrage condition in this stochastic string model, one would have to check whether the discounted bond price $e^{-rt}P(t, T)$ can be a martingale by doing stochastic calculus with respect to the Brownian sheet $B(t, x)$.

g) We have

$$\begin{aligned} & \mathbf{E} \left[\exp \left(- \int_0^T r_s ds \right) (P(T, S) - K)^+ \right] \\ &= e^{-rT} \mathbf{E} \left[\left(\exp \left(-(S-T)r - \frac{\alpha}{3}T(S-T)^3 + \sigma \int_0^{S-T} B(T, x) dx \right) - K \right)^+ \right] \\ &= e^{-rT} \mathbf{E} [(x e^{m+X} - K)^+], \end{aligned}$$

where $x = e^{-(S-T)r}$, $m = -\alpha T(S-T)^3/3$, and

$$X = \sigma \int_0^{S-T} B(T, x) dx \simeq \mathcal{N}(0, \sigma^2 t (T-t)^3 / 3).$$

Given the relation $\alpha = \sigma^2/2$, this yields

$$\begin{aligned} & \mathbf{E} \left[\exp \left(- \int_0^T r_s ds \right) (P(T, S) - K)^+ \right] \\ &= e^{-rS} \Phi \left(\sigma \sqrt{T(S-T)^3/12} + \frac{\log(e^{-(S-T)r}/K)}{\sigma \sqrt{T(S-T)^3/3}} \right) \\ &\quad - K e^{-rT} \Phi \left(-\sigma \sqrt{T(S-T)^3/12} + \frac{\log(e^{-(S-T)r}/K)}{\sigma \sqrt{T(S-T)^3/3}} \right) \\ &= P(0, S) \Phi \left(\sigma \sqrt{T(S-T)^3/12} + \frac{\log(e^{-(S-T)r}/K)}{\sigma \sqrt{T(S-T)^3/3}} \right) \\ &\quad - K P(0, T) \Phi \left(-\sigma \sqrt{T(S-T)^3/12} + \frac{\log(e^{-(S-T)r}/K)}{\sigma \sqrt{T(S-T)^3/3}} \right). \end{aligned}$$

Chapter 3

Exercise 3.1

a) We have

$$\begin{aligned} d\widehat{X}_t &= d \left(\frac{X_t}{N_t} \right) \\ &= \frac{X_0}{N_0} d \left(e^{(\sigma-\eta)B_t - (\sigma^2 - \eta^2)t/2} \right) \\ &= \frac{X_0}{N_0} (\sigma - \eta) e^{(\sigma-\eta)B_t - (\sigma^2 - \eta^2)t/2} dB_t \\ &\quad + \frac{X_0}{2N_0} (\sigma - \eta)^2 e^{(\sigma-\eta)B_t - (\sigma^2 - \eta^2)t/2} dt \\ &\quad - \frac{X_0}{2N_0} (\sigma^2 - \eta^2) e^{(\sigma-\eta)B_t - (\sigma^2 - \eta^2)t/2} dt \\ &= -\frac{X_t}{2N_t} (\sigma^2 - \eta^2) dt + \frac{X_t}{N_t} (\sigma - \eta) dB_t + \frac{X_t}{2N_t} (\sigma - \eta)^2 dt \\ &= -\frac{X_t}{N_t} \eta (\sigma - \eta) dt + \frac{X_t}{N_t} (\sigma - \eta) dB_t \end{aligned}$$

$$\begin{aligned}
&= \frac{X_t}{N_t}(\sigma - \eta)(dB_t - \eta dt) \\
&= (\sigma - \eta)\frac{X_t}{N_t}d\hat{B}_t = (\sigma - \eta)\hat{X}_t d\hat{B}_t,
\end{aligned}$$

where $d\hat{B}_t = dB_t - \eta dt$ is a standard Brownian motion under $\hat{\mathbb{P}}$.

b) By change of numéraire, we have

$$\mathbf{E}[(X_T - \lambda N_T)^+] = \hat{\mathbf{E}}\left[\frac{N_0}{N_T}(X_T - \lambda N_T)^+\right] = N_0 \hat{\mathbf{E}}[(\hat{X}_T - \lambda)^+].$$

Next, by the result of Question (a)), \hat{X}_t is a driftless geometric Brownian motion with volatility $\sigma - \eta$ under $\hat{\mathbb{P}}$, hence we have

$$\hat{\mathbf{E}}[(\hat{X}_T - \lambda)^+] = \hat{X}_0 \Phi\left(\frac{\log(\hat{X}_0/\lambda)}{\hat{\sigma}\sqrt{T}} + \frac{\hat{\sigma}\sqrt{T}}{2}\right) - \lambda \Phi\left(\frac{\log(\hat{X}_0/\lambda)}{\hat{\sigma}\sqrt{T}} - \frac{\hat{\sigma}\sqrt{T}}{2}\right),$$

by the Black-Scholes formula with zero interest rate and volatility parameter $\hat{\sigma} = \sigma - \eta$. By multiplication by N_0 and the relation $X_0 = N_0 \hat{X}_0$ we conclude to (3.5.5), i.e.

$$\begin{aligned}
\mathbf{E}[(X_T - \lambda N_T)^+] &= N_0 \hat{\mathbf{E}}[(\hat{X}_T - \lambda)^+] \\
&= N_0 \hat{X}_0 \Phi(d_+) - \lambda N_0 \Phi(d^-) \\
&= X_0 \Phi(d_+) - \lambda N_0 \Phi(d^-).
\end{aligned}$$

c) We have $\hat{\sigma} = \sigma - \eta$.

Exercise 3.2

a) By the Girsanov Theorem 3.7, the processes

$$d\hat{B}_t^{(1)} = dB_t^{(1)} - \frac{1}{N_t}dN_t \cdot dB_t^{(1)} = dB_t^{(1)} - \frac{1}{S_t^{(2)}}dS_t^{(2)} \cdot dB_t^{(1)} = dB_t^{(1)} - \eta\rho dt,$$

and

$$d\hat{B}_t^{(2)} = dB_t^{(2)} - \frac{1}{N_t}dN_t \cdot dB_t^{(2)} = dB_t^{(2)} - \frac{1}{S_t^{(2)}}dS_t^{(2)} \cdot dB_t^{(2)} = dB_t^{(2)} - \eta dt$$

are both standard Brownian motions (and martingales) under $\hat{\mathbb{P}}_2$.

b) We have

$$\begin{aligned}
d\hat{S}_t^{(1)} &= d\left(\frac{S_t^{(1)}}{S_t^{(2)}}\right) \\
&= \frac{S_0^{(1)}}{S_0^{(2)}}d(e^{\sigma B_t^{(1)} - \eta B_t^{(2)} - (\sigma^2 - \eta^2)t/2}) \\
&= \frac{S_0^{(1)}}{S_0^{(2)}}e^{\sigma B_t^{(1)} - \eta B_t^{(2)} - (\sigma^2 - \eta^2)t/2} \\
&\quad \times \left(\sigma dB_t^{(1)} + \frac{\sigma^2}{2}dt - \eta dB_t^{(2)} + \frac{\eta^2}{2}dt - \frac{\sigma^2 - \eta^2}{2}dt - \sigma\eta\rho dt\right) \\
&= \hat{S}_t^{(1)}(\sigma d\hat{B}_t^{(1)} - \eta d\hat{B}_t^{(2)}).
\end{aligned}$$

c) We note that the driftless geometric Brownian motion $(\hat{S}_t^{(1)})_{t \in \mathbb{R}}$ can be written as

$$d\hat{S}_t^{(1)} = \hat{\sigma} \hat{S}_t^{(1)} d\hat{W}_t,$$



where $(\widehat{W}_t)_{t \in \mathbb{R}}$ is a standard Brownian motion under $\widehat{\mathbb{P}}_2$. In order to determine $\widehat{\sigma}$ we note that

$$\begin{aligned}\widehat{\sigma}^2 dt &= \widehat{\sigma}^2 d\widehat{W}_t \cdot d\widehat{W}_t \\ &= \frac{d\widehat{S}_t^{(1)}}{\widehat{S}_t^{(1)}} \cdot \frac{d\widehat{S}_t^{(1)}}{\widehat{S}_t^{(1)}} \\ &= (\sigma dB_t^{(1)} - \eta d\widehat{B}_t^{(2)}) \cdot (\sigma dB_t^{(1)} - \eta d\widehat{B}_t^{(2)}) \\ &= (\sigma^2 + \eta^2 - 2\sigma\eta\rho)dt,\end{aligned}$$

hence $\widehat{\sigma}^2 = \sigma^2 + \eta^2 - 2\sigma\eta\rho$. We conclude by applying the change of numéraire formula

$$e^{-rT} \mathbf{E}^* [(S_T^{(1)} - \lambda S_T^{(2)})^+] = S_0^{(2)} \widehat{\mathbf{E}}[(\widehat{S}_T^{(1)} - \lambda)^+]$$

and the Black-Scholes formula to the the driftless geometric Brownian motion $(\widehat{S}_t^{(1)})_{t \in \mathbb{R}}$.

Exercise 3.3 We have $N_t = P(t, T)$ and from (1.4.9) and the relations $P(t, T) = F(t, r_t)$ and $P(t, S) = G(t, r_t)$ we find

$$\left\{ \begin{array}{l} \frac{dP(t, S)}{P(t, S)} = r_t dt + \sigma(t, r_t) \frac{\partial}{\partial x} \log G(t, r_t) dW_t, \\ \frac{dN_t}{N_t} = \frac{dP(t, T)}{P(t, T)} = r_t dt + \sigma(t, r_t) \frac{\partial}{\partial x} \log F(t, r_t) dW_t. \end{array} \right.$$

By the Girsanov Theorem (3.2.12) we also have

$$d\widehat{W}_t = dW_t - \frac{dN_t}{N_t} \cdot dW_t = dW_t - \sigma(t, r_t) \frac{\partial}{\partial x} \log F(t, r_t) dt,$$

hence

$$\frac{dP(t, S)}{P(t, S)} = r_t dt + \sigma^2(t, r_t) \frac{\partial}{\partial x} \log F(t, r_t) \frac{\partial}{\partial x} \log G(t, r_t) dt + \sigma(t, r_t) \frac{\partial}{\partial x} \log G(t, r_t) d\widehat{W}_t.$$

Using the relation $P(t, S) = G(t, r_t)$ we can also write

$$dP(t, S) = r_t P(t, S) dt + \sigma^2(t, r_t) \frac{\partial}{\partial x} \log F(t, r_t) \frac{\partial}{\partial x} G(t, r_t) dt + \sigma(t, r_t) \frac{\partial}{\partial x} G(t, r_t) d\widehat{W}_t.$$

Exercise 3.4 Forward contract. Taking $N_t := P(t, T)$, $t \in [0, T]$, we have

$$\begin{aligned}\mathbf{E}^* \left[\exp \left(- \int_t^T r_s ds \right) (P(T, S) - K) \mid \mathcal{F}_t \right] &= N_t \widehat{\mathbf{E}} \left[\frac{(P(T, S) - K)}{N_T} \mid \mathcal{F}_t \right] \\ &= P(t, T) \widehat{\mathbf{E}} \left[\frac{(P(T, S) - K)}{P(T, T)} \mid \mathcal{F}_t \right] \\ &= P(t, T) \widehat{\mathbf{E}}[P(T, S) - K \mid \mathcal{F}_t] \\ &= P(t, T) \widehat{\mathbf{E}}[P(T, S) \mid \mathcal{F}_t] - KP(t, T) \\ &= P(t, T) \widehat{\mathbf{E}} \left[\frac{P(T, S)}{P(T, T)} \mid \mathcal{F}_t \right] - KP(t, T) \\ &= P(t, T) \frac{P(t, S)}{P(t, T)} - KP(t, T)\end{aligned}$$

$$= P(t, S) - K P(t, T),$$

since

$$t \mapsto \frac{P(t, T)}{N_t} = \frac{P(t, S)}{P(t, T)}$$

is a martingale under the forward measure $\widehat{\mathbb{P}}$. The corresponding (static) hedging strategy is given by buying one bond with maturity S and by short selling K units of the bond with maturity T .

Remark: The above result can also be obtained by a direct argument using the tower property of conditional expectations:

$$\begin{aligned} & \mathbf{E}^* \left[\exp \left(- \int_t^T r_s ds \right) (P(T, S) - K) \middle| \mathcal{F}_t \right] \\ &= \mathbf{E}^* \left[\exp \left(- \int_t^T r_s ds \right) \left(\mathbf{E}^* \left[\exp \left(- \int_T^S r_s ds \right) \middle| \mathcal{F}_T \right] - K \right) \middle| \mathcal{F}_t \right] \\ &= \mathbf{E}^* \left[\exp \left(- \int_t^T r_s ds \right) \mathbf{E}^* \left[\exp \left(- \int_T^S r_s ds \right) - K \middle| \mathcal{F}_T \right] \middle| \mathcal{F}_t \right] \\ &= \mathbf{E}^* \left[\mathbf{E}^* \left[\exp \left(- \int_t^S r_s ds \right) - K \exp \left(- \int_t^T r_s ds \right) \middle| \mathcal{F}_T \right] \middle| \mathcal{F}_t \right] \\ &= \mathbf{E}^* \left[\exp \left(- \int_t^S r_s ds \right) - K \exp \left(- \int_t^T r_s ds \right) \middle| \mathcal{F}_t \right] \\ &= P(t, S) - K P(t, T), \quad t \in [0, T]. \end{aligned}$$

Exercise 3.5

- a) We choose $N_t := S_t$ as numéraire because this allows us to write the option payoff as $(S_T(S_T - K))^+ = N_T(S_T - K)^+$. In this case, the forward measure $\widehat{\mathbb{P}}$ satisfies

$$\frac{d\widehat{\mathbb{P}}}{d\mathbb{P}} = e^{-rT} \frac{N_T}{N_0} = e^{-rT} \frac{S_T}{S_0},$$

or

$$\frac{d\widehat{\mathbb{P}}|_{\mathcal{F}_t}}{d\mathbb{P}|_{\mathcal{F}_t}} = e^{-(T-t)r} \frac{N_T}{N_t} = e^{-(T-t)r} \frac{S_T}{S_t}, \quad 0 \leq t \leq T.$$

- b) By the change of numéraire formula of Proposition 3.5, the option price becomes

$$\begin{aligned} e^{-(T-t)r} \mathbf{E}^* [(S_T(S_T - K))^+ | \mathcal{F}_t] &= \mathbf{E}^* [e^{-(T-t)r} N_T(S_T - K)^+ | \mathcal{F}_t] \\ &= N_t \widehat{\mathbf{E}}[(S_T - K)^+ | \mathcal{F}_t] \\ &= S_t \widehat{\mathbf{E}}[(S_T - K)^+ | \mathcal{F}_t]. \end{aligned} \tag{A.11}$$

- c) In order to compute (A.11) it remains to determine the dynamics of $(S_t)_{t \in \mathbb{R}_+}$ under $\widehat{\mathbb{P}}$. Since $S_t = S_0 e^{\sigma B_t + rt - \sigma^2 t / 2}$, we have

$$\frac{d\widehat{\mathbb{P}}}{d\mathbb{P}} = e^{-rT} \frac{S_T}{S_0} = e^{\sigma B_T - \sigma^2 T / 2},$$

hence by the Girsanov Theorem, $\widehat{B}_t := B_t - \sigma t$ is a standard Brownian motion under $\widehat{\mathbb{P}}$, with

$$\begin{aligned} S_T &= S_0 e^{rT + \sigma B_T - \sigma^2 T / 2} \\ &= S_0 e^{(r + \sigma^2)T + \sigma \widehat{B}_T - \sigma^2 T / 2} \\ &= S_t e^{(r + \sigma^2)(T-t) + (\widehat{B}_T - \widehat{B}_t)\sigma - (T-t)\sigma^2 / 2}, \quad 0 \leq t \leq T. \end{aligned}$$



d) According to the above, (A.11) becomes

$$\begin{aligned}
 & e^{-(T-t)r} \mathbf{E}^* [(S_T(S_T - K))^+ | \mathcal{F}_t] = S_t \widehat{\mathbf{E}}[(S_T - K)^+ | \mathcal{F}_t] \\
 &= S_t \widehat{\mathbf{E}}[(S_0 e^{rT + \sigma^2 T + \sigma \widehat{B}_T - \sigma^2 T/2} - K)^+ | \mathcal{F}_t] \\
 &= S_t \widehat{\mathbf{E}}[(S_t e^{(r+\sigma^2)(T-t) + (\widehat{B}_T - \widehat{B}_t)\sigma - (T-t)\sigma^2/2} - K)^+ | \mathcal{F}_t] \\
 &= S_t e^{(T-t)(r+\sigma^2)} \text{Bl}(S_t, r + \sigma^2, \sigma, K, T - t), \quad 0 \leq t \leq T,
 \end{aligned}$$

since the Black-Scholes formula with interest rate $r + \sigma^2$ reads

$$\begin{aligned}
 & e^{-(T-t)(r+\sigma^2)} \widehat{\mathbf{E}}[(S_t e^{(r+\sigma^2)(T-t) + (\widehat{B}_T - \widehat{B}_t)\sigma - (T-t)\sigma^2/2} - K)^+ | \mathcal{F}_t] \\
 &= \text{Bl}(S_t, r + \sigma^2, \sigma, K, T - t), \quad 0 \leq t \leq T.
 \end{aligned}$$

Remarks:

- i) The option price can be rewritten using other Black-Scholes parametrizations, such as for example

$$S_t \text{Bl}(S_t e^{(T-t)(r+\sigma^2)}, 0, \sigma, K, T - t),$$

or

$$S_t e^{(T-t)(r+\sigma^2)} \text{Bl}(S_t, 0, \sigma, K e^{-(T-t)(r+\sigma^2)}, T - t),$$

however we prefer to choose the simplest possibility.

- ii) Deflated (or forward) processes such as $S_t/N_1 = 1$ or

$$\frac{e^{-(T-t)r}}{N_t} \mathbf{E}^* [(S_T(S_T - K))^+ | \mathcal{F}_t] = \widehat{\mathbf{E}}[(S_T - K)^+ | \mathcal{F}_t], \quad 0 \leq t \leq T,$$

are martingales under the forward measure $\widehat{\mathbb{P}}$.

- iii) This option can also be priced via an integral calculation instead of using change of numéraire, as follows:

$$\begin{aligned}
 & e^{-(T-t)r} \mathbf{E}^*[S_T(S_T - K)^+ | \mathcal{F}_t] \\
 &= e^{-(T-t)r} \mathbf{E}^*[S_t e^{(T-t)r + (B_T - B_t)\sigma - (T-t)\sigma^2/2} \\
 &\quad \times (S_t e^{(T-t)r + (B_T - B_t)\sigma - (T-t)\sigma^2/2} - K)^+ | \mathcal{F}_t] \\
 &= S_t e^{-(T-t)\sigma^2/2} \mathbf{E}^* [(S_t e^{(T-t)r + 2(B_T - B_t)\sigma - (T-t)\sigma^2/2} - K e^{(B_T - B_t)\sigma})^+ | \mathcal{F}_t] \\
 &= S_t e^{-(T-t)\sigma^2/2} \mathbf{E}^* [(x e^{(T-t)r + 2(B_T - B_t)\sigma - (T-t)\sigma^2/2} - K e^{(B_T - B_t)\sigma})^+]_{x=S_t} \\
 &= S_t e^{-(T-t)\sigma^2/2} \mathbf{E}^* [(e^{m(x)+2X} - K e^X)^+]_{x=S_t}, \quad 0 \leq t \leq T,
 \end{aligned}$$

where $X \simeq \mathcal{N}(0, v^2)$ with $v^2 = (T-t)\sigma^2$ and $m(x) = (T-t)r - (T-t)\sigma^2/2 + \log x$. Next, we note that

$$\begin{aligned}
 \mathbf{E}[(e^{m+2X} - K e^X)^+] &= \frac{1}{\sqrt{2\pi v^2}} \int_{-\infty}^{\infty} (e^{m+2x} - K e^x)^+ e^{-x^2/(2v^2)} dx \\
 &= \frac{1}{\sqrt{2\pi v^2}} \int_{-m+\log K}^{\infty} (e^{m+2x} - K e^x) e^{-x^2/(2v^2)} dx \\
 &= \frac{e^m}{\sqrt{2\pi v^2}} \int_{-m+\log K}^{\infty} e^{2x-x^2/(2v^2)} dx - \frac{K}{\sqrt{2\pi v^2}} \int_{-m+\log K}^{\infty} e^{x-x^2/(2v^2)} dx \\
 &= \frac{e^{m+2v^2}}{\sqrt{2\pi v^2}} \int_{-m+\log K}^{\infty} e^{-(2v^2-x)^2/(2v^2)} dx - \frac{K e^{v^2/2}}{\sqrt{2\pi}} \int_{-m+\log K}^{\infty} e^{-(v^2-x)^2/(2v^2)} dx \\
 &= \frac{e^{m+2v^2}}{\sqrt{2\pi v^2}} \int_{-2v^2-m+\log K}^{\infty} e^{-x^2/(2v^2)} dx - \frac{K e^{v^2/2}}{\sqrt{2\pi v^2}} \int_{-v^2-m+\log K}^{\infty} e^{-x^2/(2v^2)} dx \\
 &= \frac{e^{m+2v^2}}{\sqrt{2\pi}} \int_{(-2v^2-m+\log K)/v}^{\infty} e^{-x^2/2} dx - \frac{K e^{v^2/2}}{\sqrt{2\pi}} \int_{(-v^2-m+\log K)/v}^{\infty} e^{-x^2/2} dx
 \end{aligned}$$

$$= e^{m+2v^2} \Phi\left(2v + \frac{m - \log K}{v}\right) - K e^{v^2/2} \Phi\left(v + \frac{m - \log K}{v}\right),$$

hence

$$\begin{aligned} & e^{-(T-t)r} \mathbf{E}^* [S_T (S_T - K)^+ | \mathcal{F}_t] \\ &= S_t e^{-(T-t)\sigma^2/2} \mathbf{E}^* [(e^{m(x)+2X} - K e^X)^+]_{x=S_t} \\ &= S_t^2 e^{(T-t)(r+\sigma^2)} \Phi\left(\frac{(T-t)(r+\sigma^2) + (T-t)\sigma^2/2 + \log(S_t/K)}{\sigma\sqrt{T-t}}\right) \\ &\quad - K S_t \Phi\left(\frac{(T-t)(r+\sigma^2) - (T-t)\sigma^2/2 + \log(S_t/K)}{\sigma\sqrt{T-t}}\right) \\ &= S_t e^{(T-t)(r+\sigma^2)} \text{Bl}(S_t, r+\sigma^2, \sigma, K, T-t), \quad 0 \leq t \leq T. \end{aligned}$$

Exercise 3.6

- a) Knowing that $S_t = S_0 e^{\sigma W_t + rt - \sigma^2 t/2}$, we check that the discounted numeraire process

$$\begin{aligned} e^{-rt} N_t &:= S_t^n e^{-(n-1)n\sigma^2 t/2 - nrt} \\ &= S_0^n e^{n\sigma W_t + nrt - n\sigma^2 t/2} e^{-(n-1)n\sigma^2 t/2 - nrt} \\ &= S_0 e^{n\sigma W_t - (n\sigma)^2 t/2}, \quad 0 \leq t \leq T, \end{aligned}$$

is a martingale under \mathbb{P}^* , and

$$\frac{d\hat{\mathbb{P}}}{d\mathbb{P}^*} = e^{-rT} \frac{N_T}{N_0} = \frac{S_T^n}{S_0^n} e^{-n\sigma^2 T - nrT} = e^{n\sigma W_T - (n\sigma)^2 T/2}. \quad (\text{A.12})$$

- b) From Equation (A.12) and the Girsanov theorem, the process $\hat{W}_t := W_t - \sigma nt$ is a standard Brownian motion under $\hat{\mathbb{P}}$, and we have

$$\begin{aligned} S_t &= S_0 e^{rt + \sigma W_t - \sigma^2 t/2} \\ &= S_0 e^{rt + \sigma(\hat{W}_t + n\sigma t) - \sigma^2 t/2} \\ &= S_0 e^{(r+n\sigma^2)t + \sigma \hat{W}_t + \sigma^2 t/2}, \quad t \geq 0. \end{aligned}$$

- c) We have

$$\begin{aligned} & e^{-(T-t)r} \mathbf{E}^* [S_T^n \mathbb{1}_{\{S_T \geq K\}} | \mathcal{F}_t] = \mathbf{E}^* [e^{-(T-t)r} S_T^n \mathbb{1}_{\{S_T \geq K\}} | \mathcal{F}_t] \\ &= e^{(n-1)n\sigma^2 T/2 + (n-1)rT} \mathbf{E}^* [N_T \mathbb{1}_{\{S_T \geq K\}} | \mathcal{F}_t] \\ &= e^{(n-1)n\sigma^2 T/2 + (n-1)rT} N_t \hat{\mathbb{E}} [\mathbb{1}_{\{S_T \geq K\}} | \mathcal{F}_t] \\ &= e^{(n-1)n\sigma^2 T/2 + (n-1)rT} N_t \hat{\mathbb{P}}(S_T \geq K | \mathcal{F}_t) \\ &= e^{(n-1)n\sigma^2 T/2 + (n-1)rT} N_t \Phi\left(\frac{\log(S_t/K) + (r + (n-1/2)\sigma^2)(T-t)}{\sigma\sqrt{T-t}}\right) \\ &= S_t^n e^{(n-1)n\sigma^2(T-t)/2 + (n-1)r(T-t)} \Phi\left(\frac{\log(S_t/K) + (r + (n-1/2)\sigma^2)(T-t)}{\sigma\sqrt{T-t}}\right), \end{aligned}$$

$0 \leq t \leq T$, $n \geq 0$. Hence the power call option with payoff $(S_T^n - K^n)^+$ is priced at time $t \in [0, T]$ as

$$\begin{aligned} & e^{-(T-t)r} \mathbf{E}^* [(S_T^n - K^n)^+ | \mathcal{F}_t] \\ &= S_t^n e^{(n-1)n\sigma^2(T-t)/2 + (n-1)r(T-t)} \Phi\left(\frac{\log(S_t/K) + (r + (n-1/2)\sigma^2)(T-t)}{\sigma\sqrt{T-t}}\right) \\ &\quad - K^n e^{-(T-t)r} \Phi\left(\frac{\log(S_t/K) + (r - \sigma^2/2)(T-t)}{\sigma\sqrt{T-t}}\right), \quad n \geq 1. \end{aligned}$$



Exercise 3.7 Bond options.

a) Itô's formula yields

$$\begin{aligned} d\left(\frac{P(t,S)}{P(t,T)}\right) &= \frac{P(t,S)}{P(t,T)}(\zeta^S(t) - \zeta^T(t))(dW_t - \zeta^T(t)dt) \\ &= \frac{P(t,S)}{P(t,T)}(\zeta^S(t) - \zeta^T(t))d\hat{W}_t, \end{aligned} \quad (\text{A.13})$$

where $(\hat{W}_t)_{t \in \mathbb{R}_+}$ is a standard Brownian motion under $\hat{\mathbb{P}}$ by the Girsanov Theorem.

b) From (A.13) or (4.1.7) we have

$$\frac{P(t,S)}{P(t,T)} = \frac{P(0,S)}{P(0,T)} \exp\left(\int_0^t (\zeta^S(s) - \zeta^T(s))d\hat{W}_s - \frac{1}{2} \int_0^t |\zeta^S(s) - \zeta^T(s)|^2 ds\right),$$

hence

$$\frac{P(u,S)}{P(u,T)} = \frac{P(t,S)}{P(t,T)} \exp\left(\int_t^u (\zeta^S(s) - \zeta^T(s))d\hat{W}_s - \frac{1}{2} \int_t^u |\zeta^S(s) - \zeta^T(s)|^2 ds\right),$$

$t \in [0, u]$, and for $u = T$ this yields

$$P(T,S) = \frac{P(t,S)}{P(t,T)} \exp\left(\int_t^T (\zeta^S(s) - \zeta^T(s))d\hat{W}_s - \frac{1}{2} \int_t^T |\zeta^S(s) - \zeta^T(s)|^2 ds\right),$$

since $P(T,T) = 1$. Let $\hat{\mathbb{P}}$ denote the forward measure associated to the numéraire

$$N_t := P(t,T), \quad 0 \leq t \leq T.$$

c) For all $S \geq T > 0$ we have

$$\begin{aligned} \mathbb{E}\left[e^{-\int_t^T r_s ds}(P(T,S) - K)^+ \middle| \mathcal{F}_t\right] &= P(t,T)\hat{\mathbb{E}}\left[\left(\frac{P(t,S)}{P(t,T)} \exp\left(X - \frac{1}{2} \int_t^T |\zeta^S(s) - \zeta^T(s)|^2 ds\right) - K\right)^+ \middle| \mathcal{F}_t\right] \\ &= P(t,T)\hat{\mathbb{E}}[(e^{X+m(t,T,S)} - K)^+ \middle| \mathcal{F}_t], \end{aligned}$$

where X is a centered Gaussian random variable with variance

$$v^2(t,T,S) = \int_t^T |\zeta^S(s) - \zeta^T(s)|^2 ds$$

given \mathcal{F}_t , and

$$m(t,T,S) = -\frac{1}{2}v^2(t,T,S) + \log \frac{P(t,S)}{P(t,T)}.$$

Recall that when X is a centered Gaussian random variable with variance v^2 , the expectation of $(e^{m+X} - K)^+$ is given, as in the standard Black-Scholes formula, by

$$\mathbb{E}[(e^{m+X} - K)^+] = e^{m+v^2/2} \Phi(v + (m - \log K)/v) - K \Phi((m - \log K)/v),$$

where

$$\Phi(z) = \int_{-\infty}^z e^{-y^2/2} \frac{dy}{\sqrt{2\pi}}, \quad z \in \mathbb{R},$$

denotes the Gaussian cumulative distribution function and for simplicity of notation we dropped the indices t, T, S in $m(t,T,S)$ and $v^2(t,T,S)$.

Consequently we have

$$\begin{aligned} \mathbb{E}\left[e^{-\int_t^T r_s ds}(P(T,S) - K)^+ \middle| \mathcal{F}_t\right] &= P(t,S)\Phi\left(\frac{v}{2} + \frac{1}{v} \log \frac{P(t,S)}{KP(t,T)}\right) - KP(t,T)\Phi\left(-\frac{v}{2} + \frac{1}{v} \log \frac{P(t,S)}{KP(t,T)}\right). \end{aligned}$$

- d) The self-financing hedging strategy that hedges the bond option is obtained by holding a (possibly fractional) quantity

$$\Phi\left(\frac{\nu}{2} + \frac{1}{\nu} \log \frac{P(t, S)}{KP(t, T)}\right)$$

of the bond with maturity S , and by shorting a quantity

$$K\Phi\left(-\frac{\nu}{2} + \frac{1}{\nu} \log \frac{P(t, S)}{KP(t, T)}\right)$$

of the bond with maturity T .

Exercise 3.8

- a) The process

$$e^{-rt}S_2(t) = S_2(0)e^{\sigma_2 W_t + (\mu - r)t}$$

is a martingale if

$$r - \mu = \frac{1}{2}\sigma_2^2.$$

- b) We note that

$$\begin{aligned} e^{-rt}X_t &= e^{-rt}e^{(r-\mu)t - \sigma_1^2 t/2}S_1(t) \\ &= e^{-rt}e^{(\sigma_2^2 - \sigma_1^2)t/2}S_1(t) \\ &= e^{-\mu t - \sigma_1^2 t/2}S_1(t) \\ &= S_1(0)e^{\mu t - \sigma_1^2 t/2}e^{\sigma_1 W_t + \mu t} \\ &= S_1(0)e^{\sigma_1 W_t - \sigma_1^2 t/2} \end{aligned}$$

is a martingale, where

$$X_t = e^{(r-\mu)t - \sigma_1^2 t/2}S_1(t) = e^{(\sigma_2^2 - \sigma_1^2)t/2}S_1(t).$$

- c) By (3.5.7) we have

$$\begin{aligned} \widehat{X}(t) &= \frac{X_t}{N_t} \\ &= e^{(\sigma_2^2 - \sigma_1^2)t/2} \frac{S_1(t)}{S_2(t)} \\ &= \frac{S_1(0)}{S_2(0)} e^{(\sigma_2^2 - \sigma_1^2)t/2 + (\sigma_1 - \sigma_2)W_t} \\ &= \frac{S_1(0)}{S_2(0)} e^{(\sigma_2^2 - \sigma_1^2)t/2 + (\sigma_1 - \sigma_2)\widehat{W}_t + \sigma_2(\sigma_1 - \sigma_2)t} \\ &= \frac{S_1(0)}{S_2(0)} e^{(\sigma_1 - \sigma_2)\widehat{W}_t + \sigma_2\sigma_1 t - (\sigma_2^2 + \sigma_1^2)t/2} \\ &= \frac{S_1(0)}{S_2(0)} e^{(\sigma_1 - \sigma_2)\widehat{W}_t - (\sigma_1 - \sigma_2)^2 t/2}, \end{aligned}$$

where

$$\widehat{W}_t := W_t - \sigma_2 t$$

is a standard Brownian motion under the forward measure $\widehat{\mathbb{P}}$ defined by

$$\begin{aligned} \frac{d\widehat{\mathbb{P}}}{d\mathbb{P}} &= e^{-\int_0^T r_s ds} \frac{N_T}{N_0} \\ &= e^{-rT} \frac{S_2(T)}{S_2(0)} \end{aligned}$$



$$\begin{aligned}
&= e^{-rT} e^{\sigma_2 W_T + \mu T} \\
&= e^{\sigma_2 W_T + (\mu - r)T} \\
&= e^{\sigma_2 W_T - \sigma_2^2 t / 2}.
\end{aligned}$$

d) Given that $X_t = e^{(\sigma_2^2 - \sigma_1^2)t/2} S_1(t)$ and $\hat{X}(t) = X_t / N_t = X_t / S_2(t)$, we have

$$\begin{aligned}
e^{-rT} \mathbb{E}[(S_1(T) - \kappa S_2(T))^+] &= e^{-rT} \mathbb{E}[(e^{-(\sigma_2^2 - \sigma_1^2)t/2} X_T - \kappa S_2(T))^+] \\
&= e^{-rT} e^{-(\sigma_2^2 - \sigma_1^2)t/2} \mathbb{E}[(X_T - \kappa e^{(\sigma_2^2 - \sigma_1^2)t/2} S_2(T))^+] \\
&= S_2(0) e^{-(\sigma_2^2 - \sigma_1^2)t/2} \hat{\mathbb{E}}[(\hat{X}_T - \kappa e^{(\sigma_2^2 - \sigma_1^2)t/2})^+] \\
&= S_2(0) e^{-(\sigma_2^2 - \sigma_1^2)t/2} \hat{\mathbb{E}}[(\hat{X}_0 e^{(\sigma_1 - \sigma_2)\hat{W}_T - (\sigma_1 - \sigma_2)^2 T / 2} - \kappa e^{(\sigma_2^2 - \sigma_1^2)t/2})^+] \\
&= S_2(0) e^{-(\sigma_2^2 - \sigma_1^2)t/2} (\hat{X}_0 \Phi_+^0(T, \hat{X}_0) - \kappa e^{(\sigma_2^2 - \sigma_1^2)t/2} \Phi_-^0(T, \hat{X}_0)) \\
&= S_2(0) e^{-(\sigma_2^2 - \sigma_1^2)t/2} \hat{X}_0 \Phi_+^0(T, \hat{X}_0) \\
&\quad - \kappa S_2(0) e^{-(\sigma_2^2 - \sigma_1^2)t/2} e^{(\sigma_2^2 - \sigma_1^2)t/2} \Phi_-^0(T, \hat{X}_0) \\
&= X_0 e^{-(\sigma_2^2 - \sigma_1^2)t/2} \Phi_+^0(T, \hat{X}_0) - \kappa S_2(0) \Phi_-^0(T, \hat{X}_0) \\
&= S_1(0) e^{-(\sigma_2^2 - \sigma_1^2)t/2} \Phi_+^0(T, \hat{X}_0) - \kappa S_2(0) \Phi_-^0(T, \hat{X}_0),
\end{aligned}$$

where

$$\Phi_+^0(T, x) = \Phi \left(\frac{\log(x/\kappa)}{|\sigma_1 - \sigma_2| \sqrt{T}} + \frac{(\sigma_1 - \sigma_2)^2 - (\sigma_2^2 - \sigma_1^2)}{2|\sigma_1 - \sigma_2|} \sqrt{T} \right)$$

$$= \begin{cases} \Phi \left(\frac{\log(x/\kappa)}{|\sigma_1 - \sigma_2| \sqrt{T}} + \sigma_1 \sqrt{T} \right), & \sigma_1 > \sigma_2, \\ \Phi \left(\frac{\log(x/\kappa)}{|\sigma_1 - \sigma_2| \sqrt{T}} - \sigma_1 \sqrt{T} \right), & \sigma_1 < \sigma_2, \end{cases}$$

and

$$\Phi_-^0(T, x) = \Phi \left(\frac{\log(x/\kappa)}{|\sigma_1 - \sigma_2| \sqrt{T}} - \frac{(\sigma_1 - \sigma_2)^2 + (\sigma_2^2 - \sigma_1^2)}{2|\sigma_1 - \sigma_2|} \sqrt{T} \right)$$

$$= \begin{cases} \Phi \left(\frac{\log(x/\kappa)}{|\sigma_1 - \sigma_2| \sqrt{T}} + \sigma_2 \sqrt{T} \right), & \sigma_1 > \sigma_2, \\ \Phi \left(\frac{\log(x/\kappa)}{|\sigma_1 - \sigma_2| \sqrt{T}} - \sigma_2 \sqrt{T} \right), & \sigma_1 < \sigma_2, \end{cases}$$

if $\sigma_1 \neq \sigma_2$. In case $\sigma_1 = \sigma_2$, we find

$$\begin{aligned}
e^{-rT} \mathbb{E}[(S_1(T) - \kappa S_2(T))^+] &= e^{-rT} \mathbb{E}[S_1(T)(1 - \kappa S_2(0)/S_1(0))^+] \\
&= (1 - \kappa S_2(0)/S_1(0))^+ e^{-rT} \mathbb{E}[S_1(T)] \\
&= (S_1(0) - \kappa S_2(0)) \mathbb{1}_{\{S_1(0) > \kappa S_2(0)\}}.
\end{aligned}$$

Exercise 3.9 We have

$$\begin{aligned}
e^{-(T-t)r} \mathbb{E}^* [\mathbb{1}_{\{R_T \geq \kappa\}} | R_t] &= e^{-(T-t)r} \mathbb{P}^*(R_T \geq \kappa | R_t) \\
&= e^{-(T-t)r} \mathbb{P}^*(R_t e^{(W_T - W_t)\sigma + (r - r^f)(T-t) - (T-t)\sigma^2/2} \geq \kappa | R_t) \\
&= e^{-(T-t)r} \mathbb{P}^*(x e^{(W_T - W_t)\sigma + (r - r^f)(T-t) - (T-t)\sigma^2/2} \geq \kappa)_{x=R_t} \\
&= e^{-(T-t)r} \Phi \left(\frac{(r - r^f)(T-t) - (T-t)\sigma^2/2 - \log(\kappa/R_t)}{\sigma \sqrt{T-t}} \right),
\end{aligned}$$

after applying the hint provided, with

$$\eta^2 := (T-t)\sigma^2 \quad \text{and} \quad \mu := (r - r^f)(T-t) - (T-t)\sigma^2/2.$$

Remark: Binary options are often proposed *at the money*, i.e. $\kappa = R_t$, with a short time to maturity, for example the small value

$$T - t \simeq 30 \text{ seconds} = 0.000000951 = 9.51 \times 10^{-7} \text{ year}^{-1},$$

in which case we have

$$\begin{aligned} e^{-(T-t)r} \mathbf{E}^* [\mathbb{1}_{\{R_T \geq \kappa\}} | R_t] &= e^{-(T-t)r} \Phi \left(\left(\frac{r - r^f}{\sigma} - \frac{\sigma}{2} \right) \sqrt{T-t} \right) \\ &\simeq \Phi(0) \\ &= \frac{1}{2}. \end{aligned}$$

Taking for example $r - r^f = 0.02 = 2\%$ and $\sigma = 0.3 = 30\%$, we have

$$\left(\frac{r - r^f}{\sigma} - \frac{\sigma}{2} \right) \sqrt{T-t} = \left(\frac{0.02}{0.3} - \frac{0.3}{2} \right) \sqrt{9.51 \times 10^{-7}} = -0.000081279$$

and

$$\begin{aligned} e^{-(T-t)r} \mathbf{E}^* [\mathbb{1}_{\{R_T \geq \kappa\}} | R_t] &= e^{-(T-t)r} \Phi(-0.000081279) \\ &= e^{-r \times 0.000000951} \times 0.499968 \\ &= 0.49996801 \\ &\simeq \frac{1}{2}, \end{aligned}$$

with $r = 0.02 = 2\%$.

Exercise 3.10

a) It suffices to check that the definition of $(W_t^N)_{t \in \mathbb{R}_+}$ implies the correlation identity $dW_t^S \cdot dW_t^N = \rho dt$ by Itô's calculus.

b) We let

$$\widehat{\sigma}_t = \sqrt{(\sigma_t^S)^2 - 2\rho\sigma_t^R\sigma_t^S + (\sigma_t^R)^2}$$

and

$$dW_t^X = \frac{\sigma_t^S - \rho\sigma_t^N}{\widehat{\sigma}_t} dW_t^S - \sqrt{1 - \rho^2} \frac{\sigma_t^N}{\widehat{\sigma}_t} dW_t^R, \quad t \geq 0,$$

which defines a standard Brownian motion under \mathbb{P}^* due to the definition of $\widehat{\sigma}_t$.

Exercise 3.11

a) We have $\widehat{\sigma} = \sqrt{(\sigma^S)^2 - 2\rho\sigma^R\sigma^S + (\sigma^R)^2}$.

b) Letting $\widetilde{X}_t = e^{-rt} X_t = e^{(a-r)t} S_t / R_t$, $t \geq 0$, we have

$$\begin{aligned} \mathbf{E}^* \left[\left(\frac{S_T}{R_T} - \kappa \right)^+ \middle| \mathcal{F}_t \right] &= e^{-aT} \mathbf{E}^* \left[(X_T - e^{aT} \kappa)^+ \middle| \mathcal{F}_t \right] \\ &= e^{-(a-r)T} \mathbf{E}^* \left[\left(\widetilde{X}_T - e^{(a-r)T} \kappa \right)^+ \middle| \mathcal{F}_t \right] \\ &= e^{-(a-r)T} \left(\widetilde{X}_t \Phi \left(\frac{(r-a+\widehat{\sigma}^2/2)(T-t)}{\widehat{\sigma}\sqrt{T-t}} + \frac{1}{\widehat{\sigma}\sqrt{T-t}} \log \frac{S_t}{\kappa R_t} \right) \right. \\ &\quad \left. - \kappa e^{(a-r)T} \Phi \left(\frac{(r-a-\widehat{\sigma}^2/2)(T-t)}{\widehat{\sigma}\sqrt{T-t}} + \frac{1}{\widehat{\sigma}\sqrt{T-t}} \log \frac{S_t}{\kappa R_t} \right) \right) \end{aligned}$$



$$= \frac{S_t}{R_t} e^{(r-a)(T-t)} \Phi \left(\frac{(r-a + \widehat{\sigma}^2/2)(T-t)}{\widehat{\sigma}\sqrt{T-t}} + \frac{1}{\widehat{\sigma}\sqrt{T-t}} \log \frac{S_t}{\kappa R_t} \right) \\ - \kappa \Phi \left(\frac{(r-a - \widehat{\sigma}^2/2)(T-t)}{\widehat{\sigma}\sqrt{T-t}} + \frac{1}{\widehat{\sigma}\sqrt{T-t}} \log \frac{S_t}{\kappa R_t} \right),$$

hence the price of the quanto option is

$$e^{-(T-t)r} \mathbf{E}^* \left[\left(\frac{S_T}{R_T} - \kappa \right)^+ \middle| \mathcal{F}_t \right] \\ = \frac{S_t}{R_t} e^{-a(T-t)} \Phi \left(\frac{(r-a + \widehat{\sigma}^2/2)(T-t)}{\widehat{\sigma}\sqrt{T-t}} + \frac{1}{\widehat{\sigma}\sqrt{T-t}} \log \frac{S_t}{\kappa R_t} \right) \\ - \kappa e^{-r(T-t)} \Phi \left(\frac{(r-a - \widehat{\sigma}^2/2)(T-t)}{\widehat{\sigma}\sqrt{T-t}} + \frac{1}{\widehat{\sigma}\sqrt{T-t}} \log \frac{S_t}{\kappa R_t} \right).$$

Chapter 4

Exercise 4.1

- a) We price the floorlet at $t = 0$, with $T_1 = 9$ months, $T_2 = 1$ year, $\kappa = 4.5\%$. The LIBOR rate $(L(t, T_1, T_2))_{t \in [0, T_1]}$ is modeled as a driftless geometric Brownian motion with volatility coefficient $\widehat{\sigma} = \sigma_{1,2}(t) = 0.1$ under the forward measure \mathbb{P}_2 . The discount factors are given by

$$P(0, T_1) = e^{-9r/12} \simeq 0.970809519$$

and

$$P(0, T_2) = e^{-r} \simeq 0.961269954,$$

with $r = 3.95\%$.

- b) By (4.3.9), the price of the floorlet is

$$\mathbf{E}^* \left[e^{-\int_0^{T_2} r_s ds} (\kappa - L(T_1, T_2))^+ \right] \\ = P(0, T_2) (\kappa \Phi(-d_-(T_1)) - L(0, T_1, T_2) \Phi(-d_+(T_1))), \quad (\text{A.14})$$

where

$$d_+(T_1) = \frac{\log(L(0, T_1, T_2)/\kappa) + \sigma^2 T_1 / 2}{\sigma_1 \sqrt{T_1}},$$

and

$$d_-(T_1) = \frac{\log(L(0, T_1, T_2)/\kappa) - \sigma^2 T_1 / 2}{\sigma \sqrt{T_1}},$$

are given in Proposition 4.5, and the LIBOR rate $L(0, T_1, T_2)$ is given by

$$L(0, T_1, T_2) = \frac{P(0, T_1) - P(0, T_2)}{(T_2 - T_1) P(0, T_2)} \\ = \frac{e^{-3r/4} - e^{-r}}{0.25 e^{-r}} \\ = 4(e^{r/4} - 1) \\ \simeq 3.9695675\%.$$

Hence, we have

$$d_+(T_1) = \frac{\log(0.039695675/0.045) + (0.1)^2 \times 0.75 / 2}{0.1 \times \sqrt{0.75}} \simeq -1.404927033,$$

and

$$d_-(T_1) = \frac{\log(0.039695675/0.045) - (0.1)^2 \times 0.75 / 2}{0.1 \times \sqrt{0.75}} \simeq -1.491529573,$$

hence

$$\begin{aligned}
 & \mathbf{E}^* \left[e^{-\int_0^{T_2} r_s ds} (\kappa - L(T_1, T_1, T_2))^+ \right] \\
 &= 0.961269954 \times (\kappa \Phi(1.491529573) - L(0, T_1, T_2) \times \Phi(1.404927033)) \\
 &= 0.961269954 \times (0.045 \times 0.932089 - 0.039695675 \times 0.919979) \\
 &\simeq 0.52147141\%.
 \end{aligned}$$

Finally, we need to multiply (A.14) by the notional principal amount of \$1 million per interest rate percentage point, *i.e.* \$10,000 per percentage point or \$100 per basis point, which yields \$5214.71.

Exercise 4.2

- a) We price the swaption at $t = 0$, with $T_1 = 4$ years, $T_2 = 5$ years, $T_3 = 6$ years, $T_4 = 7$ years, $\kappa = 5\%$, and the swap rate $(S(t, T_1, T_4))_{t \in [0, T_1]}$ is modeled as a driftless geometric Brownian motion with volatility coefficient $\hat{\sigma} = \sigma_{1,4}(t) = 0.2$ under the forward swap measure $\mathbb{P}_{1,4}$. The discount factors are given by $P(0, T_1) = e^{-4r}$, $P(0, T_2) = e^{-5r}$, $P(0, T_3) = e^{-6r}$, $P(0, T_4) = e^{-7r}$, where $r = 5\%$.
- b) By Proposition 4.12 the price of the swaption is

$$(P(0, T_1) - P(0, T_4))\Phi(d_+(T_1 - t)) - \kappa\Phi(d_-(T_1))(P(0, T_2) + P(0, T_3) + P(0, T_4)),$$

where $d_+(T_1)$ and $d_-(T_1)$ are given in Proposition 4.12, and the LIBOR swap rate $S(0, T_1, T_4)$ is given by

$$\begin{aligned}
 S(0, T_1, T_4) &= \frac{P(0, T_1) - P(0, T_4)}{P(0, T_1, T_4)} \\
 &= \frac{P(0, T_1) - P(0, T_4)}{P(0, T_2) + P(0, T_3) + P(0, T_4)} \\
 &= \frac{e^{-4r} - e^{-7r}}{e^{-5r} + e^{-6r} + e^{-7r}} \\
 &= \frac{e^{3r} - 1}{e^{2r} + e^r + 1} \\
 &= \frac{e^{0.15} - 1}{e^{0.1} + e^{0.05} + 1} \\
 &= 0.051271096.
 \end{aligned}$$

By Proposition 4.12 we also have

$$d_+(T_1) = \frac{\log(0.051271096/0.05) + (0.2)^2 \times 4/2}{0.2\sqrt{4}} = 0.526161682,$$

and

$$d_-(T_1) = \frac{\log(0.051271096/0.05) - (0.2)^2 \times 4/2}{0.2\sqrt{4}} = 0.005214714,$$

Hence, the price of the swaption is given by

$$\begin{aligned}
 & (e^{-4r} - e^{-7r})\Phi(0.526161682) \\
 & - \kappa\Phi(0.005214714)(e^{-5r} + e^{-6r} + e^{-7r}) \\
 &= (0.818730753 - 0.70468809) \times 0.700612 \\
 & - 0.05 \times 0.50208 \times (0.818730753 + 0.740818221 + 0.70468809) \\
 &= 2.3058251\%.
 \end{aligned} \tag{A.15}$$

Finally, we need to multiply (A.15) by the notional principal amount of \$10 million, *i.e.* \$100,000 by interest percentage point, or \$1,000 by basis point, which yields \$230,582.51.



Exercise 4.3 Taking $t = 0$, we have $T_1 = 3$, $T_2 = 4$ and $T_3 = 5$. The LIBOR swap rate $S(t, T_1, T_3)$ is modeled as a driftless geometric Brownian motion with volatility $\sigma = 0.1$ under the forward swap measure $\widehat{P}_{i,j}$. The receiver swaption is priced using the Black-Scholes formula as

$$\begin{aligned} & \mathbb{E}^* \left[e^{-\int_t^{T_1} r_s ds} P(T_1, T_1, T_3) (\kappa - S(T_1, T_1, T_3))^+ \middle| \mathcal{F}_t \right] \\ &= \kappa \Phi(-d_-(T_1 - t)) \sum_{k=1}^2 (T_{k+1} - T_k) P(t, T_{k+1}) \\ &\quad - (P(t, T_1) - P(t, T_3)) \Phi(-d_+(T_1 - t)), \end{aligned}$$

where $\kappa = 5\%$, $r = 2\%$ and $P(t, T_1) = e^{-3r} = 0.9417$, $P(t, T_2) = e^{-4r} = 0.9231$, $P(t, T_3) = e^{-5r} = 0.9048$. Hence,

$$P(t, T_1, T_3) = P(t, T_2) + P(t, T_3) = 0.92311 + 0.90483 = 1.82794$$

and

$$S(t, T_1, T_3) = \frac{P(t, T_1) - P(t, T_3)}{P(t, T_1, T_3)} = \frac{0.9417 - 0.9048}{1.82794} = 0.02018.$$

We also have

$$\begin{aligned} d_+(T_1 - t) &= \frac{\log(S(t, T_1, T_3)/\kappa) + \sigma^2(T_1 - t)/2}{\sigma\sqrt{T_1 - t}} \\ &= \frac{\log(2.018/5) + 0.1^2 \times 3/2}{0.1\sqrt{3}} = -5.1518 \end{aligned}$$

and

$$d_-(T_1 - t) = d_+(T_1 - t) - \sigma\sqrt{T_1 - t} = -5.3250,$$

hence

$$\begin{aligned} & \mathbb{E}^* \left[e^{-\int_t^{T_1} r_s ds} P(T_1, T_1, T_3) (\kappa - S(T_1, T_1, T_3))^+ \middle| \mathcal{F}_t \right] \\ &= 0.05 \times 1.82794 \times \Phi(5.3250) - (0.9417 - 0.9048) \times \Phi(5.1518) \\ &= 0.05 \times 1.82794 \times 0.9999999 - (0.9417 - 0.9048) \times 0.9999999 \\ &= 0.054496 \\ &= 5.4496\% \\ &= 544.96 \text{ bp}, \end{aligned}$$

which yields \$54,496 after multiplication by the \$10,000 notional principal.

Exercise 4.4

a) We have

$$\begin{aligned} d \left(\frac{P(t, T_2)}{P(t, T_1)} \right) &= \frac{dP(t, T_2)}{P(t, T_1)} + P(t, T_2) d \left(\frac{1}{P(t, T_1)} \right) + dP(t, T_2) \cdot d \left(\frac{1}{P(t, T_1)} \right) \\ &= \frac{dP(t, T_2)}{P(t, T_1)} + P(t, T_2) \left(-\frac{dP(t, T_1)}{(P(t, T_1))^2} + \frac{dP(t, T_1) \cdot dP(t, T_1)}{(P(t, T_1))^3} \right) \\ &\quad - \frac{dP(t, T_1) \cdot dP(t, T_2)}{(P(t, T_1))^2} \\ &= \frac{1}{P(t, T_1)} (r_t P(t, T_2) dt + \zeta_2(t) P(t, T_2) dW_t) \end{aligned}$$

$$\begin{aligned}
& - \frac{P(t, T_2)}{(P(t, T_1))^2} (r_t P(t, T_1) dt + \zeta_1(t) P(t, T_1) dW_t) \\
& + \frac{P(t, T_2)}{(P(t, T_1))^3} ((r_t P(t, T_1) dt + \zeta_1(t) P(t, T_1) dW_t)^2) \\
& - \frac{1}{(P(t, T_1))^2} ((r_t P(t, T_1) dt + \zeta_1(t) P(t, T_1) dW_t) \cdot (r_t P(t, T_2) dt + \zeta_2(t) P(t, T_2) dW_t)) \\
& = \zeta_2(t) \frac{P(t, T_2)}{P(t, T_1)} dW_t - \zeta_1(t) \frac{P(t, T_2)}{P(t, T_1)} dW_t \\
& + (\zeta_1(t))^2 \frac{P(t, T_2)}{P(t, T_1)} dt - \zeta_1(t) \zeta_2(t) \frac{P(t, T_2)}{P(t, T_1)} dt \\
& = - \frac{P(t, T_2)}{P(t, T_1)} \zeta_1(t) (\zeta_2(t) - \zeta_1(t)) dt + \frac{P(t, T_2)}{P(t, T_1)} (\zeta_2(t) - \zeta_1(t)) dW_t \\
& = \frac{P(t, T_2)}{P(t, T_1)} (\zeta_2(t) - \zeta_1(t)) (dW_t - \zeta_1(t) dt) \\
& = (\zeta_2(t) - \zeta_1(t)) \frac{P(t, T_2)}{P(t, T_1)} d\hat{W}_t = (\zeta_2(t) - \zeta_1(t)) \frac{P(t, T_2)}{P(t, T_1)} d\hat{W}_t,
\end{aligned}$$

where $d\hat{W}_t = dW_t - \zeta_1(t)dt$ is a standard Brownian motion under the T_1 -forward measure $\hat{\mathbb{P}}$.

b) From Question (a) or (4.1.7) we have

$$\begin{aligned}
P(T_1, T_2) &= \frac{P(T_1, T_2)}{P(T_1, T_1)} \\
&= \frac{P(t, T_2)}{P(t, T_1)} \exp \left(\int_t^{T_1} (\zeta_2(s) - \zeta_1(s)) d\hat{W}_s - \frac{1}{2} \int_t^{T_1} (\zeta_2(s) - \zeta_1(s))^2 ds \right) \\
&= \frac{P(t, T_2)}{P(t, T_1)} e^{X - v^2/2},
\end{aligned}$$

where X is a centered Gaussian random variable with variance $v^2 = \int_t^{T_1} (\zeta_2(s) - \zeta_1(s))^2 ds$, independent of \mathcal{F}_t under $\hat{\mathbb{P}}$. Hence by the hint (4.5.13) with $x := P(t, T_2)/P(t, T_1)$ and $\kappa := K/x$, we find

$$\begin{aligned}
\mathbb{E}^* \left[e^{-\int_0^{T_1} r_s ds} (K - P(T_1, T_2))^+ \mid \mathcal{F}_t \right] &= P(t, T_1) \hat{\mathbb{E}} [(K - P(T_1, T_2))^+ \mid \mathcal{F}_t] \\
&= P(t, T_1) \left(K \Phi \left(\frac{v}{2} + \frac{1}{v} \log \frac{K}{x} \right) - \frac{P(t, T_2)}{P(t, T_1)} \Phi \left(-\frac{v}{2} + \frac{1}{v} \log \frac{K}{x} \right) \right) \\
&= K P(t, T_1) \Phi \left(\frac{v}{2} + \frac{1}{v} \log \frac{K}{x} \right) - P(t, T_2) \Phi \left(-\frac{v}{2} + \frac{1}{v} \log \frac{K}{x} \right).
\end{aligned}$$

Exercise 4.5

a) The forward measure $\hat{\mathbb{P}}_S$ is defined from the numéraire $N_t := P(t, S)$ and this gives

$$F_t = P(t, S) \hat{\mathbb{E}}[(\kappa - L(T, T, S))^+ \mid \mathcal{F}_t].$$

b) The LIBOR rate $L(t, T, S)$ is a driftless geometric Brownian motion with volatility σ under the forward measure $\hat{\mathbb{P}}_S$. Indeed, the LIBOR rate $L(t, T, S)$ can be written as the forward price $L(t, T, S) = \hat{X}_t = X_t/N_t$ where $X_t = (P(t, T) - P(t, S))/(S - T)$ and $N_t = P(t, S)$. Since both discounted bond prices $e^{-\int_0^t r_s ds} P(t, T)$ and $e^{-\int_0^t r_s ds} P(t, S)$ are martingales under \mathbb{P}^* , the same is true of X_t . Hence $L(t, T, S) = X_t/N_t$ becomes a martingale under the forward measure $\hat{\mathbb{P}}_S$ by Proposition 2.1, and computing its dynamics under $\hat{\mathbb{P}}_S$ amounts to removing any “ dt ” term in (4.5.14), i.e.

$$dL(t, T, S) = \sigma L(t, T, S) d\hat{W}_t, \quad 0 \leq t \leq T,$$



hence $L(t, T, S) = L(0, T, S) e^{\sigma \widehat{W}_t - \sigma^2 t / 2}$, where $(\widehat{W}_t)_{t \in \mathbb{R}_+}$ is a standard Brownian motion under $\widehat{\mathbb{P}}_S$.

c) We find

$$\begin{aligned} F_t &= P(t, S) \widehat{\mathbb{E}}[(\kappa - L(T, T, S))^+ | \mathcal{F}_t] \\ &= P(t, S) \widehat{\mathbb{E}}[(\kappa - L(t, T, S) e^{-(T-t)\sigma^2/2 + (\widehat{W}_T - \widehat{W}_t)\sigma})^+ | \mathcal{F}_t] \\ &= P(t, S)(\kappa \Phi(-d_-(T-t)) - \widehat{X}_t \Phi(-d_+(T-t))) \\ &= \kappa P(t, S) \Phi(-d_-(T-t)) - P(t, S) L(t, T, S) \Phi(-d_+(T-t)) \\ &= \kappa P(t, S) \Phi(-d_-(T-t)) - (P(t, T) - P(t, S)) \Phi(-d_+(T-t)) / (S - T), \end{aligned}$$

where $e^m = L(t, T, S) e^{-(T-t)\sigma^2/2}$, $v^2 = (T-t)\sigma^2$, and

$$d_+(T-t) = \frac{\log(L(t, T, S) / \kappa)}{\sigma \sqrt{T-t}} + \frac{\sigma \sqrt{T-t}}{2},$$

and

$$d_-(T-t) = \frac{\log(L(t, T, S) / \kappa)}{\sigma \sqrt{T-t}} - \frac{\sigma \sqrt{T-t}}{2},$$

because $L(t, T, S)$ is a driftless geometric Brownian motion with volatility σ under the forward measure $\widehat{\mathbb{P}}_S$.

Exercise 4.6

a) We have

$$P(T_i, T_i, T_j) = \sum_{k=i}^{j-1} c_{k+1} P(T_i, T_{k+1}).$$

b) It suffices to let $\tilde{c}_k = 1$, $k = i+1, \dots, j-1$. and $\tilde{c}_j = c_j + 1/\kappa$.

c) The swaption can be priced as

$$\begin{aligned} &\mathbb{E}^* \left[e^{-\int_t^{T_i} r_s ds} (P(T_i, T_i) - P(T_i, T_j) - \kappa P(T_i, T_i, T_j))^+ \mid \mathcal{F}_t \right] \\ &= \mathbb{E}^* \left[e^{-\int_t^{T_i} r_s ds} \left(1 - \kappa \sum_{k=i}^{j-1} c_{k+1} P(T_i, T_{k+1}) \right)^+ \mid \mathcal{F}_t \right] \\ &= \kappa \mathbb{E}^* \left[e^{-\int_t^{T_i} r_s ds} \left(\sum_{k=i}^{j-1} c_{k+1} F_{k+1}(T_i, \gamma_\kappa) - \sum_{k=i}^{j-1} c_{k+1} F_{k+1}(T_i, r_{T_i}) \right)^+ \mid \mathcal{F}_t \right] \\ &= \kappa \mathbb{E}^* \left[e^{-\int_t^{T_i} r_s ds} \left(\sum_{k=i}^{j-1} c_{k+1} (F_{k+1}(T_i, \gamma_\kappa) - F_{k+1}(T_i, r_{T_i})) \right)^+ \mid \mathcal{F}_t \right] \\ &= \kappa \mathbb{E}^* \left[e^{-\int_t^{T_i} r_s ds} \sum_{k=i}^{j-1} c_{k+1} \mathbb{1}_{\{r_{T_i} \leq \gamma_\kappa\}} (F_{k+1}(T_i, \gamma_\kappa) - F_{k+1}(T_i, r_{T_i})) \mid \mathcal{F}_t \right] \\ &= \kappa \mathbb{E}^* \left[e^{-\int_t^{T_i} r_s ds} \sum_{k=i}^{j-1} c_{k+1} (F_{k+1}(T_i, \gamma_\kappa) - F_{k+1}(T_i, r_{T_i}))^+ \mid \mathcal{F}_t \right] \\ &= \kappa \sum_{k=i}^{j-1} c_{k+1} \mathbb{E}^* \left[e^{-\int_t^{T_i} r_s ds} (F_{k+1}(T_i, \gamma_\kappa) - P(T_i, T_{k+1}))^+ \mid \mathcal{F}_t \right] \\ &= \kappa \sum_{k=i}^{j-1} c_{k+1} P(t, T_i) \widehat{\mathbb{E}}_i [(F_{k+1}(T_i, \gamma_\kappa) - P(T_i, T_{k+1}))^+ \mid \mathcal{F}_t], \end{aligned}$$

which is a weighted sum of bond put option prices with strike prices $F_{k+1}(T_i, \gamma_\kappa)$, $k = i, i+1, \dots, j-1$.

Exercise 4.7

a) We have

$$dX_t = \sum_{i=2}^n c_i d\widehat{P}(t, T_i) = \sum_{i=2}^n c_i \sigma_i(t) \widehat{P}(t, T_i) d\widehat{W}_t = \sigma_t X_t d\widehat{W}_t,$$

from which we obtain

$$\sigma_t = \frac{1}{X_t} \sum_{i=2}^n c_i \sigma_i(t) \widehat{P}(t, T_i) = \frac{\sum_{i=2}^n c_i \sigma_i(t) \widehat{P}(t, T_i)}{\sum_{i=2}^n c_i \widehat{P}(t, T_i)} = \frac{\sum_{i=2}^n c_i \sigma_i(t) P(t, T_i)}{\sum_{i=2}^n c_i P(t, T_i)}.$$

b) Approximating the random process $(\sigma_t)_{t \in \mathbb{R}_+}$ by the deterministic function of time

$$\widehat{\sigma}(t) := \frac{1}{X_0} \sum_{i=2}^n c_i \sigma_i(t) \widehat{P}(0, T_i) = \frac{\sum_{i=2}^n c_i \sigma_i(t) \widehat{P}(0, T_i)}{\sum_{i=2}^n c_i \widehat{P}(0, T_i)} = \frac{\sum_{i=2}^n c_i \sigma_i(t) P(0, T_i)}{\sum_{i=2}^n c_i P(0, T_i)},$$

we find

$$\begin{aligned} & P(0, T_1) \widehat{\mathbb{E}}[(X_{T_0} - \kappa)^+] \\ & \simeq P(0, T_1) \widehat{\mathbb{E}} \left[\left(X_0 e^{\int_0^{T_0} |\widehat{\sigma}(t)|^2 dW_t^{(1)} - \frac{1}{2} \int_0^{T_0} |\widehat{\sigma}(t)|^2 dt} - \kappa \right)^+ \right] \\ & = P(0, T_1) \text{Bl}_{\text{call}} \left(X_0, \kappa, \sqrt{\frac{1}{T_0} \int_0^{T_0} |\widehat{\sigma}(t)|^2 dt}, 0, T \right) \\ & = P(0, T_1) (X_0 \Phi(v/2 + (\log(X_0/\kappa))/v) - \kappa \Phi(-v/2 + (\log(X_0/\kappa))/v)) \\ & = \sum_{i=2}^n c_i P(0, T_i) \Phi \left(\frac{v}{2} - \frac{1}{v} \log \left(\sum_{i=2}^n \frac{c_i P(0, T_i)}{\kappa P(0, T_1)} \right) \right) \\ & \quad - \kappa P(0, T_1) \Phi \left(-\frac{v}{2} - \frac{1}{v} \log \left(\sum_{i=2}^n \frac{c_i P(0, T_i)}{\kappa P(0, T_1)} \right) \right), \end{aligned}$$

with $v := \sqrt{\int_0^{T_0} |\widehat{\sigma}(t)|^2 dt}$.

Exercise 4.8

a) We have

$$\frac{dP(t, T_i)}{P(t, T_i)} = r_t dt + \zeta^{(i)}(t) dB_t, \quad i = 1, 2,$$

and

$$P(T, T_i) = P(t, T_i) \exp \left(\int_t^T r_s ds + \int_t^T \zeta^{(i)}(s) dB_s - \frac{1}{2} \int_t^T |\zeta^{(i)}(s)|^2 ds \right),$$

$0 \leq t \leq T \leq T_i$, $i = 1, 2$, hence

$$\log P(T, T_i) = \log P(t, T_i) + \int_t^T r_s ds + \int_t^T \zeta^{(i)}(s) dB_s - \frac{1}{2} \int_t^T |\zeta^{(i)}(s)|^2 ds,$$

$0 \leq t \leq T \leq T_i$, $i = 1, 2$, and

$$d \log P(t, T_i) = r_t dt + \zeta^{(i)}(t) dB_t - \frac{1}{2} |\zeta^{(i)}(t)|^2 dt, \quad i = 1, 2.$$



In the present model, we have

$$dr_t = \sigma dB_t,$$

where $(B_t)_{t \in \mathbb{R}_+}$ is a standard Brownian motion under \mathbb{P} , by the [solution](#) of Exercise 1.4 and [\(1.4.9\)](#) we have

$$\zeta^{(i)}(t) = -(T_i - t)\sigma, \quad 0 \leq t \leq T_i, \quad i = 1, 2.$$

Letting

$$dB_t^{(i)} = dB_t - \zeta^{(i)}(t)dt,$$

defines a standard Brownian motion under \mathbb{P}_i , $i = 1, 2$, and we have

$$\begin{aligned} \frac{P(T, T_1)}{P(T, T_2)} &= \frac{P(t, T_1)}{P(t, T_2)} \exp \left(\int_t^T (\zeta^{(1)}(s) - \zeta^{(2)}(s)) dB_s - \frac{1}{2} \int_t^T (|\zeta^{(1)}(s)|^2 - |\zeta^{(2)}(s)|^2) ds \right) \\ &= \frac{P(t, T_1)}{P(t, T_2)} \exp \left(\int_t^T (\zeta^{(1)}(s) - \zeta^{(2)}(s)) dB_s^{(2)} - \frac{1}{2} \int_t^T (\zeta^{(1)}(s) - \zeta^{(2)}(s))^2 ds \right), \end{aligned}$$

which is an \mathcal{F}_t -martingale under \mathbb{P}_2 and under $\mathbb{P}_{1,2}$, and

$$\frac{P(T, T_2)}{P(T, T_1)} = \frac{P(t, T_2)}{P(t, T_1)} \exp \left(- \int_t^T (\zeta^{(1)}(s) - \zeta^{(2)}(s)) dB_s^{(1)} - \frac{1}{2} \int_t^T (\zeta^{(1)}(s) - \zeta^{(2)}(s))^2 ds \right),$$

which is an \mathcal{F}_t -martingale under \mathbb{P}_1 .

b) We have

$$\begin{aligned} f(t, T_1, T_2) &= -\frac{1}{T_2 - T_1} (\log P(t, T_2) - \log P(t, T_1)) \\ &= r_t + \frac{1}{T_2 - T_1} \frac{\sigma^2}{6} ((T_1 - t)^3 - (T_2 - t)^3). \end{aligned}$$

c) We have

$$\begin{aligned} df(t, T_1, T_2) &= -\frac{1}{T_2 - T_1} d \log \frac{P(t, T_2)}{P(t, T_1)} \\ &= -\frac{1}{T_2 - T_1} \left((\zeta^{(2)}(t) - \zeta^{(1)}(t)) dB_t - \frac{1}{2} (|\zeta^{(2)}(t)|^2 - |\zeta^{(1)}(t)|^2) dt \right) \\ &= -\frac{1}{T_2 - T_1} \left((\zeta^{(2)}(t) - \zeta^{(1)}(t)) (dB_t^{(2)} + \zeta^{(2)}(t) dt) - \frac{1}{2} (|\zeta^{(2)}(t)|^2 - |\zeta^{(1)}(t)|^2) dt \right) \\ &= -\frac{1}{T_2 - T_1} \left((\zeta^{(2)}(t) - \zeta^{(1)}(t)) dB_t^{(2)} - \frac{1}{2} (\zeta^{(2)}(t) - \zeta^{(1)}(t))^2 dt \right). \end{aligned}$$

d) We have

$$\begin{aligned} f(T, T_1, T_2) &= -\frac{1}{T_2 - T_1} \log \frac{P(T, T_2)}{P(T, T_1)} \\ &= f(t, T_1, T_2) - \frac{1}{T_2 - T_1} \left(\int_t^T (\zeta^{(2)}(s) - \zeta^{(1)}(s)) dB_s - \frac{1}{2} (|\zeta^{(2)}(s)|^2 - |\zeta^{(1)}(s)|^2) ds \right) \\ &= f(t, T_1, T_2) - \frac{1}{T_2 - T_1} \left(\int_t^T (\zeta^{(2)}(s) - \zeta^{(1)}(s)) dB_s^{(2)} - \frac{1}{2} \int_t^T (\zeta^{(2)}(s) - \zeta^{(1)}(s))^2 ds \right) \\ &= f(t, T_1, T_2) - \frac{1}{T_2 - T_1} \left(\int_t^T (\zeta^{(2)}(s) - \zeta^{(1)}(s)) dB_s^{(1)} + \frac{1}{2} \int_t^T (\zeta^{(2)}(s) - \zeta^{(1)}(s))^2 ds \right). \end{aligned}$$

Hence $f(T, T_1, T_2)$ has a Gaussian distribution given \mathcal{F}_t with conditional mean

$$m_1 := f(t, T_1, T_2) - \frac{1}{2} \int_t^T (\zeta^{(2)}(s) - \zeta^{(1)}(s))^2 ds$$

under \mathbb{P}_1 , resp.

$$m_2 := f(t, T_1, T_2) + \frac{1}{2} \int_t^T (\zeta^{(2)}(s) - \zeta^{(1)}(s))^2 ds$$

under \mathbb{P}_2 , and variance

$$v^2 = \frac{1}{(T_2 - T_1)^2} \int_t^T (\zeta^{(2)}(s) - \zeta^{(1)}(s))^2 ds.$$

Hence we have

$$\begin{aligned} & (T_2 - T_1) \mathbf{E}^* \left[e^{-\int_t^{T_2} r_s ds} (f(T_1, T_1, T_2) - \kappa)^+ \middle| \mathcal{F}_t \right] \\ &= (T_2 - T_1) P(t, T_2) \mathbf{E}_2 \left[(f(T_1, T_1, T_2) - \kappa)^+ \mid \mathcal{F}_t \right] \\ &= (T_2 - T_1) P(t, T_2) \mathbf{E}_2 \left[(m_2 + X - \kappa)^+ \middle| \mathcal{F}_t \right] \\ &= (T_2 - T_1) P(t, T_2) \left(\frac{v}{\sqrt{2\pi}} e^{-(\kappa - m_2)^2 / (2v^2)} + (m_2 - \kappa) \Phi((m_2 - \kappa)/v) \right). \end{aligned}$$

e) We have

$$\begin{aligned} L(T, T_1, T_2) &= S(T, T_1, T_2) \\ &= \frac{1}{T_2 - T_1} \left(\frac{P(T, T_1)}{P(T, T_2)} - 1 \right) \\ &= \frac{1}{T_2 - T_1} \left(\frac{P(t, T_1)}{P(t, T_2)} \exp \left(\int_t^T (\zeta^{(1)}(s) - \zeta^{(2)}(s)) dB_s - \frac{1}{2} \int_t^T (|\zeta^{(1)}(s)|^2 - |\zeta^{(2)}(s)|^2) ds \right) - 1 \right) \\ &= \frac{1}{T_2 - T_1} \left(\frac{P(t, T_1)}{P(t, T_2)} \exp \left(\int_t^T (\zeta^{(1)}(s) - \zeta^{(2)}(s)) dB_s^{(2)} - \frac{1}{2} \int_t^T (\zeta^{(1)}(s) - \zeta^{(2)}(s))^2 ds \right) - 1 \right) \\ &= \frac{1}{T_2 - T_1} \left(\frac{P(t, T_1)}{P(t, T_2)} \exp \left(\int_t^T (\zeta^{(1)}(s) - \zeta^{(2)}(s)) dB_s^{(1)} + \frac{1}{2} \int_t^T (\zeta^{(1)}(s) - \zeta^{(2)}(s))^2 ds \right) - 1 \right), \end{aligned}$$

and, by Itô calculus,

$$\begin{aligned} dS(t, T_1, T_2) &= \frac{1}{T_2 - T_1} d \left(\frac{P(t, T_1)}{P(t, T_2)} \right) \\ &= \frac{1}{T_2 - T_1} \frac{P(t, T_1)}{P(t, T_2)} \left((\zeta^{(1)}(t) - \zeta^{(2)}(t)) dB_t + \frac{1}{2} (\zeta^{(1)}(t) - \zeta^{(2)}(t))^2 dt \right. \\ &\quad \left. - \frac{1}{2} (|\zeta^{(1)}(t)|^2 - |\zeta^{(2)}(t)|^2) dt \right) \\ &= \left(\frac{1}{T_2 - T_1} + S(t, T_1, T_2) \right) \left((\zeta^{(1)}(t) - \zeta^{(2)}(t)) dB_t + \zeta^{(2)}(t) (\zeta^{(2)}(t) - \zeta^{(1)}(t)) dt \right) \\ &= \left(\frac{1}{T_2 - T_1} + S(t, T_1, T_2) \right) \left((\zeta^{(1)}(t) - \zeta^{(2)}(t)) dB_t^{(1)} + (|\zeta^{(2)}(t)|^2 - |\zeta^{(1)}(t)|^2) dt \right) \\ &= \left(\frac{1}{T_2 - T_1} + S(t, T_1, T_2) \right) (\zeta^{(1)}(t) - \zeta^{(2)}(t)) dB_t^{(2)}, \quad t \in [0, T_1], \end{aligned}$$

hence $t \mapsto \frac{1}{T_2 - T_1} + S(t, T_1, T_2)$ is a geometric Brownian motion, with

$$\begin{aligned} & \frac{1}{T_2 - T_1} + S(T, T_1, T_2) \\ &= \left(\frac{1}{T_2 - T_1} + S(t, T_1, T_2) \right) \exp \left(\int_t^T (\zeta^{(1)}(s) - \zeta^{(2)}(s)) dB_s^{(2)} - \frac{1}{2} \int_t^T (\zeta^{(1)}(s) - \zeta^{(2)}(s))^2 ds \right), \\ & 0 \leq t \leq T \leq T_1. \end{aligned}$$

f) We have

$$\begin{aligned} & (T_2 - T_1) \mathbf{E}^* \left[e^{-\int_t^{T_2} r_s ds} (L(T_1, T_1, T_2) - \kappa)^+ \middle| \mathcal{F}_t \right] \\ &= (T_2 - T_1) \mathbf{E}^* \left[e^{-\int_t^{T_1} r_s ds} P(T_1, T_2) (L(T_1, T_1, T_2) - \kappa)^+ \middle| \mathcal{F}_t \right] \\ &= P(t, T_1, T_2) \mathbf{E}_{1,2} \left[(S(T_1, T_1, T_2) - \kappa)^+ \middle| \mathcal{F}_t \right]. \end{aligned}$$



The forward measure \mathbb{P}_2 is defined by

$$\mathbf{E}^* \left[\frac{d\mathbb{P}_2}{d\mathbb{P}} \middle| \mathcal{F}_t \right] = \frac{P(t, T_2)}{P(0, T_2)} e^{-\int_0^t r_s ds}, \quad 0 \leq t \leq T_2,$$

and the forward swap measure is defined by

$$\mathbf{E}^* \left[\frac{d\mathbb{P}_{1,2}}{d\mathbb{P}} \middle| \mathcal{F}_t \right] = \frac{P(t, T_2)}{P(0, T_2)} e^{-\int_0^t r_s ds}, \quad 0 \leq t \leq T_1,$$

hence \mathbb{P}_2 and $\mathbb{P}_{1,2}$ coincide up to time T_1 and $(B_t^{(2)})_{t \in [0, T_1]}$ is a standard Brownian motion until time T_1 under \mathbb{P}_2 and under $\mathbb{P}_{1,2}$, consequently under $\mathbb{P}_{1,2}$ we have

$$\begin{aligned} L(T, T_1, T_2) &= S(T, T_1, T_2) \\ &= -\frac{1}{T_2 - T_1} + \left(\frac{1}{T_2 - T_1} + S(t, T_1, T_2) \right) e^{\int_t^T (\zeta^{(1)}(s) - \zeta^{(2)}(s)) dB_s^{(2)} - \frac{1}{2} \int_t^T (\zeta^{(1)}(s) - \zeta^{(2)}(s))^2 ds}, \end{aligned}$$

has same distribution as

$$\frac{1}{T_2 - T_1} \left(\frac{P(t, T_1)}{P(t, T_2)} e^{X - \text{Var}[X]/2} - 1 \right),$$

where X is a centered Gaussian random variable with variance

$$\int_t^{T_1} (\zeta^{(1)}(s) - \zeta^{(2)}(s))^2 ds$$

given \mathcal{F}_t . Hence, we have

$$\begin{aligned} (T_2 - T_1) \mathbf{E}^* \left[e^{-\int_t^{T_2} r_s ds} (L(T_1, T_1, T_2) - \kappa)^+ \middle| \mathcal{F}_t \right] \\ = P(t, T_1, T_2) \\ \times \text{Bl} \left(\frac{1}{T_2 - T_1} + S(t, T_1, T_2), \frac{\int_t^{T_1} (\zeta^{(1)}(s) - \zeta^{(2)}(s))^2 ds}{T_1 - t}, \kappa + \frac{1}{T_2 - T_1}, T_1 - t \right). \end{aligned}$$

Exercise 4.9

a) We have

$$L(t, T_1, T_2) = L(0, T_1, T_2) e^{\int_0^t \gamma_1(s) dW_s^{(2)} - \int_0^t |\gamma_1(s)|^2 ds / 2}, \quad 0 \leq t \leq T_1,$$

and $L(t, T_2, T_3) = b$. Note that we also have $P(t, T_2)/P(t, T_3) = 1 + \delta b$, hence $\widehat{P}_2 = \widehat{P}_3 = \widehat{P}_{1,2}$ up to time T_1 .

b) We use change of numéraire under the forward measure \mathbb{P}_2 .

c) We have

$$\begin{aligned} \mathbf{E}^* \left[e^{-\int_t^{T_2} r_s ds} (L(T_1, T_1, T_2) - \kappa)^+ \middle| \mathcal{F}_t \right] \\ = P(t, T_2) \widehat{\mathbf{E}}_2 [(L(T_1, T_1, T_2) - \kappa)^+ | \mathcal{F}_t] \\ = P(t, T_2) \widehat{\mathbf{E}}_2 \left[(L(t, T_1, T_2) e^{\int_t^{T_1} \gamma_1(s) dW_s^{(2)} - \int_t^{T_1} |\gamma_1(s)|^2 ds / 2} - \kappa)^+ \middle| \mathcal{F}_t \right] \\ = P(t, T_2) \text{Bl}(\kappa, L(t, T_1, T_2), \sigma_1(t), 0, T_1 - t), \end{aligned}$$

where

$$\sigma_1^2(t) = \frac{1}{T_1 - t} \int_t^{T_1} |\gamma_1(s)|^2 ds.$$

d) We have

$$\frac{P(t, T_1)}{P(t, T_1, T_3)} = \frac{P(t, T_1)}{\delta P(t, T_2) + \delta P(t, T_3)}$$

$$\begin{aligned}
&= \frac{P(t, T_1)}{\delta P(t, T_2)} \frac{1}{1 + P(t, T_3)/P(t, T_2)} \\
&= \frac{1 + \delta b}{\delta(\delta b + 2)} (1 + \delta L(t, T_1, T_2)), \quad 0 \leq t \leq T_1,
\end{aligned}$$

and

$$\begin{aligned}
\frac{P(t, T_3)}{P(t, T_1, T_3)} &= \frac{P(t, T_3)}{P(t, T_2) + P(t, T_3)} \\
&= \frac{1}{1 + P(t, T_2)/P(t, T_3)} \\
&= \frac{1}{\delta} \frac{1}{2 + \delta b}, \quad 0 \leq t \leq T_2.
\end{aligned}$$

e) We have

$$\begin{aligned}
S(t, T_1, T_3) &= \frac{P(t, T_1)}{P(t, T_1, T_3)} - \frac{P(t, T_3)}{P(t, T_1, T_3)} \\
&= \frac{1 + \delta b}{\delta(2 + \delta b)} (1 + \delta L(t, T_1, T_2)) - \frac{1}{\delta(2 + \delta b)} \\
&= \frac{1}{2 + \delta b} (b + (1 + \delta b)L(t, T_1, T_2)), \quad 0 \leq t \leq T_2.
\end{aligned}$$

We have

$$\begin{aligned}
dS(t, T_1, T_3) &= \frac{1 + \delta b}{2 + \delta b} L(t, T_1, T_2) \gamma_1(t) dW_t^{(2)} \\
&= \left(S(t, T_1, T_3) - \frac{b}{2 + \delta b} \right) \gamma_1(t) dW_t^{(2)} \\
&= S(t, T_1, T_3) \sigma_{1,3}(t) dW_t^{(2)}, \quad 0 \leq t \leq T_2,
\end{aligned}$$

with

$$\begin{aligned}
\sigma_{1,3}(t) &= \left(1 - \frac{b}{S(t, T_1, T_3)(2 + \delta b)} \right) \gamma_1(t) \\
&= \left(1 - \frac{b}{b + (1 + \delta b)L(t, T_1, T_2)} \right) \gamma_1(t) \\
&= \frac{(1 + \delta b)L(t, T_1, T_2)}{b + (1 + \delta b)L(t, T_1, T_2)} \gamma_1(t) \\
&= \frac{(1 + \delta b)L(t, T_1, T_2)}{(2 + \delta b)S(t, T_1, T_3)} \gamma_1(t).
\end{aligned}$$

f) The process $(W^{(2)})_{t \in \mathbb{R}_+}$ is a standard Brownian motion under $\hat{P}_2 = \hat{P}_{1,3}$ and $P(t, T_1, T_3) \hat{\mathbf{E}}_{1,3} [(S(T_1, T_1, T_3) - \kappa)^+ | \mathcal{F}_t]$

$$= P(t, T_2) \text{Bl}(\kappa, S(t, T_1, T_2), \tilde{\sigma}_{1,3}(t), 0, T_1 - t),$$

where $|\tilde{\sigma}_{1,3}(t)|^2$ is the approximation of the volatility

$$\frac{1}{T_1 - t} \int_t^{T_1} |\sigma_{1,3}(s)|^2 ds = \frac{1}{T_1 - t} \int_t^{T_1} \left(\frac{(1 + \delta b)L(s, T_1, T_2)}{(2 + \delta b)S(s, T_1, T_3)} \right)^2 \gamma_1(s) ds$$

obtained by freezing the random component of $\sigma_{1,3}(s)$ at time t , i.e.

$$\tilde{\sigma}_{1,3}^2(t) = \frac{1}{T_1 - t} \left(\frac{(1 + \delta b)L(t, T_1, T_2)}{(2 + \delta b)S(t, T_1, T_3)} \right)^2 \int_t^{T_1} |\gamma_1(s)|^2 ds.$$

Exercise 4.10 Bond option hedging.

a) We have

$$\begin{aligned}
&\mathbf{E}^* \left[e^{-\int_t^T r_s ds} (P(T, S) - \kappa)^+ \middle| \mathcal{F}_t \right] = V_T = V_0 + \int_0^T dV_t \\
&= P(0, T) \mathbf{E}_T [(P(T, S) - \kappa)^+] + \int_0^T \xi_s^T dP(s, T) + \int_0^T \xi_s^S dP(s, S).
\end{aligned}$$



b) We have

$$\begin{aligned}
 d\tilde{V}_t &= d\left(e^{-\int_0^t r_s ds} V_t\right) \\
 &= -r_t e^{-\int_0^t r_s ds} V_t dt + e^{-\int_0^t r_s ds} dV_t \\
 &= -r_t e^{-\int_0^t r_s ds} (\xi_t^T P(t, T) \\
 &\quad + \xi_t^S P(t, S)) dt + e^{-\int_0^t r_s ds} \xi_t^T dP(t, T) + e^{-\int_0^t r_s ds} \xi_t^S dP(t, S) \\
 &= \xi_t^T d\tilde{P}(t, T) + \xi_t^S d\tilde{P}(t, S).
 \end{aligned}$$

c) By Itô's formula we have

$$\begin{aligned}
 \mathbb{E}_T [(P(T, S) - \kappa)^+ | \mathcal{F}_t] &= C(X_t, v(t, T)) \\
 &= C(X_0, v(0, T)) + \int_0^t \frac{\partial C}{\partial x}(X_s, v(s, T)) dX_s \\
 &= \mathbb{E}_T [(P(T, S) - \kappa)^+] + \int_0^t \frac{\partial C}{\partial x}(X_s, v(s, T)) dX_s,
 \end{aligned}$$

since the process

$$t \mapsto \mathbb{E}_T [(P(T, S) - \kappa)^+ | \mathcal{F}_t]$$

is a martingale under $\tilde{\mathbb{P}}$.

d) We have

$$\begin{aligned}
 d\hat{V}_t &= d\left(\frac{V_t}{P(t, T)}\right) \\
 &= d\mathbb{E}_T [(P(T, S) - \kappa)^+ | \mathcal{F}_t] \\
 &= \frac{\partial C}{\partial x}(X_t, v(t, T)) dX_t \\
 &= \frac{P(t, S)}{P(t, T)} \frac{\partial C}{\partial x}(X_t, v(t, T)) (\sigma_t^S - \sigma_t^T) dB_t^T.
 \end{aligned}$$

e) We have

$$\begin{aligned}
 dV_t &= d\left(\frac{P(t, T)}{\hat{V}_t}\right) \\
 &= P(t, T) d\hat{V}_t + \hat{V}_t dP(t, T) + d\hat{V}_t \cdot dP(t, T) \\
 &= P(t, S) \frac{\partial C}{\partial x}(X_t, v(t, T)) (\sigma_t^S - \sigma_t^T) dB_t^T + \hat{V}_t dP(t, T) \\
 &\quad + P(t, S) \frac{\partial C}{\partial x}(X_t, v(t, T)) (\sigma_t^S - \sigma_t^T) \sigma_t^T dt \\
 &= P(t, S) \frac{\partial C}{\partial x}(X_t, v(t, T)) (\sigma_t^S - \sigma_t^T) dB_t + \hat{V}_t dP(t, T).
 \end{aligned}$$

f) We have

$$\begin{aligned}
 d\tilde{V}_t &= d(e^{-\int_0^t r_s ds} V_t) \\
 &= -r_t e^{-\int_0^t r_s ds} V_t dt + e^{-\int_0^t r_s ds} dV_t \\
 &= \tilde{P}(t, S) \frac{\partial C}{\partial x}(X_t, v(t, T)) (\sigma_t^S - \sigma_t^T) dB_t + \hat{V}_t d\tilde{P}(t, T).
 \end{aligned}$$

g) We have

$$\begin{aligned}
 d\tilde{V}_t &= \tilde{P}(t, S) \frac{\partial C}{\partial x}(X_t, v(t, T)) (\sigma_t^S - \sigma_t^T) dB_t + \hat{V}_t d\tilde{P}(t, T) \\
 &= \frac{\partial C}{\partial x}(X_t, v(t, T)) d\tilde{P}(t, S) \\
 &\quad - \frac{P(t, S)}{P(t, T)} \frac{\partial C}{\partial x}(X_t, v(t, T)) d\tilde{P}(t, T) + \hat{V}_t d\tilde{P}(t, T) \\
 &= \left(\hat{V}_t - \frac{P(t, S)}{P(t, T)} \frac{\partial C}{\partial x}(X_t, v(t, T))\right) d\tilde{P}(t, T)
 \end{aligned}$$

$$+ \frac{\partial C}{\partial x}(X_t, v(t, T)) d\tilde{P}(t, S),$$

hence the hedging strategy $(\xi_t^T, \xi_t^S)_{t \in [0, T]}$ of the bond option is given by

$$\begin{aligned}\xi_t^T &= \hat{V}_t - \frac{P(t, S)}{P(t, T)} \frac{\partial C}{\partial x}(X_t, v(t, T)) \\ &= C(X_t, v(t, T)) - \frac{P(t, S)}{P(t, T)} \frac{\partial C}{\partial x}(X_t, v(t, T)),\end{aligned}$$

and

$$\xi_t^S = \frac{\partial C}{\partial x}(X_t, v(t, T)), \quad 0 \leq t \leq T.$$

h) We have

$$\begin{aligned}\frac{\partial C}{\partial x}(x, v) &= \frac{\partial}{\partial x} \left(x\Phi\left(\frac{v}{2} + \frac{1}{v} \log \frac{x}{\kappa}\right) - \kappa\Phi\left(-\frac{v}{2} + \frac{1}{v} \log \frac{x}{\kappa}\right) \right) \\ &= x \frac{\partial}{\partial x} \Phi\left(\frac{v}{2} + \frac{1}{v} \log \frac{x}{\kappa}\right) - \kappa \frac{\partial}{\partial x} \Phi\left(-\frac{v}{2} + \frac{1}{v} \log \frac{x}{\kappa}\right) + \Phi\left(\frac{v}{2} + \frac{1}{v} \log \frac{x}{\kappa}\right) \\ &= x \frac{e^{-\frac{1}{2}\left(\frac{v}{2} + \frac{1}{v} \log \frac{x}{\kappa}\right)^2}}{\sqrt{2\pi}vx} - \kappa \frac{e^{-\frac{1}{2}\left(-\frac{v}{2} + \frac{1}{v} \log \frac{x}{\kappa}\right)^2}}{\sqrt{2\pi}vx} + \Phi\left(\frac{v}{2} + \frac{1}{v} \log \frac{x}{\kappa}\right) \\ &= \Phi\left(\frac{\log(x/\kappa) + v^2/2}{v}\right).\end{aligned}$$

As a consequence, we get

$$\begin{aligned}\xi_t^T &= C(X_t, v(t, T)) - \frac{P(t, S)}{P(t, T)} \frac{\partial C}{\partial x}(X_t, v(t, T)) \\ &= \frac{P(t, S)}{P(t, T)} \Phi\left(\frac{v^2(t, T)/2 + \log X_t}{v(t, T)}\right) \\ &\quad - \kappa \Phi\left(-\frac{v(t, T)}{2} + \frac{1}{v(t, T)} \log \frac{P(t, S)}{\kappa P(t, T)}\right) \\ &\quad - \frac{P(t, S)}{P(t, T)} \Phi\left(\frac{\log(X_t/\kappa) + v^2(t, T)/2}{\sqrt{T-t}}\right) \\ &= -\kappa \Phi\left(\frac{\log(X_t/\kappa) - v^2(t, T)/2}{v(t, T)}\right),\end{aligned}$$

and

$$\xi_t^S = \frac{\partial C}{\partial x}(X_t, v(t, T)) = \Phi\left(\frac{\log(X_t/\kappa) + v^2(t, T)/2}{v(t, T)}\right),$$

$t \in [0, T]$, and the hedging strategy is given by

$$\begin{aligned}V_T &= \mathbb{E}^* \left[e^{-\int_t^T r_s ds} (P(T, S) - \kappa)^+ \middle| \mathcal{F}_t \right] \\ &= V_0 + \int_0^t \xi_s^T dP(s, T) + \int_0^t \xi_s^S dP(s, S) \\ &= V_0 - \kappa \int_0^t \Phi\left(\frac{\log(X_t/\kappa) - v^2(t, T)/2}{v(t, T)}\right) dP(s, T) \\ &\quad + \int_0^t \Phi\left(\frac{\log(X_t/\kappa) + v^2(t, T)/2}{v(t, T)}\right) dP(s, S).\end{aligned}$$

Consequently, the bond option can be hedged by shortselling a bond with maturity T for the amount

$$\kappa \Phi\left(\frac{\log(X_t/\kappa) - v^2(t, T)/2}{v(t, T)}\right),$$



and by holding a bond with maturity S for the amount

$$\Phi\left(\frac{\log(X_t/\kappa) + \nu^2(t, T)/2}{\nu(t, T)}\right).$$

Exercise 4.11

- a) The LIBOR rate $L(t, T, S)$ is a driftless geometric Brownian motion with deterministic volatility function $\sigma(t)$ under the forward measure $\widehat{\mathbb{P}}_S$.

Explanation: The LIBOR rate $L(t, T, S)$ can be written as the forward price $L(t, T, S) = \widehat{X}_t = X_t/N_t$ where $X_t = (P(t, T) - P(t, S))/(S - T)$ and $N_t = P(t, S)$. Since both discounted bond prices $e^{-\int_0^t r_s ds} P(t, T)$ and $e^{-\int_0^t r_s ds} P(t, S)$ are martingales under \mathbb{P}^* , the same is true of X_t . Hence $L(t, T, S) = X_t/N_t$ becomes a martingale under the forward measure $\widehat{\mathbb{P}}_S$ by Proposition 2.1, and computing its dynamics under $\widehat{\mathbb{P}}_S$ amounts to removing any “ dt ” term in the original SDE defining $L(t, T, S)$, i.e. we find

$$dL(t, T, S) = \sigma(t)L(t, T, S)d\widehat{W}_t, \quad 0 \leq t \leq T,$$

hence

$$L(t, T, S) = L(0, T, S) \exp\left(\int_0^t \sigma(s)d\widehat{W}_s - \int_0^t \sigma^2(s)ds/2\right),$$

where $(\widehat{W}_t)_{t \in \mathbb{R}_+}$ is a standard Brownian motion under $\widehat{\mathbb{P}}_S$.

- b) Choosing the annuity numéraire $N_t = P(t, S)$, we have

$$\begin{aligned} \mathbb{E}^*\left[e^{-\int_t^T r_s ds} \phi(L(T, T, S)) \mid \mathcal{F}_t\right] &= \mathbb{E}^*\left[e^{-\int_t^T r_s ds} N_T \phi(L(T, T, S)) \mid \mathcal{F}_t\right] \\ &= N_t \mathbb{E}\left[\phi(L(T, T, S)) \mid \mathcal{F}_t\right] \\ &= P(t, S) \widehat{\mathbb{E}}[\phi(L(T, T, S)) \mid \mathcal{F}_t]. \end{aligned}$$

- c) Given the solution

$$\begin{aligned} L(T, T, S) &= L(0, T, S) \exp\left(\int_0^T \sigma(s)d\widehat{W}_s - \int_0^T \sigma^2(s)ds/2\right) \\ &= L(t, T, S) \exp\left(\int_t^T \sigma(s)d\widehat{W}_s - \int_t^T \sigma^2(s)ds/2\right), \end{aligned}$$

we find

$$\begin{aligned} P(t, S) \widehat{\mathbb{E}}[\phi(L(T, T, S)) \mid \mathcal{F}_t] &= P(t, S) \widehat{\mathbb{E}}\left[\phi\left(L(t, T, S) e^{\int_t^T \sigma(s)d\widehat{W}_s - \int_t^T \sigma^2(s)ds/2}\right) \mid \mathcal{F}_t\right] \\ &= P(t, S) \int_{-\infty}^{\infty} \phi(L(t, T, S) e^{x-\eta^2/2}) e^{-x^2/(2\eta^2)} \frac{dx}{\sqrt{2\pi\eta^2}}, \end{aligned}$$

because $\int_t^T \sigma(s)d\widehat{W}_s$ is a centered Gaussian variable with variance $\eta^2 := \int_t^T \sigma^2(s)ds$, independent of \mathcal{F}_t under the forward measure $\widehat{\mathbb{P}}$.

Exercise 4.12

- a) Choosing the annuity numéraire $N_t = P(t, T_i, T_j)$, we have

$$\begin{aligned} \mathbb{E}^*\left[e^{-\int_t^{T_j} r_s ds} P(T_i, T_i, T_j) \phi(S(T_i, T_i, T_j)) \mid \mathcal{F}_t\right] &= N_t \widehat{\mathbb{E}}_{i,j} \left[\frac{P(T_i, T_i, T_j)}{N_{T_i}} \phi(S(T_i, T_i, T_j)) \mid \mathcal{F}_t \right] \\ &= P(t, T_i, T_j) \widehat{\mathbb{E}}_{i,j} [\phi(S(T_i, T_i, T_j)) \mid \mathcal{F}_t]. \end{aligned}$$

- b) Since $S(t, T_i, T_j)$ is a forward price for the numéraire $P(t, T_i, T_j)$, it is a martingale under the forward swap measure $\widehat{\mathbb{P}}_{i,j}$ and we have

$$S(t, T_i, T_j) = \sigma S(t, T_i, T_j) d\widehat{W}_t^{i,j}, \quad 0 \leq t \leq T_i,$$

where $(\widehat{W}_t^{i,j})_{t \in \mathbb{R}_+}$ is a standard Brownian motion under the forward swap measure $\widehat{\mathbb{P}}_{i,j}$.

- c) Given the solution

$$S(T_i, T_i, T_j) = S(0, T_i, T_j) e^{\sigma \widehat{W}_{T_i} - \sigma^2 T_i / 2} = S(t, T_i, T_j) e^{(\widehat{W}_{T_i} - \widehat{W}_t) \sigma - (T_i - t) \sigma^2 / 2}$$

of (4.5.18), we find

$$\begin{aligned} & P(t, T_i, T_j) \widehat{\mathbb{E}}_{i,j} [\phi(S(t, T_i, T_j)) \mid \mathcal{F}_t] \\ &= P(t, T_i, T_j) \widehat{\mathbb{E}}_{i,j} \left[\phi \left(S(t, T_i, T_j) e^{(\widehat{W}_{T_i} - \widehat{W}_t) \sigma - (T_i - t) \sigma^2 / 2} \right) \mid \mathcal{F}_t \right] \\ &= P(t, T_i, T_j) \int_{-\infty}^{\infty} \phi \left(S(t, T_i, T_j) e^{\sigma x - (T_i - t) \sigma^2 / 2} \right) e^{-x^2 / (2(T_i - t))} \frac{dx}{\sqrt{2\pi(T_i - t)}}, \end{aligned}$$

because $\widehat{W}_{T_i} - \widehat{W}_t$ is a centered Gaussian variable with variance $T_i - t$, independent of \mathcal{F}_t under the forward measure $\widehat{\mathbb{P}}_{i,j}$.

- d) We find

$$\begin{aligned} & P(t, T_i, T_j) \widehat{\mathbb{E}}_{i,j} [(\kappa - S(t, T_i, T_j))^+ \mid \mathcal{F}_t] \\ &= P(t, T_i, T_j) \widehat{\mathbb{E}}_{i,j} [(\kappa - S(t, T_i, T_j) e^{-(T_i - t) \sigma^2 / 2 + (\widehat{W}_{T_i} - \widehat{W}_t) \sigma})^+ \mid \mathcal{F}_t] \\ &= P(t, T_i, T_j) (\kappa \Phi(-d_-(T_i - t)) - \widehat{X}_t \Phi(-d_+(T_i - t))) \\ &= P(t, T_i, T_j) \kappa \Phi(-d_-(T_i - t)) - P(t, T_i, T_j) S(t, T_i, T_j) \Phi(-d_+(T_i - t)) \\ &= P(t, T_i, T_j) \kappa \Phi(-d_-(T_i - t)) - (P(t, T_i) - P(t, T_j)) \Phi(-d_+(T_i - t)), \end{aligned}$$

where $e^m = S(t, T_i, T_j) e^{-(T_i - t) \sigma^2 / 2}$, $v^2 = (T_i - t) \sigma^2$, and

$$d_+(T_i - t) = \frac{\log(S(t, T_i, T_j) / \kappa)}{\sigma \sqrt{T_i - t}} + \frac{\sigma \sqrt{T_i - t}}{2},$$

and

$$d_-(T_i - t) = \frac{\log(S(t, T_i, T_j) / \kappa)}{\sigma \sqrt{T_i - t}} - \frac{\sigma \sqrt{T_i - t}}{2},$$

because $S(t, T_i, T_j)$ is a driftless geometric Brownian motion with volatility σ under the forward measure $\widehat{\mathbb{P}}_{i,j}$.

Exercise 4.13

- a) Apply the Itô formula to $d(P(t, T_1) / P(t, T_2))$.
b) We have

$$L_{T_1} = L_t \exp \left(\int_t^{T_1} \sigma(s) dB_s - \frac{1}{2} \int_t^{T_1} |\sigma(s)|^2 ds \right).$$

- c) We have

$$\begin{aligned} & P(t, T_2) \widehat{\mathbb{E}} [(L_{T_1} - \kappa)^+ \mid \mathcal{F}_t] \\ &= P(t, T_2) \widehat{\mathbb{E}} \left[\left(L_t e^{\int_t^{T_1} \sigma(s) dB_s - \frac{1}{2} \int_t^{T_1} |\sigma(s)|^2 ds} - \kappa \right)^+ \mid \mathcal{F}_t \right] \\ &= P(t, T_2) \text{Bl}(\kappa, v(t, T_1) / \sqrt{T_1 - t}, 0, T_1 - t) \\ &= P(t, T_2) \left(L_t \Phi \left(\frac{\log(x/K)}{v(t, T_1)} + \frac{v(t, T_1)}{2} \right) - \kappa \Phi \left(\frac{\log(x/K)}{v(t, T_1)} - \frac{v(t, T_1)}{2} \right) \right). \end{aligned}$$



d) Integrate the self-financing condition (4.5.23) between 0 and t .

e) We have

$$\begin{aligned} d\tilde{V}_t &= d\left(e^{-\int_0^t r_s ds} V_t\right) \\ &= -r_t e^{-\int_0^t r_s ds} dt + e^{-\int_0^t r_s ds} dV_t \\ &= -r_t e^{-\int_0^t r_s ds} \xi_t^{(1)} P(t, T_1) - r_t e^{-\int_0^t r_s ds} \xi_t^{(2)} P(t, T_2) dt \\ &\quad + e^{-\int_0^t r_s ds} \xi_t^{(1)} dP(t, T_1) + e^{-\int_0^t r_s ds} \xi_t^{(2)} dP(t, T_2) \\ &= \xi_t^{(1)} d\tilde{P}(t, T_1) + \xi_t^{(2)} d\tilde{P}(t, T_2), \quad 0 \leq t \leq T_1. \end{aligned}$$

since

$$\frac{d\tilde{P}(t, T_1)}{\tilde{P}(t, T_1)} = \zeta_1(t) dt, \quad \frac{d\tilde{P}(t, T_2)}{\tilde{P}(t, T_2)} = \zeta_2(t) dt.$$

f) We apply the Itô formula and the fact that

$$t \mapsto \widehat{\mathbb{E}}[(L_{T_1} - \kappa)^+ | \mathcal{F}_t]$$

and $(L_t)_{t \in \mathbb{R}_+}$ are both martingales under $\widehat{\mathbb{P}}$. Next we use the fact that

$$\widehat{V}_t = \widehat{\mathbb{E}}[(L_{T_1} - \kappa)^+ | \mathcal{F}_t],$$

and apply the result of Question (f).

g) Apply the Itô rule to $V_t = P(t, T_2)\widehat{V}_t$ using Relation (4.5.21) and the result of Question (f)).

h) We have

$$\begin{aligned} dV_t &= \frac{\partial C}{\partial x}(L_t, v(t, T_1))P(t, T_1)(\zeta_1(t) - \zeta_2(t))dB_t + \widehat{V}_t dP(t, T_2) \\ &= \frac{\partial C}{\partial x}(L_t, v(t, T_1))P(t, T_1)(\zeta_1(t) - \zeta_2(t))dB_t + \widehat{V}_t \zeta_2(t)P(t, T_2)dB_t \\ &= (1 + L_t) \frac{\partial C}{\partial x}(L_t, v(t, T_1))P(t, T_2)(\zeta_1(t) - \zeta_2(t))dB_t + \widehat{V}_t \zeta_2(t)P(t, T_2)dB_t \\ &= L_t \zeta_1(t) \frac{\partial C}{\partial x}(L_t, v(t, T_1))P(t, T_2)dB_t - L_t \frac{\partial C}{\partial x}(L_t, v(t, T_1))P(t, T_2)\zeta_2(t)dB_t \\ &\quad + \frac{\partial C}{\partial x}(L_t, v(t, T_1))P(t, T_2)(\zeta_1(t) - \zeta_2(t))dB_t + \widehat{V}_t \zeta_2(t)P(t, T_2)dB_t \\ &= L_t \frac{\partial C}{\partial x}(L_t, v(t, T_1))P(t, T_2)\zeta_1(t)dB_t + \left(\widehat{V}_t - L_t \frac{\partial C}{\partial x}(L_t, v(t, T_1))\right)P(t, T_2)\zeta_2(t)dB_t \\ &\quad + \frac{\partial C}{\partial x}(L_t, v(t, T_1))P(t, T_2)(\zeta_1(t) - \zeta_2(t))dB_t, \end{aligned}$$

hence

$$\begin{aligned} d\tilde{V}_t &= L_t \zeta_1(t) \frac{\partial C}{\partial x}(L_t, v(t, T_1))\tilde{P}(t, T_2)dB_t \\ &\quad + \left(\widehat{V}_t - L_t \frac{\partial C}{\partial x}(L_t, v(t, T_1))\right)\tilde{P}(t, T_2)\zeta_2(t)dB_t \\ &\quad + \frac{\partial C}{\partial x}(L_t, v(t, T_1))\tilde{P}(t, T_2)(\zeta_1(t) - \zeta_2(t))dB_t \\ &= (\tilde{P}(t, T_1) - \tilde{P}(t, T_2))\zeta_1(t) \frac{\partial C}{\partial x}(L_t, v(t, T_1))dB_t \\ &\quad + \left(\widehat{V}_t - L_t \frac{\partial C}{\partial x}(L_t, v(t, T_1))\right)d\tilde{P}(t, T_2) \\ &\quad + \frac{\partial C}{\partial x}(L_t, v(t, T_1))\tilde{P}(t, T_2)(\zeta_1(t) - \zeta_2(t))dB_t \end{aligned}$$

$$\begin{aligned}
&= (\zeta_1(t)\tilde{P}(t, T_1) - \zeta_2(t)\tilde{P}(t, T_2)) \frac{\partial C}{\partial x}(L_t, v(t, T_1)) dB_t \\
&\quad + \left(\widehat{V}_t - L_t \frac{\partial C}{\partial x}(L_t, v(t, T_1)) \right) d\tilde{P}(t, T_2) \\
&= \frac{\partial C}{\partial x}(L_t, v(t, T_1)) d(\tilde{P}(t, T_1) - \tilde{P}(t, T_2)) + \left(\widehat{V}_t - L_t \frac{\partial C}{\partial x}(L_t, v(t, T_1)) \right) d\tilde{P}(t, T_2).
\end{aligned}$$

Exercise 4.14

a) We have

$$S(T_i, T_i, T_j) = S(t, T_i, T_j) \exp \left(\int_t^{T_i} \sigma_{i,j}(s) dB_s^{i,j} - \frac{1}{2} \int_t^{T_i} |\sigma_{i,j}|^2(s) ds \right).$$

b) We have

$$\begin{aligned}
&P(t, T_i, T_j) \mathbf{E}_{i,j} [(S(T_i, T_i, T_j) - \kappa)^+ | \mathcal{F}_t] \\
&= P(t, T_i, T_j) \mathbf{E}_{i,j} \left[\left(S(t, T_i, T_j) e^{\int_t^{T_i} \sigma_{i,j}(s) dB_s^{i,j} - \frac{1}{2} \int_t^{T_i} |\sigma_{i,j}|^2(s) ds} - \kappa \right)^+ | \mathcal{F}_t \right] \\
&= P(t, T_i, T_j) \text{Bl}(\kappa, v(t, T_i) / \sqrt{T_i - t}, 0, T_i - t) \\
&= P(t, T_i, T_j) \\
&\quad \times \left(S(t, T_i, T_j) \Phi \left(\frac{\log(x/K)}{v(t, T_i)} + \frac{v(t, T_i)}{2} \right) - \kappa \Phi \left(\frac{\log(x/K)}{v(t, T_i)} - \frac{v(t, T_i)}{2} \right) \right),
\end{aligned}$$

where

$$v^2(t, T_i) = \int_t^{T_i} |\sigma_{i,j}|^2(s) ds.$$

c) Integrate the self-financing condition (4.5.28) between 0 and t .

d) We have

$$\begin{aligned}
d\tilde{V}_t &= d(e^{-\int_0^t r_s ds} V_t) \\
&= -r_t e^{-\int_0^t r_s ds} dt + e^{-\int_0^t r_s ds} dV_t \\
&= -r_t e^{-\int_0^t r_s ds} \sum_{k=i}^j \xi_t^{(k)} P(t, T_k) dt + e^{-\int_0^t r_s ds} \sum_{k=i}^j \xi_t^{(k)} dP(t, T_k) \\
&= \sum_{k=i}^j \xi_t^{(k)} d\tilde{P}(t, T_k), \quad 0 \leq t \leq T_i.
\end{aligned}$$

since

$$\frac{d\tilde{P}(t, T_k)}{\tilde{P}(t, T_k)} = \zeta_k(t) dt, \quad k = i, \dots, j.$$

e) We apply the Itô formula and the fact that

$$t \mapsto \mathbf{E}_{i,j} [(S(T_i, T_i, T_j) - \kappa)^+ | \mathcal{F}_t]$$

and $(S_t)_{t \in \mathbb{R}_+}$ are both martingales under $\mathbb{P}_{i,j}$.

f) Use the fact that

$$\widehat{V}_t = \mathbf{E}_{i,j} [(S(T_i, T_i, T_j) - \kappa)^+ | \mathcal{F}_t],$$

and apply the result of Question (e)).

g) Apply the Itô rule to $V_t = P(t, T_i, T_j) \widehat{V}_t$ using Relation (4.5.25) and the result of Question (f)).

h) From the expression (4.5.27) of the swap rate volatilities we have

$$dV_t = S_t \frac{\partial C}{\partial x}(S_t, v(t, T_i))$$



$$\begin{aligned}
& \times \left(\sum_{k=i}^{j-1} (T_{k+1} - T_k) P(t, T_{k+1}) (\zeta_i(t) - \zeta_{k+1}(t)) + P(t, T_j) (\zeta_i(t) - \zeta_j(t)) \right) dB_t \\
& + \widehat{V}_t dP(t, T_i, T_j) \\
= & S_t \frac{\partial C}{\partial x} (S_t, v(t, T_i)) \\
& \times \left(\sum_{k=i}^{j-1} (T_{k+1} - T_k) P(t, T_{k+1}) (\zeta_i(t) - \zeta_{k+1}(t)) + P(t, T_j) (\zeta_i(t) - \zeta_j(t)) \right) dB_t \\
& + \widehat{V}_t \sum_{k=i}^{j-1} (T_{k+1} - T_k) \zeta_{k+1}(t) P(t, T_{k+1}) dB_t \\
= & S_t \frac{\partial C}{\partial x} (S_t, v(t, T_i)) \sum_{k=i}^{j-1} (T_{k+1} - T_k) P(t, T_{k+1}) (\zeta_i(t) - \zeta_{k+1}(t)) dB_t \\
& + \frac{\partial C}{\partial x} (S_t, v(t, T_i)) P(t, T_j) (\zeta_i(t) - \zeta_j(t)) dB_t \\
& + \widehat{V}_t \sum_{k=i}^{j-1} (T_{k+1} - T_k) \zeta_{k+1}(t) P(t, T_{k+1}) dB_t \\
= & S_t \zeta_i(t) \frac{\partial C}{\partial x} (S_t, v(t, T_i)) \sum_{k=i}^{j-1} (T_{k+1} - T_k) P(t, T_{k+1}) dB_t \\
& - S_t \frac{\partial C}{\partial x} (S_t, v(t, T_i)) \sum_{k=i}^{j-1} (T_{k+1} - T_k) P(t, T_{k+1}) \zeta_{k+1}(t) dB_t \\
& + \frac{\partial C}{\partial x} (S_t, v(t, T_i)) P(t, T_j) (\zeta_i(t) - \zeta_j(t)) dB_t \\
& + \widehat{V}_t \sum_{k=i}^{j-1} (T_{k+1} - T_k) \zeta_{k+1}(t) P(t, T_{k+1}) dB_t \\
= & S_t \zeta_i(t) \frac{\partial C}{\partial x} (S_t, v(t, T_i)) \sum_{k=i}^{j-1} (T_{k+1} - T_k) P(t, T_{k+1}) dB_t \\
& + \left(\widehat{V}_t - S_t \frac{\partial C}{\partial x} (S_t, v(t, T_i)) \right) \sum_{k=i}^{j-1} (T_{k+1} - T_k) P(t, T_{k+1}) \zeta_{k+1}(t) dB_t \\
& + \frac{\partial C}{\partial x} (S_t, v(t, T_i)) P(t, T_j) (\zeta_i(t) - \zeta_j(t)) dB_t.
\end{aligned}$$

i) We have

$$\begin{aligned}
d\widetilde{V}_t &= S_t \zeta_i(t) \frac{\partial C}{\partial x} (S_t, v(t, T_i)) \sum_{k=i}^{j-1} (T_{k+1} - T_k) \widetilde{P}(t, T_{k+1}) dB_t \\
&+ \left(\widehat{V}_t - S_t \frac{\partial C}{\partial x} (S_t, v(t, T_i)) \right) \sum_{k=i}^{j-1} (T_{k+1} - T_k) \widetilde{P}(t, T_{k+1}) \zeta_{k+1}(t) dB_t \\
&+ \frac{\partial C}{\partial x} (S_t, v(t, T_i)) \widetilde{P}(t, T_j) (\zeta_i(t) - \zeta_j(t)) dB_t \\
= & (\widetilde{P}(t, T_i) - \widetilde{P}(t, T_j)) \zeta_i(t) \frac{\partial C}{\partial x} (S_t, v(t, T_i)) dB_t \\
&+ \left(\widehat{V}_t - S_t \frac{\partial C}{\partial x} (S_t, v(t, T_i)) \right) d\widetilde{P}(t, T_i, T_j) \\
&+ \frac{\partial C}{\partial x} (S_t, v(t, T_i)) \widetilde{P}(t, T_j) (\zeta_i(t) - \zeta_j(t)) dB_t
\end{aligned}$$

$$\begin{aligned}
&= (\zeta_i(t)\tilde{P}(t, T_i) - \zeta_j(t)\tilde{P}(t, T_j)) \frac{\partial C}{\partial x}(S_t, v(t, T_i)) dB_t \\
&\quad + \left(\widehat{V}_t - S_t \frac{\partial C}{\partial x}(S_t, v(t, T_i)) \right) d\tilde{P}(t, T_i, T_j) \\
&= \frac{\partial C}{\partial x}(S_t, v(t, T_i)) d(\tilde{P}(t, T_i) - \tilde{P}(t, T_j)) \\
&\quad + \left(\widehat{V}_t - S_t \frac{\partial C}{\partial x}(S_t, v(t, T_i)) \right) d\tilde{P}(t, T_i, T_j).
\end{aligned}$$

j) We have

$$\begin{aligned}
\frac{\partial C}{\partial x}(x, \tau, v) &= \frac{\partial}{\partial x} \left(x\Phi\left(\frac{v}{2} + \frac{1}{v} \log \frac{x}{\kappa}\right) - \kappa\Phi\left(-\frac{v}{2} + \frac{1}{v} \log \frac{x}{\kappa}\right) \right) \\
&= x \frac{\partial}{\partial x} \Phi\left(\frac{v}{2} + \frac{1}{v} \log \frac{x}{\kappa}\right) - \kappa \frac{\partial}{\partial x} \Phi\left(-\frac{v}{2} + \frac{1}{v} \log \frac{x}{\kappa}\right) + \Phi\left(\frac{v}{2} + \frac{1}{v} \log \frac{x}{\kappa}\right) \\
&= \frac{1}{v\sqrt{2\pi}} e^{-(v/2+v^{-1}\log(x/\kappa))^2/2} - \frac{\kappa}{vx\sqrt{2\pi}} e^{-(v/2+v^{-1}\log(x/\kappa))^2/2} \\
&\quad + \Phi\left(\frac{\log(x/\kappa)}{v} + \frac{v}{2}\right) = \Phi\left(\frac{\log(x/\kappa)}{v} + \frac{v}{2}\right).
\end{aligned}$$

k) We have

$$\begin{aligned}
d\tilde{V}_t &= \frac{\partial C}{\partial x}(S_t, v(t, T_i)) d(\tilde{P}(t, T_i) - \tilde{P}(t, T_j)) \\
&\quad + \left(\widehat{V}_t - S_t \frac{\partial C}{\partial x}(S_t, v(t, T_i)) \right) d\tilde{P}(t, T_i, T_j) \\
&= \Phi\left(\frac{\log(S_t/K)}{v(t, T_i)} + \frac{v(t, T_i)}{2}\right) d(\tilde{P}(t, T_i) - \tilde{P}(t, T_j)) \\
&\quad - \kappa\Phi\left(\frac{\log(S_t/K)}{v(t, T_i)} - \frac{v(t, T_i)}{2}\right) d\tilde{P}(t, T_i, T_j).
\end{aligned}$$

l) We compare the results of Questions (d) and (k)).

Chapter 5

Exercise 5.1 By absence of arbitrage we have $(1 - \alpha) e^{-r_d T} = e^{-rT}$, hence $\alpha = 1 - e^{-(r - r_d)T}$.

Exercise 5.2

- a) The bond payoff $\mathbb{1}_{\{\tau > T-t\}}$ is discounted according to the risk-free rate, before taking expectation.
- b) We have $\mathbb{E}[\mathbb{1}_{\{\tau > T-t\}}] = e^{-\lambda(T-t)}$, hence $P_d(t, T) = e^{-(\lambda+r)(T-t)}$.
- c) We have $P_M(t, T) = e^{-(\lambda+r)(T-t)}$, hence $\lambda = -r + \frac{1}{T-t} \log P_M(t, T)$.

Exercise 5.3

- a) Use the fact that $(r_t, \lambda_t)_{t \in [0, T]}$ is a Markov process.
- b) Use the tower property of the conditional expectation given \mathcal{F}_t .
- c) Writing $F(t, r_t, \lambda_t) = P(t, T)$, we have

$$\begin{aligned}
&d \left(e^{-\int_0^t (r_s + \lambda_s) ds} P(t, T) \right) \\
&= -(r_t + \lambda_t) e^{-\int_0^t (r_s + \lambda_s) ds} P(t, T) dt + e^{-\int_0^t (r_s + \lambda_s) ds} dP(t, T) \\
&= -(r_t + \lambda_t) e^{-\int_0^t (r_s + \lambda_s) ds} P(t, T) dt + e^{-\int_0^t (r_s + \lambda_s) ds} dF(t, r_t, \lambda_t)
\end{aligned}$$



$$\begin{aligned}
&= -(r_t + \lambda_t) e^{-\int_0^t (r_s + \lambda_s) ds} P(t, T) dt + e^{-\int_0^t (r_s + \lambda_s) ds} \frac{\partial F}{\partial x}(t, r_t, \lambda_t) dr_t \\
&\quad + e^{-\int_0^t (r_s + \lambda_s) ds} \frac{\partial F}{\partial y}(t, r_t, \lambda_t) d\lambda_t + \frac{1}{2} e^{-\int_0^t (r_s + \lambda_s) ds} \frac{\partial^2 F}{\partial x^2}(t, r_t, \lambda_t) \sigma_1^2(t, r_t) dt \\
&\quad + \frac{1}{2} e^{-\int_0^t (r_s + \lambda_s) ds} \frac{\partial^2 F}{\partial y^2}(t, r_t, \lambda_t) \sigma_2^2(t, \lambda_t) dt \\
&\quad + e^{-\int_0^t (r_s + \lambda_s) ds} \rho \frac{\partial^2 F}{\partial x \partial y}(t, r_t, \lambda_t) \sigma_1(t, r_t) \sigma_2(t, \lambda_t) dt + e^{-\int_0^t (r_s + \lambda_s) ds} \frac{\partial F}{\partial t}(t, r_t, \lambda_t) dt \\
&= e^{-\int_0^t (r_s + \lambda_s) ds} \frac{\partial F}{\partial x}(t, r_t, \lambda_t) \sigma_1(t, r_t) dB_t^{(1)} + e^{-\int_0^t (r_s + \lambda_s) ds} \frac{\partial F}{\partial y}(t, r_t, \lambda_t) \sigma_2(t, \lambda_t) dB_t^{(2)} \\
&\quad + e^{-\int_0^t (r_s + \lambda_s) ds} \left(-(r_t + \lambda_t) P(t, T) + \frac{\partial F}{\partial x}(t, r_t, \lambda_t) \mu_1(t, r_t) \right. \\
&\quad \left. + \frac{\partial F}{\partial y}(t, r_t, \lambda_t) \mu_2(t, \lambda_t) + \frac{1}{2} \frac{\partial^2 F}{\partial x^2}(t, r_t, \lambda_t) \sigma_1^2(t, r_t) + \frac{1}{2} \frac{\partial^2 F}{\partial y^2}(t, r_t, \lambda_t) \sigma_2^2(t, \lambda_t) \right. \\
&\quad \left. + \rho \frac{\partial^2 F}{\partial x \partial y}(t, r_t, \lambda_t) \sigma_1(t, r_t) \sigma_2(t, \lambda_t) + \frac{\partial F}{\partial t}(t, r_t, \lambda_t) \right) dt,
\end{aligned}$$

hence the bond pricing PDE is

$$\begin{aligned}
&- (x + y) F(t, x, y) + \mu_1(t, x) \frac{\partial F}{\partial x}(t, x, y) \\
&+ \mu_2(t, y) \frac{\partial F}{\partial y}(t, x, y) + \frac{1}{2} \sigma_1^2(t, x) \frac{\partial^2 F}{\partial x^2}(t, x, y) \\
&+ \frac{1}{2} \sigma_2^2(t, y) \frac{\partial^2 F}{\partial y^2}(t, x, y) + \rho \sigma_1(t, x) \sigma_2(t, y) \frac{\partial^2 F}{\partial x \partial y}(t, x, y) + \frac{\partial F}{\partial t}(t, r_t, \lambda_t) = 0.
\end{aligned}$$

d) We have

$$r_t = -a \int_0^t r_s ds + \sigma B_t^{(1)}, \quad t \geq 0,$$

hence

$$\begin{aligned}
\int_0^t r_s ds &= \frac{1}{a} (\sigma B_t^{(1)} - r_t) \\
&= \frac{\sigma}{a} \left(B_t^{(1)} - \int_0^t e^{-(t-s)a} dB_s^{(1)} \right) \\
&= \frac{\sigma}{a} \int_0^t (1 - e^{-(t-s)a}) dB_s^{(1)},
\end{aligned}$$

and

$$\begin{aligned}
\int_t^T r_s ds &= \int_0^T r_s ds - \int_0^t r_s ds \\
&= \frac{\sigma}{a} \int_0^T (1 - e^{-(T-s)a}) dB_s^{(1)} - \frac{\sigma}{a} \int_0^t (1 - e^{-(t-s)a}) dB_s^{(1)} \\
&= -\frac{\sigma}{a} \left(\int_0^t (e^{-(T-s)a} - e^{-(t-s)a}) dB_s^{(1)} + \int_t^T (e^{-(T-s)a} - 1) dB_s^{(1)} \right) \\
&= -\frac{\sigma}{a} (e^{-(T-t)a} - 1) \int_0^t e^{-(t-s)a} dB_s^{(1)} - \frac{\sigma}{a} \int_t^T (e^{-(T-s)a} - 1) dB_s^{(1)} \\
&= -\frac{1}{a} (e^{-(T-t)a} - 1) r_t - \frac{\sigma}{a} \int_t^T (e^{-(T-s)a} - 1) dB_s^{(1)}.
\end{aligned}$$

The answer for λ_t is similar.

e) As a consequence of the previous question we have

$$\mathbb{E} \left[\int_t^T r_s ds + \int_t^T \lambda_s ds \mid \mathcal{F}_t \right] = C(a, t, T) r_t + C(b, t, T) \lambda_t,$$

and

$$\begin{aligned}
& \text{Var} \left[\int_t^T r_s ds + \int_t^T \lambda_s ds \mid \mathcal{F}_t \right] = \\
&= \text{Var} \left[\int_t^T r_s ds \mid \mathcal{F}_t \right] + \text{Var} \left[\int_t^T \lambda_s ds \mid \mathcal{F}_t \right] \\
&\quad + 2 \text{Cov} \left(\int_t^T X_s ds, \int_t^T Y_s ds \mid \mathcal{F}_t \right) \\
&= \frac{\sigma^2}{a^2} \int_t^T (\mathrm{e}^{-(T-s)a} - 1)^2 ds \\
&\quad + 2\rho \frac{\sigma\eta}{ab} \int_t^T (\mathrm{e}^{-(T-s)a} - 1)(\mathrm{e}^{-(T-s)b} - 1) ds \\
&\quad + \frac{\eta^2}{b^2} \int_t^T (\mathrm{e}^{-(T-s)b} - 1)^2 ds \\
&= \sigma^2 \int_t^T C^2(a, s, T) ds + 2\rho\sigma\eta \int_t^T C(a, s, T)C(b, s, T) ds \\
&\quad + \eta^2 \int_t^T C^2(b, s, T) ds,
\end{aligned}$$

from the Itô isometry.

f) We have

$$\begin{aligned}
P(t, T) &= \mathbb{1}_{\{\tau>t\}} \mathbf{E} \left[\exp \left(- \int_t^T r_s ds - \int_t^T \lambda_s ds \right) \mid \mathcal{F}_t \right] \\
&= \mathbb{1}_{\{\tau>t\}} \exp \left(- \mathbf{E} \left[\int_t^T r_s ds \mid \mathcal{F}_t \right] - \mathbf{E} \left[\int_t^T \lambda_s ds \mid \mathcal{F}_t \right] \right) \\
&\quad \times \exp \left(\frac{1}{2} \text{Var} \left[\int_t^T r_s ds + \int_t^T \lambda_s ds \mid \mathcal{F}_t \right] \right) \\
&= \mathbb{1}_{\{\tau>t\}} \exp(-C(a, t, T)r_t - C(b, t, T)\lambda_t) \\
&\quad \times \exp \left(\frac{\sigma^2}{2} \int_t^T C^2(a, s, T) ds + \frac{\eta^2}{2} \int_t^T C^2(b, s, T) \mathrm{e}^{-(T-s)b} ds \right) \\
&\quad \times \exp \left(\rho\sigma\eta \int_t^T C(a, s, T)C(b, s, T) ds \right).
\end{aligned}$$

g) This is a direct consequence of the answers to Questions (c)) and f).

h) The above analysis shows that

$$\begin{aligned}
\mathbb{P}(\tau > T \mid \mathcal{G}_t) &= \mathbb{1}_{\{\tau>t\}} \mathbf{E} \left[\exp \left(- \int_t^T \lambda_s ds \right) \mid \mathcal{F}_t \right] \\
&= \mathbb{1}_{\{\tau>t\}} \exp \left(-C(b, t, T)\lambda_t + \frac{\eta^2}{2} \int_t^T C^2(b, s, T) ds \right),
\end{aligned}$$

for $a = 0$ and

$$\mathbf{E} \left[\exp \left(- \int_t^T r_s ds \right) \mid \mathcal{F}_t \right] = \exp \left(-C(a, t, T)r_t + \frac{\sigma^2}{2} \int_t^T C^2(a, s, T) ds \right),$$

for $b = 0$, and this implies

$$\begin{aligned}
U_\rho(t, T) &= \exp \left(\rho\sigma\eta \int_t^T C(a, s, T)C(b, s, T) ds \right) \\
&= \exp \left(\rho \frac{\sigma\eta}{ab} (T - t - C(a, t, T) - C(b, t, T) + C(a+b, t, T)) \right).
\end{aligned}$$

i) We have

$$\begin{aligned}
f(t, T) &= -\mathbb{1}_{\{\tau>t\}} \frac{\partial}{\partial T} \log P(t, T) \\
&= \mathbb{1}_{\{\tau>t\}} \left(r_t \mathrm{e}^{-(T-t)a} - \frac{\sigma^2}{2} C^2(a, t, T) + \lambda_t \mathrm{e}^{-(T-t)b} - \frac{\eta^2}{2} C^2(b, t, T) \right)
\end{aligned}$$



$$-\mathbb{1}_{\{\tau>t\}}\rho\sigma\eta C(a,t,T)C(b,t,T).$$

j) We use the relation

$$\begin{aligned}\mathbb{P}(\tau > T \mid \mathcal{G}_t) &= \mathbb{1}_{\{\tau>t\}} \mathbf{E} \left[\exp \left(- \int_t^T \lambda_s ds \right) \mid \mathcal{F}_t \right] \\ &= \mathbb{1}_{\{\tau>t\}} \exp \left(-C(b,t,T)\lambda_t + \frac{\eta^2}{2} \int_t^T C^2(b,s,T)ds \right) \\ &= \mathbb{1}_{\{\tau>t\}} e^{-\int_t^T f_2(t,u)du},\end{aligned}$$

where $f_2(t, T)$ is the Vasicek forward rate corresponding to λ_t , i.e.

$$f_2(t, u) = \lambda_t e^{-(u-t)b} - \frac{\eta^2}{2} C^2(b, t, u).$$

k) In this case we have $\rho = 0$ and

$$P(t, T) = \mathbb{P}(\tau > T \mid \mathcal{G}_t) \mathbf{E} \left[\exp \left(- \int_t^T r_s ds \right) \mid \mathcal{F}_t \right],$$

since $U_\rho(t, T) = 0$.

Bibliography

Articles

- [AL05] C. Albanese and S. Lawi. “Laplace transforms for integrals of Markov processes”. In: *Markov Process. Related Fields* 11.4 (2005), pages 677–724 (Cited on page [4](#)).
- [Bla76] F. Black. “The pricing of commodity contracts”. In: *J. of Financial Economics* 3 (1976), pages 167–179 (Cited on page [100](#)).
- [BDT90] F. Black, E. Derman, and W. Toy. “A One-Factor Model of Interest Rates and Its Application to Treasury Bond Options”. In: *Financial Analysts Journal* 46.1 (1990), pages 24–32 (Cited on page [32](#)).
- [BGM97] A. Brace, D. Gatarek, and M. Musiela. “The market model of interest rate dynamics”. In: *Math. Finance* 7.2 (1997), pages 127–155 (Cited on page [59](#)).
- [BHM18] D.C. Brody, L.P. Hughston, and D.M. Meier. “Lévy-Vasicek Models and the Long-Bond Return Process”. In: *Int. J. Theor. Appl. Finance* 21.3 (2018), page 1850026 (Cited on pages [27](#) and [137](#)).
- [CN10] C. Pintoux and N. Privault. “A direct solution to the Fokker-Planck equation for exponential Brownian functionals”. In: *Analysis and Applications* 8.3 (2010), pages 287–304 (Cited on page [24](#)).
- [CN11] C. Pintoux and N. Privault. “The Dothan pricing model revisited”. In: *Math. Finance* 21 (2011), pages 355–363 (Cited on page [22](#)).
- [Cha+92] K.C. Chan et al. “An Empirical Comparison of Alternative Models of the Short-Term Interest Rate”. In: *The Journal of Finance* 47.3 (1992). Papers and Proceedings of the Fifty-Second Annual Meeting of the American Finance Association, New Orleans, Louisiana, pages 1209–1227 (Cited on pages [27](#) and [28](#)).
- [Che+08] R.-R. Chen, X. Cheng, et al. “An Explicit, Multi-Factor Credit Default Swap Pricing Model with Correlated Factors”. In: *Journal of Financial and Quantitative Analysis* 43.1 (2008), pages 123–160 (Cited on page [129](#)).

- [Cou82] G. Courtadon. “The Pricing of Options on Default-Free Bonds”. In: *The Journal of Financial and Quantitative Analysis* 17.1 (1982), pages 75–100 (Cited on pages [5](#) and [28](#)).
- [CIR85] J.C. Cox, J.E. Ingersoll, and S.A. Ross. “A Theory of the Term Structure of Interest Rates”. In: *Econometrica* 53 (1985), pages 385–407 (Cited on pages [4](#), [6](#), [16](#), [25](#), and [28](#)).
- [Dot78] L.U. Dothan. “On the term structure of interest rates”. In: *Jour. of Fin. Ec.* 6 (1978), pages 59–69 (Cited on pages [6](#) and [22](#)).
- [EJ99] R.J. Elliott and M. Jeanblanc. “Incomplete markets with jumps and informed agents”. In: *Math. Methods Oper. Res.* 50.3 (1999), pages 475–492 (Cited on page [128](#)).
- [EJY00] R.J. Elliott, M. Jeanblanc, and M. Yor. “On models of default risk”. In: *Math. Finance* 10.2 (2000), pages 179–195 (Cited on page [127](#)).
- [FG06] R. Faff and P. Gray. “On the estimation and comparison of short-rate models using the generalised method of moments”. In: *Journal of Banking and Finance* 30 (11 2006), pages 3131–3146 (Cited on page [137](#)).
- [Fel51] W. Feller. “Two singular diffusion problems”. In: *Ann. of Math.* (2) 54 (1951), pages 173–182 (Cited on page [4](#)).
- [GK83] M.B. Garman and S.W. Kohlhagen. “Foreign Currency Option Values”. In: *J. International Money and Finance* 2 (1983), pages 231–237 (Cited on pages [78](#), [79](#), and [80](#)).
- [GER95] H. Geman, N. El Karoui, and J.-C. Rochet. “Changes of numéraire, changes of probability measure and option pricing”. In: *J. Appl. Probab.* 32.2 (1995), pages 443–458 (Cited on pages [68](#) and [85](#)).
- [HJM92] D. Heath, R. Jarrow, and A. Morton. “Bond pricing and the term structure of interest rates: a new methodology”. In: *Econometrica* 60 (1992), pages 77–105 (Cited on page [48](#)).
- [HL86] S.Y. Ho and S.B. Lee. “Term structure movements and pricing interest rate contingent claims”. In: *Journal of Finance* 41 (5 1986), pages 1011–1029 (Cited on page [6](#)).
- [HW90] J. Hull and A. White. “Pricing interest rate derivative securities”. In: *The Review of Financial Studies* 3 (1990), pages 537–592 (Cited on pages [6](#) and [48](#)).
- [Jam89] F. Jamshidian. “An exact bond option formula”. In: *The Journal of Finance* XLIV.1 (1989), pages 205–209 (Cited on pages [97](#) and [112](#)).
- [Jam96] F. Jamshidian. “Sorting out swaptions”. In: *Risk Magazine* 9.3 (1996), pages 59–60 (Cited on page [85](#)).
- [Kak15] Z. Kakushadze. “Path integral and asset pricing”. In: *Quant. Finance* 15.11 (2015), pages 1759–1771 (Cited on page [26](#)).
- [Kim02] Y.-J. Kim. “Option Pricing under Stochastic Interest Rates: An Empirical Investigation”. In: *Asia-Pacific Financial Markets* 9 (1 2002), pages 23–44 (Cited on page [98](#)).
- [Lan98] D. Lando. “On Cox processes and credit risky securities”. In: *Review of Derivative Research* 2 (1998), pages 99–120 (Cited on pages [125](#), [126](#), and [128](#)).
- [Lin07] E. Lindström. “Estimating parameters in diffusion processes using an approximate maximum likelihood approach”. In: *Annals of Operations Research* 151 (2007), pages 269–288 (Cited on page [138](#)).



- [Mam04] R.S. Mamon. “Three ways to solve for bond prices in the Vasicek model”. In: *Journal of Applied Mathematics and Decision Sciences* 8.1 (2004), pages 1–14 (Cited on page 149).
- [Mar78] W. Margrabe. “The value of an option to exchange one asset for another”. In: *The Journal of Finance* XXXIII.1 (1978), pages 177–186 (Cited on pages 83 and 87).
- [MR83] T.A. Marsh and E.R. Rosenfeld. “Stochastic Processes for Interest Rates and Equilibrium Bond Prices”. In: *The Journal of Finance* 38.2 (1983). Papers and Proceedings Forty-First Annual Meeting American Finance Association New York, N.Y., pages 635–646 (Cited on pages 5 and 28).
- [Mer73] R. Merton. “Theory of rational option pricing”. In: *Bell Journal of Economics* 4.1 (1973), pages 141–183 (Cited on page 84).
- [NS87] C.R. Nelson and A.F. Siegel. “Parsimonious Modeling of Yield Curves”. In: *Journal of Business* 60 (1987), pages 473–489 (Cited on pages 35 and 52).
- [PP17] A. Prayoga and N. Privault. “Pricing CIR yield options by conditional moment matching”. In: *Asia-Pacific Financial Markets* 24 (2017), pages 19–38 (Cited on page 25).
- [PT12] N. Privault and T.-R. Teng. “Risk-neutral hedging in bond markets”. In: *Risk and Decision Analysis* 3 (2012), pages 201–209 (Cited on pages 85, 100, and 107).
- [PU13] N. Privault and W.T. Uy. “Monte Carlo Computation of the Laplace Transform of Exponential Brownian Functionals”. In: *Methodol. Comput. Appl. Probab.* 15.3 (2013), pages 511–524 (Cited on page 24).
- [PW09] N. Privault and X. Wei. “Calibration of the LIBOR market model - implementation in PREMIA”. In: *Bankers, Markets & Investors* 99 (2009), pages 20–28 (Cited on page 108).
- [PY16] N. Privault and J.D. Yu. “Stratified approximations for the pricing of options on average”. In: *Journal of Computational Finance* 19.4 (2016), pages 95–113 (Cited on page 24).
- [Pro01] P. Protter. “A partial introduction to financial asset pricing theory”. In: *Stochastic Process. Appl.* 91.2 (2001), pages 169–203 (Cited on page 85).
- [SS01] P. Santa-Clara and D. Sornette. “The Dynamics of the Forward Interest Rate Curve with Stochastic String Shocks”. In: *The Review of Financial Studies* 14.1 (2001), pages 149–185 (Cited on page 63).
- [Vaš77] O. Vašíček. “An equilibrium characterisation of the term structure”. In: *Journal of Financial Economics* 5 (1977), pages 177–188 (Cited on pages 2, 15, 16, 39, 96, 138, and 153).
- [Wu00] X. Wu. “A New Stochastic Duration Based on the Vasicek and CIR Term Structure Theories”. In: *Journal of Business Finance and Accounting* 27 (2000) (Cited on page 33).
- [Yor92] M. Yor. “On some exponential functionals of Brownian motion”. In: *Adv. in Appl. Probab.* 24.3 (1992), pages 509–531 (Cited on page 23).

Books

- [Bou73] K.E. Boulding. In “*Energy Reorganization Act of 1973. Hearings, Ninety-third Congress, first session, on H.R. 11510*”. Washington: U.S. Government Printing Office, 1973, pages iv+422 (Cited on page 66).

- [BM06] D. Brigo and F. Mercurio. *Interest rate models—theory and practice*. Second. Springer Finance. Berlin: Springer-Verlag, 2006, pages lvi+981 (Cited on pages [16](#), [58](#), and [141](#)).
- [Cha14] A. Charpentier, editor. *Computational Actuarial Science with R*. The R Series. USA: Chapman & Hall/CRC, 2014 (Cited on pages [41](#) and [55](#)).
- [Das04] J.W. Dash. *Quantitative finance and risk management*. River Edge, NJ: World Scientific Publishing Co. Inc., 2004, pages xx+781 (Cited on page [71](#)).
- [DMM92] C. Dellacherie, B. Maisonneuve, and P.A. Meyer. *Probabilités et Potentiel*. Volume 5. Hermann, 1992. Chapter XVII-XXIV (Cited on page [127](#)).
- [DJD08] A. Downes, M.S. Joshi, and N. Denson. *Quant Job Interview Questions and Answers*. first. CreateSpace Independent Publishing Platform, 2008, pages x+316 (Cited on page [89](#)).
- [DS03] D. Duffie and K.J. Singleton. *Credit risk. Pricing, measurement, and management*. Princeton Series in Finance. Princeton, NJ: Princeton University Press, 2003, pages xvi+396 (Cited on page [128](#)).
- [Jeu80] Th. Jeulin. *Semi-martingales et grossissement d'une filtration*. Volume 833. Lecture Notes in Mathematics. Springer Verlag, 1980 (Cited on page [128](#)).
- [Pri12] N. Privault. *An Elementary Introduction to Stochastic Interest Rate Modeling (Second Edition)*. Advanced Series on Statistical Science & Applied Probability, 16. Singapore: World Scientific Publishing Co., 2012, pages xiii+228 (Cited on pages [48](#), [54](#), [56](#), [58](#), [98](#), [108](#), and [119](#)).
- [Pri14] N. Privault. *Stochastic Finance: An Introduction with Market Examples*. Financial Mathematics Series. Chapman & Hall/CRC, 2014, page 441 (Cited on pages [125](#), [127](#), [128](#), and [129](#)).
- [Pro04] P. Protter. *Stochastic integration and differential equations*. second. Volume 21. Stochastic Modelling and Applied Probability. Berlin: Springer-Verlag, 2004, pages xiv+419 (Cited on pages [14](#), [15](#), [73](#), [83](#), and [127](#)).
- [Sch05] J. Schoenmakers. *Robust LIBOR modelling and pricing of derivative products*. Chapman & Hall/CRC Financial Mathematics Series. Chapman & Hall/CRC, Boca Raton, FL, 2005, pages xvi+202 (Cited on pages [107](#) and [108](#)).
- [Shr04] S.E. Shreve. *Stochastic calculus for finance. II*. Springer Finance. Continuous-time models. New York: Springer-Verlag, 2004, pages xx+550 (Cited on page [92](#)).
- [Wat95] G. N. Watson. *A treatise on the theory of Bessel functions*. Reprint of the second (1944) edition. Cambridge: Cambridge University Press, 1995, pages viii+804 (Cited on page [24](#)).

Index

A

- absence of arbitrage 11
- abstract Bayes formula 69
- accreting swap 45
- adapted process 125
- affine model 6
- affine PDE 6, 141
- amortizing swap 45
- annuity
 - measure 102, 118, 120
 - numéraire 102, 117
- annuity numéraire 45
- approximation
 - gamma 24
- asset pricing
 - first theorem
 - continuous time, 48

B

- basis point 112
- BDT model 32
- Bermudan swaption 108
- Bessel function 4, 24
- BGM model 59, 99
- binary option 91
- Black
 - (1976) formula 100
 - caplet formula 99

- swaption formula 106
- Black-Derman-Toy model 32
- Black-Scholes
 - formula 59, 80
 - call options, 100
 - put options, 100
- bond
 - convertible 29
 - convexity 33
 - corporate 29
 - defaultable 128
 - duration 29, 33
 - immunization 144
 - ladder 102
 - option 90, 97, 163
 - pricing
 - PDE, 14, 57
 - pricing PDE 185
 - yield 21
 - zero-coupon 10
- break-even
 - rate 45, 46
- Bretton Woods 66
- bridge model 61
- Brownian
 - bridge 26
- buy back guarantee 82

C

- call
 - swaption 105
 - call-put parity 80
 - cap pricing 101
 - caplet pricing 98
 - cash-or-nothing option 91
 - CBOE 3
 - CEV model 5
 - change of numéraire 68, 80
 - Chi square distribution 4
 - Chicago Board Options Exchange 3
 - CIR model 4, 28
 - CKLS model 27
 - compounded yield to maturity 32
 - compounding
 - linear 43
 - conditional
 - survival probability 123
 - conversion rate 29
 - convertible bond 29
 - convexity 33
 - corporate bond 29
 - correlation
 - perfect 56, 63
 - problem 55
 - cost of carry 77
 - coupon
 - bond 11
 - rate 20
 - Courtadon model 5, 28
 - credit default
 - reduced-form approach 123

D

- default rate 125
- defaultable bonds 131
- deflated price 67
- Delta
 - hedging 86, 87
- derivatives
 - fixed income 93
 - interest rate 93
- diffusion
 - elasticity 6, 28
- digital option 91
- dispersion index 25
- distribution

- invariant 2, 5
- lognormal 24
- stationary 2, 5

- dollar value 32
- Dothan model 6, 22
- duration 29, 33

E

- ECB 41
- elasticity of diffusion 6, 28
- enlarge of filtration 128
- Euclidean path integral 26
- EURIBOR 43
- exchange option 83
- exponential
 - distribution 125
 - Vasicek model 5

F

- face value 10, 32
- failure rate 124
- Fano factor 25
- FED 43
- Feller condition 4
- filtration
 - enlarge of 128
- first theorem of asset pricing 48
- fixed
 - income 1
 - derivatives, 93
 - leg 44
 - rate 98
- floating
 - leg 44
 - rate 98
- floorlet 100, 113
- foreign exchange
 - option 78
- forward
 - contract 35, 89, 159
 - measure 30, 94
 - price 67
 - rate 35
 - agreement, 35
 - spot, 35–37, 98
 - swap, 44
 - swap rate 44
- forward measure 68

forward swap
measure 102, 118, 120
FRA 35

G

gamma
approximation 24
Garman-Kohlagen formula 78
Gaussian
cumulative distribution function 97
Girsanov Theorem 73
guarantee
buy back 82

H

hedge and forget 160
hedging
change of numéraire 85
static 160
HIBOR 43
HJM
condition 48
model 47, 127
Ho-Lee model 6
Hull-White model 6, 48

I

immunization 144
instantaneous forward rate 37
interest rate
derivatives 93
model
affine, 6
Constant Elasticity of Variance, 5
Courtadon, 5, 28
Cox-Ingersoll-Ross, 4
Dothan, 6, 22
exponential Vasicek, 5
Ho-Lee, 6
Hull-White, 6, 48
Marsh-Rosenfeld, 5, 28
Vasicek, 2, 7
invariant distribution 2, 5
IPython notebook 53

J

Jamshidian's trick 113

K

key lemma 127

L

Lagrangian 25
least square regression 7
leg
fixed 44
floating 44
Leibniz integral rule 49
LIBOR
model 43
rate 43
swap rate 45, 104, 106
Lipschitz function 83
lognormal
distribution 24

M

Macaulay duration 33
Margrabe formula 83
market
price of risk 14
Markov property 83, 86
Marsh-Rosenfeld model 5, 28
maturity
transformation 36
mean
reversion 1
model
affine 6
BDT 32
CEV 5
CIR 4, 28
CKLS 27
Courtadon 5, 28
Dothan 6, 22
exponential Vasicek 5
Ho-Lee 6
Hull-White 6, 48
Marsh-Rosenfeld 5, 28
Vasicek 2, 7
modified
Bessel function 4, 24
duration 33
MPoR 14
Musiela notation 47

N

- Nelson-Siegel 52, 54
- nominal value 32
- noncentral Chi square 4
- notional 45
 - principal 112, 168
- numéraire 65
 - annuity 45, 102
 - invariance 85
- numéraire invariance 85

O

- OLS 7
- option
 - binary 91
 - cash-or-nothing 91
 - digital 91
- Ornstein-Uhlenbeck process 2

P

- par value 10, 32
- parity
 - call-put 80
- path
 - freezing 114
 - integral 25, 71
 - Euclidean, 26
- payer
 - swap 44
 - swaption 103
- PDE
 - affine 6, 141
- perfect correlation 56, 63
- Planck constant 25
- Poisson
 - process 125
- power option 90
- principal amount 112
- put
 - swaption 107
- Python code 53

Q

- Quantlib 108
- quantmod 9, 41

R

- R code 3, 5, 19, 41, 108
- R package
 - quantmod 9, 41
 - RQuantLib 108
 - YieldCurve 41
- Radon-Nikodym density 68
- rate
 - default 125
 - forward 35
 - forward swap 44
 - instantaneous forward 37
 - LIBOR 43, 45
 - swap, 45
 - LIBOR swap 104, 106
 - swap 44
 - receiver swaption 107
 - recovery rate 128, 129
 - reduced-form approach 123
 - Riccati equation 18, 145
- risk
 - market price 14
- RQuantLib 108

S

- self-financing portfolio
 - change of numéraire 86
- seller swap 44
- SHIBOR 43
- SIBOR 43
 - spot forward rate 35–37, 98
- static hedging 160
- stationary distribution 2, 5
- stochastic
 - default 125
- stopping time 125
- string model 63
- survival probability 123
- Svensson parametrization 52
- swap
 - amortizing 45
 - forward measure 102, 118, 120
 - payer 44
 - rate 44, 45
 - seller 44
- swaption 103
 - Bermudan 108

T

- telescoping sum 46
- tenor structure 44, 67, 93, 129
- theorem
 - asset pricing 48
 - Girsanov 73
- TIBOR 43
- tower property 13, 69, 149, 184
- treasury note 3
- two-factor model 57

V

- variable rate 98
- Vasicek model 2, 7

Y

- yield 35, 37, 98
 - bond 21
 - compounded to maturity 32
 - curve 36
 - data, 41
 - inversion, 42
- YieldCurve (R package) 41

Z

- zero-
 - coupon bond 10

Author index

- Albanese, C. 4
Björk, T. 54
Black, F. 32, 99, 100, 106
Boulding, K.E. 66
Brace, A. 59
Brigo, D. 16, 58, 141
Brody, D.C. 27
Chan, K.C. 27
Charpentier, A. 41
Chen, R.R. 129
Cheng, X. 129
Courtadon, G. 5, 28
Cox, J.C. 4
Dash, J. 71
Dellacherie, C. 127
Denson, N. 89
Derman, E. 32
Dothan, L.U. 6, 22
Downes, A. 89
El Karoui, N. 68, 85
Elliott, R.J. 127, 128
Fabozzi, F. 129
Faff, R. 137
Feller, W. 4
Garman, M.B. 78
Gatarek, D. 59
Geman, H. 68, 85
Gray, P. 137
Guirrieri, S. 41
Guo, X. 126
Heath, D. 48
Ho, S.Y. 6
Huang, J.Z. 129
Hughston, L.P. 27
Hull, J. 6
Ingersoll, J.E. 4
Jacod, J. 128
Jamshidian, F. 85, 97, 113
Jarrow, R. 48, 126
Jeanblanc, M. 127, 128
Jeulin, Th. 128
Joshi, M.S. 89
Kakushadze, Z. 26
Karolyi, G.A. 27
Kim, Y.-J. 98
Kohlhagen, S.W. 78
Lando, D. 125
Lawi, S. 4
Lee, S.B. 6
Lindström, E. 138
Liu, B. 129
Longstaff, F.A. 27
Maisonneuve, B. 127
Mamon, R.S. 149
Margrabe, W. 83
Marsh, T.A. 5, 28
Meier, D.M. 27
Menn, C. 126
Mercurio, F. 16, 58, 141
Merton, R.C. 84
Meyer, P.A. 127
Morton, A. 48
Musielka, M. 47, 59
Nelson, C.R. 52
Pintoux, C. 22, 23
Prayoga, A. 25
Protter, P. 14, 15, 73, 83, 85, 127
Rochet, J.-C. 68, 85
Rosenfeld, E.R. 5, 28
Ross, S.A. 4
Sanders, A.B. 27
Santa-Clara, P. 63
Schoenmakers, J. 107, 108
Shreve, S. 92
Siegel, A.F. 52
Sornette, D. 63
Svensson, L.E.O. 52
Teng, T.-R. 85, 100, 107
Toy, B. 32
Uy, W.I. 23
Vašíček, O. 2, 16
Wei, X. 108
White, A. 6
Wu, X. 33
Yor, M. 23, 127, 128
Yu, J.D. 24

