

Temperfect mug

Tor-Andreas Bjone
(Dated: September 19, 2020)

We looked at the difference between a regular thermos cup (Bodum) and the Temperfect mug by measuring the temperature as a function of time. We found that the temperature was higher in the Temperfect mug because it reduced the temperature difference ∇T between the liquid in the mug and the room at the beginning by absorbing heat from the liquid and performing a phase transition between solid and liquid itself. This gave it latent heat that it could later give back to the liquid in the mug by going back to a solid. The codes can be found on my GitHub: https://github.com/Trrn13P/FYS2160/tree/master/Oblig_1

I. INTRODUCTION

In this experiment we have looked at the temperature of a liquid in two thermos cups. The Bodum cup and the temperfect cup. We pour almost boiling liquid into the cups and measure the temperature as a function of time. We did not put the lids on the cups since it would be too much of a hassle to drill two holes in the top.

The Bodum cup is just a regular thermos cup. The temperfect cup has a material that melts and thereby removing a lot of heat from the liquid in the cup, and thereafter giving that heat back by going back to a solid. This energy given up during a phase transition is called latent heat.

We will look at the temperature graphs to see if the temperfect mug preforms as advertised and discuss if it can be modelled using an Einstein crystal. You can read about the Einstein crystal in appendix 2(V).

II. THEORY

The multiplicity of an Einstein crystal is given by

$$\Omega(N, q) \approx \frac{(q + N)!}{q!N!}, \quad (1)$$

where q is the number of energy units $\epsilon = hf$, where f is the frequency and h is Planck's constant, and N is the number of particles in the crystal. We use this to find the entropy

$$S = k \ln \Omega, \quad (2)$$

where k is Boltzmann's constant. We have that the internal energy of the crystal is

$$U = q\epsilon + \frac{N}{2}\epsilon, \quad (3)$$

if we include the ground-states. This of course gives us that

$$dU = \epsilon dq, \quad (4)$$

since we vary q .

The thermodynamic identity says that

$$dU = TdS - PdV + \mu dN, \quad (5)$$

where μ is the chemical potential, P is the pressure and V is the volume. We assume that the volume, pressure and number of particles stay constant. We then get that

$$T = \frac{dU}{dS} = \epsilon \frac{dq}{dS} \Leftrightarrow \frac{T}{\epsilon} = \frac{dq}{dS}. \quad (6)$$

Numerically we find this as

$$\tilde{T}_i = \frac{T_i}{\epsilon} = \frac{q_i - q_{i-1}}{S_i - S_{i-1}}. \quad (7)$$

We have that the heat capacity

$$C_V = \frac{dU}{dT} = \epsilon \frac{dq}{dT} = \frac{dq}{dT/\epsilon} = \frac{dq}{d\tilde{T}}. \quad (8)$$

We find this numerically as

$$C_{V,i} = \frac{q_i - q_{i-1}}{\tilde{T}_i - \tilde{T}_{i-1}}. \quad (9)$$

III. METHOD

We insert temperature probes into the Bodum and Temperfect mug as shown in figure (1). We start taking the temperature. Then we pour hot water into the cups.

We use eq.7 to find the temperature numerically as a function of the number of energy units in the crystal. And when we have the temperature we use eq.9 to find the heat capacity. Both expressions will simplify a bit since $dq = 1$.

IV. RESULTS

In figure (2) we can see the temperature of the Temperfect mug and the Bodum mug plotted against the time. Since the water was not poured at the same temperature this has been adjusted in the plot as seen in figure (3).

We can see from figure (2) that the Temperfect mug absorbs a lot of heat very quickly. This is because it



Figure 1. Probes measuring temperature. Temperfect mug on the left and Bodum mug on the right.




Figure 2. Temperature against time for the two mugs.




Figure 3. Closeup of the starting points where the hot water is poured in.

goes through a phase change, from solid to liquid. It then slowly goes back to solid again and gives heat back to the liquid in the mug. That makes it so that the temperature of the liquid drops more slowly. When all of the material in the mug is solid, the liquid will drop in temperature as if it was a regular mug, like the Bodum.

The reason this is beneficial is that the heat flux is related to the ∇T , so by reducing the temperature quickly, and storing it as latent heat, you can re-introduce it to the liquid later, at a lower temperature and therefore reduce the heat flux.

The assumptions in such a system that $V, P, N = \text{const}$ is reasonable. The volume change will be minimal, the number of particles will be about the same since we are good under 100°C and the pressure is said to be constant. It will probably go as $\Delta P = C\Delta T$, where C is a constant. I assume this constant is small. This happens between ca. 0 – 2000s.

V. CONCLUSION

We can use this to model how q changes in the mug, changes the temperature of the mug material. But the Einstein solid is not valid for liquids. The Temperfect mug has a material that will melt when you put hot liquid in the mug and therefore the Einstein model is not valid at that point. After a period of time the mug material will go back into solid and here the Einstein model might be valid. As energy units is transferred from the mug material to the liquid in the mug.

APPENDIX I.

In figure (2) we can see the temperature of the Temperfect mug and the Bodum mug plotted against the time. Since the water was not poured at the same temperature this has been adjusted in the plot as seen in figure (3).

We can see from figure (2) that the Temperfect mug absorbs a lot of heat very quickly. This is because it goes through a phase change, from solid to liquid. It then slowly goes back to solid again and gives heat back to the liquid in the mug. That makes it so that the temperature of the liquid drops more slowly. When all of the material in the mug is solid, the liquid will drop in temperature as if it was a regular mug, like the Bodum.

The reason this is beneficial is that the heat flux is related to the ∇T , so by reducing the temperature quickly, and storing it as latent heat, you can re-introduce it to the liquid later, at a lower temperature and therefore reduce the heat flux.

Assume constant pressure P and volume V . This means that the work $W = 0$, so that the change in internal energy $dU = dQ$. We also assume a constant number of particles N . In this condition we can use the definition of temperature (given by the total differential of the entropy S), so that the temperature T is given by

$$T^{-1} = \left(\frac{\partial S}{\partial U} \right)_{N,V},$$

Since we only have that S is a function of T we get that

$$T^{-1} = \frac{dS}{dU} = \frac{dS}{dQ} = \frac{dS}{dT} \frac{dT}{dQ}.$$

$$T^{-1} \frac{dQ}{dT} = \frac{dS}{dT}.$$

Under constant volume the heat capacity $C_V = (\partial U / \partial T)_{V,N}$, that becomes $C_V = (dQ/dT)$ in our case. We then get that

$$\frac{dS}{dT} = \frac{C_V}{T}. \quad (10)$$

In figure (4) we see the slope of the entropy dS/dT against the temperature T , calculated by this method in H₂O. The heat capacity of H₂O is listed in I. We use this to calculate the entropy. In figure (5) we see the entropy against the temperature.

Table I. Table of the heat capacity C of H₂O in kJ/kgK .

Solid	2.108
Liquid	4.184
Gas	1.996

The feature of the graph that is relevant is the part where the phase changes are happening. That is from ca. 0s to 2000s.




Figure 4. Figure of the slope of entropy dS/dT against the temperature T .




Figure 5. Figure of the entropy S against the temperature T .

APPENDIX II.

A.

The multiplicity of an Einstein crystal is given by

$$\Omega(N, q) \approx \frac{(q + N)!}{q!N!},$$

where q is the number of energy units $\epsilon = hf$, where f is the frequency and h is Planck's constant, and N is the number of particles in the crystal. We use this to find the entropy

$$S = k \ln \Omega,$$

where k is Boltzmann's constant. We have that the internal energy of the crystal is

$$U = q\epsilon + \frac{N}{2}\epsilon,$$




Figure 6. Figure of numerically calculated entropy S/k against the number of energy units q .

if we include the ground-states. This of course gives us that

$$dU = \epsilon dq,$$

since we vary q .

The thermodynamic identity says that

$$dU = TdS - PdV + \mu dN,$$

where μ is the chemical potential, P is the pressure and V is the volume. We assume that the volume, pressure and number of particles stay constant. We then get that

$$T = \frac{dU}{dS} = \epsilon \frac{dq}{dS} \Leftrightarrow \frac{T}{\epsilon} = \frac{dq}{dS}.$$

Numerically we find this as

$$\tilde{T}_i = \frac{T_i}{\epsilon} = \frac{q_i - q_{i-1}}{S_i - S_{i-1}}.$$

We have that the heat capacity

$$C_V = \frac{dU}{dT} = \epsilon \frac{dq}{dT} = \frac{dq}{dT/\epsilon} = \frac{dq}{d\tilde{T}}.$$

We find this numerically as

$$C_{V,i} = \frac{q_i - q_{i-1}}{\tilde{T}_i - \tilde{T}_{i-1}}.$$

We see S/k as a function of q in figure (6) and $\frac{C_V}{Nk}$ as a function of q in figure (7).

B.

We have that the multiplicity of an Einstein solid is

$$\Omega(N, q) = \frac{(q + N - 1)!}{q!(N - 1)!},$$




Figure 7. Figure of numerically calculated heat capacity C_V against the number of energy units q .

where N is the number of particles and q is the number of energy units $\hbar f$. If we use $N \gg 1$, we can see that

$$\Omega(N, q) \approx \frac{(q + N)!}{q!N!}.$$

If also have $q \gg 1$ we can Stirling approximate this expression. That means that the temperature is $T > 0K$. This gives us that

$$\Omega = \frac{(q + N)!}{q!N!} \approx \frac{(q + N)^{q+N} e^{-(q+N)}}{q^q e^{-q} N^N e^{-N}} = \left(\frac{q + N}{q}\right)^q \left(\frac{q + N}{N}\right)^N$$

From the lectures we have that the multiplicity for the low temperature limit is

$$\Omega_{low\ T}(q, N) \approx \left(\frac{Ne}{q}\right)^q.$$

And the high temperature limit is

$$\Omega_{high\ T} \approx \left(\frac{qe}{N}\right)^N.$$

We see the comparison of the different Ω 's against each other in figure (8). From the figure we see that the low temperature approximation fits the Stirling approximation at low q and the high temperature approximation at high q .

We had that

$$dS = \frac{1}{U/Nk} dU = \frac{Nk}{U} dU = \frac{2Nk}{2q + N} dq.$$

This gives us that the entropy is

$$S(U, N) = Nk \int_{U_0}^U \frac{1}{U} dU = Nk \ln \frac{U}{U_0},$$




Figure 8. Figure of the low temperature approximation of the multiplicity $\Omega_{low\ T}$ vs. the high temperature approximation $\Omega_{high\ T}$ vs. the Stirling approximation $\Omega_{Stirling}$ against the number of energy units q .

where $U_0 = (1/2)N\epsilon$ is the ground state energy.

We have that the temperature

$$T(U, N) = \left(\frac{\partial S}{\partial U}\right)^{-1} = \left(Nk \frac{1}{U}\right)^{-1} = \frac{U}{Nk}.$$

And that the heat capacity

$$C_V(T, N) = \frac{dU}{dT} = \frac{d}{dT}(NkT) = Nk.$$

The specific heat capacity per. mole is

$$c = \frac{C}{n} = \frac{Nk}{n},$$

where $n = N/N_A$ is the number of moles and N_A is Avogadro's number. We then get that

$$c = \frac{Nk}{N/N_A} = N_A k = R,$$

where R is the universal gas constant.

C.

We see from figure (7) that C_V/Nk goes to one. This means that our analytical and numerical results corresponds more and more as q gets higher. And in (9) we see the heat capacity for different ϵ . We see that this does not go to $3R$ as they should and concludes that there is something wrong in the code. We can therefore not use this for a comparison to find the ϵ of a real material.




Figure 9. Heat capacity C_V for different ϵ .

APPENDIX III.

Since we have that

$$U = NkT = \epsilon q + \frac{N}{2}\epsilon$$

for the Einstein crystal, we can see that

$$dU = NkdT = \epsilon dq \Leftrightarrow dT = \frac{\epsilon}{Nk} dq.$$

We can use this to model how q changes in the mug, changes the temperature of the mug material. But the Einstein solid is not valid for liquids. The Temperfect mug has a material that will melt when you put hot liquid in the mug and therefore the Einstein model is not valid at that point. After a period of time the mug material will go back into solid and here the Einstein model might be valid. As energy units is transferred from the mug material to the liquid in the mug.

APPENDIX CODES. TERMOS TEMPERATURE

This code plots the temperatures as a function of time for the two thermos cups.

```
import matplotlib.pyplot as plt
import numpy as np
path_data = "../data_files/"

infile = open(path_data+"termokopper.txt", "r")

kopp1 = []
kopp2 = []
t = []

for line in infile:
    line = line.split()
    t.append(eval(line[0]))
    kopp1.append(eval(line[1]))
    kopp2.append(eval(line[2]))
infile.close()

t = np.asarray(t)
kopp1 = np.asarray(kopp1)
```

```
kopp2 = np.asarray(kopp2)*1.12

plt.plot(t,kopp1,label="Bodum mug")
plt.plot(t,kopp2,label="Temperfect mug")
plt.xlabel("t [s]",FontSize=15)
plt.ylabel(r"$T [\textcircled{C}]$",FontSize=15)
plt.legend()
plt.savefig("../figures/mugs.png")
plt.clf()

plt.plot(t,kopp1,label="Bodum mug")
plt.plot(t,kopp2,label="Temperfect mug")
plt.xlabel("t [s]",FontSize=15)
plt.ylabel(r"$T [\textcircled{C}]$",FontSize=15)
plt.xlim(0,150)
plt.ylim(85,90)
plt.legend()
plt.savefig("../figures/mugs_2.png")
plt.clf()
```

APPENDIX CODES. ENTROPY H2O

This code plots the entropy and heat capacity of H2O.

```
import numpy as np
import matplotlib.pyplot as plt

def C(T):
    if T<=273.15:
        return 2.03 * 1e3
    elif T>=373.15:
        return 1.998 * 1e3
    else:
        return 4.184 * 1e3

N = int(1e4)
T = np.linspace(100,500,N)
dSdT = np.zeros(N)
for i in range(N):
    dSdT[i] = C(T[i])/T[i]

S = np.zeros(N)
sum = 0

dT = (T[-1]-T[0])/N
for i in range(N):
    S[i] = sum
    sum += C(T[i])/T[i]*dT

plt.plot(T,S)
plt.xlabel(r"$T$ [K]",FontSize=15)
plt.ylabel(r"$S$ [J/K]",FontSize=15)
plt.savefig("../figures/S_mot_T")
plt.clf()

plt.plot(T,dSdT)
plt.xlabel(r"$T$ [K]",FontSize=15)
plt.ylabel(r"$dS/dT$ [J/K^2]",FontSize=15)
plt.savefig("../figures/dSdT_mot_T")
plt.clf()
```

APPENDIX CODES. ENTROPY PLOTTER

This code plots the entropy as a function of S.

```
import numpy as np
import matplotlib.pyplot as plt
from scipy.constants import k
from math import factorial

k = 1
```

```
N = int(300)
N_len = 800
q = np.linspace(1,N_len,N_len)
dq = (q[-1]-q[0])/(len(q)-1)

omega = np.zeros(N_len)
for i in range(0,N_len):
    omega[i] = factorial(int(q[i])+N-1)/(factorial(int(q[i]))*factorial(N-1))
S = k*np.log(omega)

dqdS = np.zeros(N_len)

for i in range(1,N_len):
    dqdS[i] = dq/(S[i]-S[i-1])
T = dqdS

Cv = np.zeros(N_len)
for i in range(1,N_len):
    Cv[i] = dq/(T[i]-T[i-1])

plt.plot(q,Cv/N,label=r"$N$=%g"%(N_len))
plt.xlabel("q",FontSize=15)
plt.ylabel(r"$C_V/Nk$",FontSize=15)
plt.legend()
plt.savefig("../figures/C_mot_q.png")
plt.clf()

plt.plot(q,S,label=r"$N$=%g"%(N_len))
plt.xlabel("q",FontSize=15)
plt.ylabel(r"$S/k$",FontSize=15)
plt.legend()
plt.savefig("../figures/S_mot_q.png")
plt.clf()

outfile = open("../data_files/numerical_data.txt","w+")
outfile.write("q Omega S T Cv/Nk N=%g\n"%(N_len))
for i in range(N_len):
    outfile.write("%g %.5f %.5f %.5f %.5f\n"%(q[i],omega[i],S[i],T[i],Cv[i]/N))
outfile.close()
```

APPENDIX CODES. Ω COMPARRISON.

This code plots the high T vs low T vs Stirling approx. for the Omegas.

```
import numpy as np
import matplotlib.pyplot as plt

N = 1e2
N_len = int(1e3)

q = np.linspace(1,N_len,N_len)
omega_lowT = (N*np.exp(1)/q)**q
omega_highT = (q*np.exp(1)/N)**N
omega_stirling = ((q+N)/q)**q * ((q+N)/N)**N

plt.plot(q,np.log(omega_lowT),label=r"$\Omega_{low\ T}$, N=%g"%(N))
plt.plot(q,np.log(omega_highT),label=r"$\Omega_{high\ T}$, N=%g"%(N))
plt.plot(q,np.log(omega_stirling),label=r"$\Omega_{Stirling}$, N=%g"%(N))
plt.xlabel("q",FontSize=15)
plt.ylabel("S/k",FontSize=15)
plt.legend()
plt.savefig("../figures/omegas.png")
plt.clf()
```

APPENDIX CODES. ANALYTICAL VS NUMERICAL ENTROPY

This code plots the analytical vs the numerical entropy.

```
import numpy as np
import matplotlib.pyplot as plt
```

```

k = 1;

infile = open("../data_files/numerical_data.txt", "r")
N = infile.readline().split("=")[-1]
N = eval(N)

q = []
omega_numerical = []
S_numerical = []
T_numerical = []
Cv_over_Nk_numerical = []

for line in infile:
    line = line.split()
    q.append(eval(line[0]))
    omega_numerical.append(eval(line[1]))
    S_numerical.append(eval(line[2]))
    T_numerical.append(eval(line[3]))
    Cv_over_Nk_numerical.append(eval(line[4]))
infile.close()
q = np.asarray(q)
omega_numerical = np.asarray(omega_numerical)
S_numerical = np.asarray(S_numerical)
T_numerical = np.asarray(T_numerical)
Cv_over_Nk_numerical = np.asarray(Cv_over_Nk_numerical)

S_analytical = N*k*np.log(2*q/N+1)
plt.plot(q, S_analytical)
plt.plot(q, S_numerical)
plt.show()

```

APPENDIX CODES. C COMPARRISON.

This code plots the different ϵ 's.

```

import numpy as np
import matplotlib.pyplot as plt
from scipy.constants import k
from scipy.constants import R

k = 1;

infile = open("../data_files/numerical_data.txt", "r")
N = infile.readline().split("=")[-1]
N = eval(N)

q = []
omega_numerical = []
S_numerical = []
T_numerical = []
Cv_over_Nk_numerical = []

for line in infile:
    line = line.split()
    q.append(eval(line[0]))
    omega_numerical.append(eval(line[1]))
    S_numerical.append(eval(line[2]))
    T_numerical.append(eval(line[3]))
    Cv_over_Nk_numerical.append(eval(line[4]))
infile.close()
q = np.asarray(q)
omega_numerical = np.asarray(omega_numerical)
S_numerical = np.asarray(S_numerical)
T_numerical = np.asarray(T_numerical)
Cv_over_Nk_numerical = np.asarray(Cv_over_Nk_numerical)

plt.yticks()
for epsilon in range(200, 5001, 500):
    plt.plot(T_numerical*epsilon, Cv_over_Nk_numerical*N*k/R, label="epsilon=%g"%epsilon)
plt.xlabel("T [K]", FontSize=15)
plt.legend()
plt.savefig("../figures/C_comparrison.png")
plt.clf()

```