

ECMT3150: The Econometrics of Financial Markets

2a. Conditional Heteroskedastic Models

Simon Kwok
University of Sydney

Semester 1, 2021

Outline

1. Stylized Facts of Stock Returns and Volatility
2. Testing for ARCH Effect
3. ARCH/GARCH Model
 - 3.1 Model Structure
 - 3.2 ARMA Representation
 - 3.3 Volatility Forecast
 - 3.4 Model Estimation
 - 3.5 Model Diagnostics
4. Some Extensions
 - 4.1 IGARCH
 - 4.2 GARCH-M
 - 4.3 EGARCH
 - 4.4 TGARCH
 - 4.5 SV

Volatility

Q: What is Volatility?

- ▶ Volatility is a measure of the degree of asset price/market fluctuation.
- ▶ Volatility is high during more turbulent periods/market crashes, and low during quiet or booming periods.

Q: Why do should we care about Volatility?

- ▶ It is important for pricing options and other financial derivatives.
- ▶ VIX volatility index is being traded on CBOE in the form of futures.

Some Stylized Facts

daily

Q: What are some stylized facts about stock returns?

- ▶ Stock return has mean that is close to zero.
- ▶ Stock return tends to be serially uncorrelated.¹

Q: What are some stylized facts about volatility?

- ▶ Volatility is not directly observable.
- ▶ Volatility level of stock price typically goes up fast and then decays slowly over time.
- Volatility clustering: volatility level is autocorrelated, i.e., high volatility periods tend to cluster together (and similarly for low volatility periods).
ARCH effect
- ▶ Leverage effect: volatility level tends to be inversely associated with price changes.

¹This is not true for high frequency data, however.

Model Structure

P_t : asset price at time t .

r_t : log-return of the asset at time t , given by

$$r_t = \log(P_t) - \log(P_{t-1}).$$

Let \mathcal{F}_{t-1} be the information set at time $t-1$. The conditional mean and variance of r_t are:

$$\left[\begin{array}{l} \mu_t = E(r_t | \mathcal{F}_{t-1}), \\ \sigma_t^2 = \text{Var}(r_t | \mathcal{F}_{t-1}) = E[(r_t - \mu_t)^2 | \mathcal{F}_{t-1}]. \end{array} \right. \quad \begin{array}{l} (1) \\ (2) \end{array}$$

We need a parametric model to describe the evolution of μ_t and σ_t^2 over time.

Let $a_t = r_t - \mu_t$. The dynamic regression model is:

$$r_t = \mu_t + a_t.$$

The error a_t may exhibit *conditional* heteroskedasticity, so that $\sigma_t^2 := \text{Var}(a_t | \mathcal{F}_{t-1})$ is not a constant over t .

Model Structure

- We may want to explain r_t with its lagged terms $r_{t-1}, r_{t-2}, \dots, r_{t-p}$:

$$\begin{aligned}\mu_t &= E(r_t | \mathcal{F}_{t-1}) \\ &= \phi_0 + \phi_1 r_{t-1} + \dots + \phi_p r_{t-p}\end{aligned}$$



- Sometimes we may want to include additional (weakly) exogenous regressors \mathbf{x}_t as well:

$$\begin{aligned}\mu_t &= E(r_t | \mathcal{F}_{t-1}, \mathbf{x}_t) \\ &= \phi_0 + \phi_1 r_{t-1} + \dots + \phi_p r_{t-p} + \gamma' \mathbf{x}_t\end{aligned}$$

eg r_{t-1}^2

- A popular class of models for σ_t^2 is the (generalized) autoregressive conditional heteroskedasticity model, or ARCH (GARCH) model.

Testing for ARCH Effect

Let $\hat{a}_t = r_t - \hat{\mu}_t$ be the residuals from the dynamic regression model.

data cond. mean.

Q: How to check for conditional heteroskedasticity (ARCH effect)?

A: Two approaches, both require choosing a maximum lag order m for testing.

proxy for volatility.

► Ljung-Box test $Q(m)$ on $\{\hat{a}_t^2\}$ (McLeod-Li (1983)).

→ ► Lagrange multiplier (LM) test (Engle (1982)). Carry out an F test in the linear regression ($t = m + 1, \dots, T$)

$$\hat{a}_t^2 = \alpha_0 + \alpha_1 \hat{a}_{t-1}^2 + \dots + \alpha_m \hat{a}_{t-m}^2 + e_t. \quad (3)$$

We want to test $H_0 : \alpha_i = 0$ for all $i = 1, \dots, m$ vs $H_a : \alpha_i \neq 0$ for some $i = 1, \dots, m$.

Define the restricted and unrestricted sums of squared residuals:

$SSR_r = \sum_{t=1}^T (\hat{a}_t^2 - \bar{\hat{a}^2})^2$ and $SSR_{ur} = \sum_{t=1}^T \hat{e}_t^2$, where $\bar{\hat{a}^2} = \frac{1}{T} \sum_{t=1}^T \hat{a}_t^2$, and \hat{e} is the residuals from regression (3).

The F test statistic is

$$F = \frac{(SSR_r - SSR_{ur})/m}{SSR_{ur}/(T - 2m - 1)},$$

ARCH Model

$a_t \sim \text{ARCH}(m)$ if log-return $\mu_t = 0$
model error $\rightarrow a_t = \sigma_t \varepsilon_t$

where $\varepsilon_t \sim iid(0, 1)$ (i.e., iid with mean 0 and variance 1), and

$\rightarrow \sigma_t^2 = \alpha_0 + \sum_{i=1}^m \alpha_i a_{t-i}^2$ \star proxy for vola. at $t-i$. (4)

$\sigma_t^2 = \text{Var}(a_t | \mathcal{F}_{t-1})$ is interpreted as the conditional variance at time t given the history.

$\alpha_i > 0$ due to vola. clustering.

If a_t is weakly stationary, the unconditional (long-run) variance $\sigma^2 := \text{Var}(a_t)$ exists. We may take unconditional expectations on both sides of (4) and solve for σ^2 :

$$\sigma^2 = \frac{\alpha_0}{1 - \sum_{i=1}^m \alpha_i}.$$

\neq

The necessary condition for the existence of σ^2 is $\sum_{i=1}^m \alpha_i < 1$.

GARCH Model

$a_t \sim \text{GARCH}(m, s)$ if

$$a_t = \sigma_t \varepsilon_t,$$

where $\varepsilon_t \sim \text{iid}(0, 1)$, and

$$\sigma_t^2 = \alpha_0 + \sum_{i=1}^m \alpha_i a_{t-i}^2 + \sum_{i=1}^s \beta_i \sigma_{t-i}^2. \quad (6)$$

If a_t is weakly stationary, the unconditional (long-run) variance exists and is equal to:

$$\sigma^2 = \frac{\alpha_0}{1 - \sum_{i=1}^m \alpha_i - \sum_{i=1}^s \beta_i}. \quad (7)$$

The necessary condition for the existence of σ^2 is

$$\sum_{i=1}^m \alpha_i + \sum_{i=1}^s \beta_i < 1. \quad (8)$$

$M_t = 0$ (hence pure GARCH model) ⁽⁵⁾

$$\sigma_{t+1}^2 = \alpha_0 + \sum \alpha_i a_{t+1-i}^2 + \sum \beta_i \sigma_{t+1-i}^2$$

$$\alpha_1 a_t^2 + \alpha_2 a_{t-1}^2 + \dots$$

ARMA Representation of Squared Errors GARCH(m,s)

Let $\eta_t = \underbrace{a_t^2}_{\text{error of squared log-return}} - \sigma_t^2$. Recall that $\sigma_t^2 = \text{Var}(a_t | \mathcal{F}_{t-1}) = \underline{E(a_t^2 | \mathcal{F}_{t-1})}$, so we see that $\eta_t = \underline{a_t^2 - E(a_t^2 | \mathcal{F}_{t-1})}$.

Ex: Show that $\eta_t \sim mds$, i.e., $\underline{E(\eta_t | \mathcal{F}_s) = 0}$ for all $t > s$.

We may then rewrite (6) as follows:

$$\begin{aligned} \rightarrow \quad \underline{a_t^2 - \eta_t} &= \alpha_0 + \sum_{i=1}^{\underline{m}} \alpha_i \underline{a_{t-i}^2} + \sum_{i=1}^s \beta_i (\underline{a_{t-i}^2} - \underline{\eta_{t-i}}) \\ \rightarrow \quad \underline{a_t^2} &= \alpha_0 + \underbrace{\sum_{i=1}^{\max(m,s)} (\alpha_i + \beta_i) a_{t-i}^2}_{AR} + \underbrace{\eta_t - \sum_{i=1}^s \beta_i \eta_{t-i}}_{MA}. \quad (9) \end{aligned}$$

In (9), we adopt the convention that $\alpha_i = 0$ for $i > m$, and $\beta_i = 0$ for $i > s$.

Interpretation: The squared innovation $a_t^2 \sim \underline{ARMA(\max(m,s), s)}$ with mds errors η_t .

GARCH Volatility Forecasting

At the forecast origin t , the 1-step ahead forecast of σ_{t+1}^2 is ~~known by time t~~ (due to ~~GARCH~~ **GARCH** assumption).

Optimal forecast of σ_{t+1}^2

$$\begin{aligned}\hat{\sigma}_t^2(1) &= \underline{E(\sigma_{t+1}^2 | \mathcal{F}_t)} \quad (\text{see p.20 of slides 1}) \\ &= \underline{\sigma_{t+1}^2} \quad (\text{since } \sigma_{t+1}^2 \text{ is measurable to } \mathcal{F}_t) \\ &= \underline{\alpha_0 + \sum_{i=1}^m \alpha_i a_{t+1-i}^2 + \sum_{i=1}^s \beta_i \sigma_{t+1-i}^2} \quad \leftarrow \text{from (6).}\end{aligned}$$

Q: What is the ℓ -step ahead forecast of $\sigma_{t+\ell}^2$?

$$\hat{\sigma}_t^2(1) = E(\sigma_{t+1}^2 | \mathcal{F}_t) \leftarrow \text{min. MSE.}$$

GARCH Volatility Forecasting


A: First, using (5), rewrite the recursive equation (6) as

$$\begin{aligned} \rightarrow \sigma_{t+l}^2 &= \alpha_0 + \sum_{i=1}^m \alpha_i \sigma_{t+l-i}^2 \varepsilon_{t+l-i}^2 + \sum_{i=1}^s \beta_i \sigma_{t+l-i}^2 + \sum \alpha_i \sigma_{t+l-i}^2 - \sum \alpha_i \sigma_{t+l-i}^2 \\ \sigma_{t+l-i}^2 \text{ is known with } \mathcal{F}_{t+l-i} &= \alpha_0 + \sum_{i=1}^{\max(m,s)} (\alpha_i + \beta_i) \sigma_{t+l-i}^2 + \sum_{i=1}^m \alpha_i \sigma_{t+l-i}^2 (\varepsilon_{t+l-i}^2 - 1). \end{aligned}$$

ARMA rep.

error.

Then, take expectations conditional on \mathcal{F}_t , and note that



and

$$E(\sigma_{t+l-i}^2 | \mathcal{F}_t) = \begin{cases} \hat{\sigma}_t^2(l-i) & \text{if } l-i > 0, \\ \sigma_{t+l-i}^2 & \text{otherwise;} \end{cases}$$

Tower's rule

constant as at time t .

$$\begin{aligned} E(\sigma_{t+l-i}^2 (\varepsilon_{t+l-i}^2 - 1) | \mathcal{F}_t) &= \begin{cases} E[\sigma_{t+l-i}^2 E(\varepsilon_{t+l-i}^2 - 1 | \mathcal{F}_{t+l-i-1}) | \mathcal{F}_t] & \text{if } l-i > 0, \\ \sigma_{t+l-i}^2 (\varepsilon_{t+l-i}^2 - 1) & \text{otherwise,} \end{cases} \\ \sigma_{t+l-i}^2 \text{ is known at time } t+l-i &= \begin{cases} E[\sigma_{t+l-i}^2 E(\varepsilon_{t+l-i}^2 - 1 | \mathcal{F}_t)] = 0 & \text{if } l-i > 0, \\ a_{t+l-i}^2 - \sigma_{t+l-i}^2 & \text{otherwise.} \end{cases} \end{aligned}$$

by iid of ε_t

$$E(\varepsilon_{t+l-i}^2) = \text{Var}(\varepsilon_{t+l-i}) = 1$$

GARCH Volatility Forecasting

The ℓ -step ahead forecast of $\sigma_{t+\ell}^2$ is thus given by

$$\rightarrow \hat{\sigma}_t^2(\ell) = \alpha_0 + \sum_{i=1}^{\max(m,s)} (\alpha_i + \beta_i) \hat{\sigma}_t^2(\ell-i) + \sum_{i=\ell}^m \alpha_i (a_{t+\ell-i}^2 - \sigma_{t+\ell-i}^2).$$

Handwritten notes above the equation: $E(\sigma_{t+\ell}^2 | \mathcal{F}_t)$ above the first term, $E(\sigma_{t+\ell-i}^2 | \mathcal{F}_t)$ above the recursive term, and $E(\sigma_{t+\ell-i}^2 (\sigma_{t+\ell-i}^2 - 1) | \mathcal{F}_t)$ above the summation term with a downward arrow pointing to $a_{t+\ell-i}^2 - \sigma_{t+\ell-i}^2$.

We adopt the convention that $\hat{\sigma}_t^2(\ell-i) = \sigma_{t+\ell-i}^2$ if $\ell-i \leq 0$.

The last summation is gone if $\ell > m$.

E.g.: For the GARCH(1, 1) model (i.e., $m = s = 1$), the ℓ -step ahead forecast of $\sigma_{t+\ell}^2$ is:

$$\hat{\sigma}_t^2(\ell) = \begin{cases} \alpha_0 + \alpha_1 a_t^2 + \beta_1 \sigma_t^2 & \text{if } \ell = 1, \\ \alpha_0 + (\alpha_1 + \beta_1) \hat{\sigma}_t^2(\ell-1) & \text{if } \ell > 1. \end{cases}$$

Handwritten notes: A downward arrow labeled c points to $\hat{\sigma}_t^2(\ell)$. Another downward arrow labeled c points to $\hat{\sigma}_t^2(\ell-1)$. To the right, an equation $\Rightarrow c = \frac{\alpha_0}{1 - (\alpha_1 + \beta_1)}$ is written.

Q: What happens to $\hat{\sigma}_t^2(\ell)$ as $\ell \rightarrow \infty$? \leftarrow

A: If $\sum_{i=1}^{\max(m,s)} (\alpha_i + \beta_i) < 1$, then $\hat{\sigma}_t^2(\ell)$ converges to the long-run variance $\alpha_0 / \left[1 - \sum_{i=1}^{\max(m,s)} (\alpha_i + \beta_i) \right]$. \leftarrow

Handwritten note: $m, s = 1$.

Properties of GARCH(1,1) Model

The GARCH(1,1) model is given by $E(r_t | \mathcal{F}_{t-1})$.

$$\begin{cases} r_t = \mu_t + a_t, \\ a_t = \sigma_t \varepsilon_t, \\ \sigma_t^2 = \alpha_0 + \alpha_1 a_{t-1}^2 + \beta_1 \sigma_{t-1}^2, \end{cases} \quad \leftarrow$$

with $\alpha_1, \beta_1 > 0$ and $\alpha_1 + \beta_1 < 1$, ε_t is iid noise with $E(\varepsilon_t) = 0$, and $\text{Var}(\varepsilon_t) = E(\varepsilon_t^2) = 1$.
 eg. $\varepsilon_x \sim N(0, 1) \Rightarrow K_\varepsilon = 0$

By (1) and (2), μ_t and σ_t^2 are measurable with respect to the information set \mathcal{F}_{t-1} .
 $\varepsilon_x \sim t_4 \Rightarrow K_\varepsilon > 0$

We assume for now that r_t is covariance stationary (so that its first two unconditional moments are constant over time).

Conditional Mean and Variance of GARCH(1,1)

Q: What are the conditional mean and variance of r_t ?

$$\begin{aligned}\underline{E(r_t | \mathcal{F}_{t-1})} &= E(\mu_t + a_t | \mathcal{F}_{t-1}) \\&= \mu_t + E(a_t | \mathcal{F}_{t-1}) \\&= \mu_t + \sigma_t E(\varepsilon_t | \mathcal{F}_{t-1}) \\&= \mu_t + \sigma_t E(\varepsilon_t) \\&= \underline{\mu_t}.\end{aligned}$$

$$\begin{aligned}\text{Var}(r_t | \mathcal{F}_{t-1}) &= E[(r_t - \mu_t)^2 | \mathcal{F}_{t-1}] \\&= E(a_t^2 | \mathcal{F}_{t-1}) \\&= E(\sigma_t^2 \varepsilon_t^2 | \mathcal{F}_{t-1}) \\&= \sigma_t^2 E(\varepsilon_t^2 | \mathcal{F}_{t-1}) \\&= \sigma_t^2 E(\varepsilon_t^2) = \sigma_t^2 \\&= (\alpha_0 + \alpha_1 a_{t-1}^2 + \beta_1 \sigma_{t-1}^2) \times 1 \\&= \alpha_0 + \alpha_1 \underline{a_{t-1}^2} + \beta_1 \underline{\sigma_{t-1}^2}.\end{aligned}$$

Handwritten notes: \vdots and $= \sigma_t^2$ are written in blue next to the first four lines of the variance derivation.

Unconditional Variance of GARCH(1,1)

$$E(r_t) = E(\mu_t)$$

uncond. mean

Q: What is the unconditional variance of r_t ?

A: By the law of total variance,

$$\begin{aligned} \text{Var}(r_t) &= E[\text{Var}(r_t | \mathcal{F}_{t-1})] + \text{Var}[E(r_t | \mathcal{F}_{t-1})] \\ &= E(\alpha_0 + \alpha_1 a_{t-1}^2 + \beta_1 \sigma_{t-1}^2) + \text{Var}(\mu_t) \\ &= \alpha_0 + \alpha_1 E(a_{t-1}^2) + \beta_1 E(\sigma_{t-1}^2) + 0 \\ &= \alpha_0 + \alpha_1 \text{Var}(r_{t-1}) + \beta_1 E(\sigma_{t-1}^2). \end{aligned}$$

By the stationarity of r_t , $\text{Var}(r_t) = \text{Var}(r_{t-1})$, i.e., the variance is constant over t .
Also,

$$\begin{aligned} \text{Var}(r_t) &= E(a_t^2) \\ &= E[\sigma_t^2 E(\varepsilon_t^2 | \mathcal{F}_{t-1})] \\ &= E(\sigma_t^2 E(\varepsilon_t^2)) \\ &= E(\sigma_t^2). \end{aligned}$$

By letting $V := \text{Var}(r_t) = \text{Var}(r_{t-1}) = E(\sigma_t^2)$, we have

$$V = \alpha_0 + \alpha_1 V + \beta_1 V.$$

Provided that $|\alpha_1 + \beta_1| < 1$, the solution exists, and is equal to

$$V := \text{Var}(r_t) = \frac{\alpha_0}{1 - \alpha_1 - \beta_1}. \quad \leftarrow$$

Unconditional Kurtosis of GARCH(1,1)

Assume further that r_t is stationary up to the fourth moment, that $E(\varepsilon_t^4)$ exists, and $E(\varepsilon_t^4) = 3 + K_\varepsilon$ (i.e., K_ε is the excess kurtosis of ε_t).

Q: What is the unconditional excess kurtosis of r_t ?

A: We start with the conditional fourth central moment of r_t

$$\begin{aligned} & E[(r_t - \mu_t)^4 | \mathcal{F}_{t-1}] \\ = & E(a_t^4 | \mathcal{F}_{t-1}) \\ = & E(\sigma_t^4 \varepsilon_t^4 | \mathcal{F}_{t-1}) \\ = & \sigma_t^4 E(\varepsilon_t^4) \\ = & (\alpha_0 + \alpha_1 a_{t-1}^2 + \beta_1 \sigma_{t-1}^2)^2 (3 + K_\varepsilon) \\ = & (\alpha_0^2 + \alpha_1^2 a_{t-1}^4 + \beta_1^2 \sigma_{t-1}^4 + 2\alpha_0 \alpha_1 a_{t-1}^2 + 2\alpha_0 \beta_1 \sigma_{t-1}^2 + 2\alpha_1 \beta_1 a_{t-1}^2 \sigma_{t-1}^2) (3 + K_\varepsilon). \end{aligned}$$

Let us then work out the unconditional fourth moment of a_t

$$\begin{aligned} \underline{E(a_t^4)} &= \underline{E[(r_t - \mu_t)^4]} \\ &= (3 + K_\varepsilon) [\alpha_0^2 + \alpha_1^2 E(a_{t-1}^4) + \beta_1^2 E(\sigma_{t-1}^4) \\ &\quad + 2\alpha_0 \alpha_1 E(a_{t-1}^2) + 2\alpha_0 \beta_1 E(\sigma_{t-1}^2) + 2\alpha_1 \beta_1 E(a_{t-1}^2 \sigma_{t-1}^2)]. \end{aligned} \quad (10)$$

Unconditional Kurtosis of GARCH(1,1)

By iterated expectations, the last expectation can be simplified as follows

$$\begin{aligned} E(a_{t-1}^2 \sigma_{t-1}^2) &= E[E(a_{t-1}^2 \sigma_{t-1}^2 | \mathcal{F}_{t-2})] \\ &= E[\sigma_{t-1}^2 E(a_{t-1}^2 | \mathcal{F}_{t-2})] \\ &= E[\sigma_{t-1}^2 E(\sigma_{t-1}^2 \epsilon_{t-1}^2 | \mathcal{F}_{t-2})] \\ &= E[\sigma_{t-1}^4 E(\epsilon_{t-1}^2)] = E(\sigma_{t-1}^4). \end{aligned}$$

By stationarity in the fourth moment, we have $E(a_t^4) = E(a_{t-1}^4)$. By iterated expectations,

$$\begin{aligned} E(a_t^4) &= E(\sigma_t^4 \epsilon_t^4) = E[\sigma_t^4 E(\epsilon_t^4 | \mathcal{F}_{t-1})] \\ &= E[\sigma_t^4 E(\epsilon_t^4)] = E(\sigma_t^4)(3 + K_\epsilon), \end{aligned}$$

so that $E(\sigma_t^4) = E(a_t^4)/(3 + K_\epsilon)$. We can simplify (10) as follows

$$\begin{aligned} E(a_t^4) &= (3 + K_\epsilon)[\alpha_0^2 + \alpha_1^2 E(a_t^4)] + (\beta_1^2 + 2\alpha_1\beta_1)E(a_t^4) \\ &\quad + (3 + K_\epsilon)(2\alpha_0\alpha_1 + 2\alpha_0\beta_1)E(a_t^2). \end{aligned}$$

Together with the fact that $\text{Var}(r_t) = E(a_t^2) = E(\sigma_t^2) = \frac{\alpha_0}{1 - \alpha_1 - \beta_1}$, we may then solve for $E(a_t^4)$:

$$[1 - (3 + K_\epsilon)\alpha_1^2 - \beta_1^2 - 2\alpha_1\beta_1]E(a_t^4) = (3 + K_\epsilon)[\alpha_0^2 + 2\alpha_0(\alpha_1 + \beta_1)E(a_t^2)],$$

so that

$$E(a_t^4) = \frac{(3 + K_\epsilon)[\alpha_0^2 + \frac{2\alpha_0^2(\alpha_1 + \beta_1)}{1 - \alpha_1 - \beta_1}]}{1 - (3 + K_\epsilon)\alpha_1^2 - \beta_1^2 - 2\alpha_1\beta_1} = \frac{\alpha_0^2(3 + K_\epsilon)(1 + \alpha_1 + \beta_1)}{(1 - \alpha_1 - \beta_1)[1 - 2\alpha_1^2 - K_\epsilon\alpha_1^2 - (\alpha_1 + \beta_1)^2]}.$$

Unconditional Kurtosis of GARCH(1,1)

$$E[(X-\mu)^4] = 3$$

$$X \sim N(0, 1)$$

$$K_X = 0$$

The unconditional kurtosis of r_t is thus

$$\begin{aligned} \frac{E[(r_t - \mu_t)^4]}{\{E[(r_t - \mu_t)^2]\}^2} &= \frac{(3 + K_\varepsilon)(1 + \alpha_1 + \beta_1)(1 - \alpha_1 - \beta_1)}{1 - 2\alpha_1^2 - K_\varepsilon\alpha_1^2 - (\alpha_1 + \beta_1)^2} \\ &= \frac{[1 - (\alpha_1 + \beta_1)^2](3 + K_\varepsilon)}{1 - 2\alpha_1^2 - K_\varepsilon\alpha_1^2 - (\alpha_1 + \beta_1)^2} \end{aligned}$$

↑
excess kur
:= $E[(X-\mu)^4] - 3$

By simple algebra, the unconditional excess kurtosis is given by

$$K_r := \frac{E[(r_t - \mu_t)^4]}{\{E[(r_t - \mu_t)^2]\}^2} - 3 = \frac{6\alpha_1^2 + K_\varepsilon[1 - (\alpha_1 + \beta_1)^2 + 3\alpha_1^2]}{1 - 2\alpha_1^2 - K_\varepsilon\alpha_1^2 - (\alpha_1 + \beta_1)^2}$$

↓ ARCH par

We see that K_r is zero if $\alpha_1 = 0$ and $K_\varepsilon = 0$. In other words, the ARCH parameter α_1 and the excess kurtosis of ε_t contribute to the fat-tailedness of the unconditional distribution of the log-return r_t .