

ECMT3150: The Econometrics of Financial Markets

1b. Linear Time Series Analysis

Simon Kwok
University of Sydney

Semester 1, 2021

Outline

1. MA Model
2. ARMA Model
3. Stationarity and Invertibility
4. Model Checking and Portmanteau Tests

Moving-Average (MA) Model

$MA(q)$:

$$y_t = \mu + \varepsilon_t + \theta_1 \varepsilon_{t-1} + \cdots + \theta_q \varepsilon_{t-q},$$

where $\{\varepsilon_t\} \sim wn(0, \sigma_\varepsilon^2)$. Noting that $E(y_t) = \mu$, we can demean the $MA(q)$ model into

$$\begin{aligned} u_t &= \varepsilon_t + \theta_1 \varepsilon_{t-1} + \cdots + \theta_q \varepsilon_{t-q} \\ &= [1 + \theta(L)] \varepsilon_t, \end{aligned}$$

where $\theta(x) = \sum_{i=1}^q \theta_i x^i$.

Ex: Show that for an $MA(q)$ model, the ACF is given by, for $j > 0$,

$$\begin{aligned} \rho_j &\equiv \text{Corr}(u_t, u_{t-j}) = \text{Corr}(y_t, y_{t-j}) \\ &= \begin{cases} \frac{\theta_j + \sum_{i=1}^{q-j} \theta_{j+i} \theta_i}{1 + \sum_{i=1}^q \theta_i^2} & \text{if } j \leq q, \\ 0 & \text{if } j > q, \end{cases} \end{aligned}$$

and $\rho_{-j} = \rho_j$. Plot the ACF against lag order.

MA Model Estimation

Given the time series data $\{y_t\}_{t=1}^T$, we can estimate an $MA(q)$ model by either *conditional* or *exact maximum likelihood estimation* (MLE).

- ▶ Conditional MLE: Set the initial errors (ε_t for $t = 0, -1, \dots, -q + 1$) to zero. Then assume a distribution on $\{\varepsilon_t\}_{t=1}^T$ (usually iid normal). The joint likelihood function is obtained in terms of the MA parameters $\mu, \theta_1, \dots, \theta_q$. Maximize the log-likelihood w.r.t. $\mu, \theta_1, \dots, \theta_q$.
- ▶ Exact MLE: Treat the initial errors (ε_t for $t = 0, -1, \dots, -q + 1$) as extra parameters. Maximize the log-likelihood w.r.t. $\mu, \theta_1, \dots, \theta_q, \varepsilon_0, \varepsilon_{-1}, \dots, \varepsilon_{-q+1}$.

MA Model Selection

- ▶ Use sample ACF. The asymptotic variance of $\hat{\rho}_j$ is $\frac{1}{T}$. For large T , the 95% confidence interval is roughly given by $\hat{\rho}_j \mp \frac{1.96}{\sqrt{T}}$.
- ▶ Use information criteria.

Forecasting with MA model

Suppose $\{y_t\} \sim MA(q)$ with errors $\{\varepsilon_t\} \sim wn(0, \sigma_\varepsilon^2)$. For $\ell = 1$,

$$\begin{aligned}\hat{y}_t(1) &= E[y_{t+1}|\mathcal{F}_t] \\ &= E[\mu + \varepsilon_{t+1} + \theta_1\varepsilon_t + \cdots + \theta_q\varepsilon_{t-q+1}|\mathcal{F}_t] \\ &= \mu + \theta_1\varepsilon_t + \cdots + \theta_q\varepsilon_{t-q+1},\end{aligned}$$

as $E[\varepsilon_{t+1}|\mathcal{F}_t] = 0$, and $\varepsilon_t, \dots, \varepsilon_{t-q+1}$ are measurable w.r.t. \mathcal{F}_t . The forecast error is $e_t(1) = y_{t+1} - \hat{y}_t(1) = \varepsilon_{t+1}$, with variance $Var[e_t(1)] = \sigma_\varepsilon^2$.

Forecasting with MA model

For $\ell = 2$,

$$\begin{aligned}\hat{y}_t(2) &= E[y_{t+2}|\mathcal{F}_t] \\ &= E[\mu + \varepsilon_{t+2} + \theta_2\varepsilon_{t+1} + \cdots + \theta_q\varepsilon_{t-q+2}|\mathcal{F}_t] \\ &= \mu + \theta_2\varepsilon_t + \cdots + \theta_q\varepsilon_{t-q+2}.\end{aligned}$$

The forecast error is $e_t(2) = y_{t+2} - \hat{y}_t(2) = \varepsilon_{t+2} + \theta_1\varepsilon_{t+1}$, with variance

$$\begin{aligned}\text{Var}[e_t(2)] &= \text{Var}(\varepsilon_{t+2}) + \theta_1^2 \text{Var}(\varepsilon_{t+1}) + 2\theta_1 \text{Cov}(\varepsilon_{t+2}, \varepsilon_{t+1}) \\ &= \sigma_\varepsilon^2 + \theta_1^2 \sigma_\varepsilon^2 + 0 \\ &= (1 + \theta_1^2) \sigma_\varepsilon^2.\end{aligned}$$

Forecasting with MA model

Q: What is the ℓ -step ahead forecast of an $MA(q)$ model? What are the forecast error and its variance? What happens to the forecast and its variance when ℓ increases beyond q ?

A: The ℓ -step ahead forecast is

$$\hat{y}_t(\ell) = \begin{cases} \mu + \theta_\ell \varepsilon_t + \cdots + \theta_q \varepsilon_{t-q+\ell} & \text{if } \ell \leq q, \\ \mu & \text{if } \ell > q. \end{cases} \quad (1)$$

The forecast error is

$$e_t(\ell) = \begin{cases} \varepsilon_{t+\ell} + \theta_1 \varepsilon_{t+\ell-1} + \cdots + \theta_{\ell-1} \varepsilon_{t+1} & \text{if } \ell \leq q, \\ \varepsilon_{t+\ell} + \theta_1 \varepsilon_{t+\ell-1} + \cdots + \theta_q \varepsilon_{t+\ell-q} & \text{if } \ell > q. \end{cases} \quad (2)$$

The forecast variance is

$$\text{Var}[e_t(\ell)] = \begin{cases} (1 + \theta_1^2 + \theta_2^2 + \cdots + \theta_{\ell-1}^2) \sigma_\varepsilon^2 & \text{if } \ell \leq q, \\ (1 + \theta_1^2 + \theta_2^2 + \cdots + \theta_q^2) \sigma_\varepsilon^2 & \text{if } \ell > q, \end{cases} \quad (3)$$

which is equal to $\text{Var}(y_t)$.

As ℓ increases beyond q , $\hat{y}_t(\ell)$ stays at the mean level μ , and the forecast variance remains at $\text{Var}(y_t) = (1 + \theta_1^2 + \theta_2^2 + \cdots + \theta_q^2) \sigma_\varepsilon^2$.

Autoregressive Moving-Average Model

$\{y_t\} \sim ARMA(p, q)$ if

$$\begin{aligned}y_t &= \phi_0 + \phi_1 y_{t-1} + \cdots + \phi_p y_{t-p} + \varepsilon_t + \theta_1 \varepsilon_{t-1} + \cdots + \theta_q \varepsilon_{t-q} \\[1 - \phi(L)] y_t &= \phi_0 + [1 + \theta(L)] \varepsilon_t,\end{aligned}$$

where $\{\varepsilon_t\} \sim wn(0, \sigma_\varepsilon^2)$.

After demeaning ($y_t = \mu + u_t$), the expressions become

$$\begin{aligned}u_t &= \phi_1 u_{t-1} + \cdots + \phi_p u_{t-p} + \varepsilon_t + \theta_1 \varepsilon_{t-1} + \cdots + \theta_q \varepsilon_{t-q} \\[1 - \phi(L)] u_t &= [1 + \theta(L)] \varepsilon_t.\end{aligned}\tag{4}$$

Ex: What is the ACF of a stationary $ARMA(1, 1)$ model?

Stationarity Condition

$\{y_t\}$ is said to be *stationary* if it can be expressed as an $MA(\infty)$ process of the form $u_t = [1 + \psi(L)] \varepsilon_t$.

Stationarity condition: The $ARMA(p, q)$ model is stationary iff all the roots of the polynomial equation $1 - \phi(x) = 0$ lie outside the unit circle.

In this case, $1 + \psi(L) := \frac{1 + \theta(L)}{1 - \phi(L)}$ is a well-defined infinite-order polynomial.

Q: Does the stationarity condition imply weak (covariance) stationarity?

Hint: $ARMA(p, q)$ can be expressed as $MA(\infty)$ under the stationarity condition. The Yule-Walker equations yield a unique set of autocovariances as solution: $\gamma_0, \gamma_1, \gamma_2, \dots$. Verify that the mean (by definition), the variance γ_0 and autocovariances γ_j are all time-invariant; hence the model is weakly stationary.

Stationarity Condition


Q: What is the stationarity condition for an $AR(1)$ model?

A: For $AR(1)$, we want the root of the equation $1 - \phi_1 x = 0$ to have magnitude greater than one, i.e., $|x| = \frac{1}{|\phi_1|} > 1$, so the stationarity condition is $|\phi_1| < 1$.

Q: How about $AR(2)$?

Hint: The polynomial equation is $1 - \phi_1 x - \phi_2 x^2 = 0$. Let $z = \frac{1}{x}$. The polynomial equation then becomes $z^2 - \phi_1 z - \phi_2 = 0$. Now impose the condition $|x| > 1$, which is equivalent to $|z| < 1$. Let the two roots be z_1 and z_2 , where $z_1 < z_2$. Then from the conditions $z_1 > -1$, $z_2 < 1$ and $|z_1 z_2| < 1$, we obtain $\phi_2 - \phi_1 < 1$, $\phi_1 + \phi_2 < 1$ and $|\phi_2| < 1$.

Stationarity Condition of AR(2)



Invertibility Condition

$\{y_t\}$ is said to be *invertible* if it can be expressed as an $AR(\infty)$ process of the form $[1 - \pi(L)] u_t = \varepsilon_t$.

Invertibility condition: The $ARMA(p, q)$ model is invertible iff all the roots of the polynomial equation $1 + \theta(x) = 0$ lie outside the unit circle.

In this case, $1 - \pi(L) := \frac{1 - \phi(L)}{1 + \theta(L)}$ is a well-defined infinite-order polynomial.

Q: What is the invertibility condition for an $MA(1)$ model?

A: For $MA(1)$, the invertibility condition is $|\theta_1| < 1$.

Ex: What is the invertibility condition of an $MA(2)$ model?

Ex: Is a stationary $AR(p)$ invertible?

Ex: Is an invertible $MA(q)$ stationary?

AR Representation

Given an $ARMA(p, q)$ model:

$$\begin{aligned}y_t &= \mu + u_t, \\[1 - \phi(L)] u_t &= [1 + \theta(L)] \varepsilon_t.\end{aligned}$$

Suppose the invertibility condition holds, so that $\frac{1}{1+\theta(L)}$ is a well-defined infinite-order polynomial. Define $1 + \pi(L) = \frac{1-\phi(L)}{1+\theta(L)}$. Then we can rewrite the model as $AR(\infty)$:

$$\begin{aligned}\varepsilon_t &= \frac{1 - \phi(L)}{1 + \theta(L)} u_t \\&= [1 + \pi(L)] u_t \\&= u_t + \pi_1 u_{t-1} + \pi_2 u_{t-2} + \cdots.\end{aligned}$$

MA Representation

Given an $ARMA(p, q)$ model:

$$\begin{aligned}y_t &= \mu + u_t, \\[1 - \phi(L)] u_t &= [1 + \theta(L)] \varepsilon_t.\end{aligned}$$

Suppose the stationarity condition holds, so that $\frac{1}{1-\phi(L)}$ is a well-defined infinite-order polynomial. Define $1 + \psi(L) = \frac{1+\theta(L)}{1-\phi(L)}$. Then we can rewrite the model as $MA(\infty)$:

$$\begin{aligned}u_t &= \frac{1 + \theta(L)}{1 - \phi(L)} \varepsilon_t \\&= [1 + \psi(L)] \varepsilon_t \\&= \varepsilon_t + \psi_1 \varepsilon_{t-1} + \psi_2 \varepsilon_{t-2} + \cdots.\end{aligned}\tag{5}$$

Interpretation: ψ_j as a function of j is the *impulse response function*. ψ_j is the marginal impact of the lag j shock ε_{t-j} on the current observation y_t .

MA Representation

Q: Let $\{y_t\}$ be a weakly stationary process. What happens to the ℓ -step ahead forecast and its variance as $\ell \rightarrow \infty$?

A: By weak stationarity, $\{y_t\}$ can be expressed as an $MA(\infty)$ process (5), so that

$$\begin{aligned} \text{Var}(y_t) &= \text{Var}(\varepsilon_t + \psi_1 \varepsilon_{t-1} + \psi_2 \varepsilon_{t-2} + \cdots) \\ &= \sigma_\varepsilon^2 \left(1 + \sum_{i=1}^{\infty} \psi_i^2 \right) < \infty. \end{aligned}$$

This implies that $\psi_i \rightarrow 0$ as $i \rightarrow \infty$.

It follows from (1) with $q = \infty$ that the ℓ -step ahead forecast is

$$\begin{aligned} y_t(\ell) &= \mu + \psi_\ell \varepsilon_t + \psi_{\ell+1} \varepsilon_{t-1} + \cdots \\ &\rightarrow \mu \text{ as } \ell \rightarrow \infty. \end{aligned}$$

This is the *mean-reverting property*. The forecast variance is, by (3) with $q = \infty$,

$$\begin{aligned} \text{Var}[e_t(\ell)] &\rightarrow (1 + \psi_1^2 + \psi_2^2 + \cdots) \sigma_\varepsilon^2 \\ &= \text{Var}(y_t) \text{ as } \ell \rightarrow \infty. \end{aligned}$$

Model Checking

If a model is correctly specified, the residual process $\{\hat{\varepsilon}_t\}$ should look like a white noise.

To test $H_0 : \rho_\ell = 0$ vs $H_a : \rho_\ell \neq 0$, we get the sample ACF $\hat{\rho}_\ell$ of $\{\hat{\varepsilon}_t\}$.

- Suppose $\rho_j = 0$ for all $j > \ell$ under H_0 . Then,

$$\sqrt{T}\hat{\rho}_\ell \xrightarrow{d} N\left(0, 1 + 2\sum_{j=1}^{\ell-1} \rho_j^2\right) \text{ as } T \rightarrow \infty.$$

The t ratio converges in distribution to standard normal:

$$t = \frac{\hat{\rho}_\ell}{\sqrt{\left(1 + 2\sum_{j=1}^{\ell-1} \hat{\rho}_j^2\right) / T}} \xrightarrow{d} N(0, 1) \text{ as } T \rightarrow \infty.$$

- Suppose $\{\varepsilon_t\} \sim wn(0, \sigma_\varepsilon^2)$ under H_0 . Then all $\rho_j = 0$ for all $j \neq 0$, and so

$$\sqrt{T}\hat{\rho}_\ell \xrightarrow{d} N(0, 1) \text{ as } T \rightarrow \infty.$$

This is a two-sided test (reject H_0 at α level if $|t| > z_{\alpha/2}$).

Portmanteau Tests

Set a maximum lag m . We want to test $H_0 : \rho_1 = \dots = \rho_m = 0$ vs $H_a : \rho_j \neq 0$ for some $j = 1, \dots, m$.

Assume $\{y_t\} \sim ARMA(p, q)$, with $\{\varepsilon_t\} \sim iid$ and some moment conditions.

Box-Pierce test on $\{\hat{\varepsilon}_t\}$:

$$Q^*(m) = T \sum_{\ell=1}^m \hat{\rho}_{\ell}^2 \xrightarrow{d} \chi^2(m - p - q) \text{ as } T \rightarrow \infty.$$

Ljung-Box test on $\{\hat{\varepsilon}_t\}$:

$$Q(m) = T(T+2) \sum_{\ell=1}^m \frac{\hat{\rho}_{\ell}^2}{T-\ell} \xrightarrow{d} \chi^2(m - p - q) \text{ as } T \rightarrow \infty.$$

They are one-sided tests (reject H_0 at α level if $Q(m) > \chi_{\alpha}^2$).