

# The category of representations of a concrete category as a functor category

Tibor Grün

March 22, 2021

A quiver (also called directed graph)  $q$  consists of a class of objects (or vertices)  $q_0 = \text{Obj } q$

A quiver (also called directed graph)  $q$  consists of a class of objects (or vertices)  $q_0 = \text{Obj } q$

1

3

4

5

2

A quiver (also called directed graph)  $q$  consists of a class of objects (or vertices)  $q_0 = \text{Obj } q$  and a class of morphisms (or arrows)  $q_1 = \text{Mor } q$

1


3

4

5


2

A quiver (also called directed graph)  $q$  consists of a class of objects (or vertices)  $q_0 = \text{Obj } q$  and a class of morphisms (or arrows)  $q_1 = \text{Mor } q$




together with two defining maps  $s : q_1 \longrightarrow q_0$  called source,  
and  $t : q_1 \longrightarrow q_0$  called target.

together with two defining maps  $s : q_1 \longrightarrow q_0$  called source,  
and  $t : q_1 \longrightarrow q_0$  called target.



together with two defining maps  $s : q_1 \longrightarrow q_0$  called source,  
and  $t : q_1 \longrightarrow q_0$  called target.



For example

$$s(a) = s(b) = 1$$

$$t(b) = 2$$


$$t(a) = t(c) = t(d) = t(f) = 3$$

etc.




In the finite case - which we will be dealing with most of the time here - we call  $q_0$  set of objects and  $q_1$  set of morphisms.

In the finite case - which we will be dealing with most of the time here - we call  $q_0$  set of objects and  $q_1$  set of morphisms.




In the finite case - which we will be dealing with most of the time here - we call  $q_0$  set of objects and  $q_1$  set of morphisms.



So in the quiver  $q$  we have

In the finite case - which we will be dealing with most of the time here - we call  $q_0$  set of objects and  $q_1$  set of morphisms.



So in the quiver  $q$  we have

$$q_0 = \{1, 2, 3, 4, 5\}$$

and


$$q_1 = \{a, b, c, d, e, f\}$$

Another map relates the arrows with the objects. That is the Hom-set, i.e. set of morphisms, between two objects (order matters):

$$\text{Hom} : q_0 \times q_0 \longrightarrow \mathcal{P}(q_1)$$


Another map relates the arrows with the objects. That is the Hom-set, i.e. set of morphisms, between two objects (order matters):

$$\text{Hom} : q_0 \times q_0 \longrightarrow \mathcal{P}(q_1)$$



Another map relates the arrows with the objects. That is the Hom-set, i.e. set of morphisms, between two objects (order matters):

$$\text{Hom} : q_0 \times q_0 \longrightarrow \mathcal{P}(q_1)$$



$$\text{Hom}(1, 3) = \{a\}$$


$$\text{Hom}(4, 3) = \{c, d\}$$

$$\text{Hom}(3, 4) = \{\}$$


For the same object  $X \in \mathcal{Q}_0$  we call  $\text{Hom}(X, X) = \text{End}(X)$  the endomorphism set.



For the same object  $X \in \mathcal{Q}_0$  we call  $\text{Hom}(X, X) = \text{End}(X)$  the endomorphism set.



For the same object  $X \in \mathcal{C}_0$  we call  $\text{Hom}(X, X) = \text{End}(X)$  the endomorphism set.



$$\text{End}(1) = \{a\}$$

$$\text{End}(2) = \{c\}$$

$$\text{Hom}(1, 2) = \{b\}$$

You now know what a quiver is.

A category  $\mathcal{C}$  is a quiver with two further maps:

(1) For every object  $X \in \mathcal{C}_0$  there is the identity map

$$\begin{aligned} \mathbb{1} &: \mathcal{C}_0 \longrightarrow \mathcal{C}_1 \\ X &\longmapsto \mathbb{1}_X : X \longrightarrow X \end{aligned}$$

A category  $\mathcal{C}$  is a quiver with two further maps:

( $\mathbb{1}$ ) For every object  $X \in \mathcal{C}_0$  there is the identity map

$$\begin{aligned}\mathbb{1} &: \mathcal{C}_0 \longrightarrow \mathcal{C}_1 \\ X &\longmapsto \mathbb{1}_X : X \longrightarrow X\end{aligned}$$

( $\mu$ ) For two composable morphisms  $\varphi$  and  $\psi \in \mathcal{C}_1$ , i.e. with  $t(\varphi) = s(\psi)$  there is the composition map

$$\begin{aligned}\mu &: \mathcal{C}_1 \times \mathcal{C}_1 \longrightarrow \mathcal{C}_1 \\ \varphi &: A \longrightarrow B \\ \psi &: B \longrightarrow C \\ (\varphi, \psi) &\longmapsto \mu(\varphi, \psi) := \varphi\psi : A \longrightarrow C\end{aligned}$$

The defining properties for  $\mathbb{1}$  and  $\mu$  are:

1.  $s(\mathbb{1}_M) = M = t(\mathbb{1}_M)$ , i.e.  $\mathbb{1}_M \in \text{End}(M)$ .

The defining properties for  $\mathbb{1}$  and  $\mu$  are:

1.  $s(\mathbb{1}_M) = M = t(\mathbb{1}_M)$ , i.e.  $\mathbb{1}_M \in \text{End}(M)$ .
2.  $s(\varphi\psi) = s(\varphi)$  and  $t(\varphi\psi) = t(\psi)$ , i.e. for objects  $M, L, N \in \mathcal{C}_0$  we have

$$\mu : \text{Hom}(M, L) \times \text{Hom}(L, N) \longrightarrow \text{Hom}(M, N)$$

The defining properties for  $\mathbb{1}$  and  $\mu$  are:

1.  $s(\mathbb{1}_M) = M = t(\mathbb{1}_M)$ , i.e.  $\mathbb{1}_M \in \text{End}(M)$ .
2.  $s(\varphi\psi) = s(\varphi)$  and  $t(\varphi\psi) = t(\psi)$ , i.e. for objects  $M, L, N \in \mathcal{C}_0$  we have

$$\mu : \text{Hom}(M, L) \times \text{Hom}(L, N) \longrightarrow \text{Hom}(M, N)$$

3.  $(\varphi\psi)\rho = \varphi(\psi\rho)$ , i.e. composition is associative



The defining properties for  $\mathbb{1}$  and  $\mu$  are:

1.  $s(\mathbb{1}_M) = M = t(\mathbb{1}_M)$ , i.e.  $\mathbb{1}_M \in \text{End}(M)$ .
2.  $s(\varphi\psi) = s(\varphi)$  and  $t(\varphi\psi) = t(\psi)$ , i.e. for objects  $M, L, N \in \mathcal{C}_0$  we have

$$\mu : \text{Hom}(M, L) \times \text{Hom}(L, N) \longrightarrow \text{Hom}(M, N)$$

3.  $(\varphi\psi)\rho = \varphi(\psi\rho)$ , i.e. composition is associative
4.  $\mathbb{1}_{s(\varphi)}\varphi = \varphi = \varphi\mathbb{1}_{t(\varphi)}$ , i.e. the identity is a left and right unit of the composition.

The defining properties for  $\mathbb{1}$  and  $\mu$  are:

1.  $s(\mathbb{1}_M) = M = t(\mathbb{1}_M)$ , i.e.  $\mathbb{1}_M \in \text{End}(M)$ .
2.  $s(\varphi\psi) = s(\varphi)$  and  $t(\varphi\psi) = t(\psi)$ , i.e. for objects  $M, L, N \in \mathcal{C}_0$  we have

$$\mu : \text{Hom}(M, L) \times \text{Hom}(L, N) \longrightarrow \text{Hom}(M, N)$$

3.  $(\varphi\psi)\rho = \varphi(\psi\rho)$ , i.e. composition is associative
4.  $\mathbb{1}_{s(\varphi)}\varphi = \varphi = \varphi\mathbb{1}_{t(\varphi)}$ , i.e. the identity is a left and right unit of the composition.

These properties make each endomorphism set  $\text{End}(M)$  for  $M \in \mathcal{C}$  together with the composition into a monoid, called the endomorphism monoid  $(\text{End}(M), \mu)$ .

So when you define a category, you always answer the four questions

- ▶ What are the objects?
- ▶ What are the morphisms? Especially what are the identity morphisms?
- ▶ How do you compose morphisms?
- ▶ Why is the composition associative? Why is the identity a unit for the composition?

You now know what a category is

A small example for a category: The symmetric group on two objects  $S_2$ .

$$\mathbb{1}_{\{1,2\}} \left( \begin{array}{c} \curvearrowright \\ \{1,2\} \\ \curvearrowleft \end{array} \right) (1,2)$$

The rule that the composition of  $(1,2)$  with itself results in the identity  $\mathbb{1}_{\{1,2\}}$  makes sure there are only 2 morphisms in total.

A small example for a category: The symmetric group on two objects  $S_2$ .

$$\mathbb{1}_{\{1,2\}} \left( \begin{array}{c} \curvearrowright \\ \{1,2\} \\ \curvearrowleft \end{array} \right) (1,2)$$

The rule that the composition of  $(1, 2)$  with itself results in the identity  $\mathbb{1}_{\{1,2\}}$  makes sure there are only 2 morphisms in total. This is an example of a category which is a sub-category of the category **SETS** with sets as objects and functions between sets as morphisms. Between those categories lies the category **FINSETS** in which the objects are finite sets and morphisms are functions between finite sets.

As another example of a finite subcategory of  $\mathbf{FINSETS}$ , take the category with two objects that are sets with 3 elements.

As another example of a finite subcategory of  $\mathbf{FINSETS}$ , take the category with two objects that are sets with 3 elements.

$$\{1, 2, 3\}$$



As another example of a finite subcategory of  $\mathbf{FINSETS}$ , take the category with two objects that are sets with 3 elements.

$\{1, 2, 3\}$


$\{4, 5, 6\}$

As another example of a finite subcategory of  $\mathbf{FinSets}$ , take the category with two objects that are sets with 3 elements. The endomorphisms at each object are the cyclic permutations on the 3 elements  $\{1, 2, 3\}$


$\{1, 2, 3\}$

$\{4, 5, 6\}$

As another example of a finite subcategory of  $\mathbf{FinSets}$ , take the category with two objects that are sets with 3 elements. The endomorphisms at each object are the cyclic permutations on the 3 elements  $\{1, 2, 3\}$



As another example of a finite subcategory of  $\mathbf{FINSETS}$ , take the category with two objects that are sets with 3 elements. The endomorphisms at each object are the cyclic permutations on the 3 elements  $\{1, 2, 3\}$  and on the 3 elements  $\{4, 5, 6\}$ .




As another example of a finite subcategory of  $\mathbf{FINSETS}$ , take the category with two objects that are sets with 3 elements. The endomorphisms at each object are the cyclic permutations on the 3 elements  $\{1, 2, 3\}$  and on the 3 elements  $\{4, 5, 6\}$ .




As another example of a finite subcategory of  $\mathbf{FINSETS}$ , take the category with two objects that are sets with 3 elements. The endomorphisms at each object are the cyclic permutations on the 3 elements  $\{1, 2, 3\}$  and on the 3 elements  $\{4, 5, 6\}$ . There is also a morphism from the first to the second object, mapping  $1 \mapsto 4, 2 \mapsto 5$  and  $3 \mapsto 6$ .



As another example of a finite subcategory of  $\mathbf{FINSETS}$ , take the category with two objects that are sets with 3 elements. The endomorphisms at each object are the cyclic permutations on the 3 elements  $\{1, 2, 3\}$  and on the 3 elements  $\{4, 5, 6\}$ . There is also a morphism from the first to the second object, mapping  $1 \mapsto 4, 2 \mapsto 5$  and  $3 \mapsto 6$ .




As another example of a finite subcategory of  $\mathbf{FINSETS}$ , take the category with two objects that are sets with 3 elements. The endomorphisms at each object are the cyclic permutations on the 3 elements  $\{1, 2, 3\}$  and on the 3 elements  $\{4, 5, 6\}$ . There is also a morphism from the first to the second object, mapping  $1 \mapsto 4, 2 \mapsto 5$  and  $3 \mapsto 6$ .




The endomorphism monoid on each object is the group  $C_3$ , i.e. the cyclic group on three elements. This also means that each endomorphism is invertible. We thus call this category  $C_3 C_3$ .



You may have noticed that this picture is not complete when we look back to the axioms for a category: We have three endomorphisms at the first object and three endomorphisms at the second object. Since we also have a morphism from the first to the second object, we also need all the possible compositions of those morphisms.




You may have noticed that this picture is not complete when we look back to the axioms for a category: We have three endomorphisms at the first object and three endomorphisms at the second object. Since we also have a morphism from the first to the second object, we also need all the possible compositions of those morphisms.



Of the missing two morphisms from  $\{1, 2, 3\}$  to  $\{4, 5, 6\}$ , the first one is mapping  $1 \mapsto 5, 2 \mapsto 6$  and  $3 \mapsto 4$ , where the last one is mapping  $1 \mapsto 6, 2 \mapsto 4$  and  $3 \mapsto 5$ .


You may have noticed that this picture is not complete when we look back to the axioms for a category: We have three endomorphisms at the first object and three endomorphisms at the second object. Since we also have a morphism from the first to the second object, we also need all the possible compositions of those morphisms.




Of the missing two morphisms from  $\{1, 2, 3\}$  to  $\{4, 5, 6\}$ , the first one is mapping  $1 \mapsto 5, 2 \mapsto 6$  and  $3 \mapsto 4$ , where the last one is mapping  $1 \mapsto 6, 2 \mapsto 4$  and  $3 \mapsto 5$ . So our category  $C_3 C_3$  has two objects  $\{1, 2, 3\}, \{4, 5, 6\}$  and nine morphisms in total.

The categories  $S_2$ ,  $C_3$ ,  $C_3$ , SET, FINSETS can themselves all be considered objects in a greater category, CAT, i.e. the category of categories.

The categories  $S_2$ ,  $C_3$ ,  $C_3$ ,  $SET$ ,  $FINSETS$  can themselves all be considered objects in a greater category,  $CAT$ , i.e. the category of categories.



The categories  $S_2$ ,  $C_3$ ,  $C_3$ ,  $SET$ ,  $FINSET$  can themselves all be considered objects in a greater category,  $CAT$ , i.e. the category of categories.



We know what the objects are in  $CAT$ . But what are the morphisms? What is meant with an arrow from  $SETS$  to  $FINSETS$ ?

You now know the objects in the category  $\mathbf{CAT}$  of all categories.

A functor  $F : \mathcal{C} \rightarrow \mathcal{D}$ , between categories  $\mathcal{C}$  and  $\mathcal{D}$ , consists of the following data:

- ▶ An object  $Fc \in \mathcal{D}_0$ , for each object  $c \in \mathcal{C}_0$ .
- ▶ A function  $Ff : Fc \rightarrow Fc' \in \mathcal{D}_1$ , for each morphism  $f : c \rightarrow c' \in \mathcal{C}_1$ , so that the source and target of  $Ff$  are, respectively, equal to  $F$  applied to the source or target of  $f$ , in other words,  $s(Ff) = Fs(f)$  and  $t(Ff) = Ft(f)$ .

The assignments are required to satisfy the following two functoriality axioms:

- ▶ For any composable pair  $f : M \rightarrow N, g : N \rightarrow L \in \mathcal{C}_1$ ,  $Ff \cdot Fg = F(f \cdot g)$ .
- ▶ For each object  $c \in \mathcal{C}_0$ ,  $F(1_c) = 1_{Fc}$ .



So with functors you always answer the four questions

- ▶ How does it work on objects?
- ▶ How does it work on morphisms?
- ▶ Why does it respect composition?
- ▶ Why does it respect identity morphisms?

You now know what a functor is.

You now know the objects and morphisms in the category  $\mathbf{CAT}$  of all categories.

Let us now take a look at just two categories,  $\mathcal{C}$  and  $\mathcal{D}$  as objects in  $\mathbf{CAT}$ .

The Hom-set  $\mathrm{Hom}(\mathcal{C}, \mathcal{D})$  of all functors  $F : \mathcal{C} \longrightarrow \mathcal{D}$  is itself a category, called the functor category.

This makes the functors  $F, G, H : \mathcal{C} \longrightarrow \mathcal{D}$  objects in  $\mathrm{Hom}(\mathcal{C}, \mathcal{D})$  when before they were considered morphisms.

Let us now take a look at just two categories,  $\mathcal{C}$  and  $\mathcal{D}$  as objects in  $\mathbf{CAT}$ .

The Hom-set  $\text{Hom}(\mathcal{C}, \mathcal{D})$  of all functors  $F : \mathcal{C} \longrightarrow \mathcal{D}$  is itself a category, called the functor category.

This makes the functors  $F, G, H : \mathcal{C} \longrightarrow \mathcal{D}$  objects in  $\text{Hom}(\mathcal{C}, \mathcal{D})$  when before they were considered morphisms.

$F \qquad G \qquad H$

Let us now take a look at just two categories,  $\mathcal{C}$  and  $\mathcal{D}$  as objects in  $\mathbf{CAT}$ .

The Hom-set  $\text{Hom}(\mathcal{C}, \mathcal{D})$  of all functors  $F : \mathcal{C} \longrightarrow \mathcal{D}$  is itself a category, called the functor category.

This makes the functors  $F, G, H : \mathcal{C} \longrightarrow \mathcal{D}$  objects in  $\text{Hom}(\mathcal{C}, \mathcal{D})$  when before they were considered morphisms.

$F \qquad G \qquad H$

As you can imagine, we are again looking for the morphisms in this category, i.e. what are morphisms between functors?

Let us now take a look at just two categories,  $\mathcal{C}$  and  $\mathcal{D}$  as objects in  $\mathbf{CAT}$ .

The Hom-set  $\text{Hom}(\mathcal{C}, \mathcal{D})$  of all functors  $F : \mathcal{C} \longrightarrow \mathcal{D}$  is itself a category, called the functor category.

This makes the functors  $F, G, H : \mathcal{C} \longrightarrow \mathcal{D}$  objects in  $\text{Hom}(\mathcal{C}, \mathcal{D})$  when before they were considered morphisms.

$$F \xRightarrow{\alpha} G \xRightarrow{\beta} H$$

As you can imagine, we are again looking for the morphisms in this category, i.e. what are morphisms between functors?

You now know the objects in the category  $\text{Hom}(\mathcal{C}, \mathcal{D})$  of all functors between categories  $\mathcal{C}$  and  $\mathcal{D}$ .



Given categories  $\mathcal{C}$  and  $\mathcal{D}$  and functors  $F : \mathcal{C} \rightarrow \mathcal{D}$  and  $G : \mathcal{C} \rightarrow \mathcal{D}$ , a natural transformation  $\alpha : F \Rightarrow G$  consists of:

- ▶ a morphism  $\alpha_c : Fc \rightarrow Gc \in \mathcal{D}_1$  for each object  $c \in \mathcal{C}_0$ , the collection of which define the components of the natural transformation, so that, for any morphism  $f : c \rightarrow c' \in \mathcal{C}_1$ , the following square of morphisms in  $\mathcal{D}$

$$\begin{array}{ccc} Fc & \xrightarrow{\alpha_c} & Gc \\ Ff \downarrow & & \downarrow Gf \\ Fc' & \xrightarrow{\alpha_{c'}} & Gc' \end{array}$$

commutes, i.e., has a common composite  $Fc \rightarrow Gc' \in \mathcal{D}_1$ . This means explicitly that

$$Ff\alpha_{c'} = \alpha_c Gf, \forall f : c \rightarrow c' \in \mathcal{C}_1.$$

You now know what a natural transformation is.

You now know the objects and morphisms in the category  $\text{Hom}(\mathcal{C}, \mathcal{D})$  of all functors between categories  $\mathcal{C}$  and  $\mathcal{D}$ .

Now that we know what a functor category is, we want to work towards a special kind of functor category: Given a finite concrete category  $\mathcal{C}$  whose endomorphism monoids are explicitly cyclic, we can calculate its  $\mathbb{k}$ -linear closure, i.e. the  $\mathbb{k}$ -Algebroid  $\mathcal{A}$ .

Then we can calculate the category of  $\mathbb{k}$ -linear functors from the  $\mathbb{k}$ -Algebroid  $\mathcal{A}$  into the matrix category  $\mathbb{k}\text{-Mat}$  over the same field  $\mathbb{k}$ .

$$\mathrm{Hom}_{\mathbb{k}}(\mathcal{A}, \mathbb{k}\text{-Mat})$$

category of  $\mathbb{k}$ -linear functors  $\text{Hom}_{\mathbb{k}}(\mathcal{A}, \mathbb{k}\text{-Mat})$

FinSets



finite concrete category  $\mathcal{C}$

category of  $\mathbb{k}$ -linear functors  $\text{Hom}_{\mathbb{k}}(\mathcal{A}, \mathbb{k}\text{-Mat})$

$\mathbf{FinSets}$




finite concrete category  $\mathcal{C}$


field  $\mathbb{k}$




$\mathbb{k}$ -Algebroid  $\mathcal{A}$




category of  $\mathbb{k}$ -linear functors  $\mathrm{Hom}_{\mathbb{k}}(\mathcal{A}, \mathbb{k}\text{-Mat})$











Category $\mathcal{C}$	$\text{Hom}_{\mathbb{k}}(\mathcal{A}, \mathbb{k}\text{-Mat})$	$\mathbb{k}\text{-mat}$
	$\mathbb{k}$ -linear functors $F :$	
<b>Objects</b> $\mathcal{C}_0$	$\mathcal{A} \rightarrow \mathbb{k}\text{-mat}$ $c \in \mathcal{A}_0 \mapsto Fc \in \mathbb{k}\text{-mat}_0$ $\varphi \in \mathcal{A}_1 \mapsto F\varphi \in \mathbb{k}\text{-mat}_1$	natural numbers $\mathbb{N}_0$
<b>Morphisms</b> $\mathcal{C}_1$	natural transformations, components are matrices	$m \times n$ -matrices
<b>Composition:</b>		
$\varphi : A \rightarrow B,$	$\eta : F \Rightarrow G, \eta_c : Fc \rightarrow Gc$	
$\psi : B \rightarrow C$	$\varepsilon : G \Rightarrow H, \varepsilon_c : Gc \rightarrow Hc$	
$\varphi\psi : A \rightarrow C$	$\eta\varepsilon : F \Rightarrow H, \eta_c\varepsilon_c : Fc \rightarrow Hc$	
	(component-wise)	matrix multiplication

**Category  $\mathcal{C}$**

$\text{Hom}_{\mathbb{k}}(\mathcal{A}, \mathbb{k}\text{-Mat})$

$\mathbb{k}\text{-mat}$

---

**Direct Sum:**

Let  $I = \{1, \dots, N\}$   
be a finite set

$\{F_i\}_{i \in I}$  a family  
of objects in  
 $\text{Hom}_{\mathbb{k}}(\mathcal{A}, \mathbb{k}\text{-Mat})_0$

$\{n_i\}_{i \in I}$  a family  
of objects in  $\mathbb{k}\text{-mat}_0$

$$F : c \mapsto \sum_{i=1}^N F_i(c) \quad n = \sum_{i=1}^N n_i$$

at each object the  
sum of natural  
numbers at that object

sum of natural  
numbers

Projections:

$\pi_i : F \rightarrow F_i$  with components

$$(\pi_i)_c := \begin{pmatrix} 0_{F_{<i}(c), F_i(c)} \\ 1_{F_i(c)} \\ 0_{F_{>i}(c), F_i(c)} \end{pmatrix}$$

Coprojections:

$\iota_i : F_i \rightarrow F$  with components

$$(\iota_i)_c := \begin{pmatrix} 0_{F_i(c), F_{<i}(c)} & 1_{F_i(c)} & 0_{F_i(c), F_{>i}(c)} \end{pmatrix}$$

For a morphism  $a : c \rightarrow c' \in \mathcal{A}_1$  we have a family of morphisms  $\{F_i a : F_i c \rightarrow F_i c'\}_{i \in I}$ . Then the direct sum  $F$  is defined on morphisms as

$$Fa := \sum_{i \in I} (\pi_i)_c F_i a (\iota_i)_{c'} : Fc \rightarrow Fc'$$

which satisfies

$$\begin{aligned} (\iota_i)_c Fa (\pi_i)_{c'} &= (\iota_i)_c \sum_{j \in I} (\pi_j)_c F_j a (\iota_j)_{c'} (\pi_i)_{c'} \\ &= \sum_{j \in I} (\iota_i)_c (\pi_j)_c F_j a (\iota_j)_{c'} (\pi_i)_{c'} \\ &= \sum_{j \in I} (\delta_{i,j})_c F_j a (\delta_{j,i})_{c'} \\ &= 1_{F_i c} F_i a 1_{F_i c'} \\ &= F_i a \end{aligned}$$

One step in the decomposition algorithm looks like this:

$$\begin{array}{ccccc}
 & & \eta & \rightarrow & F \\
 & \nearrow & & & \nwarrow \eta \\
 F & & & & F \\
 \uparrow \iota & & I \oplus K = F & & \uparrow \kappa \\
 I & \text{---} & \oplus & \text{---} & K
 \end{array}$$

Then we have two morphisms  $\iota, \kappa$  with target  $F$  and sources  $I, K$  such that  $I \oplus K = F$ .



---

**Algorithm 1:** DecomposeOnceByRandomEndomorphism

---

**Input :** a functor  $F$  in a functor category

**Output :** a pair  $[\iota : I \rightarrow F, \kappa : K \rightarrow F]$  of morphisms  
such that  $I \oplus K = F$  with  $I \neq 0$  and  $K \neq 0$  or  
fail if it was unable to further decompose  $F$ ;

```
1  $d := \max\{\dim_{\mathbb{k}} Fc\}_{c \in \mathcal{A}_0};$   
2 if  $d = 0$  then  
3   | return fail;      // the zero representation is  
   | indecomposable  
4 end  
5  $\mathcal{B} = [\beta_1, \dots, \beta_h]$  is a  $\mathbb{k}$ -basis of  $\text{Hom}_{\text{Hom}_{\mathbb{k}}(\mathcal{A}, \mathbb{k}\text{-Mat})}(F, F);$   
6 add  $0_{F,F}$  to  $\mathcal{B};$ 
```

---

---

---

```

7   $n := \lfloor \log_2(d) \rfloor + 1;$ 
8  for  $b \in [h + 1, h, \dots, 2]$  do
9       $\alpha := \beta_b + \text{random}(\mathbb{k}) \cdot \beta_{b-1};$            // a heuristic
        ansatz for a random endomorphism
10     for  $i \in [1, \dots, n]$  do
11          $\alpha_2 := \alpha^2;$ 
13         /* We do not expect the exponentiation to
           produce an idempotent, still this is a
           very cheap test:                                */
14         if  $\alpha = \alpha_2$  then
15             break;
16         end
17          $\alpha := \alpha_2;$ 
18     end
19     if  $\alpha = 0$  then
20         continue ;           // try another endomorphism
21     end

```

---


---

```
8 for  $b \in [h + 1, h, \dots, 2]$  do
9   ...
22    $\kappa := \text{KernelEmbedding}(\alpha)$ ;
23   if  $\kappa = 0$  then
24     | continue ;           // try another endomorphism
25   end
26    $\iota := \text{ImageEmbedding}(\alpha)$ ;
27   return  $[\iota, \kappa]$ ;
28 end

29 return fail;           // The input functor  $F$  is
    indecomposable with a high probability.
```

---

# Direct sum decomposition



---

**Algorithm 2:** WeakDirectSumDecomposition

---

**Input :** a functor  $F$  in a functor category

**Output :** a list  $[\eta_i : F_i \rightarrow F]$  of embeddings.

```
1 queue :=  $[1_F]$ ;
2 summands :=  $\emptyset$ ;
3 while queue  $\neq \emptyset$  do
4   | let  $\eta = \text{remove}(\text{queue})$ ;
5   | result :=
      DecomposeOnceByRandomEndomorphism( $s(\eta)$ );
6   | if result = fail then      //  $s(\eta)$  indecomposable
7   |   | add  $\eta$  to summands;
8   | else
9   |   |  $[\iota, \kappa] = \text{result}$ ;
10  |   | append  $[\iota\eta, \kappa\eta]$  to queue;
11  | end
12 end
13 return summands;
```

Hom-based invariants:

EmbeddingOfSumOfImagesOfAllMorphisms:

$$\begin{array}{ccc} \bigoplus_{i \in \text{hom}} M & \xrightarrow{u_{\text{out}}(\text{hom})} & N \\ & \searrow \pi & \nearrow \iota \\ & I & \end{array}$$

