

Assignment 3

Meriam Elabor, 1076589

March 22, 2018

Question 1.1

1-KNN		
Record	Distance from the record	Label of the record
6	1.29007933	-1
Prediction		-1

Question 1.2

3-KNN		
Record	Distance from the record	Label of the record
26	1.36993215	1
21	1.65639809	1
39	2.09803532	1
Prediction		1

Question 1.3

5-KNN		
Record	Distance from the record	Label of the record
29	3.23194266	1
8	5.03028237	-1
36	5.2278505	1
37	5.23353427	1
39	5.61435592	1
Prediction		1

Question 1.4

Testing Error Rates								
1-NN Rate	Test	Error	3-NN Rate	Test	Error	5-NN Rate	Test	Error
0.025			0.05			0.075		

Question 2.1

$$h_{\Theta}(x) \in (1, -1)$$

Question 2.2

let $J(\Theta) = \frac{1}{N} \sum_{n=1}^N (h_{\Theta}(x^n) - y^n)^2$, $h_{\Theta}(x^n) = \frac{2}{1+e^{\Theta^T \cdot x}} - 1$, $g_{\Theta}(x^n) = \frac{1}{1+e^{\Theta^T \cdot x}}$,
then $\forall k = \{0, \dots, 4\}$

$$\begin{aligned} \frac{\partial J(\Theta)}{\partial \theta_k} &= \frac{\partial J(\Theta)}{\partial h_{\Theta}(x^n)} \cdot \frac{\partial h_{\Theta}(x^n)}{\partial \theta_k} \\ \frac{\partial J(\Theta)}{\partial h_{\Theta}(x^n)} &= \frac{2}{N} \sum_{n=1}^N (h_{\Theta}(x^n) - y^n) \\ \frac{\partial h_{\Theta}(x^n)}{\partial \theta_k} &= \frac{2x_k e^{\Theta^T \cdot x}}{(1+e^{\Theta^T \cdot x})^2} = 2x_k g_{\Theta}(x^n)(1 - g_{\Theta}(x^n)) \\ \text{thus } \frac{\partial J(\Theta)}{\partial \theta_k} &= 4x_k \frac{1}{N} \sum_{n=1}^N (h_{\Theta}(x^n) - y^n) g_{\Theta}(x^n)(1 - g_{\Theta}(x^n)) \end{aligned}$$

$$\text{and } \theta_k \rightarrow \theta_k - \alpha 4x_k \frac{1}{N} \sum_{n=1}^N (h_{\Theta}(x^n) - y^n) g_{\Theta}(x^n)(1 - g_{\Theta}(x^n)), \forall k = \{0, \dots, 4\}$$

Question 2.3

Question 4			
Testing record	Classifier $h_{\Theta}(x^{(n)})$	Output	Final Output
5	-0.949479		-1
10	-0.991887		-1
15	-0.996641		-1
20	-0.992353		-1
25	0.893525		1
30	0.968980		1
35	0.991332		1
40	0.983889		1

Question 3.1

let $f(X) = \frac{1}{1+e^X}$, $X \in R$ then
 $h_{\Theta}(x) = g_2 = f(g_1\theta_6 + x_0\theta_5)$, $h_{\Theta}(x) \in (0, 1)$
 $g_1 = f(x_0\theta_0 + x_1\theta_1 + x_2\theta_2 + x_3\theta_3 + x_4\theta_4)$

Question 3.2

let $f(X) = \frac{1}{1+e^X}$, $X \in R$ and
 $J(\Theta) = (h_\Theta(x) - y)^2$ then
 $\frac{d}{dX}f(X) = f(X)(1 - f(X))$
and $\forall k = \{5, 6\}$

$$\begin{aligned}\frac{\partial J(\Theta)}{\partial \theta_k} &= \frac{\partial J(\Theta)}{\partial h_\Theta(x)} \cdot \frac{\partial h_\Theta(x)}{\partial \theta_k} \\ \frac{\partial J(\Theta)}{\partial h_\Theta(x)} &= 2(h_\Theta(x) - y) \\ \frac{\partial h_\Theta(x)}{\partial \theta_k} &= 2x_k h_\Theta(x)(1 - h_\Theta(x)) \\ \text{thus } \frac{\partial J(\Theta)}{\partial \theta_k} &= 4x_k(h_\Theta(x) - y)h_\Theta(x)(1 - h_\Theta(x))\end{aligned}$$

$$\text{and } \theta_k \rightarrow \theta_k - \alpha 4x_k(h_\Theta(x) - y)h_\Theta(x)(1 - h_\Theta(x)), \forall k = \{5, 6\}$$

$$\forall k = \{0, \dots, 4\}$$

$$\begin{aligned}\frac{\partial J(\Theta)}{\partial \theta_k} &= \frac{\partial J(\Theta)}{\partial h_\Theta(x)} \cdot \frac{\partial h_\Theta(x)}{\partial g1} \cdot \frac{\partial g1}{\partial \theta_k} \\ \frac{\partial J(\Theta)}{\partial h_\Theta(x)} &= 2(h_\Theta(x) - y) \\ \frac{\partial h_\Theta(x)}{\partial g1} &= 2\theta_6 h_\Theta(x)(1 - h_\Theta(x)) \\ \frac{\partial g1}{\partial \theta_k} &= x_k g1(1 - g1) \\ \text{thus } \frac{\partial J(\Theta)}{\partial \theta_k} &= 4x_k \theta_6 (h_\Theta(x) - y)h_\Theta(x)(1 - h_\Theta(x))g1(1 - g1)\end{aligned}$$

$$\text{and } \theta_k \rightarrow \theta_k - \alpha 4x_k \theta_6 (h_\Theta(x) - y)h_\Theta(x)(1 - h_\Theta(x))g1(1 - g1), \forall k = \{0, \dots, 4\}$$