




ALMA-MATER OF QUAID-E-AZAM MOHAMMAD ALI JINNAH
SINDH MADRESSATUL ISLAM UNIVERSITY

CALCULUS

Dr. Muhammad Ali



In this lecture we will discuss about Roll's Theorem,
Cauchy mean value theorem, Lagrange's Mean Value
Theorem and power series

Roll's Theorem

Suppose that $y = f(x)$ is continuous at every point of the closed interval $[a, b]$ and differentiable at every point of its interior (a, b) . If

$$f(a) = f(b) = 0,$$

then there is at least one number c in (a, b) at which

$$f'(c) = 0.$$

Proof:

There are two cases arise

Case: I

$$\text{If } f(x) = k$$

(1)

$$f(a) = k$$

$$f(b) = k$$

$$\Rightarrow f(a) = f(b)$$

From (1)

$$f'(x) = 0$$

at $x = c$

$$f'(c) = 0 \quad \text{Hence proved}$$

Case: II


Consider a function which is continuous on $[a, b]$ and differentiable (a, b) where

$$f(a) = f(b).$$

From Fig.

$$f(c - h) - f(c) \leq 0 \quad (2)$$

Dividing both sides by $-h$ and taking limit as $h \rightarrow 0$ we have



$$\lim_{h \rightarrow 0} \frac{f(c-h) - f(c)}{-h} \geq 0$$

$$\Rightarrow f'(c) \geq 0$$

$$\Rightarrow f'(c) > 0, f'(c) = 0 \quad (A)$$

Again from Fig.

$$f(c+h) - f(c) \leq 0 \quad (3)$$


Dividing both sides by h and taking limit as $h \rightarrow 0$ we have

$$\lim_{h \rightarrow 0} \frac{f(c+h) - f(c)}{h} \leq 0$$

$$\text{i.e. } f'(c) < 0, f'(c) = 0 \quad (B)$$

From (A) and (B) $f'(c) > 0$ and $f'(c) < 0$ at a time it is not possible i.e. only possible is

$$f'(c) = 0$$



Exercise

Discuss the validity of Roll's theorem. Find value of c (whenever possible).

1. $f(x) = x^2 - 3x + 2$ *on* $[1, 2]$

2. $f(x) = \sin^2 x$ *on* $[0, \pi]$

3. $f(x) = x^2 - 7x + 12$ *on* $[3, 4]$

4. $f(x) = x^2 + 2x - 1$ *on* $[0, 1]$

1. Solution:

$$f(x) = x^2 - 3x + 2 \quad (1)$$

Now, at $x = 1$

$$(1) \Rightarrow f(1) = 1^2 - 3(1) + 2 = 3 - 3 = 0$$

at $x = 2$

$$(1) \Rightarrow f(2) = 2^2 - 3(2) + 2 = 4 - 6 + 2 = 6 - 6 = 0$$

$$\Rightarrow f(1) = f(2)$$

Hence, Roll's theorem is valid. Therefore, by Roll's theorem

$$f'(c) = 0 \quad (2)$$

$$\text{From (1)} \quad f'(x) = 2x - 3$$

$$\text{at } x = c, \quad f'(c) = 2c - 3 \quad (3)$$

Comapring (2) and (3)

$$2c - 3 = 0$$

$$\Rightarrow 2c = 3 \Rightarrow c = \frac{3}{2} = 1.5$$

2. Solution:

$$f(x) = \sin^2 x \quad (1)$$

Now, at $x = 0$

$$(1) \Rightarrow f(0) = \sin^2(0) = 0$$

at $x = \pi$

$$(1) \Rightarrow f(\pi) = \sin^2(\pi) = 0$$

$$\Rightarrow f(0) = f(\pi)$$

Hence, Roll's theorem is valid. Therefore, by Roll's theorem

$$f'(c) = 0 \quad (2)$$

$$\text{From (1)} \quad f'(x) = 2\sin x \cos x = \sin(2x)$$

$$\text{at } x = c, \quad f'(c) = \sin(2c) \quad (3)$$

Comapring (2) and (3)

$$\sin(2c) = 0$$

$$\Rightarrow 2c = 0, \pi \Rightarrow c = \frac{\pi}{2}$$

3. Solution:

$$f(x) = x^2 - 7x + 12 \quad (1)$$

Now, at $x = 3$

$$(1) \Rightarrow f(3) = 3^2 - 7(3) + 12 = 9 - 21 + 12 = 0$$

at $x = 4$

$$(1) \Rightarrow f(4) = 4^2 - 7(4) + 12 = 16 - 28 + 12 = 0$$

$$\Rightarrow f(3) = f(4)$$

Hence, Roll's theorem is valid. Therefore, by Roll's theorem

$$f'(c) = 0 \quad (2)$$

$$\text{From (1)} \quad f'(x) = 2x - 7$$

$$\text{at } x = c, \quad f'(c) = 2c - 7 \quad (3)$$

Comapring (2) and (3)

$$2c - 7 = 0$$

$$\Rightarrow 2c = 7 \Rightarrow c = \frac{7}{2} = 3.5$$

4. Solution:

$$f(x) = x^2 + 2x - 1 \quad (1)$$

Now, at $x = 0$

$$(1) \Rightarrow f(0) = 0^2 + 2(0) - 1 = -1$$

at $x = 1$

$$(1) \Rightarrow f(1) = 1^2 + 2(1) - 1 = 3 - 1 = 2$$

$$\Rightarrow f(0) \neq f(1)$$

Hence, Roll's theorem is not valid.

Lagrange's Mean Value Theorem

Statement: Let $f(x)$ be a function such that

(i) $f(x)$ is continuous on $[a, b]$, $b > a$

(ii) $f(x)$ is differentiable on $]a, b[$

Such that there must be a number $c \in (a, b)$ such that

$$f(b) - f(a) = f'(c)(b - a)$$

Or

$$\frac{f(b) - f(a)}{b - a} = f'(c)$$

Proof:

Let $f(x)$ is a continuous function on $[a, b]$ and differentiable on (a, b) then

$\phi(x)$ be a new function which is also continuous on $[a, b]$ and

differentiable on (a, b) and this can be written as


$$\phi(x) = Mf(x) + Nx \quad (1)$$

Where M and N are constants

By Roll's theorem

$$\phi(a) = \phi(b)$$

$$\Rightarrow Mf(a) + Na = Mf(b) + Nb$$



$$\text{slope of } AB = \frac{f(b) - f(a)}{b - a} = f'(c) = \text{slope of tangent at } C$$

$$\begin{aligned}
&\Rightarrow Mf(a) - Mf(b) = Nb - Na \\
&\Rightarrow M[f(a) - f(b)] = (b - a)N \\
&\Rightarrow -M[f(b) - f(a)] = (b - a)N \\
&\Rightarrow \frac{f(b) - f(a)}{b - a} = -\frac{N}{M} \quad (2)
\end{aligned}$$

Again by Roll's theorem

$$\phi'(c) = 0$$

From (1)

$$\phi'(x) = Mf'(x) + N$$

put $x = c$

$$\Rightarrow \phi'(c) = Mf'(c) + N \text{ but } \phi'(c) = 0$$

$$\Rightarrow Mf'(c) + N = 0$$

$$\Rightarrow f'(c) = -\frac{N}{M} \quad (3) \text{ Comparing (2) and (3)}$$

$$\frac{f(b) - f(a)}{b - a} = f'(c)$$

Example: 1 If $f(x) = x^3 - 3x - 1$ on $\left[-\frac{11}{7}, \frac{13}{7}\right]$. Find the value of C by LMVT.

Solution:

$$f(x) = x^3 - 3x - 1 \quad (1)$$

Put $x = -\frac{11}{7}$ in (1)

$$(1) \Rightarrow f\left(-\frac{11}{7}\right) = \left(-\frac{11}{7}\right)^3 - 3\left(-\frac{11}{7}\right) - 1 = -0.17$$

Put $x = \frac{13}{7}$ in (1)

$$(1) \Rightarrow f\left(\frac{13}{7}\right) = \left(\frac{13}{7}\right)^3 - 3\left(\frac{13}{7}\right) - 1 = -0.17$$

Now, differentiate (1) w.r.t. x

$$f'(x) = 3x^2 - 3$$

$$\text{at } x = c, f'(c) = 3c^2 - 3,$$

By LMVT

$$f'(c) = \frac{f(b) - f(a)}{b - a} = \frac{(-0.17) - (-0.17)}{\frac{13}{7} - \left(-\frac{11}{7}\right)} = \frac{0}{\frac{13}{7} + \frac{11}{7}} = 0$$

$$\Rightarrow 3c^2 - 3 = 0,$$

$$\Rightarrow 3c^2 = 3,$$

$$\Rightarrow c^2 = 1,$$

$$\Rightarrow c = \pm 1$$

Practice Problem: *If $f(x) = x^3 - 5x^2 + 4x - 2$ on $[2, 4]$. Find the value of C by LMVT*

Practice Problem: If $f(x) = x^3 - 5x^2 + 4x - 2$ on $[2,4]$. Find the value of C by LMVT

Solution:

$$f(x) = x^3 - 5x^2 + 4x - 2 \quad (1)$$

Put $x = 2$ in (1)

$$(1) \Rightarrow f(2) = (2)^3 - 5(2)^2 + 4(2) - 2 = 8 - 20 + 8 - 2 = -6$$

Put $x = 4$ in (1)

$$(1) \Rightarrow f(4) = (4)^3 - 5(4)^2 + 4(4) - 2 = 64 - 80 + 16 - 2 = -2$$

Now, differentiate (1) w.r.t. x

$$f'(x) = 3x^2 - 10x + 4$$

$$\text{at } x = c, f'(c) = 3c^2 - 10c + 4,$$

By LMVT

$$f'(c) = \frac{f(b) - f(a)}{b - a} = \frac{(-2) - (-6)}{4 - 2} = \frac{4}{2} = 2$$

$$\Rightarrow 3c^2 - 10c + 4 = 2,$$

$$\Rightarrow 3c^2 - 10c + 2 = 0,$$

$$c = \frac{-b \pm \sqrt{b^2 - 4ac}}{2a} = \frac{-4 \pm \sqrt{(4)^2 - 4(3)(2)}}{2(3)}$$

$$c = \frac{-4 \pm \sqrt{16 - 24}}{6} = \frac{-4 \pm \sqrt{-8}}{6} = \frac{-4 \pm 2\sqrt{2}i}{6} = \frac{-2 \pm \sqrt{2}i}{3}$$

Practice Problem: If $f(x) = x^2 - 7x^2 + 12$ on $[3,4]$. Find the value of C .

Cauchy Mean Value Theorem

Statement:

Let $f(x)$ and $g(x)$ be two functions that satisfying following conditions.

- (i) $f(x)$ and $g(x)$ both are continuous on $[a, b]$,*
- (ii) $f(x)$ and $g(x)$ both are differentiable on $]a, b[$*
- (iii) $g'(x) \neq 0, \quad \forall x$ then*

$$\frac{f'(c)}{g'(c)} = \frac{f(b) - f(a)}{g(b) - g(a)}$$

Example:1 If $f(x) = e^x$ and $g(x) = e^{2x}$ on $[0, \ln 2]$.

Find value of C by Cauchy mean value theorem.

Solution:

$$f(x) = e^x \quad (1)$$

$$g(x) = e^{2x} \quad (2)$$

Since, by Cauchy mean value theorem

$$\frac{f'(c)}{g'(c)} = \frac{f(b) - f(a)}{g(b) - g(a)} \quad (3)$$

at $x = a = 0$,

$$(1) \Rightarrow f(0) = e^0 = 1,$$

$$(2) \Rightarrow g(0) = e^0 = 1$$

at $x = b = \ln 2$,

$$(1) \Rightarrow f(\ln 2) = e^{\ln 2} = 2,$$

$$(2) \Rightarrow g(\ln 2) = e^{2\ln 2} = e^{\ln 4} = 4$$

Again from equation (1) and (2)

$$(1) \Rightarrow f'(x) = e^x,$$

$$(2) \Rightarrow g'(x) = 2e^{2x}$$

At $x = c$,

$$f'(c) = e^c,$$

$$g'(x) = 2e^{2c}$$

Putting all above values in (3)

$$(3) \Rightarrow \frac{e^c}{2e^{2c}} = \frac{2 - 1}{4 - 1} = \frac{1}{3} \Rightarrow \frac{1}{2e^c} = \frac{1}{3} \Rightarrow 2e^c = 3 \Rightarrow c = \ln\left(\frac{3}{2}\right)$$

Example:2 If $f(x) = x^2$ and $g(x) = x^3$.

Verify, Cauchy mean value theorem on $[1, 2]$.

Also find value of C .

Solution:

$$f(x) = x^2 \quad (1)$$

$$g(x) = x^3 \quad (2)$$

Since, by Cauchy mean value theorem

$$\frac{f'(c)}{g'(c)} = \frac{f(b) - f(a)}{g(b) - g(a)} \quad (3)$$

at $x = a = 1$,

$$(1) \Rightarrow f(1) = (1)^2 = 1,$$

$$(2) \Rightarrow g(1) = (1)^3 = 1$$

at $x = b = 2$,

$$(1) \Rightarrow f(2) = (2)^2 = 4,$$

$$(2) \Rightarrow g(2) = (2)^3 = 8$$

Again from equation (1) and (2)

$$(1) \Rightarrow f'(x) = 2x,$$

$$(2) \Rightarrow g'(x) = 3x^2$$

At $x = c$,

$$f'(c) = 2c,$$

$$g'(c) = 3c^2$$

Putting all above values in (3)

$$(3) \Rightarrow \frac{2c}{3c^3} = \frac{4 - 1}{8 - 1} = \frac{3}{7} \Rightarrow \frac{1}{c^2} = \frac{9}{14} \Rightarrow c^2 = \frac{14}{9} \Rightarrow c = \sqrt{\left(\frac{14}{9}\right)}$$

Power Series :-

A series $\sum_{n=0}^{\infty} C_n (x-\alpha)^n = C_0 + C_1(x-\alpha)$

$+ C_2(x-\alpha)^2 + C_3(x-\alpha)^3 + \dots$ is a power series

around α and C_n 's are called coefficients.

The series $\sum_{n=0}^{\infty} C_n x^n$ is a power series around '0'.

e.g. $\sum_{n=0}^{\infty} n^2 x^n$, $\sum_{n=0}^{\infty} \frac{x^n}{n!}$, $\sum_{n=0}^{\infty} n^3 (x-1)^n$ etc.

Power Series

The equation of the form

$$\sum_{n=0}^n a_n(x-a)^n = a_0 + a_1(x-a) + a_2(x-a)^2 + \dots,$$

is called a power series.

Consider,

$$f(x) = a_0 + a_1(x-a) + a_2(x-a)^2 + \dots \quad (1)$$

then

$$f'(x) = a_1 + 2a_2(x-a) + 3a_3(x-a)^2 + \dots.$$

$$f''(x) = 2 \cdot 1a_2 + 3 \cdot 2a_3(x-a) + 4 \cdot 3a_4(x-a)^2 + \dots.$$

$$f'''(x) = 3! a_3 + 4! a_4(x-a) + \dots.$$

\vdots

Put $x = a$,

$$f(a) = a_0$$

$$f'(a) = a_1$$

$$f''(a) = 2! a_2 \text{ or } a_2 = \frac{f''(a)}{2!}$$

$$f'''(a) = 3! a_3 \text{ or } a_3 = \frac{f'''(a)}{3!}$$

\vdots

Putting these values of a_0, a_1, a_2, \dots in eq. (1)

$$(1) \Rightarrow f(x) = f(a) + f'(a)(x - a) + \frac{f''(a)}{2!}(x - a)^2 + \frac{f'''(a)}{3!}(x - a)^3 + \dots \quad (2)$$

This equation is called Taylor series for $f(x)$ at $x = a$.

Put $a = 0$ in equation (2) then

$$f(x) = f(0) + f'(0)x + \frac{f''(0)}{2!}x^2 + \frac{f'''(0)}{3!}x^3 + \dots \quad (3)$$

The equation (3) is called **Machaurin series** for $f(x)$.

Example: 3 Find Machaurian's series for $f(x) = e^x$.

Solution:

The Machaurin's series $f(x)$ is

$$f(x) = f(0) + f'(0)x + \frac{f''(0)}{2!}x^2 + \frac{f'''(0)}{3!}x^3 + \dots \quad (1)$$

Here, $f(x) = e^x$

Now,

$$f(x) = e^x \quad \Rightarrow f(0) = 1$$

$$f'(x) = e^x \quad \Rightarrow f'(0) = e^0 = 1$$

$$f''(x) = e^x \quad \Rightarrow f''(0) = e^0 = 1$$

\vdots

$$(1) \Rightarrow e^x = 1 + x + \frac{1}{2!}x^2 + \frac{1}{3!}x^3 + \dots$$

$$e^x = \sum_{n=0}^n \frac{x^n}{n!} = 1 + x + \frac{1}{2!}x^2 + \frac{1}{3!}x^3 + \dots$$

Practice Problems: Find Manchurian series for

(i) $\cos x$

(ii) $\frac{1}{1-x}$

Q1 Apply Maclaurin's series to prove That

$$i) \ln(1+x) = x - \frac{x^2}{2} + \frac{x^3}{3} - \frac{x^4}{4} + \dots$$

Sol $f(x) = \ln(1+x)$
Diff w.r.t x

$$f(0) = \ln(1+0) = \ln 1 = 0$$

$$\boxed{f(0) = 0}$$

$$f'(x) = \frac{1}{1+x} \frac{d}{dx}(1+x)$$

$$f'(x) = \frac{1}{1+x}$$

$$\boxed{f'(0) = 1}$$

$$f'(x) = (1+x)^{-1} \text{ Diff w.r.t. } x$$

$$f''(x) = (-1)(1+x)^{-2} (0+1)$$

$$f''(x) = (-1)(1+x)^{-2}$$

Diff again w.r.t. x

$$f'''(x) = (-1)(-2)(1+x)^{-3} (0+1)$$

$$= 2(1+x)^{-3}$$

Diff aga w.r.t. x

$$f'''(0) = 2(1+0)^{-3}$$

$$\boxed{f'''(0) = 2}$$

$$f^{(4)}(x) = 2(-3)(1+x)^{-4}$$

$$= -6(1+x)^{-4}$$

$$f^{(4)}(0) = -6(1+0)^{-4}$$

$$\boxed{f^{(4)}(0) = -6}$$

As we know By Maclaurin's Series

$$f(x) = f(0) + x f'(0) + \frac{x^2}{2!} f''(0) + \frac{x^3}{3!} f'''(0) + \frac{x^4}{4!} f^{(4)}(0) + \dots$$

But $f(x) = \ln(1+x)$

$$\ln(1+x) = 0 + x(1) + \frac{x^2}{2!}(-1) + \frac{x^3}{3!}(2) + \frac{x^4}{4!}(-6) + \dots$$

$$\ln(1+x) = x - \frac{x^2}{2} + \frac{x^3}{3} - \frac{x^4}{4} + \dots$$

$$\frac{2}{3!} = \frac{2}{3 \cdot 2 \cdot 1}$$

$$\frac{6}{4!} = \frac{6}{4 \cdot 3 \cdot 2 \cdot 1}$$

Apply Maclaurin's series to prove that
 vi) $\ln(1-x) = -x - \frac{x^2}{2} - \frac{x^3}{3} - \frac{x^4}{4} - \dots$

Sol $f(x) = \ln(1-x)$
 Diff w.r.t x
 $f(0) = \ln(1-0) = \ln 1 = 0$
 $\boxed{f(0) = 0}$

$$f'(x) = \frac{1}{1-x} \frac{d}{dx}(1-x)$$

$$f'(x) = \frac{-1}{1-x}$$

$$\boxed{f'(0) = -1}$$

$$f'(x) = -(1-x)^{-1} \text{ Diff w.r.t. } x$$

$$f''(x) = (+1)(1-x)^{-2}(0-1)$$

$$f''(x) = (-1)(1-x)^{-2}$$

$$f''(0) = -1(1-0)^{-2} = -1$$

$$\boxed{f''(0) = -1}$$

Diff again w.r.t. x

$$f'''(x) = (-1)(-2)(1-x)^{-3}(0-1)$$

$$= -2(1-x)^{-3}$$

Diff again w.r.t. x

$$f'''(0) = -2(1-0)^{-3}$$

$$\boxed{f'''(0) = -2}$$

$$f^{(4)}(x) = -2(-3)(1-x)^{-4}(0-1)$$

$$= -6(1-x)^{-4}, f(0) = -6(1-0)^{-4}$$

$$\boxed{f^{(4)}(0) = -6}$$

As we know By Maclaurin's Series

$$f(x) = f(0) + x f'(0) + \frac{x^2}{2!} f''(0) + \frac{x^3}{3!} f'''(0) + \frac{x^4}{4!} f^{(4)}(0) + \dots$$

But $f(x) = \ln(1-x)$

$$\ln(1-x) = 0 + x(-1) + \frac{x^2}{2!}(-1) + \frac{x^3}{3!}(-2) + \frac{x^4}{4!}(-6) + \dots$$

$$= -x - \frac{x^2}{2} - \frac{x^3}{3} - \frac{x^4}{4} - \dots$$

$$e^x = 1 + x + \frac{x^2}{2!} + \frac{x^3}{3!} + \dots$$

$$\sin x = \dots$$

$$\cos x = \dots$$

$$\ln(1+x) = \dots$$

$$\ln(1-x) = \dots$$

Example: 1 Find Taylor's series for $\sin x$ at $x = \frac{\pi}{2}$.

Solution:

The Taylor's series at $x = a$ is

$$f(x) = f(a) + f'(a)(x - a) + \frac{f''(a)}{2!}(x - a)^2 + \frac{f'''(a)}{3!}(x - a)^3 + \dots \quad (1)$$

Here, $f(x) = \sin x$ and $a = \frac{\pi}{2}$

Now,

$$f(x) = \sin x \quad \Rightarrow f\left(\frac{\pi}{2}\right) = 1$$

$$f'(x) = \cos x \quad \Rightarrow f'\left(\frac{\pi}{2}\right) = 0$$

$$f''(x) = -\sin x \quad \Rightarrow f''\left(\frac{\pi}{2}\right) = -1$$

$$f'''(x) = -\cos x \quad \Rightarrow f'''\left(\frac{\pi}{2}\right) = 0$$

$$(1) \Rightarrow \sin x = 1 - \frac{1}{2}\left(x - \frac{\pi}{2}\right)^2 + \frac{1}{4!}\left(x - \frac{\pi}{2}\right)^4 - \frac{1}{6!}\left(x - \frac{\pi}{2}\right)^6 + \dots$$

Example: 2 Find Taylor's series for $f(x) = \frac{1}{\sqrt{x}}$ at $x = 9$.

Solution:

The Taylor's series at $x = a$ is

$$f(x) = f(a) + f'(a)(x - a) + \frac{f''(a)}{2!}(x - a)^2 + \frac{f'''(a)}{3!}(x - a)^3 + \dots \quad (1)$$

Here, $f(x) = \frac{1}{\sqrt{x}}$ and $a = 9$

Now,

$$f(x) = \frac{1}{\sqrt{x}} \quad \Rightarrow f(9) = \frac{1}{3}$$

$$f'(x) = -\frac{1}{2}x^{-\frac{3}{2}} \quad \Rightarrow f'(9) = -\frac{1}{2}(9)^{-\frac{3}{2}} = \frac{1}{2 \cdot 3^3}$$

$$f''(x) = \frac{3}{4}x^{-\frac{5}{2}} \quad \Rightarrow f''(9) = \frac{3}{4}(9)^{-\frac{5}{2}} = \frac{3}{2^2 \cdot 3^5}$$

$$f'''(x) = -\frac{15}{8}x^{-\frac{7}{2}} \quad \Rightarrow f'''(9) = -\frac{15}{8}(9)^{-\frac{7}{2}} = -\frac{15}{2^3 \cdot 3^7}$$

$$(1) \Rightarrow \frac{1}{\sqrt{x}} = \frac{1}{3} - \frac{1}{2 \cdot 3^3}(x - 9) + \frac{3}{2^2 \cdot 2! \cdot 3^5}(x - 9)^2 - \frac{15}{2^3 \cdot 3! \cdot 3^7}(x - 9)^3 + \dots$$

TAILOR THEOREM

$$f(x+h) = f(x) + h f'(x) + \frac{h^2}{2!} f''(x) + \frac{h^3}{3!} f'''(x) + \dots + \frac{h^n}{n!} f^{(n)}(x) + \dots$$

Q2 Show That $\cos(x+h) = \cos x - h \sin x - \frac{h^2}{2!} \cos x + \frac{h^3}{3!} \sin x + \dots$
and Evaluate $\cos 61^\circ$

Sol-

$$f(x) = \cos x$$

Diff w.r.t. x

$$f'(x) = -\sin x$$

Diff w.r.t. x

$$f''(x) = -\cos x$$

Diff again w.r.t. x

$$f'''(x) = -(-\sin x) = \sin x$$

\vdots

$$f(x+h) = \cos(x+h)$$

As we know $f(x+h) = f(x) + h f'(x) + \frac{h^2}{2!} f''(x) + \frac{h^3}{3!} f'''(x) + \dots$

$$\cos(x+h) = \cos x + h(-\sin x) + \frac{h^2}{2!}(-\cos x) + \frac{h^3}{3!}(\sin x) + \dots$$

$$\cos(x+h) = \cos x - h \sin x - \frac{h^2}{2!} \cos x + \frac{h^3}{3!} \sin x + \dots$$

Let $x = 60^\circ$, $h = 1^\circ = 0.01745$

$$\cos(60^\circ + 1^\circ) = \cos 60^\circ - (0.01745) \sin 60^\circ - \frac{(0.01745)^2}{2} \cos 60^\circ + \dots$$

$$\cos 61^\circ \approx .4848$$

$\cos 59^\circ$
 $\cos 31^\circ$
 $\cos 29^\circ$
 $\cos 46^\circ$
 $\cos 44^\circ$

EXERCISE No 2.8

Q3 Show that: $2^{x+h} = 2^x \left[1 + h(\ln 2) + \frac{h^2}{2!} (\ln 2)^2 + \frac{h^3}{3!} (\ln 2)^3 + \dots \right]$

Sol $f(x) = 2^x$, $f(x+h) = 2^{x+h}$

Diff w.r.t. x

$$f'(x) = \frac{d}{dx} (2^x) \quad \text{using } \frac{d}{dx} (a^x) = a^x \cdot \ln a$$

$$f'(x) = 2^x \cdot \ln 2$$

Diff w.r.t. x

$$f''(x) = \ln 2 \cdot \frac{d}{dx} 2^x \\ = \ln 2 \cdot (2^x) \cdot \ln 2$$

$$f''(x) = (\ln 2)^2 \cdot 2^x$$

Diff w.r.t. x

$$f'''(x) = (\ln 2)^2 \cdot 2^x \cdot \ln 2 \\ = (\ln 2)^3 \cdot 2^x$$

\vdots

By Taylor Theorem

$$f(x+h) = f(x) + hf'(x) + \frac{h^2}{2!} f''(x) + \frac{h^3}{3!} f'''(x) + \dots \\ 2^{x+h} = 2^x + h(2^x \ln 2) + \frac{h^2}{2!} (\ln 2)^2 2^x + \frac{h^3}{3!} (\ln 2)^3 2^x + \dots \\ = 2^x \left[1 + h(\ln 2) + \frac{h^2}{2!} (\ln 2)^2 + \frac{h^3}{3!} (\ln 2)^3 + \dots \right]$$