

Tropical representation of Weyl groups associated with certain rational varieties

Teruhisa Tsuda and Tomoyuki Takenawa

Dedicated to Professor Kazuo Okamoto on his sixtieth birthday

Abstract

Starting from certain rational varieties blown-up from $(\mathbb{P}^1)^N$, we construct a tropical, i.e., subtraction-free birational, representation of Weyl groups as a group of pseudo isomorphisms of the varieties. We develop an algebro-geometric framework of τ -functions as defining functions of exceptional divisors on the varieties. In the case where the corresponding root system is of affine type, our construction yields a class of (higher order) q -difference Painlevé equations and its algebraic degree grows quadratically.

1 Introduction

The aim of the present work is to develop the theory of birational representation of Weyl groups associated with algebraic varieties.

At the beginning of the twentieth century, it was discovered by Coble and Kantor, and later by Du Val, that certain types of Cremona transformations act on the configuration space of point sets [Cob29]. Let $X_{m,n}$ be the configuration space of n points in general position in the projective space \mathbb{P}^{m-1} . Then, the Weyl group $W(T_{2,m,n-m})$ corresponding to the Dynkin diagram $T_{2,m,n-m}$ (see Figure 1) acts birationally on $X_{m,n}$ and is generated by permutations of n points and the standard Cremona transformation with respect to each m points. An algebro-geometric and modern interpretation of this theory is due to Dolgachev and Ortland [DO88]; they showed that the Cremona action of $W(T_{2,m,n-m})$ induces a *pseudo isomorphism*, i.e., an isomorphism except for subvarieties of codimension two or higher, between certain rational varieties blown-up from \mathbb{P}^{m-1} at generic n points, which they call *generalized Del Pezzo varieties*. It is worth mentioning that if $(m, n) = (3, 9)$, the affine Weyl group of type $E_8^{(1)}$ appears and its lattice part gives rise to an important discrete dynamical system, i.e., the elliptic-difference Painlevé equation [Sak01]; see also [ORG01, KMNOY03, KMNOY06].

In this paper, starting from a certain rational variety blown-up from $(\mathbb{P}^1)^N$ along appropriate subvarieties that are not only point sets, we construct a birational representation of Weyl groups corresponding to the Dynkin diagram T_ℓ^k . Here T_ℓ^k refers to the tree given in Figure 1, specified by a

2000 *Mathematics Subject Classification* 14E07, 14L30, 20F55, 34M55, 37K10, 39A13.

Keywords: Cremona transformation, τ -function, Painlevé equation, rational variety, Schur function, tropical representation, universal character, Weyl group.




Figure 1: Dynkin diagrams $T_{2,m,n-m}$ and T_ℓ^k

pair of sequences $\mathbf{k} = (k_1, \dots, k_N), \ell = (\ell_1, \dots, \ell_N) \in (\mathbb{Z}_{>0})^N$. It is remarkable that T_ℓ^k includes all of the simply-laced affine cases $A_n^{(1)}, D_n^{(1)}$ and $E_n^{(1)}$, which are relevant to a class of higher-dimensional discrete dynamical systems of Painlevé type. For example if $N = 3$ and $\mathbf{k} = \ell = (1, 2, 1)$, we have $T_{(1,2,1)}^{(1,2,1)} = D_4^{(1)}$. This representation of Weyl groups is *tropical*, i.e., given in terms of subtraction-free birational mappings [Kir01] and, interestingly enough, possesses a geometric framework of τ -functions.

In the next section, we begin with blowing-up $(\mathbb{P}^1)^N$ along certain subvarieties of codimension three. Let X be the rational variety thus obtained. These X 's constitute a family. Applying a cohomological technique (cf. [DO88]), we construct the root and coroot lattices of type T_ℓ^k included in the Néron-Severi bilattice $N(X) \simeq (H^2(X, \mathbb{Z}), H_2(X, \mathbb{Z}))$. The associated Weyl group $W = W(T_\ell^k)$ naturally acts on $N(X)$ as isometries (Lemma 2.1). We see that this linear action of W on $N(X)$ leads to a birational representation of W on the family of varieties itself as a group of pseudo isomorphisms (Theorem 2.2). An element of W naturally induces an appropriate permutation among the set of exceptional divisors on X , as similar to the classical topic: 27 lines (or exceptional curves) on a cubic surface and a Weyl group of type E_6 . In § 3, for the purpose of describing the action of W at the level of defining polynomials of exceptional divisors, we introduce a geometric framework of τ -functions (Definition 3.2 and Theorems 3.1 and 3.3). One important advantage of our τ -functions is that we can trace the resulting value for any $w \in W$ by using only the defining polynomials of suitable divisors, although it is generally difficult to compute iterations of rational mappings. In particular, our representation in an affine case provides a discrete dynamical system arising from the lattice part of the affine Weyl group. Such a discrete dynamical system is equipped with a set of commuting discrete time evolutions and its algebraic degree grows in the quadratic order. And it is regarded as a (higher order) q -difference Painlevé equation; in § 4, the $A_n^{(1)}$ and $D_n^{(1)}$ cases are demonstrated as typical examples. In § 5, we briefly indicate (from a soliton-theoretic point of view) some interesting relationships between τ -functions and the *character polynomials* appearing in representation theory of classical groups, i.e., the Schur functions or the universal characters.

2 Tropical Weyl group actions on rational varieties

In this section, we consider a certain rational variety X blown-up from $(\mathbb{P}^1)^N$ and construct appropriate root and coroot lattices included in the Néron-Severi bilattice $N(X)$. Moreover, the action of the corresponding Weyl group is realized as pseudo isomorphisms of X and is shown to possess a tropical (or subtraction-free birational) representation.

2.1 Rational variety and root system

Denote by $f = (f_1, f_2, \dots, f_N) \in (\mathbb{P}^1)^N$ the inhomogeneous coordinate and consider the following subvarieties:

$$\begin{aligned} C_n^i &= \{f_{n-1} = 0, f_n = -u_n^i, f_{n+1} = \infty\}, \quad i = 1, \dots, k_n, \\ C_n^{-j} &= \{f_{n-1} = \infty, f_n = -1/v_n^{-j}, f_{n+1} = 0\}, \quad j = 1, \dots, \ell_n, \end{aligned}$$

for $n = 1, 2, \dots, N$, where u_n^i and v_n^{-j} are nonzero parameters. Let $\epsilon : X \rightarrow (\mathbb{P}^1)^N$ be the blowing-up along $\{C_n^i, C_n^{-j}\}$. Since X is a rational variety, we have

$$H^2(X, \mathbb{Z}) \simeq NS(X) = \bigoplus_{n=1}^N \left(\mathbb{Z}H_n \oplus \bigoplus_{i=1}^{k_n} \mathbb{Z}E_n^i \oplus \bigoplus_{j=1}^{\ell_n} \mathbb{Z}E_n^{-j} \right),$$

where $NS(X)$ is the Néron-Severi group of X ; we denote by H_n the divisor class of hyperplanes $\{f_n = \text{const.}\}$ and by E_n^i the class of exceptional divisors $\epsilon^{-1}(C_n^i)$. The Poincaré duality guarantees $H_2(X, \mathbb{Z}) \simeq (H_2(X, \mathbb{Z}))^*$. We can choose a basis $\{h_n, e_n^i, e_n^{-j}\}$ of $H_2(X, \mathbb{Z})$, where h_n corresponds to a line of degree $(0, \dots, 0, 1, 0, \dots, 0)$ and e_n^i to a line restricted in a fibre ($\simeq \mathbb{P}^2$) of the exceptional divisor $\epsilon^{-1}(C_n^i)$. Thus the intersection pairing $\langle \cdot, \cdot \rangle : H^2(X, \mathbb{Z}) \times H_2(X, \mathbb{Z}) \rightarrow \mathbb{Z}$ is defined by $\langle H_m, h_n \rangle = \delta_{m,n}$, $\langle E_m^i, e_n^j \rangle = -\delta_{m,n} \delta_{i,j}$ and $\langle \text{otherwise} \rangle = 0$.


Introduce the root lattice Q and coroot lattice \check{Q} as follows:

$$Q = \bigoplus_{n=1}^N \bigoplus_{-\ell_n+1 \leq i \leq k_n-1} \mathbb{Z}\alpha_n^i \subset H^2(X, \mathbb{Z}) \quad \text{and} \quad \check{Q} = \bigoplus_{n=1}^N \bigoplus_{-\ell_n+1 \leq i \leq k_n-1} \mathbb{Z}\check{\alpha}_n^i \subset H_2(X, \mathbb{Z}),$$

where

$$\begin{aligned} \alpha_n^0 &= H_n - E_n^1 - E_n^{-1}, & \check{\alpha}_n^0 &= h_{n-1} + h_{n+1} - e_n^1 - e_n^{-1}, \\ \alpha_n^i &= E_n^i - E_n^{i+1}, & \check{\alpha}_n^i &= e_n^i - e_n^{i+1} \quad (i = 1, \dots, k_n - 1), \\ \alpha_n^{-j} &= E_n^{-j} - E_n^{-j-1}, & \check{\alpha}_n^{-j} &= e_n^{-j} - e_n^{-j-1} \quad (j = 1, \dots, \ell_n - 1). \end{aligned}$$

The Dynkin diagram of the canonical root basis forms the tree T_ℓ^k :



The simple reflection s_n^i associated with a root α_n^i naturally acts on the Néron-Severi bilattice $N(X) \simeq (H^2(X, \mathbb{Z}), H_2(X, \mathbb{Z}))$ as

$$s_n^i(\Lambda) = \Lambda + \langle \Lambda, \check{\alpha}_n^i \rangle \alpha_n^i, \quad \Lambda \in H^2(X, \mathbb{Z}), \quad (2.1a)$$

$$s_n^i(\lambda) = \lambda + \langle \alpha_n^i, \lambda \rangle \check{\alpha}_n^i, \quad \lambda \in H_2(X, \mathbb{Z}). \quad (2.1b)$$

One can easily check that these reflections indeed satisfy the fundamental relations (see [Kac90]) of the Weyl group $W = W(T_\ell^k) = \langle s_n^i \rangle$.

The half of the anti-canonical class $-\frac{1}{2}K_X = \sum_{n=1}^N (H_n - \sum_{i=1}^{k_n} E_n^i - \sum_{j=1}^{\ell_n} E_n^{-j}) \in H^2(X, \mathbb{Z})$ can be decomposed in two ways

$$-\frac{1}{2}K_X = \sum_{n=1}^N D_n^0 = \sum_{n=1}^N D_n^\infty,$$

where $D_n^0 = H_n - \sum_{i=1}^{k_{n+1}} E_{n+1}^i - \sum_{j=1}^{\ell_{n-1}} E_{n-1}^{-j}$ and $D_n^\infty = H_n - \sum_{i=1}^{k_{n-1}} E_{n-1}^i - \sum_{j=1}^{\ell_{n+1}} E_{n+1}^{-j}$. Note that each D_n^0 and D_n^∞ corresponds to the strict transform of hyperplanes $\{f_n = 0\}$ and $\{f_n = \infty\}$, respectively. In parallel, we shall *formally* define an element $-\frac{1}{2}k_X \in H_2(X, \mathbb{Z})$ by

$$-\frac{1}{2}k_X = \sum_{n=1}^N \left(2h_n - \sum_{i=1}^{k_n} e_n^i - \sum_{j=1}^{\ell_n} e_n^{-j} \right) = \sum_{n=1}^N d_n^0 = \sum_{n=1}^N d_n^\infty,$$

where $d_n^0 = h_{n-1} + h_{n+1} - \sum_{i=1}^{k_{n+1}} e_{n+1}^i - \sum_{j=1}^{\ell_{n-1}} e_{n-1}^{-j}$ and $d_n^\infty = h_{n-1} + h_{n+1} - \sum_{i=1}^{k_{n-1}} e_{n-1}^i - \sum_{j=1}^{\ell_{n+1}} e_{n+1}^{-j}$. We see that $Q \subset \{d_n^0, d_n^\infty\}^\perp$ and $\check{Q} \subset \{D_n^0, D_n^\infty\}^\perp$. Hence we have the

Lemma 2.1. *All elements $w \in W \subset \text{Aut}(N(X))$ leave the intersection pairing $\langle \cdot, \cdot \rangle$, $\frac{1}{2}K_X$ and $\frac{1}{2}k_X$ invariant.*

2.2 Birational representation of Weyl groups

We summarize below the linear action of generators s_n^i on the basis of $H^2(X, \mathbb{Z})$:

$$\begin{aligned} s_n^0(H_{n\pm 1}) &= H_{n\pm 1} + H_n - E_n^1 - E_n^{-1}, & s_n^0(E_n^{\pm 1}) &= H_n - E_n^{\mp 1}, \\ s_n^i(E_n^{\{i, i+1\}}) &= E_n^{\{i+1, i\}} & \text{for } 1 \leq i \leq k_n - 1, \\ s_n^{-j}(E_n^{\{-j, -j-1\}}) &= E_n^{\{-j-1, -j\}} & \text{for } 1 \leq j \leq \ell_n - 1. \end{aligned}$$

Now, let us extend the above linear action of the Weyl group $W = \langle s_n^i \rangle$ to the level of birational transformations on the rational variety X . We introduce the *multiplicative root variables* $a_n^i \in \mathbb{C}^\times$ attached to the canonical roots α_n^i and fix the action of s_n^i on them as

$$s_n^i(a_n^i) = \frac{1}{a_n^i}, \quad s_n^i(a_n^{i\pm 1}) = a_n^i a_n^{i\pm 1}, \quad s_n^0(a_{n\pm 1}^0) = a_n^0 a_{n\pm 1}^0. \quad (2.2)$$

Using the root variables, we fix the parameterization of subvarieties C_n^i as follows:

$$u_n^1 = u_n = \frac{(a_n^0 \prod_{j=1}^{\ell_n-1} (a_n^{-j})^{1-j/\ell_n})^{\ell_n/(k_n+\ell_n)}}{(\prod_{i=1}^{k_n-1} (a_n^i)^{1-i/k_n})^{k_n/(k_n+\ell_n)}}, \quad v_n^1 = v_n = \frac{(a_n^0 \prod_{i=1}^{k_n-1} (a_n^i)^{1-i/k_n})^{k_n/(k_n+\ell_n)}}{(\prod_{j=1}^{\ell_n-1} (a_n^{-j})^{1-j/\ell_n})^{\ell_n/(k_n+\ell_n)}}, \quad (2.3)$$

and $u_n^i = a_n^{i-1} u_n^{i-1}$, $v_n^{-j} = a_n^{-j+1} v_n^{-j+1}$ for $i, j \geq 2$. Namely, a projective equivalence class of the arrangement of subvarieties $\{C_n^i\}$ in $(\mathbb{P}^1)^N$ can be identified with a point of the Cartan subalgebra associated with the Dynkin diagram T_ℓ^k . In parallel, the rational varieties X 's under consideration constitute a family parameterised by the multiplicative root variables $a = (a_n^i)$; so we shall write clearly as $X = X_a$. Let $\mathbb{K}(f)$ be the field of rational functions in $f = (f_1, f_2, \dots, f_N)$, where the coefficient field $\mathbb{K} = \mathbb{C}(a^{1/p})$ is generated by a suitable fractional power $(a_n^i)^{1/p}$ of a_n^i . We then have the following birational representation of W over $\mathbb{K}(f)$, acting on the family of varieties $\mathcal{X} = \cup_a X_a$.

Theorem 2.2. *The birational transformations s_n^i defined by*

$$s_n^0(f_{n-1}) = (a_n^0)^{\ell_{n-1}/(k_{n-1}+\ell_{n-1})} f_{n-1} \frac{f_n + v_n^{-1}}{f_n + u_n}, \quad (2.4a)$$

$$s_n^0(f_{n+1}) = (a_n^0)^{-k_{n+1}/(k_{n+1}+\ell_{n+1})} f_{n+1} \frac{f_n + u_n}{f_n + v_n^{-1}}, \quad (2.4b)$$

and $s_n^i(f_m) = f_m$ (otherwise), together with (2.2), realize the Weyl group $W = W(T_\ell^k)$ over the field $\mathbb{K}(f)$. Moreover, each s_n^i maps X_a to $X_{s_n^i(a)}$ as a pseudo isomorphism of rational varieties and actually induces the linear action (2.1) on $N(X)$.

Proof. First we notice that $s_n^0 : u_n \leftrightarrow 1/v_n$, $s_n^i : u_n^i \leftrightarrow u_n^{i+1}$ ($1 \leq i \leq k_n - 1$) and $s_n^{-j} : v_n^{-j} \leftrightarrow v_n^{-j-1}$ ($1 \leq j \leq \ell_n - 1$). It is easy to verify by direct computation that $\langle s_n^i \rangle$ satisfies the fundamental relations of $W(T_\ell^k)$.

For $i \neq 0$, s_n^i is trivial and acts as an isomorphism $X_a \rightarrow X_{s_n^i(a)}$, so we have only to check that s_n^0 actually induces a pseudo isomorphism. The points of indeterminacy of the rational map $s_n^0 : (\mathbb{P}^1)^N \rightarrow (\mathbb{P}^1)^N$ are given by the following subvarieties:

$$\begin{aligned} I_1(a) &= \{f_n = -u_n, f_{n-1} = 0\}, & I_2(a) &= \{f_n = -u_n, f_{n+1} = \infty\}, \\ I_3(a) &= \{f_n = -v_n^{-1}, f_{n+1} = 0\}, & I_4(a) &= \{f_n = -v_n^{-1}, f_{n-1} = \infty\}, \end{aligned}$$

of codimension two. Since the same argument is also valid for the other I_n ($n = 2, 3, 4$), in what follows we shall only treat I_1 . We see that s_n^0 maps I_1 to I_3 generically, except for $C_n^1 = I_1 \cap I_2$; furthermore, C_n^1 equals the inverse image $(s_n^0)^{-1}(\{f_n = -1/v_n\})$. By the blowing-up $\epsilon : X_a \rightarrow (\mathbb{P}^1)^N$, the indeterminacy at C_n^1 is resolved and s_n^0 is extended holomorphically to $\epsilon^{-1}(C_n^1)$. Thus we have proved that s_n^0 acts on X_a as a pseudo isomorphism, i.e., an isomorphism except for subvarieties of codimension two or higher. \square

Remark 2.3. Our representation given in Theorem 2.2 above is obviously tropical and hence admits a combinatorial counterpart via the ultra-discretization [TTMS96]:

$$a \times b \rightarrow a + b, \quad a/b \rightarrow a - b, \quad a + b \rightarrow \min(a, b).$$

3 τ -Functions

An element of the Weyl group $W = W(T_\ell^k)$ naturally induces a permutation among the exceptional divisors on the rational variety X . In this section, we introduce a geometric framework of τ -functions, which describes the above permutations and therefore governs our tropical representation of W .

3.1 Representation over the field of τ -variables

We set

$$\theta_n^0 = k_{n+1} + \ell_{n-1} \quad \text{and} \quad \theta_n^\infty = k_{n-1} + \ell_{n+1},$$

which are equal to the numbers of centers of the blowing-up included in hyperplanes $\{f_n = 0\}$ and $\{f_n = \infty\}$, respectively. From now on, we assume for simplicity that

$$k_{n-1}k_{n+1} = \ell_{n-1}\ell_{n+1}.$$

In this case, (2.4) can be equivalently rewritten into

$$s_n^0(f_{n-1}) = f_{n-1} \frac{v_n^{\omega_n} f_n + v_n^{-1+\omega_n}}{u_n^{-\omega_n} f_n + u_n^{1-\omega_n}}, \quad (3.1a)$$

$$s_n^0(f_{n+1}) = f_{n+1} \frac{u_n^{-\omega_n} f_n + u_n^{1-\omega_n}}{v_n^{\omega_n} f_n + v_n^{-1+\omega_n}}, \quad (3.1b)$$

where $\omega_n = \theta_n^0 / (\theta_n^0 + \theta_n^\infty)$. Now we consider the decomposition of variables

$$f_n = \frac{x_{n+1}}{x_{n-1}}, \quad (3.2)$$

and take a birational transformation

$$s_n^i(x_n) = x_n \frac{v_n^{\omega_n} x_{n+1} + v_n^{-1+\omega_n} x_{n-1}}{u_n^{-\omega_n} x_{n+1} + u_n^{1-\omega_n} x_{n-1}}. \quad (3.3)$$

One can easily verify that (3.3) with (2.2) also defines a birational representation of W on the field $\mathbb{K}(x)$. Moreover, the representation (3.1) at the level of the f -variables is of course derived from (3.3) via (3.2). Introducing new variables τ_n^i attached to the centers C_n^i of the blowing-up, we shall consider a further decomposition of variables in terms of the τ -variables

$$x_n = \frac{\xi_n}{\eta_n}, \quad \xi_n = \prod_{i=1}^{k_n} \tau_n^i, \quad \eta_n = \prod_{j=1}^{\ell_n} \tau_n^{-j}. \quad (3.4)$$

Theorem 3.1. Define the birational transformations s_n^i by

$$s_n^0(\tau_n^1) = \frac{v_n^{\omega_n} \xi_{n+1} \eta_{n-1} + v_n^{-1+\omega_n} \xi_{n-1} \eta_{n+1}}{\tau_n^{-1}}, \quad (3.5a)$$

$$s_n^0(\tau_n^{-1}) = \frac{u_n^{-\omega_n} \xi_{n+1} \eta_{n-1} + u_n^{1-\omega_n} \xi_{n-1} \eta_{n+1}}{\tau_n^1}, \quad (3.5b)$$

$$s_n^i(\tau_n^{\{i, i+1\}}) = \tau_n^{\{i+1, i\}} \quad (1 \leq i \leq k_n - 1), \quad (3.5c)$$

$$s_n^{-j}(\tau_n^{\{-j, -j-1\}}) = \tau_n^{\{-j-1, -j\}} \quad (1 \leq j \leq \ell_n - 1), \quad (3.5d)$$

$$s_n^i(\tau_m^j) = \tau_m^j \quad (\text{otherwise}), \quad (3.5e)$$

where $\omega_n = \theta_n^0/(\theta_n^0 + \theta_n^\infty)$, $\xi_n = \prod_{i=1}^{k_n} \tau_n^i$ and $\eta_n = \prod_{j=1}^{\ell_n} \tau_n^{-j}$. Then (3.5) with (2.2) realizes the Weyl group $W(T_\ell^k)$ over the field $L = \mathbb{K}(\tau)$ of rational functions in indeterminates τ_n^i .

We remark that, conversely, both realizations in terms of f -variables and x -variables are deduced from Theorem 3.1 through (3.2) and (3.4).

3.2 τ -Function and its geometric meaning

By means of Theorem 3.1, we now introduce the concept of τ -functions; cf. [KMNOY03, Tsu06]. Let us consider the Weyl group orbit $M = W\{E_n^i\} \subset H^2(X, \mathbb{Z})$ of the classes of exceptional divisors. Note that $\dim |\Lambda| = 0$ for any $\Lambda \in M$.

Definition 3.2. The τ -function $\tau : M \rightarrow L$ is defined by

- (i) $\tau(E_n^i) = \tau_n^i$;
- (ii) $\tau(w\Lambda) = w\tau(\Lambda)$ for any $\Lambda \in M$ and $w \in W$.

By applying Theorem 3.1, we can determine inductively the explicit formula of $\tau(\Lambda)$ as a rational function in τ_n^i . Alternatively below, we shall present a geometric (and *a priori*) characterization of our τ -functions. Let us first prepare some notations; let $\zeta = (\zeta_1^0 : \zeta_1^\infty, \dots, \zeta_N^0 : \zeta_N^\infty)$ be the homogeneous coordinate of $(\mathbb{P}^1)^N$ such that

$$f_n = \frac{\zeta_n^0}{\zeta_n^\infty}. \quad (3.6)$$

For a multiple index $m = (m_1^0, m_1^\infty, \dots, m_N^0, m_N^\infty) \in (\mathbb{Z}_{\geq 0})^{2N}$, we set $\zeta^m = \prod_n (\zeta_n^0)^{m_n^0} (\zeta_n^\infty)^{m_n^\infty}$.

An element $\Lambda = \sum d_n H_n - \sum \mu_n^i E_n^i \in M$ corresponds to a unique hypersurface of degree $d = (d_1, \dots, d_N)$ passing through C_n^i with multiplicity μ_n^i . Let

$$\Phi_\Lambda(\zeta) = \sum_m A_m \zeta^m \in \mathbb{K}[\zeta] \quad (3.7)$$

be the defining polynomial of the hypersurface, where $A_m \in \mathbb{K}$ is a monic monomial in multiplicative root variables. Normalize Φ_Λ as its coefficients satisfy the following condition:

$$\prod_m A_m^{(1/\theta)^m} = 1, \quad (3.8)$$

with $(1/\theta)^m = \prod_n (1/\theta_n^0)^{m_n^0} (1/\theta_n^\infty)^{m_n^\infty}$. A few examples of the normalized defining polynomial $\Phi_\Lambda(\zeta)$ ($\Lambda \in M$) are

$$\begin{aligned} \Phi_{E_n^i} &= 1, \\ \Phi_{H_n - E_n^1} &= (u_n)^{-\omega_n} \zeta_n^\infty (f_n + u_n), \\ \Phi_{H_n + H_{n+1} - E_n^1 - E_{n+1}^{-1} - E_{n+1}^{-1}} &= u_n^{(1-\omega_n)\omega_{n+1}} v_n^{-(1-\omega_n)(1-\omega_{n+1})} v_{n+1}^{-1+\omega_{n+1}} \\ &\quad \times \zeta_n^\infty \zeta_{n+1}^\infty \left(1 + \frac{f_n}{u_n} + v_{n+1} f_{n+1} + v_n v_{n+1} f_n f_{n+1} \right). \end{aligned}$$

Now suppose that

$$\zeta_n^0 = \xi_{n+1} \eta_{n-1}, \quad \zeta_n^\infty = \xi_{n-1} \eta_{n+1}. \quad (3.9)$$

We then have an expression of the τ -function $\tau(\Lambda)$ in terms of the normalized defining polynomial $\Phi_\Lambda(\zeta)$ of the corresponding divisor Λ .

Theorem 3.3. For any $\Lambda = \sum_n d_n H_n - \sum_{n,i} \mu_n^i E_n^i \in M$, we have the equality

$$\tau(\Lambda) \prod_{n,i} (\tau_n^i)^{\mu_n^i} = \Phi_\Lambda(\zeta). \quad (3.10)$$

Hence, one can in principle trace the resulting value with respect to any iteration of Weyl group actions by means of the defining polynomials of appropriate divisors. We see from (3.10) that $\tau(\Lambda)$ is a Laurent polynomial in the indeterminates τ_n^i , though, from the definition of τ -functions, we can only state that $\tau(\Lambda)$ is rational. This *regularity* is an interesting feature of our τ -functions and should be relevant to the theory of infinite-dimensional integrable systems such as the KP hierarchy, UC hierarchy, etc.; see § 5. Note that the τ -function can be considered as a counterpart of height functions in the sense that the original inhomogeneous coordinate $f = (f_1, f_2, \dots, f_N)$ of $(\mathbb{P}^1)^N$ is recovered as a ratio of τ -functions (see (3.2) and (3.4)).

Proof of Theorem 3.3. We will prove the theorem by induction. If $\Lambda = H_n - E_n^{\pm 1} = s_n^0(E_n^{\mp 1}) \in M$, then (3.10) follows immediately from (3.5a) and (3.5b). We assume that (3.10) is true for $\Lambda \in M$. Then it is enough to verify for $\Lambda' = w(\Lambda)$ such that $w \in \{s_n^i\}$ is a generator of $W = W(T_\ell^k)$. The action of s_n^i ($i \neq 0$) is just a permutation of τ -variables, so we need only to concentrate upon the nontrivial case $w = s_n^0$.

For $\Lambda = \sum_n d_n H_n - \sum_{n,i} \mu_n^i E_n^i \in M$, let us introduce a polynomial $\varphi_\Lambda(f) \in \mathbb{K}[f]$ in the inhomogeneous coordinate $f = (f_1, \dots, f_N)$ of degree $d = (d_1, \dots, d_N)$ defined by $\varphi_\Lambda(f) = \Phi_\Lambda(\zeta) \prod_n (\zeta_n^\infty)^{-d_n}$.

Claim. We have

$$s_n^0(\varphi_\Lambda(f)) = c(f_n + u_n)^{-d_{n-1} + \mu_n^{-1}} (f_n + v_n^{-1})^{-d_{n+1} + \mu_n^1} \varphi_{s_n^0 \Lambda}(f), \quad (3.11)$$

where the normalizing constant c is given by $c = (u_n^{d_{n-1} - \mu_n^{-1}} v_n^{-d_{n+1} + \mu_n^1})^{\omega_n}$.

To prove this, first we notice that $s_n^0 \Lambda = \Lambda + \langle \Lambda, \check{\alpha}_n^0 \rangle \alpha_n^0 = \Lambda + (d_{n-1} + d_{n+1} - \mu_n^1 - \mu_n^{-1}) \alpha_n^0$; accordingly the degree $d' = (d'_1, \dots, d'_N)$ of $\varphi_{s_n^0 \Lambda}(f)$ reads $d'_n = d_{n-1} + d_n + d_{n+1} - \mu_n^1 - \mu_n^{-1}$ and $d'_i = d_i$ ($i \neq n$). The multiplicity of $C_n^1 = \{f_{n-1} = 0, f_n = -u_n, f_{n+1} = \infty\}$ on the hypersurface $\{\varphi_\Lambda(f) = 0\} \subset (\mathbb{P}^1)^N$ is given by $\text{ord}_{C_n^1}(\varphi_\Lambda) = \mu_n^1$. Hence the defining polynomial $\varphi_\Lambda(f)$ is of the form

$$\varphi_\Lambda(f) = \sum_{i \geq 0} \psi_{\mu_n^1 - d_{n+1} + i},$$

where $\psi_i = \psi_i(f)$ is homogeneous of degree i in $(f_{n-1}, f_n + u_n, f_{n+1}^{-1})$. In the same way, from $\text{ord}_{C_n^{-1}}(\varphi_\Lambda) = \mu_n^{-1}$, we can also describe it as $\varphi_\Lambda = \sum_{i \geq 0} \tilde{\psi}_{\mu_n^{-1} - d_{n-1} + i}$ where $\tilde{\psi}_i$ is homogeneous of degree i in $(f_{n-1}^{-1}, f_n + v_n^{-1}, f_{n+1})$. By applying (3.1), we see that $s_n^0(\varphi_\Lambda) \times (f_n + u_n)^{d_{n-1} - \mu_n^{-1}} (f_n + v_n^{-1})^{d_{n+1} - \mu_n^1}$ is a polynomial of degree at most d' and therefore equals $(\text{const.}) \times \varphi_{s_n^0 \Lambda}$. Here we used implicitly the fact that s_n^0 is a pseudo isomorphism (Theorem 2.2) and transforms $\{\varphi_\Lambda(f) = 0\}$ generically into $\{\varphi_{s_n^0 \Lambda}(f) = 0\}$. Since (3.1) keeps the normalization (3.8) invariant, we can immediately calculate the constant and arrive at (3.11).

Applying s_n^0 to both sides of (3.10) and using the claim above, we obtain

$$\begin{aligned}
(\text{LHS}) &\xrightarrow{s_n^0} \tau(\Lambda') \prod_{i,j} (\tau_j^i)^{\mu_j^i} \times \left(\frac{s_n^0(\tau_n^1)}{\tau_n^1} \right)^{\mu_n^1} \left(\frac{s_n^0(\tau_n^{-1})}{\tau_n^{-1}} \right)^{\mu_n^{-1}}, \\
(\text{RHS}) &= \varphi_\Lambda(f) \prod_j (\zeta_j^\infty)^{d_j} \\
&\xrightarrow{s_n^0} s_n^0(\varphi_\Lambda(f)) \prod_j (\zeta_j^\infty)^{d_j} \times \left(\frac{s_n^0(\tau_n^1)}{\tau_n^1} \right)^{d_{n+1}} \left(\frac{s_n^0(\tau_n^{-1})}{\tau_n^{-1}} \right)^{d_{n-1}} \\
&= \varphi_{\Lambda'}(f) \prod_j (\zeta_j^\infty)^{d'_j} \times \left(\frac{s_n^0(\tau_n^1)}{\tau_n^1} \right)^{\mu_n^1} \left(\frac{s_n^0(\tau_n^{-1})}{\tau_n^{-1}} \right)^{\mu_n^{-1}} (\tau_n^1 \tau_n^{-1})^{-d_{n-1}-d_{n+1}+\mu_n^1+\mu_n^{-1}}.
\end{aligned}$$

Here we have used the formulae $f_n + u_n = u_n^{\omega_n} s_n^0(\tau_n^{-1}) \tau_n^1 / \zeta_n^\infty$ and $f_n + v_n^{-1} = v_n^{-\omega_n} s_n^0(\tau_n^1) \tau_n^{-1} / \zeta_n^\infty$. Thus, we get

$$\Phi_{\Lambda'}(\zeta) = \varphi_{\Lambda'}(f) \prod_j (\zeta_j^\infty)^{d'_j} = \tau(\Lambda') \prod_{i,j} (\tau_j^i)^{\mu_j^i} \times (\tau_n^1 \tau_n^{-1})^{d_{n-1}+d_{n+1}-\mu_n^1-\mu_n^{-1}},$$

which is exactly (3.10) for $\Lambda' = s_n^0 \cdot \Lambda$. The proof of the theorem is complete. \square

Remark 3.4. We can naturally extend the definition of a τ -function also for the divisors $D_n^0 = H_n - \sum_i E_{n+1}^i - \sum_j E_{n-1}^{-j}$ and $D_n^\infty = H_n - \sum_i E_{n-1}^i - \sum_j E_{n+1}^{-j}$ by

$$\tau(D_n^0) \xi_{n+1} \eta_{n-1} = \zeta_n^0, \quad \tau(D_n^\infty) \xi_{n-1} \eta_{n+1} = \zeta_n^\infty.$$

Noticing that D_n^0 and D_n^∞ are invariant under the action of Weyl group $W = W(T_\ell^k)$, we set $\tau(D_n^0) = \tau(D_n^\infty) \equiv 1$. Therefore the assumption (3.9) consists with (3.10).

4 Examples of affine case and q -Painlevé equations

We assume that the Dynkin diagram T_ℓ^k is of affine type. Then the lattice part of the affine Weyl group $W(T_\ell^k)$ provides an interesting class of discrete dynamical systems of Painlevé type. This is based on the following facts:

(i) (Existence of commuting time evolutions) For each root $\alpha \in Q$, we have the translation t_α acting on the Néron-Severi bilattice $N(X) \simeq (H^2(X, \mathbb{Z}), H_2(X, \mathbb{Z}))$ by the Kac formula [Kac90, §6.5]:

$$t_\alpha(\Lambda) = \Lambda - \langle \Lambda, \check{\delta} \rangle \alpha + \left(\langle \Lambda, \check{\alpha} \rangle - \frac{1}{2} \langle \alpha, \check{\alpha} \rangle \langle \Lambda, \check{\delta} \rangle \right) \delta, \quad (4.1)$$

$$t_\alpha(\lambda) = \lambda - \langle \delta, \lambda \rangle \check{\alpha} + \left(\langle \alpha, \lambda \rangle - \frac{1}{2} \langle \alpha, \check{\alpha} \rangle \langle \delta, \lambda \rangle \right) \check{\delta}, \quad (4.2)$$

for $\Lambda \in H^2(X, \mathbb{Z})$ and $\lambda \in H_2(X, \mathbb{Z})$. Here $\delta = -\frac{1}{2}K_X$ and $\check{\delta} = -\frac{1}{2}k_X$ are $W(T_\ell^k)$ -invariants (Lemma 2.1) and play the role of the null vector and its dual, respectively. The additivity property of translations $t_\alpha t_\beta = t_\beta t_\alpha = t_{\alpha+\beta}$ holds. This implies that the discrete dynamical system

$$t_\alpha(f_i) = F_{\alpha,i}(a, f) \quad (i = 1, \dots, N)$$

corresponding to a root $\alpha \in Q$ has a set of commuting discrete time evolutions $\{t_\beta | \beta \in Q\}$, where $F_{\alpha,i}(a, f) \in \mathbb{K}(f)$ is a rational function explicitly determined by Theorem 2.2; cf. [NY98].

(ii) (**Quadratic degree growth**) From (i), we have

$$\begin{aligned} (t_\alpha)^n(H_i) &= t_{n\alpha}(H_i) \\ &= H_i - n\langle H_i, \check{\delta} \rangle \alpha + \left(n\langle H_i, \check{\alpha} \rangle - \frac{n^2}{2} \langle \alpha, \check{\alpha} \rangle \langle H_i, \check{\delta} \rangle \right) \delta. \end{aligned}$$

Therefore the value of pairing $\langle (t_\alpha)^n(H_i), h_j \rangle$ is in the order n^2 , that is, the degree of $(t_\alpha)^n(f_i)$ with respect to f_j is at most n^2 .

Generally, a quadratic degree growth asserts integrability of the dynamical system under consideration; in [Tak01], it was proved by a similar technique as above that the degree growth of every *second order* difference Painlevé equation (in Sakai's classification [Sak01]) is quadratic. Also, as shown in [Tak04], a certain class of integrable higher dimensional dynamics defined on generalized Del Pezzo varieties has a quadratic degree growth. More elementarily, the degree of the n^{th} iteration of an addition formula of an elliptic curve grows quadratically as a rational mapping; see, e.g., [Sil86].

In this section, we shall demonstrate the $A_r^{(1)}$ and $D_r^{(1)}$ cases as examples.

4.1 The case of $A_{N-1}^{(1)}$

Let us consider the case $k_n = \ell_n = 1$ ($n \in \mathbb{Z}/N\mathbb{Z}$), the corresponding Dynkin diagram T_ℓ^k is of type $A_{N-1}^{(1)}$. We have, from Theorems 2.2 and 3.1, a tropical representation of affine Weyl group $W(A_{N-1}^{(1)})$ as pseudo isomorphisms of a rational variety X_a . Moreover, we can extend $W(A_{N-1}^{(1)})$ to $W(A_{N-1}^{(1)}) \times W(A_1^{(1)})$ in the following manner.

First we reformulate the parameters u_n and v_n as follows:

$$u_n = \frac{a_n b_1}{b_0}, \quad v_n = \frac{a_n b_0}{b_1}, \quad (4.3)$$

where a_n and b_i ($i = 0, 1$) are nonzero parameters such that

$$\prod_{n=1}^N a_n = (b_0 b_1)^N = q^N. \quad (4.4)$$

Let $\gamma_{1,n}$ and $\check{\gamma}_{1,n}$ be a pair of vectors defined by

$$\begin{aligned} \gamma_{1,n} &= -\frac{K_X}{2} - H_n - E_n^1 + E_{n-1}^{-1} + E_n^{-1} + E_{n+1}^{-1} \in H^2(X, \mathbb{Z}), \\ \check{\gamma}_{1,n} &= h_n - e_n^1 \in H_2(X, \mathbb{Z}), \end{aligned}$$

for $n \in \mathbb{Z}/N\mathbb{Z}$. We see that these are mutually orthogonal (-2) -vectors, namely, $\langle \gamma_{1,i}, \check{\gamma}_{1,j} \rangle = -2\delta_{i,j}$. Define the action of the reflection $r_{1,n}$ associated with the pair $(\gamma_{1,n}, \check{\gamma}_{1,n})$ on $N(X) = (H^2(X, \mathbb{Z}), H_2(X, \mathbb{Z}))$ in the same way as (2.1). Let $\beta_1 = \sum_{n=1}^N \gamma_{1,n}$ and $\check{\beta}_1 = \sum_{n=1}^N \check{\gamma}_{1,n}$ and define the associated reflection r_1 by $r_1 = r_{1,1} \circ \cdots \circ r_{1,N}$. In parallel, we consider an involution $\iota : E_n^1 \leftrightarrow E_n^{-1}, e_n^1 \leftrightarrow e_n^{-1}$ of $N(X)$; let $(\beta_0, \check{\beta}_0) = \iota(\beta_1, \check{\beta}_1)$ and define $r_0 = \iota \circ r_1 \circ \iota$. Then we

have $\langle r_0, r_1 \rangle \simeq W(A_1^{(1)})$. Note that $(\beta_i, \check{\beta}_i)$ ($i = 0, 1$) is orthogonal to $(\alpha_n, \check{\alpha}_n)$, where $\alpha_n = \alpha_n^0 = H_n - E_n^1 - E_n^{-1}$ and $\check{\alpha}_n = \check{\alpha}_n^0 = h_{n-1} + h_{n+1} - e_n^1 - e_n^{-1}$ are the (co-)root bases of $A_{N-1}^{(1)}$. Furthermore, we have in fact

$$\begin{aligned} Q \oplus \mathbb{Z}\beta_1 &= \left(\bigoplus_{n=1}^N \mathbb{Z}\alpha_n \right) \oplus \mathbb{Z}\beta_1 = \{d_n^0, d_n^\infty\}^\perp \subset H^2(X, \mathbb{Z}), \\ \check{Q} \oplus \mathbb{Z}\check{\beta}_1 &= \left(\bigoplus_{n=1}^N \mathbb{Z}\check{\alpha}_n \right) \oplus \mathbb{Z}\check{\beta}_1 = \{D_n^0, D_n^\infty\}^\perp \subset H_2(X, \mathbb{Z}). \end{aligned}$$

In order to realize r_i ($i = 0, 1$) as birational transformations, we give the action of r_1 on the cohomology basis of $H^2(X, \mathbb{Z})$:

$$r_1(E_n^1) = E_n^1 + \gamma_{1,n} = \sum_{i \neq n} H_i - \sum_{j=1}^N E_j^1 - \sum_{k \neq n, n \pm 1} E_k^{-1},$$

$r_1(E_n^{-1}) = E_n^{-1}$ and $r_1(H_n) = H_n + \gamma_{1,n}$. For $\Lambda = r_1(E_n^1)$, the normalized defining polynomial reads

$$\Phi_\Lambda(\zeta) = G_n(\zeta) := \left(1 + \sum_{j=1}^{N-1} \prod_{i=1}^j \frac{u_{n+i}}{f_{n+i}} \right) \prod_{k=1}^{N-1} (u_{n+k})^{-1+k/N} \zeta_{n+k}^0.$$

By virtue of Theorem 3.3, we thus obtain a birational action of r_1 on τ -variables:

$$r_1(\tau_n^1) = \frac{G_n(\zeta)}{\prod_j \tau_j^1 \prod_{k \neq n, n \pm 1} \tau_k^{-1}}, \quad r_1(\tau_n^{-1}) = \tau_n^{-1},$$

where we recall that $\zeta_n^0 = \tau_{n+1}^1 \tau_{n-1}^{-1}$, $\zeta_n^\infty = \tau_{n-1}^1 \tau_{n+1}^{-1}$ and $f_n = \zeta_n^0 / \zeta_n^\infty$.

Now, we fix the action on the multiplicative root variables a_n and b_i as

$$\begin{aligned} s_n(a_n) &= a_n^{-1}, & s_n(a_{n \pm 1}) &= a_n a_{n \pm 1}, \\ r_i(b_i) &= b_i^{-1}, & r_i(b_j) &= b_i^2 b_j \quad (i \neq j), \\ \pi(a_n) &= a_{n+1}, & \iota(b_{\{0,1\}}) &= b_{\{1,0\}}. \end{aligned} \tag{4.5}$$

The following birational transformations $\langle s_n, \pi, r_i, \iota \rangle$ together with (4.5) realize the extended affine Weyl group $W = \widetilde{W}(A_{N-1}^{(1)}) \times \widetilde{W}(A_1^{(1)})$:

$$s_n(\tau_n^1) = \frac{v_n^{1/2} \tau_{n+1}^1 \tau_{n-1}^{-1} + v_n^{-1/2} \tau_{n-1}^1 \tau_{n+1}^{-1}}{\tau_n^{-1}}, \tag{4.6a}$$

$$s_n(\tau_n^{-1}) = \frac{u_n^{-1/2} \tau_{n+1}^1 \tau_{n-1}^{-1} + u_n^{1/2} \tau_{n-1}^1 \tau_{n+1}^{-1}}{\tau_n^1}, \tag{4.6b}$$

$$r_1(\tau_n^1) = \frac{G_n(\zeta)}{\prod_j \tau_j^1 \prod_{k \neq n, n \pm 1} \tau_k^{-1}}, \tag{4.6c}$$

and $r_0 = \iota \circ r_1 \circ \iota$ with $\pi(\tau_n^{\pm 1}) = \tau_{n+1}^{\pm 1}$ and $\iota(\tau_n^{\pm 1}) = \tau_n^{\mp 1}$. This is in fact equivalent to the representation due to Kajiwara et al. [KNY02a, KNY02b], which was originally discovered in connection with a q -analogue of KP hierarchy, without any algebro-geometric setup. The birational action $f_n =$


$(\tau_{n+1}^1 \tau_{n-1}^{-1})/(\tau_{n-1}^1 \tau_{n+1}^{-1}) \mapsto \overline{f_n}$ of the translation $r_1 \circ \iota : (b_1, b_0; a_n) \mapsto (qb_1, b_0/q; a_n)$ is the q -Painlevé equation of type $A_{N-1}^{(1)}$:

$$\overline{f_n} = f_{n+1} \frac{g_{n-1}(f)}{g_{n+1}(f)}, \quad (4.7)$$

where $g_n(f) = (1 + \sum_{j=1}^{N-1} \prod_{i=1}^j u_{n+i}/f_{n+i}) \prod_{k=1}^{N-1} (u_{n+k})^{-1+k/N}$. The family $\mathcal{X} = \cup_a X_a$ of rational varieties provides a sort of phase space for the dynamics of (4.7).

4.2 The case of $D_{N+3}^{(1)}$

For $k_1 = \ell_1 = k_N = \ell_N = 2$ and $k_i = \ell_i = 1$ ($2 \leq i \leq N-1$), the Dynkin diagram T_ℓ^k is of type $D_{N+3}^{(1)}$ and looks as follows:



We fix the action of simple reflections s_i on the multiplicative root variables $a = (a_0, \dots, a_{N+3})$ as $s_i(a_j) = a_j a_i^{-C_{ij}}$, where C_{ij} is the Cartan matrix of type $D_{N+3}^{(1)}$. We define

$$\begin{aligned} u_1 &= a_1 a_0^{-1/2} a_{N+3}^{1/2}, & v_1 &= a_1 a_0^{1/2} a_{N+3}^{-1/2}, \\ u_i &= v_i = a_i \quad (2 \leq i \leq N-1), \\ u_N &= a_N a_{N+1}^{-1/2} a_{N+2}^{1/2}, & v_N &= a_N a_{N+1}^{1/2} a_{N+2}^{-1/2}. \end{aligned}$$

From Theorem 3.1, we have the following tropical representation of the affine Weyl group $W(D_{N+3}^{(1)})$ over the field $L = \mathbb{C}(a^{1/4})(\tau_n^i)$:

$$\begin{aligned} s_n(\tau_n^1) &= \frac{v_n^{1/2} \xi_{n+1} \eta_{n-1} + v_n^{-1/2} \xi_{n-1} \eta_{n+1}}{\tau_n^{-1}}, \\ s_n(\tau_n^{-1}) &= \frac{u_n^{-1/2} \xi_{n+1} \eta_{n-1} + u_n^{1/2} \xi_{n-1} \eta_{n+1}}{\tau_n^1}, \end{aligned}$$

for $1 \leq n \leq N$, $s_0(\tau_1^{\{1,2\}}) = \tau_1^{\{2,1\}}$, $s_{N+3}(\tau_1^{\{-1,-2\}}) = \tau_1^{\{-2,-1\}}$, $s_{N+1}(\tau_N^{\{1,2\}}) = \tau_N^{\{2,1\}}$ and $s_{N+2}(\tau_N^{\{-1,-2\}}) = \tau_N^{\{-2,-1\}}$. Here we set $\xi_n = \prod_{i=1}^{k_n} \tau_n^i$ and $\eta_n = \prod_{j=1}^{\ell_n} \tau_n^{-j}$ for $1 \leq n \leq N$ and $\xi_n = \eta_n \equiv 1$ for $n = 0, N+1$. When $N = 2$ ($D_5^{(1)}$ case), the above representation in τ -variables coincides with that of [TM06, §1]. So, the birational action on f -variables of a translation part of the Weyl group yields the q -Painlevé VI equation [JS96, Sak01]. For general $N \geq 3$, one may regard it as a (higher order) generalization of the q -Painlevé VI equation equipped with $W(D_{N+3}^{(1)})$ symmetry.

Remark 4.1. We shall make a slightly technical remark. In order to describe the (co)-root lattice of $D_{N+3}^{(1)}$ appropriately inside the Néron-Severi bilattice $N(X)$, we need to consider rational variety X blown-up from $(\mathbb{P}^1)^{N+1}$. However, such X turns out to be redundant for the dynamical system under consideration, because each $W(D_{N+3}^{(1)})$ -orbit of a generic point is confined in a certain hypersurface of X . Hence the essential dimension of the space of initial values for the $W(D_{N+3}^{(1)})$ -dynamics is N ; for instance, recall that the q -Painlevé VI equation ($D_5^{(1)}$ case) is of second order.

5 τ -Functions and character polynomials

In this section, for the case of $A_{N-1}^{(1)}$, we try to interpret from a soliton-theoretic point of view the relationship between our τ -functions and character polynomials, i.e., the Schur functions or the universal characters.

5.1 Universal characters

For a pair of partitions $\lambda = (\lambda_1, \lambda_2, \dots, \lambda_r)$ and $\mu = (\mu_1, \mu_2, \dots, \mu_{r'})$, the *universal character* $S_{[\lambda, \mu]}(x, y)$ is a polynomial in $(x, y) = (x_1, x_2, \dots, y_1, y_2, \dots)$ defined by the determinant formula of *twisted* Jacobi-Trudi type [Koi89]:

$$S_{[\lambda, \mu]}(x, y) = \det \left(\begin{array}{cc} p_{\mu_{r'-i+1}+i-j}(y), & 1 \leq i \leq r' \\ p_{\lambda_{i-r'}-i+j}(x), & r' + 1 \leq i \leq r + r' \end{array} \right)_{1 \leq i, j \leq r+r'}, \quad (5.1)$$

where p_n is a polynomial defined by the generating function $\sum_{k \in \mathbb{Z}} p_k(x) z^k = \exp(\sum_{n=1}^{\infty} x_n z^n)$ and $p_n(x) = 0$ for $n < 0$. The Schur function $S_{\lambda}(x)$ (see, e.g., [Mac95]) is regarded as a special case of the universal character:

$$S_{\lambda}(x) = \det(p_{\lambda_i-i+j}(x)) = S_{[\lambda, \emptyset]}(x, y).$$

If we count the degree of the variables as $\deg x_n = n$ and $\deg y_n = -n$, then $S_{[\lambda, \mu]}(x, y)$ is a weighted homogeneous polynomial of degree $|\lambda| - |\mu|$, where $|\lambda| = \lambda_1 + \dots + \lambda_r$. The universal character $S_{[\lambda, \mu]}$ describes the irreducible character of a rational representation of the general linear group $GL(n; \mathbb{C})$ corresponding to a pair of partitions $[\lambda, \mu]$, while the Schur function S_{λ} does that of a polynomial representation corresponding to a partition λ ; see [Koi89] for details.

Example 5.1. For $\lambda = (2, 1)$ and $\mu = (1)$, we have

$$S_{[(2,1),(1)]}(x, y) = \begin{vmatrix} p_1(y) & p_0(y) & p_{-1}(y) \\ p_1(x) & p_2(x) & p_3(x) \\ p_{-1}(x) & p_0(x) & p_1(x) \end{vmatrix} = \left(\frac{x_1^3}{3} - x_3 \right) y_1 - x_1^2.$$

5.2 Notations

Let us recall some terminology relevant to the case at hand. A subset $\mathbf{m} \subset \mathbb{Z}$ is said to be a *Maya diagram* if $i \in \mathbf{m}$ (for $i \ll 0$) and $i \notin \mathbf{m}$ (for $i \gg 0$). Each Maya diagram $\mathbf{m} = \{\dots, m_3, m_2, m_1\}$ corresponds to a unique partition $\lambda = (\lambda_1, \lambda_2, \dots)$ such that $m_i - m_{i+1} = \lambda_i - \lambda_{i+1} + 1$. For a sequence of integers $\nu = (\nu_1, \nu_2, \dots, \nu_N) \in \mathbb{Z}^N$, we consider the Maya diagram

$$\mathbf{m}(\nu) = (N\mathbb{Z}_{<\nu_1} + 1) \cup (N\mathbb{Z}_{<\nu_2} + 2) \cup \dots \cup (N\mathbb{Z}_{<\nu_N} + N),$$

and denote by $\lambda(\nu)$ the corresponding partition. We call a partition of the form $\lambda(\nu)$ an *N-core partition*. It is well-known that a partition λ is *N-core* if and only if λ has no hook with length of a multiple of N . See [Nou04].

Example 5.2. If $N = 3$ and $\nu = (2, 0, 3)$ then the resulting partition reads $\lambda(\nu) = (4, 2, 1, 1)$.

Also, we prepare the following standard notations of q -analysis [GR90].
 q -shifted factorials:

$$(z; q)_\infty = \prod_{i=0}^{\infty} (1 - zq^i), \quad (z; p, q)_\infty = \prod_{i,j=0}^{\infty} (1 - zp^i q^j),$$

and $(z_1, \dots, z_r; q)_\infty = (z_1; q)_\infty \cdots (z_r; q)_\infty$, etc.

Elliptic gamma function:

$$\Gamma(z; p, q) = \frac{(pqz^{-1}; p, q)_\infty}{(z; p, q)_\infty}.$$

5.3 τ -Functions in terms of character polynomials

In what follows, we shall only treat the case of $A_{N-1}^{(1)}$ (see § 4.1). Introduce the lattice $P(A_{\ell-1}) = \mathbb{Z}^\ell / \mathbb{Z}(\mathbf{e}_1 + \cdots + \mathbf{e}_\ell)$ of rank $\ell - 1$, called the *weight lattice* of type $A_{\ell-1}$. Here $\{\mathbf{e}_i\}$ refers to the canonical basis of \mathbb{Z}^ℓ . Note that $P(A_1) \simeq \mathbb{Z}$. For an element $\Lambda \in M = W.\{E_n^\pm\}$ of the Weyl group orbit, we can associate a sequence of integers $\nu = (\nu_1, \dots, \nu_N) \in \mathbb{Z}^N$ and an integer $\kappa \in \mathbb{Z}$ through the formulae

$$\nu_i - \nu_{i+1} = \langle \Lambda, \check{\alpha}_i \rangle \quad (i \neq N), \quad \nu_N - \nu_1 + 1 = \langle \Lambda, \check{\alpha}_N \rangle \quad \text{and} \quad \kappa = \langle \Lambda, \check{\beta}_0 \rangle.$$

Since $\nu \in \mathbb{Z}^N$ is uniquely determined modulo $\mathbb{Z}(\mathbf{e}_1 + \cdots + \mathbf{e}_N)$, we can regard (ν, κ) as a point of $P(A_{N-1}) \times P(A_1)$. In fact, this is a one-to-one correspondence between M and $P(A_{N-1}) \times P(A_1)$. Now let us index the τ functions as

$$\tau(\Lambda) =: \sigma_\nu^\kappa. \tag{5.2}$$

For example, we see that $\tau(E_n^1) = \sigma_{\mathbf{e}_1 + \cdots + \mathbf{e}_n}^0$ and $\tau(E_n^{-1}) = \sigma_{\mathbf{e}_1 + \cdots + \mathbf{e}_n}^1$. We have in general

$$T_n(\sigma_\nu^\kappa) = \sigma_{\nu + \mathbf{e}_n}^\kappa \quad (n \in \mathbb{Z}/N\mathbb{Z}), \quad \widetilde{T}(\sigma_\nu^\kappa) = \sigma_\nu^{\kappa+1},$$

where $T_n = \pi \circ s_{n+N-2} \circ \cdots \circ s_{n+1} \circ s_n$ and $\widetilde{T} = r_1 \circ \iota$ are the translations of affine Weyl groups $\widetilde{W}(A_{N-1}^{(1)})$ and $\widetilde{W}(A_1^{(1)})$, respectively. As previously seen in § 3, the τ -function σ_ν^κ is expressed as a rational function in the indeterminates τ_i^\pm by means of the defining polynomial of the corresponding divisor. However, substituting appropriate special values for the indeterminates τ_i^\pm , we have another expression of the τ -function in terms of the character polynomials.

Theorem 5.3 (cf. Kajiwara-Noumi-Yamada [KNY02b]). *Define*

$$F(t) = \frac{(q^{2N} t^{2N}; q^{2N}, q^{2N})_\infty}{(q^2 t^2; q^2, q^2)_\infty} \Gamma(-q^{N-1} t^{N-1}; q^{N-1}, q^{N-1}),$$

$$H_\nu = \prod_{(i,j) \in \lambda(\nu)} (q^{h_{ij}} - q^{-h_{ij}}) q^{(i-j)/2},$$

where h_{ij} is the hook-length, that is, $h_{ij} = \lambda_i + \lambda'_j - i - j + 1$. Under the specialization $a_i = q$, $\tau_i^1 = F(1/b_0)$ and $\tau_i^{-1} = F(b_1)$ for all $i \in \mathbb{Z}/N\mathbb{Z}$, the τ -function σ_ν^κ is expressed in terms of the Schur function attached to the N -core partition $\lambda(\nu)$ as

$$\sigma_\nu^\kappa = F(t) H_\nu S_{\lambda(\nu)}(x), \tag{5.3}$$

where $t = q^\kappa/b_0$ and

$$x_n = \frac{t^n + t^{-n}}{n(q^n - q^{-n})} \quad (n = 1, 2, \dots).$$

This theorem is based on the principle that appropriate reductions of infinite-dimensional integrable systems, such as the KP hierarchy, yield Painlevé-type equations. In fact, the Weyl group actions (4.6) lead us to the bilinear relation among τ -functions $\sigma_\nu^\kappa \in \mathbb{K}(\tau_i^\pm)$ of the form

$$q^{N\nu_i - |\nu| + i - 1} c_i (t^N - t^{-N}) \sigma_\nu^\kappa \sigma_{\nu+e_i}^\kappa = t \sigma_\nu^{\kappa-1} \sigma_{\nu+e_i}^{\kappa+1} - t^{-1} \sigma_\nu^{\kappa+1} \sigma_{\nu+e_i}^{\kappa-1}, \quad (5.4)$$

where $t = q^\kappa/b_0$ and $c_i = \prod_{j=1}^{N-1} (a_{i+j-1}/q)^{-1+j/N}$. The bilinear relation (5.4) above is exactly the similarity reduction of the q -KP hierarchy. Since the Schur function S_λ is a *homogeneous* polynomial solution of the q -KP hierarchy and is thus compatible with the similarity reduction, we can immediately verify Theorem 5.3; see [KNY02b] for details. Moreover, when N is even, (5.4) can also be obtained, alternatively, from the q -UC hierarchy by a similarity reduction [Tsu04, Tsu05a]. Therefore, the following theorem is a direct consequence of the fact that the universal character $S_{[\lambda, \mu]}$ is homogeneous and solves the q -UC hierarchy.

Theorem 5.4. *In the case $N = 2g + 2$ ($g = 1, 2, \dots$), define*

$$\begin{aligned} \widetilde{F}(a, t) &= \frac{(-aq^2t^2, -a^{-1}q^2t^2; q^2, q^4)_\infty (q^{4g+4}t^{4g+4}; q^{4g+4}, q^{4g+4})_\infty}{(q^4t^4; q^4, q^4)_\infty} \Gamma(-a^{1/2}qt, -a^{-1/2}qt; q, q^2), \\ \widetilde{H}_\nu &= a_\nu^{(|\lambda(\nu_{\text{even}})| - |\lambda(\nu_{\text{odd}})|)/2} \prod_{(i,j) \in \lambda(\nu_{\text{odd}})} (q^{2h_{ij}} - q^{-2h_{ij}}) \prod_{(i,j) \in \lambda(\nu_{\text{even}})} (q^{-2h_{ij}} - q^{2h_{ij}}), \end{aligned}$$

with $a_\nu = aq^{2(|\nu_{\text{even}}| - |\nu_{\text{odd}}|)}$, $\nu_{\text{even}} = (\nu_2, \nu_4, \dots, \nu_{2g+2})$ and $\nu_{\text{odd}} = (\nu_1, \nu_3, \dots, \nu_{2g+1})$. Under the specialization $a_{2i+1} = a$, $a_{2i} = a^{-1}q^2$, $\tau_{2i}^1 = \widetilde{F}(a, 1/b_0)$, $\tau_{2i}^{-1} = \widetilde{F}(a, b_1)$, $\tau_{2i+1}^1 = \widetilde{F}(aq^{-2}, 1/b_0)$ and $\tau_{2i+1}^{-1} = \widetilde{F}(aq^{-2}, b_1)$ for all $i \in \mathbb{Z}/(g+1)\mathbb{Z}$, the τ -function σ_ν^κ is expressed in terms of the universal character attached to a pair of $(g+1)$ -core partitions as

$$\sigma_\nu^\kappa = \widetilde{F}(a_\nu, t) \widetilde{H}_\nu S_{[\lambda(\nu_{\text{odd}}), \lambda(-\nu_{\text{even}})]}(x, y), \quad (5.5)$$

where $t = q^\kappa/b_0$ and

$$x_n = \frac{t^n + t^{-n} - (-a_\nu)^n (q^n + q^{-n})}{n(q^{2n} - q^{-2n})}, \quad y_n = \frac{t^n + t^{-n} - (-a_\nu)^{-n} (q^n + q^{-n})}{n(q^{-2n} - q^{2n})} \quad (n = 1, 2, \dots).$$

Taking into account the origin of our realization of Weyl groups, Theorems 5.3 and 5.4 exemplify a quite curious coincidence between algebraic geometry of particular rational varieties and representation theory. Similar results have been established also for the $D_5^{(1)}$ and $E_6^{(1)}$ cases [Tsu05b, TM06] by the use of the q -UC hierarchy. The authors expect there should exist another geometric understanding of such a relationship between τ -functions and character polynomials, without employing the theory of infinite-dimensional integrable systems.

Acknowledgement. The authors wish to thank Ralph Willox for his helpful comments on the first draft of this article. They have also benefited from discussions with colleagues in Kobe: Tetsu Masuda, Masatoshi Noumi, Yasuhiro Ohta, Masa-Hiko Saito and Yasuhiko Yamada. This work was partially supported by a grant-in-aid from the Japan Society for the Promotion of Science (JSPS).

References

- [Cob29] A. B. Coble, *Algebraic geometry and theta functions*, (Amer. Math. Soc., Providence, RI, 1929).
- [DO88] I. Dolgachev and D. Ortland, *Point sets in projective spaces and theta functions*, Astérisque **165** (1988).
- [GR90] G. Gasper and M. Rahman, *Basic hypergeometric series*, Encyclopedia of Mathematics and its Applications **35** (Cambridge University Press, Cambridge, 1990).
- [JS96] M. Jimbo and H. Sakai, *A q -analog of the sixth Painlevé equation*, Lett. Math. Phys. **38** (1996), 145–154.
- [Kac90] V. G. Kac, *Infinite dimensional Lie algebras*, 3rd ed. (Cambridge University Press, Cambridge, 1990).
- [KMNOY03] K. Kajiwara, T. Masuda, M. Noumi, Y. Ohta and Y. Yamada, *${}_{10}E_9$ solution to the elliptic Painlevé equation*, J. Phys. A: Math. Gen. **36** (2003), L263–L272.
- [KMNOY06] K. Kajiwara, T. Masuda, M. Noumi, Y. Ohta and Y. Yamada, *Point configurations, Cremona transformations and the elliptic difference Painlevé equation*, Séminaires et Congrès **14** (2006), 175–204.
- [KNY02a] K. Kajiwara, M. Noumi and Y. Yamada, *Discrete dynamical systems with $W(A_{m-1}^{(1)} \times A_{n-1}^{(1)})$ symmetry*, Lett. Math. Phys. **60** (2002), 211–219.
- [KNY02b] K. Kajiwara, M. Noumi and Y. Yamada, *q -Painlevé systems arising from q -KP hierarchy*, Lett. Math. Phys. **62** (2002), 259–268.
- [Kir01] A. N. Kirillov, *Introduction to tropical combinatorics* (Physics and combinatorics, 2000, eds. A. N. Kirillov and N. Liskova, Singapore: World Scientific, 2001, pp. 82–150).
- [Koi89] K. Koike, *On the decomposition of tensor products of the representations of the classical groups: By means of the universal characters*, Adv. Math. **74** (1989), 57–86.
- [Mac95] I. G. Macdonald, *Symmetric Functions and Hall Polynomials*, 2nd ed. Oxford Mathematical Monographs (Oxford University Press Inc., New York, 1995).
- [Nou04] M. Noumi, *Painlevé equations through symmetry*, (American Mathematical Society, 2004).
- [NY98] M. Noumi and Y. Yamada, *Affine Weyl groups, discrete dynamical systems and Painlevé equations*, Comm. Math. Phys. **199** (1998), 281–295.
- [ORG01] Y. Ohta, A. Ramani and B. Grammaticos, *An affine Weyl group approach to the eight-parameter discrete Painlevé equation*, J. Phys. A: Math. Gen. **34** (2001), 10523–10532.
- [Sak01] H. Sakai, *Rational surfaces associated with affine root systems and geometry of the Painlevé equations*, Comm. Math. Phys. **220** (2001), 165–229.
- [Sil86] J. H. Silverman, *The arithmetic of elliptic curves*, (Springer-Verlag, New York, 1986).
- [Tak01] T. Takenawa, *Algebraic entropy and the space of initial values for discrete dynamical systems*, J. Phys. A: Math. Gen. **34** (2001), 10533–10545.

- [Tak04] T. Takenawa, *Integrability of n -dimensional dynamical systems of type $E_7^{(1)}$ and $E_8^{(1)}$* , (Preprint, arXiv:nlin/0409051).
- [TTMS96] T. Tokihiro, D. Takahashi, J. Matsukidaira and J. Satsuma, *From soliton equations to integrable cellular automata through a limiting procedure*, Phys. Rev. Lett. **76** (1996), 3247–3250.
- [Tsu04] T. Tsuda, *Universal characters and an extension of the KP hierarchy*, Comm. Math. Phys. **248** (2004), 501–526.
- [Tsu05a] T. Tsuda, *Universal characters and q -Painlevé systems*, Comm. Math. Phys. **260** (2005), 59–73.
- [Tsu05b] T. Tsuda, *Universal character and q -difference Painlevé equations with affine Weyl groups*, (Preprint, UTMS 2005–21).
- [Tsu06] T. Tsuda, *Tropical Weyl group action via point configurations and τ -functions of the q -Painlevé equations*, Lett. Math. Phys. **77** (2006), 21–30.
- [TM06] T. Tsuda and T. Masuda, *q -Painlevé VI equation arising from q -UC hierarchy*, Comm. Math. Phys. **262** (2006), 595–609.

Teruhisa Tsuda tudateru@math.kyushu-u.ac.jp
 Faculty of Mathematics, Kyushu University, Hakozaki, Fukuoka 812-8581, Japan

Tomoyuki Takenawa takenawa@kaiyodai.ac.jp
 Faculty of Marine Technology, Tokyo University of Marine Science and Technology, Echujiima, Tokyo 135-8533, Japan