

Discovery of a Bipolar Outflow from 2MASSW J1207334-393254 a 24 M_{jup} Brown Dwarf

E.T. Whelan¹

T.P. Ray¹

S. Randich²

F. Bacciotti³

R. Jayawardhana²

L. Testi²

A. Natta²

S. Mohanty⁴

ABSTRACT

The 24 M_{jup} brown dwarf 2MASS1207-3932 has for some time been known to show clear signs of classical T Tauri-like accretion. Through analysis of its oxygen forbidden emission we have discovered that it is driving a bipolar outflow. Blue and red-shifted components to the [OI] λ 6300 forbidden emission line are seen at velocities of -8kms^{-1} and $+4\text{kms}^{-1}$. Spectro-astrometry recovers the position of both components relative to the BD, at $\sim 0''.08$ (in opposing directions). A position velocity diagram of the line region supports the spectro-astrometric results. The $\text{H}\alpha$ and $\text{HeI}\lambda$ 6678 lines were also analysed. These line regions are not offset with respect to the continuum ruling out the presence of spectro-astrometric artifacts and underlining the validity of the [OI] λ 6300 results. The low radial velocity

¹School of Cosmic Physics, Dublin Institute for Advanced Studies

²Osservatorio Astrofisico di Arcetri

³Department of Astronomy and Astrophysics, University of Toronto

⁴Harvard-Smithsonian CfA

of the outflow, and relatively large offsets, are consistent with 2MASS1207-3932 having a near edge-on disk, as proposed by Scholz et al. (2005). 2MASS1207-3932 is now the smallest mass galactic object known to drive an outflow. The age of the TW Hydrae Association (~ 8 Myr) also makes this one of the oldest objects with a resolved jet. This discovery not only highlights the robustness of the outflow mechanism over an enormous range of masses but also suggests that it may even be feasible for young giant planets with accretion disks to drive outflows.

Subject headings: 2MASSWJ1207334-393254 — stars: low mass, brown dwarfs — stars: formation — ISM: jets and outflows

1. Introduction

Recent studies have highlighted the numerous ways in which young brown dwarfs (BDs) bear a striking resemblance to classical T Tauri stars (CTTSs) (Jayawardhana et al. 2003a; Natta et al. 2004; Whelan et al. 2005; Scholz & Jayawardhana 2006). This comparison not only provides information relevant to theories of BD formation, but also allows the validity of the accretion/ejection models for the formation of solar-mass stars, to be tested in the sub-stellar domain. Emphasis has been on probing accretion (Jayawardhana et al. 2003b; Natta & Testi 2001). While such studies point to T Tauri like infall in BDs, relatively little is known about their outflow activity. Outflows in CTTSs are directly related to magnetospheric infall (Hartigan et al. 1994; Konigl & Pudritz 2000), hence it is feasible that BDs demonstrating strong accretion will drive jets.

The $H\alpha$ 10% line width criterion (full width of the line at 10% of the peak emission) used to distinguish accreting T Tauri stars from those where the line emission is primarily due to chromospheric activity, has been successfully applied to BDs (Jayawardhana et al. 2003a). A BD with a $H\alpha$ 10% width $> 200 \text{ km s}^{-1}$ is accepted as a strong accretor. High resolution spectroscopic observations subsequently found that many of the accreting BDs also have forbidden emission lines (FELs) (Fernández & Comerón 2001; Muzerolle et al. 2003). The FELs of CTTSs are important outflow tracers and their discovery provided the first hint that accreting BDs could also power outflows. Direct imaging has proven very difficult as expected on theoretical grounds (Masciadri & Raga 2004). An alternative approach adopted by Whelan et al. (2005) is to search for sub-stellar outflows through the spectro-astrometric analysis of FELs. Using this method the first BD outflow, driven by ρ -Oph 102, was discovered. Here the detection of a faint bipolar jet from a second strongly accreting BD, 2MASS1207-3932 (hereafter 2MASS1207) using the ESO VLT, is reported.

These observations are part of an important on-going study to search for outflows from BDs.

That such outflows are present is physically reasonable since it seems unlikely that the outflow mechanism has anything to do with the eventual triggering of fusion in the core. Moreover jets from the very low mass stars Par-Lup3-4 (Fernández & Comerón 2005) and ESO-H α 574 (Comerón & Reipurth 2006) have been found recently. In addition a molecular outflow from the L1014 dense low luminosity core and possible BD precursor has been found by Bourke et al. (2005).

2. Target, Observations and Analysis

2MASS1207 was identified by Gizis (2002) and Mohanty et al. (2003) as a sub-stellar member of the TW Hydrae Association. Given the age of the association (~ 8 Myr), the evolutionary tracks of Chabrier et al. (2000) were then used to estimate a mass of $\sim 35 M_{jup}$ for this BD. More recently the mass has been revised downwards to $24M_{jup} \pm 6M_{jup}$ (Chauvin et al. 2005; Mohanty et al. 2006). The nature of this BD as a strong accretor was clearly evident from early observations. Mohanty et al. (2003) reported bright asymmetric H α emission, with a full width at 10% of the peak flux, of $> 200 \text{ kms}^{-1}$. The new optical spectra presented here, lead to a value of $\sim 320 \text{ kms}^{-1}$. Mohanty et al. (2005) were the first to observe the [OI] λ 6300 line in the spectrum of 2MASS1207 suggesting it could be driving an outflow.

2MASS1207 is known for its line variability. Scholz et al. (2005) studied its H α emission over a period of 6 weeks and reported a change by a factor of 5-10 in the accretion rate during that time. They also used the shape and variability of the H α line to infer a near edge-on orientation for the accretion disk. The prominent red-shifted absorption feature in the H α line (see Figure 2b) is strongly dependent on the inclination i , between the rotational axis and the line of sight (Hartmann et al. 1994). The absorption feature should only be observed when the disk is seen close to edge-on i.e. $i \geq 60^\circ$. Refer to Scholz et al. (2005) and Scholz & Jayawardhana (2006) for further details. Such a geometry suggests that an outflow, will be in the plane of the sky. Lastly 2MASS1207 is a binary with a planetary mass companion (Chauvin et al. 2004, 2005). To summarise 2MASS1207 is an ideal candidate to include in our search for outflows driven by young BDs. It is a strong accretor with known [OI] λ 6300 emission.

High resolution ($R=40,000$) cross-dispersed spectra of 2MASS1207 (CD3 disperser, spectral range 4810-6740 Å) were obtained with UV-Visual Echelle Spectrometer (UVES) (Dekker et al. 2000) on the ESO VLT UT2, on May 16 2006. The results presented in this

letter represent the sum of two 45 min exposures with a $1''$ slit. The seeing was $0''.6$ and the spatial sampling was $0''.182$ per pixel. The data was reduced using standard IRAF routines, and the 2D spectra obtained were smoothed using an elliptical Gaussian filter of FWHM $0.12 \text{ \AA} \times 0''.22$. The Gaussian smoothing improved the signal to noise and effectively decreased the spectral and spatial resolution to $R \sim 8000$ and $0''.7$ respectively. The night sky lines were then removed. All velocities are quoted with respect to the systemic velocity of 2MASS1207 which is measured from the $\text{LiI}\lambda 6708$ line to be $+2 \text{ km s}^{-1}$, in the LSR frame. The final stage in the analysis was to apply the technique of spectro-astrometry to the smoothed spectra.

For the majority of protostars with jets, the FEL regions of their spectra are very obviously spatially extended, particularly at high velocities, e.g. the FELs of the CTTS DG Tau trace a $\sim 12''$ knotty jet (Whelan et al. 2004). When however the emission is very compact and close to the source, specialised methods have to be used to probe its nature. In particular spectro-astrometry can investigate through Gaussian fitting, the offset of emission line features smaller than the seeing disc of the observation. The result of this technique is an offset-velocity diagram with the displacement of a spectral feature, e.g. the $[\text{OI}]\lambda 6300$, line shown as a function of velocity and relative to the continuum centroid of the object (Bailey 1998; Takami et al. 2001; Whelan et al. 2004, 2005). Figure 2 shows the results of the spectro-astrometry and see Whelan et al. (2005) for a thorough discussion of the method.

3. Results

The $[\text{OI}]\lambda\lambda 6300, 6363$ lines are the only “traditional” FEL tracers seen in the UVES spectrum. A position velocity (PV) diagram of the smoothed $[\text{OI}]\lambda 6300$ spectrum is presented in Figure 1. Blue and red-shifted emission is observed at velocities of $\sim -8 \text{ km s}^{-1}$ and $\sim +4 \text{ km s}^{-1}$ and a relative displacement between the two parts of the line is apparent. Also the relatively very small radial velocity of the outflow is consistent with the near edge-on disk hypothesis for 2MASS1207. We observe the $[\text{OI}]\lambda 6300$ line at the same velocity as reported by Mohanty et al. (2005). However due to the use of a larger telescope (8m as opposed to 6.5m) and a longer exposure time we retrieved more detailed kinematic information. Note also that Scholz et al. (2005) failed to detect forbidden emission in the spectrum of this BD. Again this is due to the use of a smaller telescope and the larger spectral resolution of their observations ($R \sim 60,000$). The $[\text{OI}]$ night sky line has been removed here and there is emission (at $\sim -35 \text{ km s}^{-1}$) on the farside of the night sky line position. This is probably also associated with the blue-shifted outflow.

The results of a spectro-astrometric analysis of the smoothed spectrum is shown in Figure 2(a). The first step in this analysis is to map the position of the BD through Gaussian




Fig. 1.— Continuum subtracted Position Velocity diagram of the $[\text{OI}]\lambda 6300$ line in 2MASS1207. Here the spectrum was smoothed in both the spectral and spatial directions using an elliptical Gaussian function of FWHM $0.12 \text{ \AA} \times 0''.22$. The contours begin at 3 times the r.m.s noise and increase in intervals of the r.m.s noise. The zero spatial offset line is measured from an accurate mapping of the continuum position using spectro-astrometry. The red and blue-shifted components to the line are obvious and an opposing offset in these components is suggested. This offset is recovered using spectro-astrometry (see Figure 2). The $[\text{OI}]\lambda 6300$ night sky line has been removed from the spectrum and the feature blueward of this sky line, at $\sim -35 \text{ km s}^{-1}$ is probably also part of the blue-shifted outflow.

fitting of the spatial profile of the continuum. As the optical continuum emission from a BD, is by definition much weaker than for a CTTS say, it is necessary to bin or smooth the BD continuum in order to increase the spectro-astrometric accuracy. The error in the spectro-astrometric analysis depends strongly on the S/N and is given by $\sigma_{centroid} = \frac{seeing(mas)}{2.3548\sqrt{N_p}}$ where N_p is the number of detected photons. Once the continuum is mapped it is subtracted and the spatial profile of the pure line emission analysed. An approach first adopted by Whelan et al. (2005) was to bin the continuum and line emission in such a way that as to ensure that a comparable number of photons are sampled at each point. In other words the spectro-astrometric accuracy achieved in the line and continuum is comparable.

Here in the same way the continuum and [OI] λ 6300 line emission are smoothed so that the spectro-astrometric error in both regions are the same. The blue and red-shifted components to the [OI] λ 6300 line are found to be offset in opposing directions to an absolute distance of ~ 80 mas (see Figure 2a). Hence it is clear that the [OI] λ 6300 emission originates in a faint bipolar outflow driven by 2MASS1207. The $1-\sigma$ error in the measurements, of 18 mas is marked, by the dashed line in Figure 2a. Also as in Whelan et al. (2005) the presence or not of the [OI] λ 6300 night sky line was found to have no effect on the spectro-astrometric results.

Other lines of interest are [OI] λ 6363, H α and HeI λ 6678. The [OI] λ 6363 is ~ 3 times fainter than the [OI] λ 6300 line and hence is just below the detection limit in the raw spectrum. By smoothing the spectrum using a large gaussian filter (FWHM in dispersion direction = 0.35 \AA) the [OI] λ 6363 is revealed at a similar velocity to the [OI] λ 6300 line, although detailed kinematic data is lost. The primary origin of the H α and HeI λ 6678 lines is in the accretion flow. As this occurs on a very small scale we expect to measure no spectro-astrometric offset and as can be seen in Figure 2, none is found ruling out the presence of spectro-astrometric artifacts (Brannigan et al. 2006). The H α line is often seen in CTTS to have an outflow component. However at these scales any outflow component would certainly be extremely weak relative to the bulk of the H α emission and therefore again very difficult to detect. As the H α line is much brighter than any other part of the spectrum, and as any outflow signature would be found in the wings, the line and the continuum are smoothed in such a way that the error in the continuum and the line-wings are comparable. All that is seen is a tightening of the H α position (due to the higher S/N) as one moves towards the peak of the line (Figure 2b).




Fig. 2.— Offset velocity diagrams in the vicinity of the [OI]λ6300, Hα and He I lines. The line profiles are an average of three pixel rows in the spatial direction centred on the continuum. Note the continuum has not been subtracted. The green dashed lines delineate the $\pm 1 \sigma$ error envelope for the centroid position of the continuum. As explained in the text the line and continuum centroid errors are the same in all cases. In particular for Hα the spectrum is smoothed so that error in the centroid position of the wing emission is comparable to that for the continuum. The error decreases for the line peak region as is clear in (b). The red-shifted absorption feature in the Hα line is a signature of infall. The bipolar offset in the [OI]λ6300 reveals the presence of an outflow. No offsets are measured in the Hα and He I lines as expected ruling out the possibility of spectro-astrometric artifacts. Lastly the gaps in the position of the continuum blueward of the [OI]λ6300 and Hα lines, mark the position of night sky lines. The night sky lines have been removed from the line profiles and as found by Whelan et al. (2005) the presence or not of these sky lines has no effect on the spectro-astrometric results.

4. Scaling Down from T Tauri Jets

While it is accepted that accretion activity in BDs is simply a scaled down version of in-fall in CTTSs (Jayawardhana et al. 2003a), the presumption that BDs also drive scaled-down versions of T Tauri jets and outflows while credible, has yet to be generally confirmed. Here the evidence gathered so far supporting the continuation of the CTT paradigm for outflow activity, into the young BD mass regime, is summarised. Table 1 presents a comparison between the three BDs whose high resolution optical spectra have been analysed by us to date. LS-RCrA 1 probably has the strongest FELs of any BD (Fernández & Comerón 2001) and it is very likely that these FELs are excited in an jet. Yet direct imaging and a preliminary spectro-astrometric study failed to confirm such an origin for these features (Fernández & Comerón 2005; Whelan et al. 2006). In particular spectro-astrometry is difficult due to the faintness of this object which in turn is almost certainly caused by an edge-on disk (Barrado y Navascués et al. 2004). The relatively low radial velocities of the FELs of LS-RCrA 1 support this picture. From Table 1 ρ -Oph 102, 2MASS1207 and LS-RCrA 1 appear to launch outflows of comparable velocities. Also for 2MASS1207 and ρ -Oph 102 the [OI] λ 6300 line emitting region is offset by similar distances from the BD. Hence these three sub-stellar objects present a consistent picture of outflow activity in BDs, but how do their FEL regions (FELRs) compare to those of CTTSs?

The most notable features of the FELRs of CTTSs are their blue-shifted and often multi-component nature (Hirth et al. 1997; Davis et al. 2003). The lack of red-shifted FE is accounted for by the obscuring effect of the accretion disk. Only blue-shifted emission is observed for ρ -Oph 102 and the scale of the blue-shifted offset ($0''.085$) reported by Whelan et al. (2005) for ρ -Oph 102 would suggest a minimum (projected) disk radius of $0''.1$ (≥ 15 AU at the distance to the ρ -Ophiuchi cloud) in order to hide any red-shifted component. For 2MASS1207 the fact that both blue and red-shifted FE is detected is consistent with the near edge-on disk hypothesis.

Lastly Whelan et al. (2005) discuss the distance from a typical BD at which the critical density (n_{cr}) for forbidden emission is expected to be reached, if it is assumed that BD jets are similar in temperature, line excitation and opening angle to T Tauri jets. This distance is derived by considering how n_{cr} ($2 \times 10^6 \text{ cm}^{-3}$ for [OI] λ 6300) is expected to scale with mass flux through the jet. Using the predictions of Masciadri & Raga (2004), the distance for forbidden emission (and thus measured offsets) is estimated to be 3-10 times closer to a BD than a CTTS. The [OI] λ 6300 offsets measured for 2MASS1207 lie within this mass range. These offsets are maximised due to the probable edge-on disk geometry of 2MASS1207. In conclusion there is much evidence in existence to support the continuation T Tauri like outflow activity into the mass range of young BDs.

5. Conclusions

Our analysis of the high spatial resolution spectrum of 2MASS1207 clearly uncovers the existence of a bipolar, T-Tauri like outflow. This result is in agreement with the classification of 2MASS1207 as a strong accretor with forbidden emission and its proposed near edge-on disk geometry. As 2MASS1207 is only a $\sim 24M_{jup}$ object it now ranks as the least massive galactic object known to drive a jet. Given the age of the TW Hydrae Association it is also one of the oldest galactic objects with a resolved jet. The age of 2MASS1207 highlights that outflows can persist for a relatively long time, even when an object is sub-stellar, and the derived mass accretion rates are relatively low. Overall this discovery is not only intriguing but important as it emphasises the robustness of the outflow mechanism over an enormous range of masses (up to $10^{8-9}M_{\odot}$ in Active Galactic Nuclei) and it adds weight to the idea that at least, massive planets may also be outflow sources when they form.

The present work was supported in part by the European Community’s Marie Curie Actions - Human Resource and Mobility within the JETSET (Jet Simulations, Experiments and Theory) network under contract MRTN-CT-2004 005592 and by Science Foundation Ireland (contract no. 04/BRG/P02741). This work is based on observations that were taken at the European Southern Observatory, Chile (077.C-0771)

REFERENCES

- Bailey, J. A. 1998, *Proc. SPIE*, 3355, 932
- Barrado y Navascués, D., Mohanty, S., & Jayawardhana, R. 2004, *ApJ*, 604, 284
- Bourke, T. L., Crapsi, A., Myers, P. C., Evans, N. J., II, Wilner, D. J., Huard, T. L., Jørgensen, J. K., & Young, C. H. 2005, *ApJ*, 633, L129
- Brannigan, E., Takami, M., Chrysostomou, A., & Bailey, J. 2006, *MNRAS*, 367, 315
- Comerón, F., & Reipurth, B. 2006, *A&A*, 458, L21
- Chabrier, G., Baraffe, I., Allard, F., & Hauschildt, P. 2000, *ApJ*, 542, 464
- Chauvin, G., Lagrange, A.-M., Dumas, C., Zuckerman, B., Mouillet, D., Song, I., Beuzit, J.-L., & Lowrance, P. 2004, *A&A*, 425, L29
- Chauvin, G., Lagrange, A.-M., Dumas, C., Zuckerman, B., Mouillet, D., Song, I., Beuzit, J.-L., & Lowrance, P. 2005, *A&A*, 438, L25

- Davis, C. J., Whelan, E., Ray, T. P., & Chrysostomou, A. 2003, *A&A*, 397, 693
- Dekker, H., D’Odorico, S., Kaufer, A., Delabre, B., & Kotzlowski, H. 2000, *Proc. SPIE*, 4008, 534
- Fernández, M., & Comerón, F. 2001, *A&A*, 380, 264
- Fernández, M., & Comerón, F. 2005, *A&A*, 440, 1119
- Gizis, J. E. 2002, *ApJ*, 575, 484
- Hartigan, P., Morse, J. A., & Raymond, J. 1994, *ApJ*, 436, 125
- Hirth, G. A., Mundt, R., & Solf, J. 1997, *A&AS*, 126, 437
- Hartmann, L., Hewett, R., & Calvet, N. 1994, *ApJ*, 426, 669
- Jayawardhana, R., Mohanty, S., & Basri, G. 2003, *ApJ*, 592, 282
- Jayawardhana, R., Ardila, D. R., Stelzer, B., & Haisch, K. E., Jr. 2003, *AJ*, 126, 1515
- Konigl, A., & Pudritz, R. E. 2000, *Protostars and Planets IV*, 759
- Masciadri, E., & Raga, A. C. 2004, *ApJ*, 615, 850
- Mohanty, S., Jayawardhana, R., & Barrado y Navascués, D. 2003, *ApJ*, 593, L109
- Mohanty, S., Jayawardhana, R., & Basri, G. 2005, *ApJ*, 626, 498
- Mohanty, S., Jayawardhana, R., Huelamo, N., & Mamajek, E. 2006, *ApJ* in press, astro-ph/0610550
- Muzerolle, J., Hillenbrand, L., Calvet, N., Briceño, C., & Hartmann, L. 2003, *ApJ*, 592, 266
- Natta, A., & Testi, L. 2001, *A&A*, 376, L22
- Natta, A., Testi, L., Muzerolle, J., Randich, S., Comerón, F., & Persi, P. 2004, *A&A*, 424, 603
- Scholz, A., Jayawardhana, R., & Brandeker, A. 2005, *ApJ*, 629, L41
- Scholz, A., & Jayawardhana, R. 2006, *ApJ*, 638, 1056
- Takami, M., Bailey, J., Gledhill, T. M., Chrysostomou, A., & Hough, J. H. 2001, *MNRAS*, 323, 177

Whelan, E. T., Ray, T. P., & Davis, C. J. 2004, *A&A*, 417, 247

Whelan, E. T., Ray, T. P., Bacciotti, F., Natta, A., Testi, L., & Randich, S. 2005, *Nature*, 435, 652

Whelan, E. T., Ray, T. P., Bacciotti, F., & Jayawardhana, R. 2006, *New Astronomy Review*, 49, 582

Source	2MASS1207	ρ -Oph 102	LS-RCrA 1
Mass (M_{jup})	24	60	35
\dot{M}_{acc} ($M_{\odot}yr^{-1}$)	10^{-11}	1.25×10^{-9}	10^{-9}
[OI] λ 6300(kms $^{-1}$, ")	+4/-8,+0.08/-0.08	-40, 0.085	-9, –
[SII] λ 6731(kms $^{-1}$, ")	–,–	-45, 0.085	-15,–
[NII] λ 6583	–,–	-45,–	-19,–

Table 1: Table comparing the mass (in Jupiter masses), mass accretion rates and typical FEL velocities of 2MASS1207, ρ -Oph 102 and LS RCrA 1. The accretion rate for ρ -Oph 102 was derived using the H α emission line and is taken from Natta et al. (2004). For 2MASS1207 and LS RCrA 1 Mohanty et al. (2005) used the CaII λ 8662 line fluxes to infer the mass accretion rates. The mass estimate for LS-RCr A1 is taken from Barrado y Navascués et al. (2004).