

GloBAM Data Management Plan

Peter Desmet

Cecilia Nilsson

Adriaan Dokter

2019-09-09

Contents

Colophon	2
1 Introduction	3
1.1 Research purpose	3
1.2 Data manager	3
1.3 How this DMP is maintained	3
Source data	3
2 European radar data (pvol)	3
2.1 Source	3
2.2 License	4
2.3 Geographical scope	4
2.4 Temporal scope	5
2.5 Format	5
2.6 File name conventions	5
2.7 Quality assurance	6
2.8 Storage & backup	6
3 US radar data (pvol)	6
3.1 Source	6
3.2 License	6
3.3 Geographical scope	6
3.4 Temporal scope	8
3.5 Format	8
3.6 File name conventions	8
3.7 Quality assurance	8
3.8 Storage & backup	8
4 Other external data	9
4.1 Wind/weather data	9
4.2 Land use	9
4.3 Artificial light	9
4.4 Wind energy installations	9
Produced data	10
5 Processing pipeline	10
5.1 European pvol	10
5.2 European pvol to vp	10
5.3 vp archiving	10

5.4	vp to vpts	10
6	Vertical profiles of aerial migrants data (vp)	11
6.1	Source	11
6.2	License	11
6.3	Geographical scope	11
6.4	Temporal scope	14
6.5	Format	14
6.6	File name conventions	14
6.7	Quality assurance	14
6.8	Storage & backup	14
7	Time series of vertical profiles data (vpts)	14
7.1	Source	14
7.2	License	14
7.3	Geographical scope	14
7.4	Temporal scope	14
7.5	Format	15
7.6	File name conventions	15
7.7	Quality assurance	15
7.8	Storage & backup	15
	Publishing & preservation	15
8	Open data	15
8.1	Data	15
8.2	Repository	15
8.3	License	15
8.4	Metadata	15
8.5	Standards	16
8.6	Frequency	16

Colophon

Distributions of this data management plan:

- Latest version (website): <https://enram.github.io/globam-dmp/>
- Latest version (pdf): <https://enram.github.io/globam-dmp/globam-dmp.pdf>
- Latest version (source files): <https://github.com/enram/globam-dmp>
- Periodically archived version (pdf): *to be deposited on Zenodo.org*

Suggested citation for the latest version:

Desmet P, Nilsson C, Dokter A (2019) GloBAM Data Management Plan. Version 2019-09-09.
<https://enram.github.io/globam-dmp/>

All versions are licensed under a Creative Commons Attribution License (CC BY 4.0).

1 Introduction

1.1 Research purpose

Migratory animals play significant roles in shaping ecosystems through a variety of transport and trophic effects that also represent services and disservices to human infrastructure, agriculture and welfare. Their aerial and terrestrial habitats have changed dramatically over the past decades and are expected to change further, particularly due to rapid climate change, increased urbanization, wind energy installations, and habitat fragmentation.

Within *GloBAM: Towards monitoring, understanding and forecasting global biomass flows of aerial migrants*, we aim to use weather radar data to quantify the biomass flows of aerial migrants (birds, insects and bats) from regional to continental scales across Europe and North America, over time-scales from days to years. We are particularly interested in identifying the drivers of migrant movements and abundances and will relate the timing and intensity of movements to a suite of atmospheric, climatic and landscape variables, exploring the implications for aerial migrants in a changing world.

For more information on GloBAM, see the project website.

1.2 Data manager

Peter Desmet is responsible for data management and DMP maintenance in GloBAM. He is also leading *WP1 - Data infrastructure*. Peter works as open data coordinator for the Open science lab for biodiversity at the Research Institute for Nature and Forest (INBO). He and his team have extensive expertise and experience with data management and open data publication meeting FAIR principles.

1.3 How this DMP is maintained

1. This DMP is maintained and versioned on GitHub at <https://github.com/enram/globam-dmp/>.
2. Each chapter is an R Markdown file (`Rmd`) in the `src` directory of that GitHub repository. You can access it directly by clicking the pencil icon in the top navigation of the website version of this DMP.
3. Changes to the R Markdown files can be made by contributors to the GitHub repository or suggested by anyone as pull requests. Textual changes can be done directly on GitHub, code changes are better tested in RStudio first.
4. Accepted changes (i.e. changes to the `master` branch) will trigger an automatic build procedure that will generate a new version of the DMP using the R package bookdown. The date of the build is used as the version number.

Source data

2 European radar data (pvol)

This chapter describes polar volume (`pvol`) data from European weather radars. It is one the main data sources for GloBAM.

2.1 Source

European `pvol` data are collected by weather radars from **national weather services** in Europe. They have different approaches in processing, archiving and providing access to `pvol` data (from restricted access to

open data). Even though some GloBAM partners (e.g. the University of Amsterdam) have archived subsets of these data for research use, the scattered scope and access to European `pvol` data is a serious barrier to GloBAM, which ideally requires access to all European `pvol` data using the same protocol.

The best source for consolidated `pvol` data is therefore **BALTRAD**, managed by SMHI (contact person: Günther Haase). `pvol` data are archived there, but access to that server is restricted to SMHI. See 5.1 for details on how `pvol` data are submitted to this archive.

The rest of this chapter discusses the specifics of the BALTRAD archive.

2.2 License

European radar data exchange is coordinated by the *European Operational Program for Exchange of Weather Radar Information (EUMETNET/OPERA)*. GloBAM has access to these data via the *EIG EUMETNET license agreement for Research use of OPERA data*, which grants a specific list of people (involved in ENRAM and GloBAM) access to radar data from selected countries under the following conditions (excerpt):

The Licensor grants the Licensees a non-Exclusive license to use intellectual property belonging to the EUMETNET members as defined below [...]

The OPERA Members Data will be provided for use within the ENRAM Project for the purpose of extracting animal migration information for scientific research.

The Grant of this License does not permit use of the OPERA Members Data licensed to be used for commercial purposes or exploitation for profit.

The license agreement allows us to access European `pvol` data for the sole purpose of extracting animal migration information for scientific research, which is what GloBAM intends to do. Storing, sharing or other use of these data is **restricted** and requires prior written consent of the Licensor.

2.3 Geographical scope

OPERA manages a list of radars, their unique codes (e.g. `odimcode`) and associated metadata. The list currently contains 232 radars. This **OPERA database** is available as an interactive map derived from a `json` file containing the metadata. To keep track of changes, we archive this `json` on GitHub every time we update the DMP.

The license agreement allows access to data from 19 countries:

##	country	iso_code	radars	operational
##	Austria	AT	5	5
##	Belgium	BE	3	3
##	Croatia	HR	5	2
##	Czechia	CZ	2	2
##	Denmark	DK	5	5
##	Estonia	EE	2	2
##	Finland	FI	10	10
##	France	FR	31	25
##	Germany	DE	20	20
##	Netherlands	NL	3	2
##	Norway	NO	11	11
##	Poland	PL	8	8
##	Portugal	PT	4	3
##	Slovakia	SK	4	4
##	Slovenia	SI	2	2
##	Spain	ES	15	15

##	Sweden	SE	12	12
##	Switzerland	CH	5	5
##	United Kingdom	GB	16	16
##	Total	-	163	152

The BALTRAD archive does not contain `pvol` data for all of these radars/countries however (see 5.1). Its precise geographical scope is currently **unknown**, but being gathered.

2.4 Temporal scope

The temporal scope of the BALTRAD archive is currently **unknown**, but being gathered. GloBAM needs a **2-3 year archive** of European `pvol` and derived `vp` data to tackle its research questions. By starting from a `pvol` archive, the generation of `vp` data can be reproduced if need be, e.g. to make sure adequate `vol2bird` settings/versions are used.

The `pvol` archive at BALTRAD is likely to be the best source for this archive, but its scope and quality need to be assessed before we can proceed, see 2.7.

2.5 Format

The delivery of European `pvol` data is defined in the license agreement:

The Licensor will make the OPERA Members Data available to the Licensees in accordance to the following data description and technical specifications:

- Single-site polar volumes containing reflectivity, optionally also Doppler velocity, uncorrected reflectivity and dual-pol parameters. Data model ODIM as described in OPERA pages of the EUMETNET website (<http://www.eumetnet.eu>)
- Update frequency – 15 minutes
- Issue time – up to 120 minutes after data time
- Format: HDF5
- Delivery method: FTP via Internet using an ad hoc server. A username and password will be created for ENRAM.
- Availability of OPERA Members Data will be on the basis of reasonable endeavour
- Support: OPERA documents are available from the EUMETNET website’s OPERA pages (<http://www.eumetnet.eu>). OPERA Program Manager will manage technical questions related to the OPERA Products but may address more complicated issues to a team member who may charge a fee.

Data is thus provided for every 15 minutes as `hdf5` in the OPERA ODIM_h5 format. These files can be read in `bioRad` with `read_pvolfile()`.

2.6 File name conventions

File names are generally of the format `bejab_pvol_20190620T055500Z_0x9.h5`, where:

- `be`: two-letter code (ISO 3166-1 alpha-2)
- `jab`: three-letter code (last 3 letters from the ODIM code, see 2.3)
- `_pvol_`: indication that this is a `pvol` data file
- `20190620`: date in `yyyymmdd` format
- `T055500Z`: time in `hhmmss` format + timezone (Z for UTC time)
- `_0x9`: suffix (to be ignored)

2.7 Quality assurance

The scope and quality of the BALTRAD archive need to be assessed. Tasks for this are listed here and include:

1. Get a file listing for `pvol` archive
2. Select subsets: first 2 days of data per radar/month for 2016, 2017, 2018 (72 days): either `pvol` or pre-merged scans
3. Transfer subsets to accessible FTP server
4. Merge to `pvol` (test 1)
5. Process with `vol2bird` to `vp` (test 2)
6. Store output `vp` files
7. Visual control of `vp` (test 3)

2.8 Storage & backup

European `pvol` data are stored on the BALTRAD archive, managed and financed by SMHI. The archive is a restricted FTP server, only accessible to SMHI (see 5.1).

3 US radar data (`pvol`)

This chapter describes polar volume (`pvol`) data from weather radars in the United States. It is one the main data sources for GloBAM.

3.1 Source

US `pvol` data are collected by the National Oceanic and Atmospheric Administration (NOAA), which operates a network of approximately 160 Next Generation Weather Radar (NEXRAD), WSR-88D sites in the US and associated territories. Archived polar volumes (`pvol`s) (called “level II data” in the US) are stored on various services:

1. Amazon S3, see NEXRAD on AWS. Archived data are hosted in the `noaa-nexrad-level2` Amazon S3 bucket in the `us-east-1` AWS region. These data are made available 10-20 minutes lag relative to real-time. There is also a real-time data stream, for which scans have not been merged yet into polar volumes.
2. Google Cloud.

3.2 License

There are no restrictions on the use of these data.

3.3 Geographical scope

The NEXRAD network covers continental US, Alaska, Hawaii, Guam, South Korea and Puerto Rico. Sites in use vary slightly over time, see here for more information, maps here.

VCP12 Coverage

- 4,000 ft above ground level*
- 6,000 ft above ground level*
- 10,000 ft above ground level*

* Bottom of beam height (assuming Standard Atmospheric Refraction).
Terrain blockage indicated where 50% or more of beam blocked.

0 125 250 500 750 Miles

NOAA
NATIONAL OCEANIC AND ATMOSPHERIC ADMINISTRATION
U.S. DEPARTMENT OF COMMERCE

7

3.4 Temporal scope

The archive spans from June 1991 to present. However, during this time significant changes have been made. A resolution increase was implemented in 2008-2009 (so-called superresolution data). Dual polarization capabilities were added in 2010-2013.

3.5 Format

See this page for available decoding tools of the radar format. US pvol data can be read in bioRad with `read_pvolfile()`.

3.6 File name conventions

From the AWS open data project documentation for archive data:

Each volume scan file of archival data is available as an object in Amazon S3. The basic data format is:

`/<Year>/<Month>/<Day>/<NEXRAD Station>/<filename>`

Where:

- `<Year>` is the year the data was collected
- `<Month>` is the month of the year the data was collected
- `<Day>` is the day of the month the data was collected
- `<NEXRAD Station>` is the NEXRAD ground station (map of ground stations)
- `<filename>` is the name of the file containing the data. These are compressed files (compressed with gzip). The file name has more precise timestamp information.

All files in the archive use the same compressed format (.gz). The data file names are, for example, `KAKQ20010101_080138.gz`. The file naming convention is:

`GGGGYYYYMMDD_TTTTTT`

Where:

- `GGGG` = Ground station ID (map of ground stations)
- `YYYY` = year
- `MM` = month
- `DD` = day
- `TTTTTT` = time when data started to be collected (GMT)

Note that the 2015 files have an additional field on the file name. It adds “_V06” to the end of the file name. An example is `KABX20150303_001050_V06.gz`.

3.7 Quality assurance

Unknown.

3.8 Storage & backup

US pvol data are stored on AWS and other cloud services, see 3.1.

4 Other external data

This chapter describes other external data sources required by GloBAM.

4.1 Wind/weather data

4.1.1 Users

Needed for at least WP3, WP4, WP5 and maybe WP2. The type of weather data (wind, precipitation) will differ between analyses.

4.1.2 Sources

- World: to be defined
- US: NCEP/NCAR Reanalysis data from NOAA has been used before.
- Europe: to be defined, see <https://github.com/enram/globam-dmp/issues/21>

4.2 Land use

To be defined, see <https://github.com/enram/globam-dmp/issues/25>

4.3 Artificial light

4.3.1 Users

Needed for WP3.

4.3.2 Sources

Visible Infrared Imaging Radiometer Suite (VIIRS) Day/Night Band provided by NOAA. See that page for a description, filenaming conventions, data types/formats and providing credit.

We may also use the new Black Marble product from NASA (not yet released).

4.4 Wind energy installations

4.4.1 Users

Needed for WP4 (Judy Shamoun-Baranes).

4.4.2 Sources

Depends on the exact research questions and what is available. Will be updated in DMP once known.

Produced data

5 Processing pipeline

This chapter describes the processing pipeline from `pvol` over `vp` to `vpts` data.

5.1 European `pvol`

BALTRAD collects `pvol` data for Europe. For `pvol` data to be available in BALTRAD, several conditions need to be met:

1. A country should send both reflectivity and radial velocity data to the OPERA data centre, which is called ODYSSEY. While many countries are sending reflectivity data, radial velocities are still unavailable for many countries.
2. ODYSSEY should forward these data to the BALTRAD datahub. It is standard policy to do so, but in practice data is not yet forwarded correctly for some radars/countries.
3. Currently BALTRAD and ODYSSEY store data at a 15 minute interval, higher resolution data is not yet available.

5.2 European `pvol` to `vp`

A server at BALTRAD processes any incoming `pvol` data with `vol2bird`. This pipeline is maintained by Günther Haase (SMHI).

The resulting `vp` files are stored for 2 days on a private FTP server to which we have access. The used `vol2bird` version is stored in the metadata of the `vp` files and can differ over time. The latest files are created with `vol2bird` 0.3.20 (last tested April 11, 2019).

5.3 `vp` archiving

European `vp` data are archived daily by a pipeline running on Amazon Web Services (AWS) which transfers `vp` files from the BALTRAD FTP server to a public S3 bucket. This pipeline is maintained by the Open science lab for biodiversity and its code and documentation are available in this repository. Note that any issues in the pipeline that are not resolved within 2 days (the time `vp` data are kept on the FTP server) **can result in data loss**.

The pipeline also updates a coverage file and bundles `vp` files in monthly zips per radar (e.g. `bejab201904.zip`). A public website allows easier file navigation of the S3 bucket.

United States `vp` data are currently not archived. See <https://github.com/enram/globam-dmp/issues/19>

5.4 `vp` to `vpts`

Not yet defined, but likely to be part of the ENRAM pipeline (see 5.3).

6 Vertical profiles of aerial migrants data (vp)

This chapter describes vertical profiles of aerial migrants/biological signals (**vp**) data for Europe and the United States. Vertical profiles contain the speed, direction and density of aerial migrants at different altitudes for a specific radar location and time, and form the source data for GloBAM specifically and aeroecological research in general.

6.1 Source

European vp data are generated daily from **pvo1** data on the BALTRAD server (see 5.2) and then transferred to the public ENRAM data repository (see ??). **United States vp** data are being generated for the 25 year **pvo1** archive, but not yet publicly archived.

vp files can be downloaded from the ENRAM data repository individually, as monthly zips, or automatically in bioRad with `download_vpfiles()`.

6.2 License

Data in the ENRAM data repository are available as **open data** under a Creative Commons Zero waiver.

For the **European data** it is recommended to acknowledge EUMETNET/OPERA in publications resulting from the use of these data as follows:

We acknowledge the European Operational Program for Exchange of Weather Radar Information (EUMETNET/OPERA) for providing access to European radar data, facilitated through a research-only license agreement between EUMETNET/OPERA members and ENRAM.

6.3 Geographical scope

The coverage of the ENRAM data repository is recorded daily in `coverage.csv` and summarized here:

##	radar	2016	2017	2018	2019	Total
##	bejab	21	NA	194	235	450
##	bewid	19	12	205	233	469
##	bezav	21	NA	NA	NA	21
##	bgvar	21	NA	NA	NA	21
##	ctcdv	21	NA	NA	NA	21
##	ctpda	21	NA	NA	NA	21
##	czbrd	21	NA	286	234	541
##	czska	21	NA	286	236	543
##	deasb	NA	NA	221	93	314
##	deboo	31	17	305	93	446
##	dedrs	31	21	304	93	449
##	deeis	31	21	304	93	449
##	deemd	NA	20	49	NA	69
##	deess	31	21	304	93	449
##	defbg	31	NA	NA	NA	31
##	defld	31	21	303	92	447
##	deflg	NA	21	49	NA	70
##	dehnr	31	21	304	93	449
##	deisn	31	NA	NA	NA	31
##	demem	31	21	301	93	446
##	deneu	31	21	301	93	446

##	denhb	31	21	300	90	442
##	deoft	31	21	304	93	449
##	depro	31	21	302	93	447
##	deros	25	21	294	93	433
##	desna	NA	21	303	93	417
##	detur	29	21	303	93	446
##	deumd	31	21	304	93	449
##	dkbor	NA	34	85	238	357
##	dkrom	NA	38	85	237	360
##	dksin	NA	38	85	238	361
##	dkste	NA	38	85	237	360
##	dkvir	NA	27	85	238	350
##	eehar	NA	20	35	1	56
##	eesur	NA	20	35	NA	55
##	esalm	NA	38	287	180	505
##	esbad	NA	38	302	180	520
##	esbar	NA	38	303	181	522
##	escor	NA	38	287	180	505
##	eslid	NA	38	303	180	521
##	eslpa	NA	NA	14	5	19
##	esmad	NA	38	301	180	519
##	esmal	NA	38	303	181	522
##	esmur	NA	38	302	180	520
##	espma	NA	38	303	179	520
##	essan	NA	38	302	180	520
##	essev	NA	36	300	177	513
##	essse	NA	38	303	174	515
##	esval	NA	38	294	180	512
##	eszar	NA	38	303	176	517
##	fianj	22	154	38	NA	214
##	fiika	22	153	39	NA	214
##	fikes	22	151	38	NA	211
##	fikor	22	146	38	NA	206
##	fikuo	22	156	38	NA	216
##	filuo	22	153	38	NA	213
##	fipet	22	99	NA	NA	121
##	fiuta	22	157	38	NA	217
##	fivan	22	157	39	NA	218
##	fivim	22	132	34	NA	188
##	frabb	21	38	320	221	600
##	fraja	NA	NA	NA	6	6
##	frale	21	26	321	227	595
##	frave	21	NA	283	238	542
##	frbla	21	38	321	229	609
##	frbol	21	38	279	185	523
##	frbor	21	36	321	238	616
##	frbou	21	38	173	235	467
##	frcae	21	38	320	235	614
##	frche	21	38	317	228	604
##	frcol	21	38	312	237	608
##	frgre	21	38	320	233	612
##	frlep	21	38	320	225	604
##	frmcl	21	38	304	237	600
##	frmom	21	38	320	238	617

##	frmtc	21	35	302	218	576
##	frnan	21	38	317	237	613
##	frnim	21	38	320	238	617
##	frniz	21	NA	281	238	540
##	fropo	21	38	241	59	359
##	frpla	21	38	315	238	612
##	frtou	21	37	305	238	601
##	frtra	21	38	320	238	617
##	frtre	6	38	316	238	598
##	frtro	21	NA	282	237	540
##	hrbil	NA	38	35	NA	73
##	hrosi	NA	38	35	NA	73
##	nldbl	22	4	NA	NA	26
##	nldhl	21	125	295	237	678
##	nlhrw	NA	21	294	237	552
##	plbrz	21	3	261	236	521
##	plgda	21	3	261	235	520
##	plleg	21	3	259	236	519
##	plpas	NA	3	259	236	498
##	plpoz	21	3	260	236	520
##	plram	21	3	260	236	520
##	plrze	21	3	222	237	483
##	plswi	21	3	260	236	520
##	ptfar	119	NA	NA	52	171
##	ptlis	NA	NA	NA	31	31
##	ptliz	122	NA	NA	NA	122
##	ptprt	113	NA	NA	109	222
##	seang	21	137	310	194	662
##	searl	21	153	318	123	615
##	sease	21	16	NA	NA	37
##	seatv	NA	NA	NA	73	73
##	sebaa	NA	NA	NA	20	20
##	sehem	NA	43	314	227	584
##	sehud	21	NA	NA	NA	21
##	sehuv	NA	36	295	228	559
##	sekaa	NA	NA	NA	132	132
##	sekir	21	144	318	238	721
##	sekkrr	21	157	171	NA	349
##	selek	21	118	208	222	569
##	sella	NA	NA	NA	131	131
##	selul	21	148	183	NA	352
##	seoer	NA	36	307	228	571
##	seosd	NA	142	311	225	678
##	seosu	13	NA	NA	NA	13
##	seovi	21	59	NA	NA	80
##	sevar	21	NA	NA	NA	21
##	sevox	NA	133	312	223	668
##	sevil	21	153	318	29	521
##	silis	21	35	304	227	587
##	sipas	21	35	311	196	563
##	skjav	NA	38	301	237	576
##	skkoj	NA	38	301	230	569
##	skkub	NA	NA	280	236	516
##	sklaz	NA	NA	280	238	518

6.4 Temporal scope

Data transfer to the ENRAM data repository become more or less operational in March 2018 (with a gap in July 2018). Data from 2016 were uploaded for the European flyway study Nilsson et al. 2019.

6.5 Format

A `vp` file is generated for each originating `pvo1` file and thus has the same granularity. `vp` data are stored as `hdf5` files in the ODIM bird profile format specification. These files can be read in `bioRad` with `read_vpfiles()`.

6.6 File name conventions

File names follow the same format as `pvo1` files (see 2.5), but with `_vp_` (e.g. `bejab_vp_20190620T055500Z_0x9.h5`). The directory structure of the ENRAM data repository is documented here.

6.7 Quality assurance

To be defined.

6.8 Storage & backup

European `vp` data are stored on AWS (see 5.3). Each file transferred to the S3 bucket is also backed up in a second S3 bucket.

7 Time series of vertical profiles data (`vpts`)

This chapter describes time series of vertical profiles (`vpts`). `vpts` are `vp` data bundled in time series (without data loss) and are a more convenient way for downloading and reading this type of data.

7.1 Source

These files are not yet generated, but will be stored in the ENRAM data repository.

7.2 License

To be defined, but likely as **open data** under a Creative Commons Zero waiver.

7.3 Geographical scope

To be defined, but likely the same as the `vp` data.

7.4 Temporal scope

To be defined, but likely the same as the `vp` data.

7.5 Format

Not yet defined, but likely text format (`txt`, `json`, `csv`) and tabular. See this issue.

7.6 File name conventions

To be defined.

7.7 Quality assurance

To be defined.

7.8 Storage & backup

To be defined, but likely the same as the `vp` data.

Publishing & preservation

8 Open data

This chapter describes how we will publish data of long term value that GloBAM will use and generate.

8.1 Data

The data of long term value that GloBAM will generate are the `vp/vpts` data for Europe and the United States (see the respective chapters).

8.2 Repository

As there is no specific research repository for `vp/vpts` data, we will publish these data on the general research repository Zenodo. This has already been done for Nilsson et al. 2018 and will likely look similar for data published for GloBAM. Data packages published on Zenodo are assigned a Digital Object Identifier (DOI) for easier referencing (both a generic one and one for each updated version).

8.3 License

Data will be published under a Creative Commons Zero waiver.

8.4 Metadata

Data will be documented with metadata describing the contributors, provenance, resolution, temporal and spatial coverage, and how to use these with open source software developed or contributed to by GloBAM.

8.5 Standards

Data will be kept in their source format, i.e. the ODIM bird profile format specification (see 6.5).

8.6 Frequency

Data will be published on Zenodo every two years. Closer to real-time data are already publicly available in the ENRAM data repository.