

Journal of Oral Implantology

Fifteen years of platelet rich fibrin (PRF) in dentistry and oromaxillofacial surgery: How high is the level of scientific evidence?

--Manuscript Draft--

Manuscript Number:	aaid-joi-D-17-00179R3
Full Title:	Fifteen years of platelet rich fibrin (PRF) in dentistry and oromaxillofacial surgery: How high is the level of scientific evidence?
Short Title:	Level of scientific evidence in clinical PRF Research
Article Type:	Review Paper
Keywords:	platelet rich fibrin; Scientific evidence; Dentistry; maxillofacial surgery; LSCC; I-PRF, A-PRF
Corresponding Author:	Shahram Ghanaati, MD DMD PhD Goethe University Frankfurt Frankfurt, Hessen GERMANY
Corresponding Author Secondary Information:	
Corresponding Author's Institution:	Goethe University Frankfurt
Corresponding Author's Secondary Institution:	
First Author:	Shahram Ghanaati, MD DMD PhD
First Author Secondary Information:	
Order of Authors:	Shahram Ghanaati, MD DMD PhD Carlos Herrera-Vizcaino, DMD Sarah Al-Maawi, DMD Jonas Lorenz, DMD Richard J. Miron, DDS Katja Nelson, DMD, PhD Frank Schwarz, DMD, PhD Joseph Choukroun, MD Robert Sader, MD, DMD, PhD
Order of Authors Secondary Information:	
Abstract:	<p>Abstract:</p> <p>Platelet-rich fibrin is a blood concentrate system used for soft tissue and bone tissue regeneration. In the last decade, platelet rich fibrin (PRF) has been widely used in different indication fields, particularly in oral and maxillofacial surgery. This review is aimed to investigate the level of scientific evidence of published articles related to the use of PRF for bone and soft tissue regeneration in dentistry and maxillofacial surgery. An electronic literature research using the biomedical search engine "National Library of Medicine" (PubMed-MEDLINE) was performed in May 2017. A total of 392 articles were found, 72 of which were classified for each indication field. When comparing PRF with biomaterials vs biomaterial alone in sinus lift (5 studies; IIa), no statistically significant differences were detected. Socket preservation and ridge augmentation using PRF significantly enhanced new bone formation compared to healing without PRF (seven studies Ib, IIa, IIb). Reepithelialization and bone regeneration was achieved in 96 of 101 patients diagnosed with medication-related osteonecrosis of the jaw (5 studies, III). In periodontology, PRF alone (six studies; Ib, IIa, IIb) or its combination with biomaterials (six studies; Ib, IIa, IIb) significantly improved the pocket depth and attachment loss compared to a treatment without PRF. Over 70% of the</p>

	patients were part of studies with a high level of scientific evidence (randomized and controlled prospective studies). This published evidence, with a high scientific level, showed that PRF (38 articles) is a beneficial tool that significantly improves bone and soft tissue regeneration. However, the clinical community requires a standardization of PRF protocols to further examine the benefit of PRF in bone and soft tissue regeneration in reproducible studies, with a higher scientific level of evidence.
--	--

Fifteen years of platelet-rich fibrin (PRF) in dentistry and oromaxillofacial surgery: How high is the level of scientific evidence?

Abstract:

Platelet-rich fibrin is a blood concentrate system used for soft tissue and bone tissue regeneration. In the last decade, platelet rich fibrin (PRF) has been widely used in different indication fields, particularly in oral and maxillofacial surgery. This review is aimed to investigate the level of scientific evidence of published articles related to the use of PRF for bone and soft tissue regeneration in dentistry and maxillofacial surgery. An electronic literature research using the biomedical search engine “National Library of Medicine” (PubMed-MEDLINE) was performed in May 2017. A total of 392 articles were found, 72 of which were classified for each indication field. When comparing PRF with biomaterials vs biomaterial alone in sinus lift (5 studies; IIa), no statistically significant differences were detected. Socket preservation and ridge augmentation using PRF significantly enhanced new bone formation compared to healing without PRF (seven studies Ib, IIa, IIb). Reepithelialization and bone regeneration were achieved in 96 of 101 patients diagnosed with medication-related osteonecrosis of the jaw (5 studies, III). In periodontology, PRF alone (six studies; Ib, IIa, IIb) or its combination with biomaterials (six studies; Ib, IIa, IIb) significantly improved the pocket depth and attachment loss compared to a treatment without PRF. Over 70% of the patients were part of studies with a high level of scientific evidence (randomized and controlled prospective studies). This published evidence, with a high scientific level, showed that PRF (38 articles) is a beneficial tool that significantly improves bone and soft tissue regeneration. However, the clinical community requires a standardization of PRF protocols to further examine the benefit of PRF in bone and soft tissue regeneration in reproducible studies, with a higher scientific level of evidence.

Keywords: Platelet rich fibrin · Scientific evidence · Dentistry · Maxillofacial surgery · LSCC · I-PRF

Formatted: Font color: Auto

1 Introduction:

2 The development and application of biomaterials has increased over the past years. In dentistry and
3 maxillofacial surgery, biomaterials of autologous, allogenic and ~~xenogeneic~~xenogeneic sources are
4 widely used¹⁻⁴. There are advantages and disadvantages that can be associated with each of them⁵.

5 Surgical techniques performed to obtained autografts can be difficult and associated with higher
6 morbidity. Despite the difficulties, ~~of some surgical techniques in obtaining autografts and the~~
7 ~~accompanying patient morbidity and risks~~, they are considered to be the gold-standard method in
8 bone and soft tissue regeneration, particularly because of their autologous regenerative capacity⁶.

9 However, alternative autologous techniques introduced platelet concentrates (PC), autologous blood
10 derivatives characterized by the ease of the technique used to obtain them⁷. The history of PC can be

11 traced back to 1940. Young and Medawar described that they successfully reunited peripheral nerves
12 in animals, sealing them in blood plasma. Based on these findings, Helena Matras developed the
13 fibrin sealant technique during experiments on rat skin and later through clinical applications in
14 maxillofacial surgery in 1982⁷. This fibrin sealant product consisted of a mixture of fibrinogen and
15 thrombin, which formed a clot that was used for wound covering⁸. The natural evolution turned fibrin
16 sealants into PC when platelets were added to the formula⁹. The first PC protocols were described by

17 the so-called autologous platelet-derived wound healing factors (PDWHF) and were published in
18 1986 by Knighton et al. and in 1998, also by Marx, with the name platelet-rich plasma (PRP),
19 emphasizing platelets and growth factor content¹⁰⁻¹². This PC system requires the addition of
20 anticoagulants during its processing procedure and aims to concentrate platelets but eliminate blood
21 derived leukocytes. In 2001, as an alternative to PRP with numerous advantages, Choukroun
22 developed platelet-rich fibrin (PRF), a second generation of PCs that does not require the addition of
23 anticoagulants¹³.

24 PRF is a one-step standardized method of obtaining PCs. Through centrifugation of peripheral blood,
25 physiologic clot formation and fractioning are induced without the requirement of additives, such as

Field Code Changed

Field Code Changed

Field Code Changed

Field Code Changed

Field Code Changed

Field Code Changed

Field Code Changed

Field Code Changed

Field Code Changed

1 anticoagulants, thus, PRF is the only PC system that is fully autologous¹⁴. Using specific tubes with
2 a glass surface initiates the coagulation cascade and activates platelets during centrifugation. The
3 resulting PRF consists of a fibrin scaffold that contains platelets, leukocytes and plasma proteins.
4 After centrifugation, the resulted 3D matrix of the PRF clot serves as a reservoir of growth factors¹⁵.
5 Bioactive molecules (growth factor, cytokines, fibrinogen, fibronectin, thrombospondin, adhesive
6 proteins, coagulation factor) are primarily released from the alpha and dense platelet's granules¹⁶.
7 Additionally, cell interaction between activated platelets and the included leukocytes and their
8 subfamilies, such as neutrophils, seems to substantially increase the degranulation of inflammatory
9 cytokines (IL-1 β , IL-8) and chemokine (MCP-1)¹⁷.

Field Code Changed

Field Code Changed

Field Code Changed

Field Code Changed

10 The initial protocol to obtain PRF described the need of the application of high relative centrifugal
11 forces (RCF)^{18,19}. As a first attempt to understand PRF, our group reduced RCF through a further
12 histological analysis of the PRF clot and found that reducing the RCF increases the number platelets
13 and leukocytes, with a balanced distribution of cells within the matrix²⁰. In that manner, a systematic
14 analysis (gradual reduction approach) of the influence of RCF on the PRF-based matrices was
15 conducted and ~~showed-found~~ that by further reducing the RCF, an even higher increase of cell
16 concentration is obtained, as well as the release of growth factors²¹.

Field Code Changed

Field Code Changed

Field Code Changed

17 Basic science and translational research investigated the influence of PRF in soft and bone
18 regeneration in vitro. The behavior of osteoblasts treated with PRF in vitro ~~showed-found~~ a
19 significantly improved activity compared to the non-treated osteoblasts²². In addition, a significantly
20 higher migration rate, collagen formation and growth factor release activity ~~were-was~~ observed in
21 gingival fibroblasts treated with PRF compared to fibroblasts treated with PRP or without PRF
22 treatment in vitro²³.

Field Code Changed

Field Code Changed

23 Based on the previous results, the low-speed centrifugation concept (LSCC) was introduced as a
24 possible tool to create solid and liquid matrices of PRF, thereby generating more bioactive PRF

1 matrices. ~~These matrices that~~ contain high leukocyte and platelet concentrations, as well as advanced
2 growth factor release, according to the clinical requirements²¹.

Field Code Changed

3 Since PRF was first introduced in 2001 as a second-generation PC, it has found its way into different
4 fields of dentistry and oral maxillofacial surgery. In the last decade, there has been a constant increase
5 in the number of published papers related to the clinical applications of PRF with a great diversity of
6 experimental methods and results. This diversity has caused uncertainty about the level of scientific
7 evidence of the published articles and whether the results are reproducible. ~~Th~~~~is~~ ~~review~~ is focused
8 on, ~~we aimed to~~ individually evaluation on of these published articles with a certain level of scientific
9 evidence and reports their contributions to dentistry and maxillofacial surgery.

11 2. Materials and Methods

12 2.1 Literature research:

13 An electronic literature research was conducted using the biomedical search engine “National Library
14 of Medicine” (PubMed-MEDLINE) in May 2017. To obtain results that involved the whole scope of
15 dentistry and PRF, the keywords used for the search were “PRF”, “PRF and bone” and “PRF and soft
16 tissue”. The search resulted in a total of 392 articles. The articles were selected by reviewing the titles
17 and abstracts of the articles; 72 were selected, according to the following inclusion criteria:

- 18 - articles with the word platelet rich fibrin (PRF) in the title,
- 19 - articles related only to clinical applications in general dentistry and its different application fields
- 20 - human studies, and
- 21 - articles published in English.

22 Systematic reviews obtained through the electronic literature research were not included in the study
23 but were used to identify potentially relevant information. If needed, the references of the selected

articles were examined to search for the description of the used methodology. The full texts of the selected articles were examined by three authors (SG, CH and SA), and data were extracted using a standardized Excel data sheet (as a collection tool). The following information was recorded: name of the first author, publication year, field of dentistry (endodontics, implantology, maxillofacial surgery, orthodontics and periodontology), follow-up, number of patients involved, use of biomaterials, country, and level of scientific evidence. All collected data were arranged in tables, bars or maps using Microsoft® Excel power view® 2013 (Redmond, Washington, USA) and analyzed. The results were grouped and presented in each field (fig. 1, table 1).

Field Code Changed

2.2 Classification of the level of evidence:

All articles were independently examined by three authors (SG, CH and SA), and after a consensus, a level of scientific evidence was assigned to each article, according to the US Agency for Healthcare and Quality (table 2). The result of the search was 21 randomized clinical trials (Ib), 21 non-randomized controlled prospective studies (IIa), 13 quasi-experimental studies (IIb) and 17 case control studies (III). The method of assigning a level of scientific evidence was initiated in 1979 by the Canadian Task Force on the Periodic Health Examination, with the intention of supporting their literature recommendations on evidence-based medicine. Since the first description, randomized clinical trials have been placed at the highest level of the scientific hierarchy because they are designed to be unbiased and create less risk of systematic errors. Evaluation of the published clinical articles of PRF by their level of scientific evidence could help clinicians make selecting their best treatment options decisions as to which treatment is likely to be indicated for their patients.²⁴⁻²⁸.

Field Code Changed

2.3. Paper evaluation:

In the second step, the most used evaluation parameters in the field of periodontology (i.e., pocket depth [PD] and clinical attachment level [CAL]) were selected, and the results between the control and test groups (level of scientific evidence Ib, IIa and IIb) were extracted as the mean values of PD and CAL and calculated as the difference between the baseline and the reevaluation after six, nine or 12 months for each study separately. The data were presented graphically using Graph-Pad Prism version 6.0 (GraphPad Software, La Jolla, California). The results from each paper were reported individually without comparison or statistical analysis between the studies. Statistical differences were only reported in this manuscript when the article being examined ~~showed~~ demonstrated a significant difference in their results. The results from the studies with a level III of scientific evidence (case reports) were extracted and reported in the corresponding field. In addition, the biomaterials being evaluated in the studies and the ratio (when included) in which they were mixed with PRF were arranged and cross-referenced, as shown in table 3.

3. Results

The flowchart demonstrates the number of examined articles grouped by their level of scientific evidence and fields of dentistry (fig. 2). The rated 72 articles selected for the review showed that in the last decade, most papers were published in 2015 and 2016. By adding all patients found in the rated 72 articles, we obtained a result of 1861 patients involved in PRF studies; 728 (39.1%) participated in randomized level Ib studies, 603 (32.4%) participated in well design controlled prospect level IIa studies, 293 (15.7%) participated in quasi-experimental level IIb studies, and 237 (12.7%) participated in level III case control studies (fig. 1). The field of dentistry that showed a higher development in PRF was oral and maxillofacial surgery (38.3%), followed by periodontology (36.9%) (fig. 1)^{19,29-99}. In recent years, the use of PRF in different fields has expanded, according to the selection of published papers in the last two years (fig. 3). There were no complications reported in 28 dental pathologies treated with PRF (table 1). Furthermore, as seen in the extracted information, India is the country with the highest number of published PRF articles, particularly in the field of

Formatted: Font color: Auto

Field Code Changed

periodontology. The world map proportionally shows the number of papers published in each country, represented by different circle sizes (fig. 4).

3.1 PRF in endodontics:

Six studies were associated with the endodontic field. Five articles were case control studies (level III)^{29,31,32,33,93}, while one was a randomized study (level Ib)³⁰. In this field, a total of 55 patients were treated with PRF by placing it into the dental canals or in the peri-radicular lesions. One study reported five patients with incomplete healing. Two patients did not respond to treatment at all³⁰; in the cases of the remaining 48 patients, complete resolution and bone regeneration of the apical lesions could be achieved. The treated pathologies were immature teeth with necrotic pulps, acute chronic apical abscess and suppurative chronic apical periodontitis. Apical closure and root lengthening were observed in all cases diagnosed with immature necrotic pulps of the teeth^{29,31-33,93}.

3.2 PRF in implantology:

A total of 255 patients participated in 10 studies related to the use of PRF in implantology, with the level of scientific evidence Ib^{39,42}, IIa^{34,35,36,38,41}, IIb³⁷ and III^{9,44,0}. The treated pathologies were severe maxillary or mandibular resorption and peri-implant bone defects^{36,37,39}. Several treatment options were also described, in which PRF was used as a sole biomaterial or in combination with other biomaterials^{40,34}. Follow-up care ranged from six months⁴² to six years without implant loss³⁶. The articles reported successful osseointegration⁴², enhancement of biomechanical quality^{41,94}, safe and reliable sinus floor elevation⁴⁰, significant crestal bone gain compared to a control group without the use of PRF³⁹, and an increase of the secondary stability of implants³⁷. An additional study reported dehiscence in 31 patients after performing implant placement and tissue augmentation with PRF using a double layer technique. ~~Nevertheless, A~~after six months, all implants were osseointegrated, without any signs of infections⁴². Furthermore, rinsing dental implants prior to treatment significantly increased secondary stability over time³⁷ and significantly reduced bone resorption when PRF was used to cover the implant site³⁹. A total 19 patients presenting with (Schwarz et al. 2007¹⁰⁰: defect

Field Code Changed

Field Code Changed

Field Code Changed

Field Code Changed

Field Code Changed

Field Code Changed

Field Code Changed

Field Code Changed

Field Code Changed

Field Code Changed

Field Code Changed

Field Code Changed

Field Code Changed

Field Code Changed

Field Code Changed

Field Code Changed

Field Code Changed

Field Code Changed

Field Code Changed

classes Ib+II, Ic+II, Id+II, Ie+II) peri-implantitis (38 implants) were treated with PRF and ~~showed~~ demonstrated a survival rate of 100% after six months of follow up³⁶.

Field Code Changed

3.3 PRF in maxillofacial surgery

In the field of oral and maxillofacial surgery, four groups were identified: sinus lift, socket preservation, bone regeneration/augmentation and medication related osteonecrosis of the jaw (MRONJ).

3.3.1 Sinus lift treatment (maxillofacial surgery/implantology)

Eight articles and 198 patients were identified in this group, with IIa^{19,34,35,38,50,62}, Iib⁹⁵, and III levels of evidence⁴⁰. Two studies evaluated PRF as a sole filling biomaterial using the lateral or crestal approach^{34,40}, and the results ~~showed~~ demonstrated that the regenerated bone was sufficient for implant insertion and without bone resorption for up to six years; the final bone gain was significant, between 8.5 and 12 mm of bone gain (10.4 ± 1.2)^{34,40}. After a one-year prospective study, PRF as a sole graft material was also said to be reliable and stable⁴⁰. In a case control study, PRF was mixed with deproteinized bovine bone, and the study reported a 31% increase in the peri-implant bone density⁹⁵. Five studies evaluated PRF in combination with biomaterial versus biomaterial alone. In the PRF group, accelerated regeneration was observed; however, no statistically significant~~ce~~ between-group differences ~~were was~~ observed in new bone formation^{19,35,38,50,62}.

Field Code Changed

Field Code Changed

Field Code Changed

Field Code Changed

Field Code Changed

Field Code Changed

Field Code Changed

Field Code Changed

Field Code Changed

Field Code Changed

Field Code Changed

Field Code Changed

Field Code Changed

Field Code Changed

Field Code Changed

Field Code Changed

Field Code Changed

Field Code Changed

Field Code Changed

Field Code Changed

3.3.2 Socket preservation

A total of 14 articles investigated the effect of PRF in the treatment of socket preservation with the level of scientific evidence Ib^{51,66}, IIa^{43,44,46,48,54,65}, Iib^{45,55,58}, III^{47,53,56}. The evaluation parameters focused on bone regeneration and pain assessment. Nine studies, with a total of 343 patients, evaluated the influence of PRF on bone regeneration (PRF vs. blood clot)^{43-45,54-56,58,65,66}. Six studies showed significant improvement of socket bone fill^{56,65,66}, vertical gain of oral cortical plate⁵⁸, alveolar

Field Code Changed

ridge contour⁵³ and bone density⁵⁵ in the groups using PRF compared to the control group without PRF. Three studies showed improved bone regeneration in the PRF group but were not statistically significant^{45,43,56}.

Pain was evaluated using the visual analog scale score (PRF vs natural healing); the group treated with PRF showed significantly reduced pain compared to natural alveolus healing⁴⁵. However, Abhishek Singh et al. ~~has~~ reported that although pain could be reduced in the study group, it was not statistically significant⁴³.

A further study with a IIa level of scientific evidence evaluated the use of PRF as a tool to manage hemorrhagic complications. Fifty patients with records of heart surgery and ~~undergoing~~ anticoagulant therapy were treated by placing PRF in 168 post-extraction sockets; only two complications were reported in patients, with an international normalized ratio (INR) of 3.7. The study showed that PRF could serve as a sealing material to avoid hemorrhagic complications⁴⁸. Furthermore, in a case study, the stability of implants inserted in PRF's preserved sockets versus no socket preservation was evaluated. Significantly higher implant stability and significantly lower bone resorption were achieved compared to implants inserted in non-preserved sockets⁴⁷.

3.3.3 Bone regeneration and ridge augmentation

One study with a Ib level of scientific evidence involved 24 patients diagnosed ~~ed~~ with cleft alveolar ridge; ~~it~~ reported new bone formation to be statistically significant after the treatment with iliac crest graft combined with PRF (test group) compared to iliac crest grafts without PRF (control group)⁴⁹. In an additional study, 20 patients with cyst lesions were treated with PRF; the bone defects were filled with PRF after cyst enucleation (evidence level III, without controls). The bone defects partially regenerated after 3 months, with complete bone healing after six months⁶⁸.

3.3.4 Medication Related Osteonecrosis of the Jaw (MRONJ)

Field Code Changed

Field Code Changed

Field Code Changed

Field Code Changed

Field Code Changed

Field Code Changed

Field Code Changed

Field Code Changed

Field Code Changed

Field Code Changed

Field Code Changed

1 In total, five articles investigated the effect of PRF in the treatment of MRONJ (level of scientific
2 evidence IIa, IIb, III). Two studies with a IIa and IIb level of evidence reported statistically significant
3 improvement of wound healing compared to the control group. Epithelization was observed in the
4 PRF group within 2–4 weeks and in the control group within 2–8 weeks^{59,64}. The remaining 3 case
5 control studies with a level of evidence IIb and III reported positive clinical results^{53,52,61}. A total of
6 101 patients with MRONJ were treated using PRF as a multilayer coverage of the bone; 3 studies
7 reported a complete healing of all the cases^{53,61,64}; the remaining two articles reported no resolution
8 in 1/15 cases⁵² and 4/55 of the cases⁵⁹. The studies showed a fast epithelialization within four weeks
9 to 3 months and a total bone closure of the defect in 96 patients (during antiresorptive treatment or
10 with an interrupted treatment prior to surgery). A group of 25 patients were treated with a mixture of
11 PRF plus recombinant human bone morphogenetic proteins 2 (rhBMP2) and compared to a control
12 group of 30 patients treated with PRF only; the results showed statistically significant improvements
13 in healing in the PRF+BMP2 group⁵⁹.

14 3.5 Orthodontics

15 Only one article was found in this field with a level III of scientific evidence. An orthodontic-surgical
16 treatment using Wilcko's modified periodontally accelerated osteogenic orthodontics (PAOO) was
17 performed in 11 patients; PRF was minced or applied as a membrane combined with mineralized
18 human cancellous bone allograft + deproteinized bovine bone material (3:2) and metronidazole (500
19 mg). All patients healed without any problems; orthodontic treatment took, on average, 9.3 months,
20 with stable treatment after two years⁶⁹.

21 3.6 Periodontology

22 A total of 26 articles were selected with different levels of scientific evidence Ib^{71–75,79,81,83,85,89–92,96,97},
23 IIa^{76,82,84,86,98}, IIb^{70,77,87,88} and III^{78,99}. More randomized studies with control and test groups were found
24 in this field. The pathologies treated were chronic periodontitis with different levels of bone defects,
25 gingival recession and gingiva hyperpigmentation. A total of 13 of the 26 articles included a PRF

Field Code Changed

Field Code Changed

Field Code Changed

Field Code Changed

Field Code Changed

Field Code Changed

Field Code Changed

Field Code Changed

Field Code Changed

Field Code Changed

Field Code Changed

Field Code Changed

Field Code Changed

Field Code Changed

Field Code Changed

Field Code Changed

study-group as a sole treatment compared to conventional treatments (e.g., open flap debridement, connective tissue graft, and autogenous bone graft)^{70,73,74,79,81–83,85,87,90–92}. Ten of these studies reported that the use of PRF improved equally (4 articles)^{70,83,91,97} or significantly (six articles)^{73,74,79,81,85,90} for PD and/or CAL compared to the conventional surgical treatments. An additional study showed no significant differences in PD reduction and CAL gain⁸² (fig. 5 and 6). As opposed to measuring PD and CAL, the previously mentioned articles and two others obtained improvements in pain reduction, healing index^{83,87,92,97} and bone-defect fill^{79,70}.

A total of 14 studies investigated the combination of PRF with different biomaterials (e.g., hydroxyapatite bone graft^{71,99}, freeze-dried bone allograft [DFDBA]^{72,75,77,90}, autogenous bone graft⁷⁶, amnion allograft membrane⁸⁴, bioactive glass calcium phosphosilicate^{86,88}, anorganic bovine bone mineral⁸⁹, enamel matrix derivative⁹⁷, bovine porous bone mineral⁹⁸ and bioactive Gengigel hyaluronic acid)⁷⁸. The most frequent combination used was PRF+DFDBA; the used ratio of the mixture was reported only in six articles (1:1 or 1:2); (table 3). Three studies reported statistically significant improvements in PD and CAL compared to the sole use of biomaterial or PRF⁹⁰. Six studies combining PRF with biomaterials of allogenic or xenogeneic source showed significantly improved CAL and PD in the PRF group compared to the control group (biomaterial without PRF). Two studies did not show significant results between the evaluated groups⁹⁷, and 1 reported better root coverage in the control group⁹⁶. Furthermore, the use of autogenous bone graft was compared to the use of PRF as a sole filling material for bone regeneration, and no statistically significant improvement was observed between the evaluated groups after the intervention⁷⁶. Two studies observed that the use of bioactive Gengigel or hydroxyapatite graft combined with PRF seems to have advantages in terms of the bone regeneration of furcation intrabony defects^{78,99}.

4. Discussion

In the present review, a level of scientific evidence was assigned to each article^{see in} to establish a parameter of comparison and to give guidance to clinicians. To avoid bias in the selection of the

Field Code Changed

Field Code Changed

Field Code Changed

Field Code Changed

Field Code Changed

Field Code Changed

Field Code Changed

Field Code Changed

Field Code Changed

Field Code Changed

Field Code Changed

Field Code Changed

Field Code Changed

Field Code Changed

Field Code Changed

Field Code Changed

Field Code Changed

Field Code Changed

Field Code Changed

Field Code Changed

Field Code Changed

Field Code Changed

Field Code Changed

1 articles, the initial inclusion criteria were based on the title of the articles and, second, on the content
2 of the abstract to verify its relationship to dentistry and its different fields. The extracted data were
3 reported based on the articles in table 1.

4 As seen in the results of this review, over 70% of the patients were part of a randomized or controlled
5 prospective study. The subsequent data can serve as a compendium for clinical practitioners
6 combining dental pathologies and treatments where PRF is used (table 1). Note that 1861 patients
7 participated in 72 articles included in this review, resulting in 28 different diagnosis and pathologies.
8 In recent years, the application of PRF in different fields has expanded, as the selection of published
9 papers in the last two years (fig. 3) indicates. This result may be primarily because of the absence of
10 complications, positive results, autologous source and the simplified and clinically applicable
11 preparation method.

12 The World of PRF map illustrates the presence throughout the world of PRF to give an idea of how
13 research in the clinical environment has developed (fig. 4). All fields of dentistry are involved,
14 however, periodontology and oral-maxillofacial surgery remain the primary areas of application. The
15 properties of PRF to accelerate the healing process of soft and hard tissue, while controlling pain and
16 inflammation, opens up a broad area of application to clinicians.

17 The essence of PRF may rely on the delivery of blood components at an early stage of healing, which
18 diminishes the noxious phases of stress-induced hyper-metabolism in response to injury¹⁰¹.
19 Additionally, PRF from humans can regulate inflammation, promote vascularization, provide a
20 matrix for cells and improve the healing of tissue and bone²⁰.

21 Continuous research in the field of PRF has shown that PRF, which is a bioactive autologous system,
22 can be considered as a drug delivery system²⁰. Current studies aim to optimize the regenerative
23 capacity of PRF using the low-speed centrifugation concept (LSCC). In this context, PRF matrices
24 that are centrifuged using a low relative centrifugation force (RCF) show higher leukocyte and platelet

Field Code Changed

Field Code Changed

Field Code Changed

1 concentrations compared to PRF matrices that are centrifuged using a high RCF²¹. Consequently,
2 matrices generated in the low RCF range release a higher concentration of growth factors (VEGF,
3 EGF, TGF β -1) compared to those generated in a high RCF range^{21,102}.

Field Code Changed

4 The results of the clinical application of PRF in different fields of dentistry and oral-maxillofacial
5 surgery ~~showed-demonstrated~~ that, presently, there is a considerable group of published articles with
6 a high scientific evidence of the effectivity of PRF in bone and soft tissue regeneration.

Field Code Changed

7 In the field of periodontology, 23 studies with a scientific evidence of Ib, IIa or IIb have shown that
8 the application of PRF led to a significantly improved pocket depth (PD) and clinical attachment level
9 (CAL) compared to the groups treated without PRF, and one study with a scientific evidence of Ib
10 reported better root coverage in the control group. Additionally, the results in this review highlight
11 the fact that in the field of periodontology, the use of biomaterials in combination with PRF seems to
12 significantly improve the regeneration potential of the used biomaterials, which may be related to the
13 “biologization” of the acellular biomaterial prior to their application and could facilitate the cell-cell
14 communication and biomaterial integration in the application region¹⁰³.

Field Code Changed

15 ~~Nevertheless, Most~~ studies regarding the surgical-periodontal treatment have focused on
16 periodontitis with different diagnosis. ~~Whereas-Only~~ one study with a scientific evidence of IIa has
17 evaluated the effect of PRF application in peri-implantitis (i.e., 100% survival rate of the implants)³⁶,
18 with significantly improved PD compared to treatment without PRF. A recent Cochrane review
19 focused on the effectiveness of current treatments for peri-implantitis; it associated the use of bovine-
20 derived xenograft with PD and CAL improvement¹⁰⁴. Interestingly, six of the studies included in this
21 review obtained statistically significant improvement of PD and CAL using PRF + biomaterial
22 compared to the sole use of a biomaterial. Furthermore, the most frequent used biomaterial was found
23 to be demineralized freeze-dried bone allograft (DFDBA), showing significant positive results. In
24 particular, it has been described that biomaterials derived from natural bone sources, such as DFDBA,
25 contain BMPs⁷⁵. An additional study included in the review combined PRF+BMPs and obtained

Field Code Changed

Field Code Changed

Field Code Changed

1 significant clinical improvement compared to sole PRF. The observed interaction between PRF and
2 BMPs could help explain how PRF could enhance the regenerative potential in the case of
3 biomaterials derived from natural bone sources. Some points still need to be clarified (e.g., if the
4 mixing ratio of PRF and the autologous grafts and/or biomaterials has any effect over the obtained
5 clinical results). As observed in our review, only six of the included articles reported the mixing ratio
6 (1:1 or 1:2) (table 3)³⁸⁷⁰⁷¹⁷⁵⁹¹⁹⁸. Today, due to the increased application of dental implants in the last
7 decades, peri-implantitis has become a key area in periodontology, as the primary long-term
8 complication after implant placement. The results of the present review could encourage clinicians to
9 use PRF for peri-implantitis treatment, and this process requires further research in this area.

10 Interestingly, a further application field, such as bone regeneration within the sinus cavity (sinus-lift),
11 showed no statistically significant difference between the new bone regeneration using biomaterials
12 in combination with PRF compared to biomaterials alone in studies with a scientific evidence of Ib,
13 IIa or IIb. ~~Whereas~~ PRF as a sole material was shown to support bone regeneration in the sinus lift.
14 However, the level of scientific evidence of these studies is low ~~because due to the they~~ lack of a
15 control group. Moreover, there is a general lack of research investigating the bone regeneration in
16 this field.

17 ~~However, Bb~~ bone regeneration after socket preservation with PRF was shown to accelerate/improve
18 bone density and tissue healing compared to physiological defect healing. These results were
19 underlined by 14 of the studies with a level of scientific evidence of Ib, IIa or IIb. ~~The However, the~~
20 contribution of PRF to bone healing should be further evaluated in critical bone defects and more
21 complex augmentations, such as ~~the -challenging of~~ vertical and horizontal bone augmentation. In
22 this manner, a higher evidence of the benefits in bone and tissue regeneration can be obtained.
23 ~~Furthermore, Aa~~ comparison between the combination of PRF with biomaterials and biomaterials
24 alone has not yet been investigated. Therefore, further studies are required to evaluate the role of PRF
25 in combination with biomaterials and their bioactivity.

Field Code Changed

1 These studies all ~~used~~utilized clinical and radiological methods to investigate regeneration capacity
2 and bone stability. However, only a few single studies ~~that~~ have used histological methods to analyze
3 bone regeneration in humans.

4 In addition the application of PRF in socket of extracted third molars led to significantly reduced pain
5 compared to conventional socket healing^{45,47,51}. These observations could be explained by the ability
6 of PRF's ~~of~~ release of different pro- and anti-inflammatory cytokines, such as IL-4, that modulate
7 the inflammatory response and are involved in the cascade of pain.

Field Code Changed

8 In addition, the quality of new bone formation and stability seemed to be improved by the PRF
9 treatment, ~~and~~ thus, the mechanical stability and osteointegration of implants inserted in the bone
10 regions treated with PRF or covered by PRF have been shown to exhibit significantly higher
11 secondary stability and experienced significantly less bone resorption compared to implants inserted
12 in non-treated bones^{37,60}.

Field Code Changed

13 Additionally, five studies have shown that PRF is beneficial in the treatment of medication related to
14 osteonecrosis of the jaw. Reepithelialization and bone regeneration ~~were~~was achieved in 96 of the
15 101 treated cases. In this context, PRF provided a less invasive treatment option for multimorbid
16 patients. Among others, this disease pattern is related to impaired vascularization and wound
17 healing⁵⁹. Thereby, these results may be achieved due to the bioactivity of PRF and its capability of
18 releasing VEGF and EGF, which primarily contribute to angiogenesis and epithelialization¹⁰².

Field Code Changed

Field Code Changed

19 In other fields of dentistry, such as endodontics and orthodontics, little research has been performed,
20 with a low level of scientific evidence. ~~In addition,~~ There is a general lack of long term or
21 retrospective studies evaluating the long-term results of implants inserted in bones treated with PRF
22 compared to non-treated bones.

23 Our results provide published evidence of high scientific level that PRF (38 articles) is a beneficial
24 tool that significantly enhances bone and soft tissue regeneration. Furthermore, 17 articles have

reported that with the use of PRF as a sole biomaterial, similar results were obtained compared to the conventional treatment. In contrast, only one study reported statistically significant root coverage in the control group (conventional treatment) compared to the test group (PRF). There may be other cases with similar results in this field, which have remained unpublished thus far. The remaining 16 articles, with a scientific evidence level III, observed clinical resolution of the treated cases with PRF. In summary, this review serves as an overview of the level of scientific evidence of studies that analyzed the role of PRF in dentistry and oral ~~and~~, maxillofacial surgery. To promote studies with a higher scientific level of evidence, the clinical community requires a standardization of PRF protocols and more well-designed and conducted studies.

5. Conclusion

The present review provides an overview of the literature regarding the existing clinical research of PRF and its level of scientific evidence. Over 70% of patients participated in studies with a high level of scientific evidence (randomized and controlled prospect studies). In nine of the studies, the application of PRF in periodontology showed significantly improved pocket depth and clinical attachment level compared to the conventional treatment. ~~Moreover, S~~significantly enhanced new bone formation was reported during socket preservation and ridge augmentation procedures (seven articles) comparing the used of PRF to the normal bone healing without PRF. ~~Furthermore, N~~no statistically significant differences were found in the addition of PRF to biomaterial in sinus lift compared to sole biomaterials. The results obtained when patients were diagnosed with medication-related osteonecrosis of the jaw (MRONJ) treated with PRF are also relevant, resulting in 96 of 101 of the patients healing uneventfully. Thus, the present review highlights the existing studies with a high scientific level of evidence that equally show the contribution of PRF in bone and soft tissue regeneration. From the 72 articles involved in the review, only one reported statistically significant results favoring the control group against the PRF group. Although 16 articles with a level III of scientific evidence reported good outcomes, they have the lowest level of scientific evidence. To

promote reproducible studies with a higher scientific level of evidence, the clinical community requires a standardization of PRF protocols to show the benefit of PRF in tissue regeneration.

References

References

1. Ghanaati S, Kovács A, Barbeck M, et al. Bilayered, non-cross-linked collagen matrix for regeneration of facial defects after skin cancer removal: a new perspective for biomaterial-based tissue reconstruction. *J Cell Commun Signal*. 2016;10(1):3-15.
2. Tsaryk R, Gloria A, Russo T, et al. Collagen-low molecular weight hyaluronic acid semi-interpenetrating network loaded with gelatin microspheres for cell and growth factor delivery for nucleus pulposus regeneration. *Acta Biomater*. 2015;20:10-21.
3. Schlee M, Ghanaati S, Willershausen I, Stimmlmayr M, Sculean A, Sader RA. Bovine pericardium based non-cross linked collagen matrix for successful root coverage, a clinical study in human. *Head Face Med*. 2012;8(1):6
4. Lorenz J, Blume M, Barbeck M, et al. Expansion of the peri-implant attached gingiva with a three-dimensional collagen matrix in head and neck cancer patients—results from a prospective clinical and histological study. *Clin Oral Investig*. 2017;21(4):1103-1111.
5. Duttonhoefer F, Hieber SF, Stricker A, Schmelzeisen R, Gutwald R, Sauerbier S. Follow-up of implant survival comparing ficoll and bone marrow aspirate concentrate methods for hard tissue regeneration with mesenchymal stem cells in humans. *Biores Open Access*. 2014;3(2):75-76.
6. Sakkas A, Wilde F, Heufelder M, Winter K, Schramm A. Autogenous bone grafts in oral implantology-is it still a "gold standard? A consecutive review of 279 patients with 456 clinical procedures. *Int J Implant Dent*. 2017;3(1):23.
7. Matras H. The use of fibrin sealant in oral and maxillofacial surgery. *J Oral Maxillofac Surg*. 1982;40(10):617-622.
8. Laurens N, Koolwijk P, de Maat MP. Fibrin structure and wound healing. *J Thromb Haemost*. 2006;4(5):932-939.
9. Cieslik-Bielecka A, Choukroun J, Guillaume O, Dohan Ehrenfest DM. L-PRP/L-PRF in Esthetic Plastic Surgery, Regenerative Medicine of the Skin and Chronic Wounds. *Curr Pharm Biotechnol*. 2012;13:1266-1277.
10. Knighton DR, Ciresi KF, Fiegel VD, Austin LL, Butler EL. Classification and treatment of chronic nonhealing wounds. Successful treatment with autologous platelet-derived wound healing factors (PDWHF). *Ann Surg*. 1986;204(3):322-330.

Formatted: Font color: Auto

11. Perez AGM, Lana JFSD, Rodrigues AA, Luzo ACM, Belangero WD, Santana MHA. Relevant Aspects of Centrifugation Step in the Preparation of Platelet-Rich Plasma. *ISRN Hematol*. 2014.
12. Marx RE. Platelet-Rich Plasma: Evidence to Support Its Use. *J Oral Maxillofac Surg*. 2004;62(4):489-496.
13. Dohan DM, Choukroun J, Diss A, et al. Platelet-rich fibrin (PRF): A second-generation platelet concentrate. Part I: Technological concepts and evolution. *Oral Surgery, Oral Med Oral Pathol Oral Radiol Endodontology*. 2006;101(3).
14. Choukroun J, Adda F, Schoeffler C, Vervelle A. Uneopportunit  en paro-implantologie: le PRF. *Implantodontie*. 2001;(42):55-62.
15. Fujioka-Kobayashi M, Miron RJ, Hernandez M, Kandalam U, Zhang Y, Choukroun J. Optimized Platelet Rich Fibrin With the Low Speed Concept: Growth Factor Release, Biocompatibility and Cellular Response. *J Periodontol*. 2016:1-17.
16. Fern ndez-Delgado N, Hern ndez-Ram rez P, Forrellat-Barrios M. Platelet functional spectrum: from hemostasis to regenerative medicine. *Rev Cuba Hematol Inmunol y Hemoter*. 2012;28(3):200-216.
17. Neumann FJ, Marx N, Gawaz M, et al. Induction of cytokine expression in leukocytes by binding of thrombin-stimulated platelets. *Circulation*. 1997;95(10):2387-2394.
18. Choukroun J, Diss A, Simonpieri A, et al. Platelet-rich fibrin (PRF): A second-generation platelet concentrate. Part IV: Clinical effects on tissue healing. *Oral Surgery, Oral Med Oral Pathol Oral Radiol Endodontology*. 2006;101(3):56-60.
19. Choukroun J, Diss A, Simonpieri A, et al. Platelet-rich fibrin (PRF): A second-generation platelet concentrate. Part V: Histologic evaluations of PRF effects on bone allograft maturation in sinus lift. *Oral Surgery, Oral Med Oral Pathol Oral Radiol Endodontology*. 2006;101(3):299-303.
20. Ghanaati S, Booms P, Orlowska A, et al. Advanced Platelet-Rich Fibrin: A New Concept for Cell-Based Tissue Engineering by Means of Inflammatory Cells. *J Oral Implantol*. 2014;40(6):679-689.
21. J. Choukroun and S. Ghanaati. Reduction of relative centrifugation force within PRF- (Platelet-Rich-Fibrin) concentrates advances patients' own inflammatory cells and platelets: First introduction of the Low Speed Centrifugation Concept. *Eur J Trauma Emerg Surg*. 2017.
22. Dohle E, El Bagdadi K, Sader R, Choukroun J, James Kirkpatrick C, Ghanaati S. PRF-based matrices to improve angiogenesis in an in vitro co-culture model for bone tissue engineering. *J Tissue Eng Regen Med*. 2017.
23. Wang X, Zhang Y, Choukroun J, Ghanaati S, Miron RJ. Behavior of Gingival Fibroblasts on Titanium Implant Surfaces in Combination with either Injectable-PRF or PRP. *Int J Mol Med*. 2017;18(331).
24. McCarthy CM, Collins ED, Pusic AL. Where do we find the best evidence? *Plast Reconstr Surg*. 2008;122:1942-1947-1951.

Formatted: Font color: Auto, English (United States)

Formatted: Font color: Auto

- 1 25. Burns P, Rohrich R, Chong K. The Levels of Evidence and their role in Evidence-Based
2 Medicine. *Plast Reconstr Surg*. 2011;128(1):305-310.
- 3 26. Bhandari M, Devereaux PJ, Montori V, Cina C. User guide to the surgical literature How to
4 assess a randomized controlled trial in surgery. *Can J Surg*. 2004;47(3).
- 5 27. Guyatt GH, Sackett DL, Cook DJ. User's Guides to the Medical Literature Therapy or
6 Prevention Results and Will They Help Me? *J Am Med Assoc*. 1994;271(1):59-63.
- 7 28. Healthcare MOF, Affairs C. *Clinical Practice Guideline for Schizophrenia and Incipient*
8 *Psychotic Disorder Clinical Practice Guideline for Schizophrenia and Incipient Psychotic Disorder*. First
9 edit. Barcelona: Catalan Agency for Health Technology Assessment and Research; 2009.
- 10 29. Geeta IB, Galagali G, Sangeeta K, Pushpa S, Noushin F. A natuaral meliorate: Revolutionary
11 tissue engineering in endodontics. *J Clin Diagnostic Res*. 2013;7(11):2644-2646.
- 12 30. Dhiman M, Kumar S, Duhan J, Sangwan P, Tewari S. Effect of Platelet-rich Fibrin on Healing
13 of Apicomarginal Defects: A Randomized Controlled Trial. *J Endod*. 2015;41(7):985-991.
- 14 31. Singh S, Singh A, Singh S, Singh R. Application of PRF in surgical management of periapical
15 lesions. *Natl J Maxillofac Surg*. 2013;4(1):94-99.
- 16 32. Subash D, Shoba K, Aman S, Bharkavi SKI. Revitalization of an immature permanent
17 mandibular molar with a necrotic pulp using platelet-rich fibrin: A case report. *J Clin Diagnostic Res*.
18 2016;10(11):ZD21-ZD23.
- 19 33. Bakhtiar H, Esmaeili S, Fakhr Tabatabayi S, Ellini MR, Nekoofar MH, Dummer PMH. Second-
20 generation Platelet Concentrate (Platelet-rich Fibrin) as a Scaffold in Regenerative Endodontics: A
21 Case Series. *J Endod*. 2017;43(3):401-408.
- 22 34. Simonpieri A, Choukroun J, Del Corso M, Sammartino G, Dohan Ehrenfest DM. Simultaneous
23 sinus-lift and implantation using microthreaded implants and leukocyte- and platelet-rich fibrin as
24 sole grafting material: a six-year experience. *Implant Dent*. 2011;20(1):2-12.
- 25 35. Zhang Y, Tangl S, Huber CD, Lin Y, Qiu L, Rausch-Fan X. Effects of Choukroun's platelet-rich
26 fibrin on bone regeneration in combination with deproteinized bovine bone mineral in maxillary
27 sinus augmentation: A histological and histomorphometric study. *J Cranio-Maxillofacial Surg*.
28 2012;40(4):321-328.
- 29 36. Hamzacebi B, Oduncuoglu B, Alaaddinoglu EE. Treatment of Peri-implant Bone Defects with
30 Platelet-Rich Fibrin. *Int J Periodontics Restorative Dent*. 2015;35(3):414-422 9p.
- 31 37. Öncü E, Alaaddinoğlu E. The Effect of Platelet-Rich Fibrin on Implant Stability. *Int J Oral*
32 *Maxillofac Implants*. 2015;30(3):578-582.
- 33 38. Bolukbasi N, Ersanlı S, Keklikoglu N, Basegmez C, Ozdemir T. Sinus Augmentation With
34 Platelet-Rich Fibrin in Combination With Bovine Bone Graft Versus Bovine Bone Graft in
35 Combination With Collagen Membrane. *J Oral Implantol*. 2015;41(5):586-595.

- 1 39. Boora P, Rathee M, Bhoria M. Effect of Platelet Rich Fibrin (PRF) on peri-implant soft tissue
2 and crestal bone in one-stage implant placement: A randomized controlled trial. *J Clin Diagnostic*
3 *Res.* 2015;9(4):ZC18-ZC21.
- 4 40. Kanayama T, Horii K, Senga Y, Shibuya Y. Crestal Approach to Sinus Floor Elevation for
5 Atrophic Maxilla Using Platelet-Rich Fibrin as the Only Grafting Material: A 1-Year Prospective Study.
6 *Implant Dent.* 2015:32-38.
- 7 41. Angelo T, Marcel W. Biomechanical Stability of Dental Implants in Augmented Maxillary
8 Sites: Results of a Randomized Clinical Study with Four Different Biomaterials and PRF and a
9 Biological View on Guided Bone Regeneration. *Biomed Res Int.* 2015;2015:17.
- 10 42. Hehn J, Schwenk T, Striegel M, Schlee M. The effect of PRF (platelet-rich fibrin) inserted with
11 a split-flap technique on soft tissue thickening and initial marginal bone loss around implants: results
12 of a randomized, controlled clinical trial. *Int J Implant Dent.* 2016;2(1):13.
- 13 43. Singh A, Kohli M, Gupta N. Platelet rich fibrin: a novel approach for osseous regeneration. *J*
14 *Maxillofac Oral Surg.* 2012;11(4):430-434.
- 15 44. Doiphode AM, Hegde P, Mahindra U, Santhosh Kumar SM, Tenglikar PD, Tripathi V.
16 Evaluation of the efficacy of platelet-rich plasma and platelet-rich fibrin in alveolar defects after
17 removal of impacted bilateral mandibular third molars. *J Int Soc Prev Community Dent.* 2016;6(Suppl
18 1):S47-52.
- 19 45. Emanuele Ruga, DDS, Cesare Gallesio, MD D, Paolo Boffano M. Platelet-Rich Fibrin and
20 Piezoelectric Surgery: A Safe Technique for the Prevention of Periodontal Complications in Third
21 Molar Surgery. *J Craniofac Surg.* 2011;22(5):1951-1955.
- 22 46. Marenzi G, Riccitiello F, Tia M, di Lauro A, Sammartino G. Influence of Leukocyte- and
23 Platelet-Rich Fibrin (L-PRF) in the Healing of Simple Postextraction Sockets: A Split-Mouth Study.
24 *Biomed Res Int.* 2015;2015:369-273.
- 25 47. Kotsakis G, Prasad H, Rohrer M, et al. Extraction socket management utilizing Platelet-Rich-
26 Fibrin : A proof-of-principle study of the accelerated-early implant placement concept. *J Oral*
27 *Implantol.* 2015.
- 28 48. Sammartino G, Ehrenfest DMD, Carile F, Tia M, Bucci P. Prevention of Hemorrhagic
29 Complications After Dental Extractions Into Open Heart Surgery Patients Under Anticoagulant
30 Therapy: The Use of Leukocyte- and Platelet-Rich Fibrin. *J Oral Implantol.* 2011;37(6):681-690.
- 31 49. Shawky H, Seifeldin SA. Does Platelet-Rich Fibrin Enhance Bone Quality and Quantity of
32 Alveolar Cleft Reconstruction? *Cleft Palate Craniofac J.* 2015;0(0).
- 33 50. Tatullo M, Marrelli M, Cassetta M, et al. Platelet rich fibrin (P.R.F.) in reconstructive surgery
34 of atrophied maxillary bones: Clinical and histological evaluations. *Int J Med Sci.* 2012;9(10):872-
35 880.
- 36 51. Kumar N, Prasad K, Ramanujam L, K R, Dexith J, Chauhan A. Evaluation of Treatment
37 Outcome After Impacted Mandibular Third Molar Surgery With the Use of Autologous Platelet-Rich
38 Fibrin: A Randomized Controlled Clinical Study. *J Oral Maxillofac Surg.* 2014;(February):1-8.

Formatted: Font color: Auto, English (United States)

Formatted: Font color: Auto

- 1 52. Nørholt SE, Hartlev J. Surgical treatment of osteonecrosis of the jaw with the use of platelet-
2 rich fibrin: a prospective study of 15 patients. *Int J Oral Maxillofac Surg*. 2016.
- 3 53. Soydan SS, Uckan S. Management of bisphosphonate-related osteonecrosis of the jaw with
4 a platelet-rich fibrin membrane: Technical report. *J Oral Maxillofac Surg*. 2014;72(2):322-326.
- 5 54. Suttapreyasri S, Leepong N. Influence of platelet-rich fibrin on alveolar ridge preservation. *J*
6 *Craniofac Surg*. 2013;24(4):1088-1094.
- 7 55. Yelamali T, Saikrishna D. Role of Platelet Rich Fibrin and Platelet Rich Plasma in Wound
8 Healing of Extracted Third Molar Sockets: A Comparative Study. *J Maxillofac Oral Surg*.
9 2015;14(2):410-416.
- 10 56. Kumar YR, Mohanty S, Verma M, et al. Platelet-rich fibrin: The benefits. *Br J Oral Maxillofac*
11 *Surg*. 2016;54(1):57-61.
- 12 57. Kapustecki M, Niedzielska I, Marek HB, Ranowski B. Alternative method to treat oroantral
13 communication and fistula with autogenous bone graft and platelet rich fibrin. *Med Oral Patol Oral*
14 *Cir Bucal*. 2016;21(5):608-613.
- 15 58. Anwandter A, Bohmann S, Nally M, Castro AB, Quirynen M, Pinto N. Dimensional changes
16 of the post extraction alveolar ridge, preserved with Leukocyte- and Platelet Rich Fibrin: A clinical
17 pilot study. *J Dent*. 2016;52:23-29.
- 18 59. Park JH, Kim JW, Kim SJ. Does the Addition of Bone Morphogenetic Protein 2 to Platelet-
19 Rich Fibrin Improve Healing After Treatment for Medication-Related Osteonecrosis of the Jaw? *J*
20 *Oral Maxillofac Surg*. 2016;75(6):1176-1184.
- 21 60. Öncü E und AA in E. Enhancement of Immediate Implant Stability and Recovery Using
22 Platelet-Rich Fibrin. *Int J Periodontics Restorative Dent*. 2017.
- 23 61. Gönen ZB, Yılmaz Asan C. Treatment of bisphosphonate-related osteonecrosis of the jaw
24 using platelet-rich fibrin. *Cranio - J Craniomandib Pract*. 2016;9634(July):1-5.
- 25 62. Gokhan GURLER CD. Effects of leukocyte-platelet rich fibrin on postoperative complications
26 of direct sinus lifting. *Minerva Stomatol*. 2016;65(4):207-212.
- 27 63. Cortese A, Pantaleo G, Borri A, Caggiano M, Amato M. Platelet-rich fibrin (PRF) in implant
28 dentistry in combination with new bone regenerative technique in elderly patients. *Int J Surg Case*
29 *Rep*. 2016;28:52-56.
- 30 64. Asaka T, Ohga N, Yamazaki Y, Sato J, Satoh C, Kitagawa Y. Platelet-rich fibrin may reduce the
31 risk of delayed recovery in tooth-extracted patients undergoing oral bisphosphonate therapy: a trial
32 study. *Clin Oral Investig*. 2016:1-8.
- 33 65. Varghese MP, Manuel S, Kumar L. K. S. Potential for Osseous Regeneration of Platelet-Rich
34 Fibrin-A Comparative Study in Mandibular Third Molar Impaction Sockets. *J Oral Maxillofac Surg*.
35 2016:1-8.

Formatted: Font color: Auto, English (United States)

Formatted: Font color: Auto

66. Temmerman A, Vandessel J, Castro A, et al. The use of leucocyte and platelet-rich fibrin in socket management and ridge preservation: a split-mouth, randomized, controlled clinical trial. *J Clin Periodontol.* 2016;43(11):990-999.
67. Bilginaylar K. Uncommon Odontogenic Orocutaneous Fistula of the Jaw Treated with Platelet-Rich Fibrin. *Case Rep Dent.* 2017;2017:1-5.
68. Dar M, Hakim T, Shah A, Najar L, Yaqoob G, Lanker F. Use of autologous platelet-rich fibrin in osseous regeneration after cystic enucleation: A clinical study. *J Oral Biol Craniofacial Res.* 2016;6:S29-S32.
69. Munoz F, Jiménez C, Espinoza D, Vervelle A, Beugnet J, Haidar Z. Use of leukocyte and platelet-rich fibrin (L-PRF) in periodontally accelerated osteogenic orthodontics (PAOO): Clinical effects on edema and pain. *J Clin Exp Dent.* 2016;8(2):e119-24.
70. Pradeep AR, Nagpal K, Karvekar S, Patnaik K, Naik SB, Guruprasad CN. Platelet Rich Fibrin with 1% Metformin for the Treatment of Intrabony Defects in Chronic Periodontitis: A Randomized Controlled Clinical Trial. *J Periodontol.* 2015;12:1-14.
71. Pradeep R, Karvekar S, Nagpal K, Patnaik K, Raju A, Singh P. Rosuvastatin 1.2 mg in Situ Gel Combined With 1:1 Mixture of Autologous Platelet-Rich Fibrin and Porus- Hydroxyapatite Bone Graft in Surgical Treatment of Mandibular Degree II Furcation Defects: A Randomized Clinical Control Trial. *J Periodontol.* 2015;(January):1-15.
72. Gamal AY, Abdel Ghaffar KA, Algezwy OA. Crevicular Fluid Growth Factors Release Profile Following the Use of Platelets Rich Fibrin (PRF) and Plasma Rich Growth Factors (PRGF) in Treating Periodontal Intrabony Defects (Randomized Clinical Trial). *J Periodontol.* 2016:1-13.
73. Sharma A, Pradeep A. Autologous Platelet-Rich Fibrin in the Treatment of Mandibular Degree II Furcation Defects: A Randomized Clinical Trial. *J Periodontol.* 2011;82(October):1396-1403.
74. Sharma A, Pradeep a. R. Treatment of 3-Wall Intrabony Defects in Patients With Chronic Periodontitis With Autologous Platelet-Rich Fibrin: A Randomized Controlled Clinical Trial. *J Periodontol.* 2011;82(12):1705-1712.
75. Agarwal A, Gupta ND. Platelet-rich plasma combined with decalcified freeze-dried bone allograft for the treatment of noncontained human intrabony periodontal defects: a randomized controlled split-mouth study. *Int J Periodontics Restorative Dent.* 2014;34(5):705-711.
76. Mathur A, Bains VK, Gupta V, Jhingran R, Singh RP. Evaluation of intrabony defects treated with platelet-rich fibrin or autogenous bone graft: A comparative analysis. *Eur J Dent.* 2015;9(1):100-108.
77. Bharti CB and V. Evaluation of efficacy of autologous platelet rich fibrin with demineralized-freeze dried bone allograft in the treatment of periodontal intrabony defects. *J Indian Soc Periodontol.* 2013;17(3):361-366.

78. Sandhu GK, Khinda PK, Gill AS, Kalra HS. Surgical re-entry evaluation of regenerative efficacy of bioactive Gengigel and platelet-rich fibrin in the treatment of grade II furcation: A novel approach. *Contemp Clin Dent*. 2015;6(4):570-573.
79. Ajwani H, Shetty S, Gopalakrishnan D, et al. Comparative evaluation of platelet-rich fibrin biomaterial and open flap debridement in the treatment of two and three wall intrabony defects. *J Int oral Heal JIOH*. 2015;7(4):32-37.
80. Moussa M, El-Dahab OA, El Nahass H. Anterior Maxilla Augmentation Using Palatal Bone Block with Platelet-Rich Fibrin: A Controlled Trial. *Int J Oral Maxillofac Implants*. 2016;31(3):708-715.
81. Bajaj P, Pradeep AR, Agarwal E, et al. Comparative evaluation of autologous platelet-rich fibrin and platelet-rich plasma in the treatment of mandibular degree II furcation defects: A randomized controlled clinical trial. *J Periodontal Res*. 2013;48(5):573-581.
82. Rosamma Joseph V, Sam G, Vijay Amol N. Clinical evaluation of autologous platelet rich fibrin in horizontal alveolar bony defects. *J Clin Diagnostic Res*. 2014;8(11):ZC43-ZC47.
83. Jankovic S, Aleksic Z. Use of Platelet-Rich Fibrin Membrane Following Treatment of Gingival Recession : A Randomized Clinical Trial. *Int J Periodontics Restorative Dent*. 2012;32(2):41-50.
84. Agarwal S, Jhingran R, Bains V, Srivastava R, Madan R, Rizvi I. Patient-centered evaluation of microsurgical management of gingival recession using coronally advanced flap with platelet-rich fibrin or amnion membrane: A comparative analysis. *Eur J Dent*. 2016;10(1):121.
85. Thorat M, Pradeep AR, Pallavi B. Clinical effect of autologous platelet-rich fibrin in the treatment of intra-bony defects: A controlled clinical trial. *J Clin Periodontol*. 2011;38(10):925-932.
86. Chang YC, Zhao JH. Effects of platelet-rich fibrin on human periodontal ligament fibroblasts and application for periodontal infrabony defects. *Aust Dent J*. 2011;56(4):365-371.
87. Bansal M, Kumar A, Puri K, Khatri M, Geeti G, Vi H. Clinical and Histologic Evaluation of Platelet-Rich Fibrin Accelerated Epithelization of Gingival Wound. *J Cutan Aesthet Surg*. 2016;9(3):196-200.
88. Biswas S, Sambashivaiah S, Kulal R, Bilichodmath S, Kurtzman GM. Comparative Evaluation of Bioactive Glass (Putty) and Platelet Rich Fibrin in Treating Furcation Defects. *J Oral Implantol*. 2016;42(5):411-415.
89. Sezgin Y, Uraz A, Taner IL, Çulhaoglu R. Effects of platelet-rich fibrin on healing of intra-bony defects treated with anorganic bovine bone mineral. *Braz Oral Res*. 2017;31(1):1-11.
90. Nikhil D., Chandradas Shivamurthy, Ravindra Vivekananda M. Rangaraju Sheetal J, Shivaprasad D. Efficacy of platelet rich fibrin in the treatment of human intrabony defects. *J Adv Pharm Technol Res*. 2015;4(1):25-30.
91. Kanoriya D, Pradeep AR, Garg V, Singhal S. Mandibular Degree II Furcation Defects Treatment With Platelet Rich Fibrin and 1% Alendronate Gel Combination: A Randomized Controlled Clinical Trial. *J Periodontol*. 2016;88(3):1-13.

92. Ustaoglu G, Ercan E, Tunali M. The role of titanium-prepared platelet-rich fibrin in palatal mucosal wound healing and histoconduction. *Acta Odontol Scand.* 2016;74(7):558-564.
93. Shivashankar VY, Johns DA, Kumar MR. Platelet Rich Fibrin in the revitalization of tooth with necrotic pulp and open apex. *J Conserv Dent.* 2017;15(4):395-398.
94. Peck MT, Marnewick J, Stephen L. Case Report Alveolar Ridge Preservation Using Leukocyte and Platelet-Rich Fibrin : A Report of a Case. *Case Rep Dent.* 2011;2011(Figure 1):1-5.
95. F. Inchingolo, M. Tatullo, M. Marrelli, A.M. Inchingolo, S. Scacco, A.D. Inchingolo G. Dipalma, D. Vermesan°, A. Abbinante R. Cagiano. Trial with Platelet-Rich Fibrin and Bio-Oss used as grafting materials in the treatment of the severe maxillar bone atrophy : clinical and radiological evaluations. *Eur Rev Med Pharmacol Sci.* 2010;14:1075-1084.
96. Aroca S, Keglevich T, Barbieri B, Gera I, Etienne D. Clinical Evaluation of a Modified Coronally Advanced Flap Alone or in Combination With a Platelet-Rich Fibrin Membrane for the Treatment of Adjacent Multiple Gingival Recessions : A 6-Month Study. *J Periodontol.* 2009;80(2).
97. Jankovic S. The coronally advanced flap in combination with platelet-rich fibrin (PRF) and enamel matrix derivative in the Treatment of Gingival Recession : A Comparative Study. *Eur J Esthet Dent.* 2010;5(3).
98. Lekovic V, Milinkovic I, Aleksic Z, Jankovic S, Stankovic P, Eb K. Platelet-rich fibrin and bovine porous bone mineral vs . platelet-rich fibrin in the treatment of intrabony periodontal defects. *J Periodontal Res.* 2012;47:409-417.
99. Bains R, Bains VK, Loomba K, Verma K, Nasir A. Management of pulpal floor perforation and grade II Furcation involvement using mineral trioxide aggregate and platelet rich fibrin : A clinical report. *Contemp Clin Dent.* 2012;3(Suppl 2):223-227.
100. Schwarz F, Sculean A. Comparison of naturally occurring and ligature-induced peri-implantitis bone defects in humans and dogs. *Clin Oral Impl Res.* 2007;18:161-170.
101. Demling RH. The role of anabolic hormones for wound healing in catabolic states. *J Burns Wounds.* 2005;4:46-59.
102. El Bagdadi K, Kubesch A, Yu X, et al. Reduction of relative centrifugal forces increases growth factor release within solid platelet-rich-fibrin (PRF)-based matrices: a proof of concept of LSCC (low speed centrifugation concept). *Eur J Trauma Emerg Surg.* 2017.
103. Al-maawi S, Orlowska A, Sader R, Kirkpatrick CJ, Ghanaati S. In vivo cellular reactions to different biomaterials — Physiological and pathological aspects and their consequences. *Semin Immunol J.* 2017.
104. Esposito M, Grusovin MG, Esposito M. Treatment of peri-implantitis: what interventions are effective? A Cochrane systematic review. *Eur J Oral Implantol.* 2012;5.

1
2
3
4
5
6
7
8
9
10
11
12
13
14
15
16
17
18
19
20
21
22

Figures

Fig. 1: Distribution of all included articles by field of application and the level of scientific evidence.

Fig. 2: Flowchart of the study design and results.

Fig. 3: The number of published PRF articles in clinical dentistry over the last decade. The literature review was conducted in May 2017.

Fig. 4: The world map shows the tendency towards research on PRF. The circle sizes correspond to the sum of articles collected in this literature review, related to each country.

Fig. 5: The use of PRF as sole filling material vs. conventional open access flap.


Fig. 6: The use of PRF with biomaterial vs. biomaterials alone.

Tables



Table 1: All collected data were organized by field of dentistry. The numbers in bracket [] match with references included in the article. *Indications: (BR/A) Bone regeneration and augmentation, (PC) pain control, (PRA) periodontal attachment regeneration, (R/A) revascularization/apexification of non-vital tooth, (SD) surgical depigmentation, (SL) sinus lift, (SP) socket preservation and (WH) wound healing.


Table 2: Classification of the level of scientific evidence, according to the US Agency for Healthcare Research and Quality.


Table 3: The mixing ratio of PRF and the autologous grafts and/or biomaterials cross-referenced.




Click here to access/download
Rebuttal Letter (for revisions)
cover letter (002).docx














Pocket depth (PD) improvement




Clinical attachment level (CAL) improvement



Pocket depth (PD) improvement



Clinical attachment level (CAL) improvement



Name of the First Author	Dental Field *(indication)	Diagnosis	Treatment	Results	Patients	Follow up Months	Level of Scientific Evidence
I.B. GEETA (2013) India [29]	Endodontics (R/A) of non-vital tooth	Chronic periapical abscess	PRF was placed into the canals. 1. Revascularization of non-vital immature teeth, 2. Regenerative pulpotomy, 3. Apical barrier for apexification of non-vital immature tooth, 4. Periapical surgery	1. Induce faster periapical healing 2. Ideal scaffold in revascularization of immature permanent teeth with necrotic pulps	4	18	III
MEENU DHIMAN (2016) India [30]	Endodontics (R/A) of non-vital tooth	Suppurative chronic apical periodontitis and apicomarginal communication	Peri-radicular curettage enucleation Test group: PRF gel was carefully placed into the cavity. Control group: without PRF	PRF group showed a statistically significant ($P = .046$) better improvement in PD than the control group. Within the limits of this study, it can be concluded that a high success rate may be attained in apicomarginal defects with endodontic microsurgery. PRF may not necessarily improve the outcome.	30	12	Ib
SMITA SINGH (2016) India [31]	Endodontics (BR/A)	Periradicular lesions of endodontic origin	Peri-radicular curettage enucleation PRF gel was carefully placed into the cavity.	Radiographically all patients showed complete bone regeneration at the end of six months. It requires around 1 year for complete healing to occur after the periapical surgery while with the use of PRF, healing is fastened and requires approx. 6 months for complete regeneration of bone.	15	6	III
DAYALAN SUBASH (2016) India [32]	Endodontics (R/A) of non-vital tooth	necrotic pulp and symptomatic apical periodontitis	The fragmented PRF membrane was placed in the pulp chamber and pushed apically with the help of endodontic pluggers.	Radiographic examination revealed resolution of periapical lesion, thickening of the dentinal walls, root lengthening with apical closure showing a continuous lamina dura.	1	12	III
HENGAMEH BAKHTIAR (2016) UK [33]	Endodontics (R/A) of non-vital tooth	Immature teeth with necrotic pulps. acute apical abscess	The liquid that was obtained by compression of the PRF clot was used to irrigate the root canal. The PRF membrane was inserted inside the root canal.	The tooth was asymptomatic at the six-month follow-ups. Parallel radiographic examination revealed no periapical lesion at the one-month follow-up. The initiation of apical closure was obvious by the three-month follow-up. Signs of apical closure were observed after 6 months. Thickening of dentinal walls and complete apical closure were noted over 18 months.	4	18	III
Vasundara Yayathi Shivashankar (2013) India [93]	Endodontics (R/A)	Necrotic pulp and open apex	The PRF was condensed into the canal using a finger plugger till the level the cemento-enamel junction. Grey MTA was placed directly over the PRF to a thickness of 3mm. The patient was recalled after 3 days and the setting of MTA was confirmed. The access cavity was then double sealed with GIC and Composite restoration	The tooth #8 showed negative response to percussion and palpation tests and responded positive to CO2 ice or an electric pulp tester (EPT). Radiograph revealed continued thickening of the dentinal walls, root lengthening, regression of the periapical lesion and apical closure.	1	12	III
ALAIN SIMONPIERI (2011) France [34]	Implantology (SL)	Severe maxillary resorption	Sinus lift -PRF clots and membranes as sole filling material during lateral sinus-lift with immediate implantation	A total of 52 implants were placed. The main results in this case series were that no implant was lost during this 6-year experience and that the vertical bone gain was always substantial and stable. The final bone gain was always very significant, between 8.5 and 12 mm bone gain (10.4 ± 1.2).	20	72	Ila
YU ZHANG (2012) China [35]	Implantology (SL)	Maxillary bone atrophy with a residual crest height of less than 5mm	Sinus lift – Test group: A mixture of deproteinized bovine bone mineral and PRF preparation was inserted in sinuses. Control group: the sinuses were grafted with deproteinized bovine bone alone.	The percentage of newly formed bone in PRF group was about 1.4 times that of the control group (18.35%, 5.62% vs. 12.95% 5.33%). No statistically significant difference between groups.	10	6	Ila
BURAK HAMZACEBI (2015) Turkey [36]	Implantology (BR/A)	Peri-implant bone defect	Flap surgery group (control) and flap surgery + PRF group (PRF). Full thickness mucoperiosteal flap, debridement, implant surface decontamination and filling with PRF membranes.	100% survival of all implants. After 6 months, the PRF group showed a statistically significantly higher reduction in PD and gain in CAL than the control group. Clinically more effective than open flap alone regardless of the defect configuration.	19	6	Ila
ELIF ONCU (2015) Turkey [37]	Implantology (BR/A)	Partially or fully edentulous	Test: implants, PRF+ group Control: implants without PRF. PRF membranes were applied to one of the implant osteotomies. The related implant was thoroughly rinsed with PRF liquid and inserted.	At the end of the first month, the mean ISQ in the PRF+ group in type 2 bone was significantly higher. Secondary stability increases over time. PRF enhanced the stability of implants	20	1	Ilb
NILUFER BOLUKBASI (2015) Turkey [38]	Implantology (SL)	Atrophic posterior maxilla	Sinus lift – Test group: Bovine bone graft and PRF. Control group: resorbable collagen membrane and bovine bone graft	Overall implant survival rate was 100% in both test and control groups. Bone height gain and resorption were calculated radiologically in the test and control groups, the differences were not statistically significant ($p=0.093$).	25	24	Ila
PRIYANKA BOORA (2015) India [39]	Implantology (BR/A)	Partially edentulous in anterior maxilla	Group I: PRF group (The immediately prepared PRF was placed over the surgical site) and Group II: Non-PRF group. One stage non-functional immediate prosthetic protocol.	In PRF group, statistically significant crestal bone level changes were noted within three months. The amount of crestal bone level changes as exhibited in study group had statistically significant lesser mean value than control group. Intergroup comparison for probing depth and bleeding and probing at one month and three months was statistically insignificant.	20	8	Ib
TROEDHAN ANGELO (2015) Germany [41]	Implantology (BR/A)	Maxillary atrophy	Group 1: Particles of 40% beta-tricalcium phosphate (beta-TCP) and 60% hydroxyapatite (HA). Group 2: Microporous particles of pure beta-TCP, group 3: beta-TCP and Platelet Rich Fibrin placed subperiosteal	All subperiosteal tunnel augmentation sites presented sufficient even buccal bone gain in radiographic follow-up. The results of the current study back the experimental findings concerning the advantages for more reliable bone regeneration when A-PRF is used and suggest A-PRF to enhance biomechanical bone quality to a constant higher level in the clinical routine in conjunction with Piezotome surgery and BiSHB grafts. No statistical significant differences were found ITVs.	82	7	Ila
JULIA HEHN (2016) Germany [42]	Implantology (BR/A)	Patients aged 18+, who required an implant in the posterior mandible, were eligible for this study	Bone augmentation and tissue thickening – titanium implants were inserted at bone level with primary stability. In the test group, the tissue was augmented with a PRF membrane using a double layered technique. In the control group, the implant treatment was realized without mucosa thickening.	Though surgical flaps were all sutured completely free of tension, a post-operative dehiscence above the implant could be observed in all test patients within the first week. This process resulted in a complete loss of mucosal and augmented tissues above the implant. The open areas were healed by secondary intention. All implants were clinically osseointegrated and stable and showed no sign of infection. The comparison of the differences in bone loss showed no significance between the control and test groups.	31	6	Ib
TAKEO KANAYAMA (2016) Japan [40]	Implantology (SL)	Atrophic posterior maxilla	Sinus lift – Floor elevation procedures by the crestal approach. PRF as sole filling material	This 1-year prospective study showed that the use of PRF as the sole grafting material during simultaneous sinus lift and implantation is a safe and reliable method.	27	12	III

Mogammad Thabit Peck (2011) South Africa [94]	Implantology (BR/A)	Partially edentulous	Alveolar ridge preservation - Tooth roots were extracted atraumatically. A total of three L-PRF membranes were formed and inserted into the extraction socket site.	The extraction site showed signs of healing with no evidence of residual inflammation. The site was free of infection and the L-PRF membrane was still clearly visible. Even though it remained exposed to the oral environment, there were no signs of membrane disintegration or infection. The patient also reported minimal pain during the postoperative period. At implant insertion, the quality of the newly formed bone was such that it allowed for the implant to be inserted at an insertion torque of more than 35 Ncm.	1	3	III
JOSEPH CHOUKROU N (2006) France [21]	Maxillofacial surgery (SL)	Maxillary atrophy	Sinus lift - Control group (FDBA without PRF). Test group (FDBA with PRF) filling with freeze-dried bone allograft (FDBA) + PRF	The histomorphometric results of control group (FDBA without PRF) after 8 months appear equivalent to those of the test group (FDBA with PRF) after 4 months. It possible to consider sinus floor augmentation surgery with a shorter healing period before implant placement (4 months instead of 8 months). PRF membranes appear to be able to treat sinus membrane perforation and permit the surgery to be completed.	9	8	Ila
EMANUELE RUGA (2011) Italy [45]	Maxillofacial surgery (SP) (PC)	Impacted third molar	Socket preservation - Control: No PRF test: PRF was placed into the extraction socket	In the study group less pain was referred and hygienic maintenance was facilitated by a more rapid alveolar socket fulfillment; consequently, distal periodontal condition was improved also. The healing processes seemed to be accelerated, and the patient's postoperative comfort seemed to be positively influenced	14	6	Iib
GILBERTO SAMMARTIN O (2011) Italy [48]	Maxillofacial surgery (SP)	Records of mechanical heart valve substitution. Tooth extraction	Socket preservation - PRF was placed into the extraction socket	There were only 2 hemorrhagic complications (4%); these occurred in 2 patients with an INR of 3.7, a 61-year-old man and a 54-year-old woman.	50	7 days	Ila
ABHISHEK SINGH (2012) India [43]	Maxillofacial surgery (SP)	Impacted third molar	Socket preservation - Test group: PRF was placed into the extraction socket. Control: Without PRF	Healing Index of Soft Tissue: There was significant difference between study and control site in all 20 patients. Assessment of bone density after 12 weeks showed that average gray scale value for PRF (study) site was comparatively higher than non-PRF (control) site. There was no significant difference between study and control site.	20	3	Ila
MARCO TATULLO (2012) Italy [50]	Maxillofacial surgery (SL)	Maxillary atrophy	Sinus lift - Control: Deproteinized bovine bone and test: amorphous and membranous PRF together with deproteinized bovine bone.	PRF leads to the production of new bone, already 106 days after reconstructive surgery. No statistically significant differences (p > 0.05) regarding ISQ mean values were found between test groups and control groups in each of protocol we performed.	60	36	Ila
SIDIKA SINEM SOYDAN (2013) Turkey [53]	Maxillofacial surgery (WH) (SP)	Medication-Related Osteonecrosis of the Jaw (MRONJ)	Socket preservation - PRF was placed into the extraction socket	Total bone closure was achieved and new mucosa was visible. No gingival loss, inflammation, or infection was detected at the postoperative 6-month follow-up, and the patient was able to use his dentures 1 month after the procedure.	1	6	III
SRISURANG SUTTAPREY ASRI (2013) Thailand [54]	Maxillofacial surgery (SP)	Post extraction socket	Socket preservation - Tooth extraction. Test group: The PRF was filled in the socket. Control group: The alveoli were filled with natural blood clot.	Soft-tissue healing: No statistically significant differences were detected between the groups. Alveolar Ridge Contour: statistical significance at the first week (T1) between the PRF and the control group. Radiographic Evaluation: no statistically significant differences were detected among the groups.	8	2	Ila
GAETANO MARENZI (2014) Italy [46]	Maxillofacial surgery (SP)	Root fracture, residual root, periapical granuloma, and orthodontic reason.	Socket preservation - Control: Natural healing. Study: The sockets were filled with L-PRF	All patients completed the study. No cases of bleeding, infection, alveolar osteitis, or other surgical complications were reported. Comparisons between values relative to the study and control sides showed better healing and faster socket closure for the side treated with L-PRF, with differences statistically significant at days 3 and 7.	26	1	Ila
NILIMA KUMAR (2014) India [51]	Maxillofacial surgery (SP) (PC)	Impacted third molar	Socket preservation - Test: PRF was placed into the extraction socket, followed by flap approximation. Control: Only primary closure	Difference between the two groups was statistically significant (p value 0.017) indicating that the application of PRF in the extraction socket aided in reducing the patient's post-operative pain, swelling and interincisal distance.	31	3	Ib
TEJESH YELAMALI (2014) India [55]	Maxillofacial surgery (SP)	Impacted third molar	Socket preservation - Split mouth experiment. PRF was placed on the right side and PRP on the left side.	Healing capability for PRF group were significantly higher as compared to PRP group. The mean values of bone density for PRF groups were significantly higher as compared to PRP groups.	20	4	Iib
AMOL M DOIPHODE (2015) India [44]	Maxillofacial surgery (SP)	Impacted third molar	Socket preservation - Group I: control. Group IIa: filled with PRP gel. Group IIb: filled with PRF	Group I showed dehiscence in 5 (33.3%) out of 15 cases. Group IIa showed dehiscence in 3 (20%) cases whereas Group IIb (PRF) did not show any sign of wound dehiscence. The difference in the decrease in ABH at 2, 4, and 6 months postoperatively in the group IIb was statistically significant.	30	6	Ila
GEORGIOS KOTSAKIS (2015) USA [47]	Maxillofacial surgery (SP) (PC)	Maxillary uperright first premolar. class I mobility	Socket preservation - Tooth extraction, placement of a PRF plug and membrane in the socket. 8 weeks later implants were places for 'early accelerated' implant placement"	The implant remained successfully in function at the 48-month follow-up, with no signs of mobility or inflammation and no subjective symptoms, such as pain, or altered sensation.	1	48	III
HAZEM SHAWKY (2015) Egypt [49]	Maxillofacial surgery (BR/A)	Alveolar cleft	Bone regeneration - Group A: Maxillary alveolar cleft reconstruction with anterior iliac crest bone graft combined with PRF. Group B: Autogenous anterior iliac crest bone graft.	No recurrence of the oronasal fistulae and no signs of infection or evidence of bone graft loss. Group A: The mean amount of newly formed bone volume was 0.78 cm ³ and the percentage of newly formed bone was 82.6%. Group B The mean amount of newly formed bone volume was 0.62 cm ³ , and the mean percentage of newly formed bone was 68.38%. There was a statistically significant increase in the percentage of newly formed bone in group A.	24	6	Ib
S.E NORHOLT (2016) Denmark [52]	Maxillofacial surgery (WH)	Medication Related Osteonecrosis of the Jaw (MRONJ)	Elevation of a mucoperiosteal flap. Necrotic bone was removed and bone surface was smoothed. Bone covered with PRF	Successful outcome was defined as complete mucosal healing and no symptoms from the jaw. The outcome of the surgical treatment was successful in 14 of the 15 patients (93%). In conclusion, 14 of the 15 patients in this study had a successful outcome following the surgical treatment of ONJ with the use of PRF membranes to ensure a multilayered closure	15	6	Iib
YUVIKA RAJ KUMAR (2016) India [56]	Maxillofacial surgery (SP)	Impacted third molar	Socket preservation - Test: PRF was placed into the extraction socket. Control: Without PRF	These numbers show that platelet-rich fibrin gradually increases the osseous healing at every follow-up, and the mean value remains consistently higher in the experimental group. The values for fractal analysis were 0.03 (95% CI -0.02 to 0.08) higher in the experimental group across the time points compared with the control, but the difference was not significant.	34	6	III
MICHAL KAPUSTECKI (2016) Poland [57]	Maxillofacial surgery (BR/A)	Oroantral communication and fistula	Bone regeneration - Collection of monocortical bone blocks from mental protuberance or mandible oblique line. The graft was stabilized using a bicortical screw. The graft and surrounding bone were covered with a PRF membrane.	Complete healing of the operated area in 18 patients. In the study group in all cases closure of the oroantral communication was observed. Mean measurement alveolar width values in a group of 20 patients= Post-extraction alveolus (mm): 13.5 and control 13. Mean measurement alveolar height values in a group of 20 patients = Post-extraction alveolus (mm): 13 and control: 12.5.	20	6	III

ANDREAS ANWANDTER (2016) Chile [58]	Maxillofacial surgery (SP)	un-restorable caries lesions and advanced periodontal disease	Socket preservation – The teeth were extracted. The socket was completely filled with several L-PRF clots. Finally L-PRF membranes were positioned over the clots, inside the socket.	The mean horizontal bone loss was 1.56 mm. A total horizontal resorption of 1.53 mm was reported at the alveolar crest. During the first 4 months after tooth extraction the sockets reduced 4.5 mm in depth. The center of the socket had a significant filling of 5.72±3.6 mm (p = 0.0001) and the oral cortical plate had a mean vertical gain of 0.09 mm 1.57 mm (p = 0.9). This would bring L-PRF's effectiveness at the same level as these osseous substitutes.	18	4	I Ib
ZEYNEP BURÇIN GÖNEN (2016) Turkey [61]	Maxillofacial surgery (WH)	Medication-related osteonecrosis of the jaw Stage-3 (MRONJ)	Bone regeneration – One layer of PRF was laid into the alveolar bone cavity. The second layer was placed superficially and sutured to the surrounding gingiva. Acellular plasma was injected submucosally around the wound after the procedure.	Epithelization occurred in the second week postoperatively, and there was no infection or inflammation of the gingival tissues. New gingival tissue formation was observed after four weeks, and healing was uneventful without any paresthesia. Three months after the surgical procedure, whole closure of the exposed bone with new gingival tissue was achieved.	1	12	III
GOKHAN GÜRLER (2016) Turkey [62]	Maxillofacial surgery (SL) (PC)	Posterior maxillary bone height of less than <5 mm.	Sinus lift – Study group: allogeneic freeze dried corticocancellous bone chips mixed with L-PRF. Bony sinus windows were covered with two layers of PRF membranes. Control: Allogeneic bone graft and bony sinus windows were covered with a resorbable collagen membrane	There were gradual improvements in postoperative pain, swelling, sleeping, eating, phonetics, activities of daily living, and missed work days in the L-PRF group, but the differences between the groups were not significant. HI scores of the L-PRF group (4.2±0.9) were higher than that of the control group (3.6±0.7) on the 7th (4.7±0.4) and 14th postoperative days (4.4±0.5); differences were not statistically significant.	24	14	I Ia
ANTONIO CORTESE (2016) Italy [63]	Maxillofacial surgery (BR/A)	Reabsorbed edentulous ridge secondary to previous extraction	Bone augmentation – Split crest flapless new technique for implant insertion. Test group: autologous platelet rich fibrin (PRF) has been used to fill the osteotomy gap or simply as regenerative material. Control: traditional technique	All patients underwent an uneventful implant surgery. At T1 mean height loss was of 0 mm and of 1.2 mm at T3 for group 1 (new flapless technique) while for group 2 patients (traditional implant insertion technique) mean height bone loss was 2.4 mm at T1 and of 3.4 at T3. Mean difference for height bone loss between the two groups of patients was 2.4 mm at T1 and 2.2 mm at T3.	10	6	I Ib
TAKUYA ASAKA (2016) Japan [64]	Maxillofacial surgery (WH)	Medication-related osteonecrosis of the jaw (MRONJ)	Socket preservation – PRF group (prospective) and a control group (retrospective). PRF was laid directly over the bone to fill the socket, which was then sutured using 4.0 nylon sutures	Postoperative infections and the onset of MRONJ, characterized by exposed necrotic bone or fistulae, were also absent at the time of follow-up in all patients. All sockets were filled with granulation tissue within 2 weeks and completely epithelized by 4 weeks. Radiographic bone regeneration as a radiopaque rise was observed after 3 months in all PRF cases. Early epithelization was confirmed in all PRF patients including patients receiving steroids and immunosuppressants. The prevalence of delayed recovery was significantly higher in the control group (P < 0.05) than in the PRF group.	102	3	I Ib
ANDY TEMMERMAN (2016) Chile [66]	Maxillofacial surgery (SP)	symmetrical bilateral tooth extractions in the maxilla or mandible	Socket preservation – At the test site 2–5 PRF clots, depending on the size of the socket, were inserted and compressed with a large plunger. The site was covered with 2–3 L-PRF membranes. At the control site, a cross-suture was performed to stabilize the coagulum	Statically significant differences (p = 0.004) were found in the percentage of socket fill between test [94.7% (26.9)] and control sites [63.3% (31.9)].	22	3	I b
MANZOOR DAR (2016) India [68]	Maxillofacial surgery (BR/A)	Cystic lesions	Bone regeneration – All cystic lesions were enucleated and PRF was placed in the bony defects	Radiographically all patients showed that PRF promotes faster osseous regeneration within the 3rd postoperative month, and within 6th postoperative month, complete bone regeneration was seen.	20	6	III
MAHMOUD MOUSSA (2016) Egypt [80]	Maxillofacial surgery (BR/A)	Horizontally deficient edentulous maxillary anterior ridges (≤ 4.5 mm)	Bone augmentation – Test group: Palatal Bone Block covered with PRF. Control group: only bone block graft.	PRF showed superior results when used to cover the palatal bone block, decreasing the block's surface resorption when compared with only using bone block graft. The test group showed statistically significantly lower mean graft resorption than the control group (test, 0.8 ± 0.6 mm; control, 1.6 ± 0.9 mm; P = .006)	12	4	I Ia
ELIF ÖNCÜ (2017) Turkey [60]	Maxillofacial surgery (BR/A)	Indications for dental extraction	Bone regeneration – After the extraction, using split-mouth design, test sockets were coated with L-PRF and control sockets without L-PRF.	Results showed a statistically significant difference between the stability of L-PRF+ and L-PRF– implants at 1 week and at 1 month. Mean marginal bone resorption was higher in the control group at 1 year.	26	12	I Ib
JUNG-HYUN PARK (2017) Korea [59]	Maxillofacial surgery (WH)	Medication-related osteonecrosis of the jaw (MRONJ)	Bone regeneration and healing of tissue – Group 1: L-PRF. Group 2: L-PRF+rhBMP2 L-PRF was placed at the bone surface and primary closure of the mucoperiosteal flap was performed.	L-PRF (n = 25): Complete resolution, n 9 (36.0%), Delayed resolution 13 (52.0%), No resolution n 3 (12.0%) L-PRF Plus+ BMP-2 (n = 30) Complete resolution: 18 (60.0%), Delayed resolution: 11 (36.7%), No resolution 1 (3.3%). statistically significant compared with that of the therapy using L-PRF alone (P = .028)	55	6	I Ia
MATHEW P. VARGHESE (2017) India [65]	Maxillofacial surgery (SP)	impacted mandibular third molars	Socket preservation – Randomization was performed after extraction. On one side the socket was sutured primarily (control site); on the other side, autologous PRF gel was placed and then the socket was sutured (test site).	The average percentage of bone fill in the PRF group was 57.90 (SD, 26.789) and that of the non-PRF group was 46.74 (SD, 17.713; P < .05). Soft tissue healing as evaluated by the healing index of Landry et al also was found to be better at the PRF test site and it was statistically significant (P < .05).	30	3	I Ia
KANI BILGINAYLAR (2017) Turkey [67]	Maxillofacial surgery (BR/A)	Odontogenic Orocutaneous Fistula	Bone regeneration – The treatment started with root canal treatment. 2 days after the filling of the root canal, apical resection was performed and platelet-rich fibrin (PRF) was administered into the bone cavity and a PRF membrane was used to close the site as well.	At the third month follow-up, radiographically, intraoral, and extra oral examinations revealed complete healing of bone at periapical lesion area and spontaneous healing of orocutaneous fistula.	1	3	III
F.INCHINGOLO (2010) Italy [95]	Maxillofacial surgery (SL) (BR/A)	Severe maxillar bone atrophy	Surgical procedure of “one-stage sinus lift” (implant insertion occurred concurrently with sinus lift, and the healing and integration time was 6-9 months). Before inserting the implant, a small quantity of filling material was placed in the cavity (Bio-Oss and PRF).	In all the cases included in this protocol, the Authors observed a successful implant-prosthetic rehabilitation. All patients reported no pain to percussion, no sign of tissue suffering to the soft peri-implant tissues and the presence of an optimal primary stability of the inserted implants. Sn average increase in the peri-implant bone density of 31%	23	6	I Ib

FRANCISCO MUÑOZ (2016) Chile [69]	Orthodontics (BR/A)	Moderate to severe crowding, Class II or III malocclusions	PAOO + (PRF+ Puros®/Bio-Oss® graft+ Metronidazole)	Our results support the wound healing and osteogenic properties of L-PRF, yet also suggest that the healing capacity of L-PRF may exceed the limits reported. This combinatorial approach seems to reduce the risk of tear/dehiscence of the flap with subsequent graft exposure and contamination, providing an optimal environment for wound healing.	11	24	III
ANUJ SHARMA (2011) India [72]	Periodontology (PAR)	Intra-Bony defects Grade II furcation defect	Randomly allocated and treated either with (test group) autologous PRF and OFD or (control group) OFD.	Test sites presented with a significantly greater reduction in PD than control sites, with a difference of 2.17 mm. The gain in attachment level (CAL) significantly greater in the test sites. Test sites presented with a significantly greater vertical defect fill (50.8 – 6.24) than the control sites (16.7 – 6.42) at 9 months	18	9	Ib
ANUJ SHARMA (2011) India [74]	Periodontology (PAR)	Chronic Periodontitis	(Test group) Autologous PRF with OFD or (control group) OFD alone in the treatment of 3-wall IBDs.	Test sites presented with a significantly greater PD reduction (4.55 ± 1.87 mm) than control site. The PAL gain was also greater in test sites (3.31 ± 1.76) compared to the control group (2.77 ± 1.44 mm). Test sites (48.2% – 5.72%) presented with a significantly greater IBD fill than did control sites.	32	9	Ib
THORAT M.K. (2011) India [85]	Periodontology (PAR)	Intra-Bony defects	Conventional flap surgery with autologous PRF (test group), OFD alone (control group). PRF in the treatment of intra bony defects in chronic periodontitis patients	The mean PD and CAL at baseline and after 9 month in control and test group was 6.75 ± 1.69 , 6.50 ± 1.75 and 3.19 ± 1.52 , 4.38 ± 2.16 mm, respectively, and was found to be significant ($p<0.01$). IBD at baseline (4.40 ± 1.04) and test group (4.52 ± 1.11) was compared, it was found to be non-significant at baseline ($p<0.05$) and was significant at 9-month comparison ($p<0.05$).	32	9	Ib
Y-C CHANG (2011) Taiwan [86]	Periodontology (PAR)	Chronic periodontitis Infrabony defects	Minced PRF with bioactive glass were applied to the defect walls and root surfaces. The PRF membrane was adapted over the grafted defect and above the CEJ	These results indicated that PRF may contribute to the differentiation of human PDLFs into osteoblasts. In this study, PRF enhanced p-ERK, OPG and ALP expression. The application of PRF for periodontal osseous defects achieves probing depth reduction, clinical attachment gain and radiographic defect fills.	6	12	Ila
SASHA JANKOVIC (2012) Serbia [82]	Periodontology (PAR) (PC)	Gingival recession Miller Class I or II	On one side, the gingival recession was treated with a CAF and PRF membrane (PRF group). The other side was treated with a CTG in combination with a CAF (control).	No side effects were reported. VRD, CAL and PD differences between the groups were not statistically significant at baseline or 6 months. Healing index: Results obtained in the PRF group were statistically superior in reference with data recorded in the CTG group ($P < .05$). The pain intensity was statistically different between groups for the first 7 days, favoring the PRF group.	15	6	Ib
CHHAYA BANSAL (2013) India [77]	Periodontology (PAR)	Chronic periodontitis Interproximal intrabony defects	Mucoperiosteal flap elevation followed by the placement of Group I: DFDBA alone. In Group II, mixture of PRF with DFDBA was done.	A combination of PRF with DFDBA demonstrated better results in probing pocket depth reduction and clinical attachment level gain as compared to DFDBA alone in the treatment of periodontal intrabony defects. On comparison, the difference between the two groups was statistically no significant at all time intervals.	10	6	Iib
PAVAN BAJAJ (2013) India [81]	Periodontology (PAR)	Chronic periodontitis Mandibular degree II furcation defects.	PRF with OFD or autologous PRP with OFD, or OFD alone. Treatment of degree II furcation defects	Both PRF and PRP sites presented with a significantly greater PD reduction (4.29 ± 1.04 mm, 3.92 ± 0.93 mm respectively) than control site (1.58 ± 1.02 mm) at 9 month postoperatively ($p < 0.05$). RVCAL gain was also greater in the PRF (2.87 ± 0.85 mm) and PRP (2.71 ± 1.04 mm) sites as compared to control site (1.37 ± 0.58 mm). There was no significant difference between PRP and PRF groups.	42	9	Ib
ROSAMMA JOSHEP (2014) India [82]	Periodontology (PAR)	Chronic periodontitis	PRF gel (experimental group I). PRF gel and PRF membrane (experimental group II). OFD alone (Control group).	The magnitude of changes were higher for both the experimental groups compared to control ($p<0.05$) and there was no significance difference between the experimental groups ($p>0.05$). radiographic bone levels RCH (-1 ± 2.8 mm in control, 1 ± 4.7 mm in group I, 0.33 ± 2.28 mm in group II) at baseline and nine months postsurgery showed no significant differences in all groups ($p>0.05$). The negative value for RCH in control group implies resorption of alveolar bone postoperatively compared to baseline.	15	9	Ila
A R PRADEEP (2015) India [70]	Periodontology (PAR)	Chronic periodontitis Interproximal intrabony defects	Four treatment groups: OFD alone, OFD with PRF, OFD with 1% MF and OFD + PRF+1% MF. Treatment of 2/3 wall intrabony defects (IBDs) with OFD + PRF +1% MF gel	All treated cases showed uneventful wound healing and the drug was well tolerated by all the patients. Greater PD reduction and RAL gain seen in PRF+1% MF group. IBD depth reduction in MF group and PRF group was not statistically significant when compared to each other.	126	9	Iib
A R PRADEEP (2015) India [71]	Periodontology (PAR)	Chronic periodontitis	OFD + Placebo gel (group 1), PRF+HA with OFD (group 2), RSV 1.2mg gel+PRF+HA with OFD (group 3).	Results of our study suggest that, furcation defects when treated simultaneously with RSV 1.2 mg+HA+PRF (group 3) showed significantly positive correlation with PD reduction (4.62 ± 1.03 mm) and RVAL, RHAL gain (4.17 ± 0.70 and 4.05 ± 0.76 mm, respectively) with greater defect depth reduction (3.68 ± 0.32 mm) and % bone defect fill (61.94 ± 3.54 %).	120	9	Ib
ASHISH AGARWAL (2015) India [75]	Periodontology (PAR)	Chronic periodontitis Interproximal intrabony defects	Test group: PRF/DFDBA and control group: the DFDBA/saline. DFDBA was mixed with PRF at a proportion of 1:1 (v/v) and filled into the defect. A membrane of compressed PRF was trimmed and adapted over the grafted defects	PRF/DFDBA group exhibited statistically significantly greater changes compared with the DFDBA/saline group in PD (4.15 ± 0.84 vs 3.60 ± 0.51 mm), CAL (3.73 ± 0.74 vs 2.61 ± 0.68 mm), REC (0.47 ± 0.56 vs 1.00 ± 0.61 mm), bone fill (3.50 ± 0.67 vs 2.49 ± 0.64 mm) and defect resolution (3.73 ± 0.63 vs 2.75 ± 0.57 mm)	30	12	Ib
ASHISH MATUR (2015) India [76]	Periodontology (PAR)	Generalized chronic periodontitis	Treated with OFD and filled with PRF (OFD with PRF) or OFD with ABG.	There was significant mean PPD reduction and CAL gain, in both the treated groups; PRF (2.67 ± 1.29 mm, 2.53 ± 1.06 mm) and the ABG treated group (2.4 ± 1.06 mm, 2.67 ± 1.63 mm). No significant difference in mean change after intervention between the two groups was observed for clinical parameters (PI, GI, PPD, CAL, AC, and BD level)	25	6	Ila
GURKIRAT KAUR SHANDU (2015) India [78]	Periodontology (PAR)	Chronic periodontitis Interproximal intrabony defects Grade II furcation defects	The debrided defect was slightly overfilled with Gengigel. PRF was filled into the furcation defect. A GTR membrane was used to cover the furcation. The reflected flap was repositioned over the PRF membrane.	These studies support the fact that the combined use of Gengigel in conjunction with PRF seems to have an added benefit in the regeneration of Class II furcation, as described in this case report.	1	6	III
HIMANSHU AJWANI (2015) India [79]	Periodontology (PAR)	Chronic periodontitis Two and three-wall intrabony defects ≥ 3 mm	The control group consisted of sites treated with OFD alone, whereas test-group sites were treated with OFD with autologous PRF	Intra group and Inter group comparisons showed statistical significant reduction with PD and RAL and no difference was observed with GML levels. Statistically significant improvements were seen with the mean defect fill (cement enamel junction to base of the defect [CEJ-BOD] and AC-BOD) ($P = 0.003^*$) when intra group and inter group comparisons were made. PRF + OFD stimulated a significant improvement in the clinical and radiographic parameters and increase in bone fill compared to OFD alone at 9 months.	20	9	Ib
AHMED Y. GAMAL (2016) Egypt [72]	Periodontology (PAR)	Severe chronic periodontitis	Group 1: bone substitute grafting DBM control group. Group 2: PRGF combined with DBM. Group 3: PRF combined with DBM. Papillary preservation flap, filling with PRF and xenograft in intrabony periodontal defects.	PRF and control group GF levels were found also none significantly different. All groups showed statistically significant improvement in PD reduction, CAL gain and IBD fill compared to baseline data. No statistically significant differences were found between groups.	30	9	Ib

SUMIT KUMAR AGARWAL (2016) India [84]	Periodontology (PAR)	Gingival recession Miller Class I or II	CAF procedures under the microsurgical approach in the management of Miller's Class I and II gingival recession defects with the use of either PRF or amnion membrane (AM) in comparison to CAF alone.	Root coverage was obtained in 52.3% patients, our results reported superior root coverage in PRF treated site as compared to control group. At 6 months follow-up, significant increase in GT measurements in group I (CAF with PRF), whereas, nonsignificant increase for group II (CAF with AM) and no change or decrease for group III (CAF alone) as compared to baseline was observed.	23	6	Ila
MANSI BANSAL (2016) India [87]	Periodontology (SD) (WH) (PC)	Gingiva hyperpigmentation	split mouth design: 1 quadrant was covered with PRF membrane and 2 quadrant was covered with a periodontal dressing (Coe-Pack)	Healing index (HI) After both 3 and 5 days postoperatively, CoePack sites had greater erythematous area as compared to PRF. At 7 days postoperatively, both the sites exhibited normal tissue architecture. After 5 days, two patients complained of slight pain at CoePack site whereas on PRF site none of the patient reported with pain.	5	7 days	Iib
SHRIPARNA BISWAS India (2016) [88]	Periodontology (PAR)	mandibular molar class II furcation defects	Group I: 10 furcation defects were treated using bioactive glass (NovaBone) bone graft putty material. Group II: 10 furcation defects were treated using platelet rich fibrin (PRF).	Both treatments showed improvement in gingival index (GI) and plaque index (PI). There was an overall reduction in both vertical and horizontal probing depth in both groups; however, vertical probing depth reduction in group II showed better statistically significant results than did group I at the end of 3 months ($P=0.0004$), 6 months ($P<0.00001$), and 9 months ($P=0.0004$).	15	9	Iib
YASEMIN SEZGIN (2016) Turkey [89]	Periodontology (PAR)	Generalized, severe, chronic periodontal disease	Split-mouth design. The selected sites were randomly divided into control (ABBM alone) and test (ABBM-PRF) groups. The defect was filled to the existing alveolar crest. PRF was used to cover the defect as a barrier.	Significantly greater CAL gain was detected in the test group than in the control group ($p < 0.05$). In addition, increase in GR was not statistically significant at 6 months after treatment compared to baseline ($p > 0.05$) in the test group, whereas this increase was statistically significant in the control group ($p < 0.05$); Moreover, PD decreased significantly compared to baseline at 6 months after treatment in both groups ($p < 0.05$), but the inter-group differences were not statistically significant at any evaluation time point ($p > 0.05$)	21	6	Ib
NIKHIL D. CHANDRAD AS (2016) India [90]	Periodontology (PAR)	chronic periodontitis – intrabony defect	Group A (PRF with DBM), group B (PRF alone) and group C (OFD). The mixture was delivered into the defect.	Mean PD reduction and RAL gain were greater in group A (4.25 ± 1.48 , 3.92 ± 0.90) and group B (3.82 ± 0.75 , 3.27 ± 0.65) than control (3.00 ± 1.21 , 2.25 ± 0.62). Furthermore, Statistically significant improvement in LBG and %BF was found in group A (3.47 ± 0.53 , 61.53 ± 4.54) compared to group B (2.55 ± 0.61 , 49.60 ± 14.08) and group C (1.21 ± 0.80 , 24.69 ± 15.59).	38	9	Ib
DHARMEND RA KANORIYA (2016) India [91]	Periodontology (PAR)	Mandibular Degree II Furcation Defects	Group 1 treated with open-flap debridement; group 2 treated with access therapy with autologous PRF; and group 3 treated with access therapy and autologous PRF + 1% ALN gel	Reduction in PD was greater in group 3 (4.4 ± 0.57 mm) when compared with group 2 ($3.69 - 0.76$ mm) and group 1 ($2.41 - 0.77$ mm). RVAL and RHAL gain was greater in group 3 ($4.12 - 0.6$ mm and $3.64 - 0.90$ mm) when compared with group 2 ($3.39 - 0.49$ mm and $2.86 - 0.062$ mm) and group 1 ($2.33 - 0.48$ mm and $2.04 - 0.35$ mm). Group 3 showed greater significant radiographic bone fill ($56.01\% - 2.64\%$) as opposed to group 2 ($49.43\% - 3.70\%$) and group 1 ($10.25\% - 3.66\%$).	72	9	Ib
GÜLBAHAR USTA OĞLU (2016) Turkey [92]	Periodontology (WH)	Inadequate attached gingiva and gingival recession defects.	Free gingival graft (FGG) donor sites were treated with T-PRF and compared with an untreated control group. An FGG was obtained and then sutured at the prepared recipient bed. The T-PRF membrane was then prepared to be compatible with the FGG donor site	The T-PRF group showed a significantly higher CWE than did the control group on day 14 (68.7% for the test group; 16.7% for the control group) ($p<0.001$). CWE was observed in all patients in both groups on day 21. PSTT in the T-PRF group was significantly thicker than that in the control group after 6 months of healing (4.51 ± 0.58 mm in the T-PRF group and 3.93 ± 0.69 mm in the control group) ($p<0.05$)	40	21 days	Ib
Sofia Aroca (2009) France [96]	Periodontology (PAR)	Miller Class I and II recession defects	CAFs were performed on both sides of the mouth, either in conjunction with a PRF membrane (test side) or without (contralateral control side)	At 6 months, the difference in root coverage between the two groups was statistically significant: 80.7% – 14.7% and 91.5% – 11.4% for test and control sites, respectively. A significant increase in GTH between baseline and 6 months was observed only in the test group (from 1.1 – 0.4 mm to 1.4 – 0.5 mm.	20	6	Ib
Sasha Jankovic (2010) Serbia [97]	Periodontology (PAR)	Bilateral Miller Class I or II maxillary gingival recession.	Two treatment groups: CAF combine with EMD or CAF combined with PRF. The flap was advanced so as to cover the PRF membrane and sutured.	The mean decrease of GR was not statistically significant. The WKT increase at 12 month was statistically significant favoring the EMD group. PD measurements were not significant. The pain intensity was statistically significantly between groups for the first 5 days, favoring the PRF group. No clinical advantages in between PRF and EMD.	20	12	Ib
Lekovic V. (2012) USA [98]	Periodontology (PAR)	Intrabony defects	PRF–BPBM group and PRF alone group. A membrane of compressed PRF was trimmed and adapted over the grafted defect	Mean pocket reduction in the PRF group was 3.35 ± 0.68 mm on buccal and 3.24 ± 0.73 mm on lingual sites and in the PRF–BPBM group 4.47 ± 0.78 mm on buccal and 4.29 ± 0.82 mm on lingual sites. The differences observed between the two groups were statistically significant in favor of the PRF–BPBM group. The PRF group presented with a clinical attachment gain of 2.24 ± 0.73 mm on buccal sites and 2.12 ± 0.78 mm on lingual sites, while the gain for the PRF–BPBM group was of 3.82 ± 0.78 mm on buccal and 3.71 ± 0.75 mm on lingual sites. The differences in attachment gain observed between the two groups were significantly better in the PRF–BPBM group.	17	6	Ila
Rhythm Bains (India) 2012 [99]	Periodontology (PAR)	Retrograde periodontitis along with Grade II furcation	The furcation was filled with PRF gel mixed with hydroxyapatite graft and over it PRF membrane was placed.	The patient was asymptomatic when he last reported 1½ year after the treatment with significant disappearance of radiolucency on radiographs and probing depth reduction from 10 mm to 3 mm.	1	18	III


ABH: Alveolar bone height	ITVs: Insertion-torque-value
ABBM: Inorganic bovine bone mineral	LBG: Linear bone growth
ABG: Autogenous bone graft	L-PRF: Leukocytes platelet rich fibrin
AC: Alveolar crest	MF: Metformin
ALP: alkaline phosphatase	MRONJ: Medication related osteonecrosis of the jaw
AM: amnion membrane	mSBI: Modified sulcus bleeding index
BD: Bony defects	OFD: Open flap debridement
BF: Bone fill	ONJ: Osteonecrosis of the jaw
BiSHB: Biphasic self-hardening biomaterial	OPG: Osteoprotegerin
BMP: Bone morphogenetic protein	PAOO: Periodontally accelerated osteogenic orthodontics
BPBM: Bovine porous bone mineral	PAL: Periodontal attachment level
CAF: Coronally advanced flap	PD: Pocket depth
CAL: Clinical attachment level	PDGF: Platelet-derived growth factor
CEJ: Cementoenamel junction	PDLF: Periodontal ligament fibroblasts
CTG: Connective tissue graft	p-ERK: Phosphorylated extracellular signal-regulated protein kinase
CWE: Complete wound epithelialization	PI: Plaque index
DBM: Demineralized bone matrix	PPD: Probing pocket depth
DFDBA: Demineralized freeze-dried bone allografts	PRF: Platelet rich fibrin
EMD: Enamel matrix derivative	PRGF: Preparation rich in growth factors
FDDBA: Freeze-dried bone allografts	PRP: Platelet rich plasma
FGG: Free gingival graft	PSTT: palatal soft-tissue thickness
GI: Gingival index	PSTT: phone Crest Height
GI: Gingival index	REC: Mucosal recession

GT: Gingival thickness	RHAL: Relative horizontal attachment level
GTH: Gingival thickness	RHCAL: Relative horizontal clinical attachment level
GTR: Guided tissue regeneration	RVCAL: Relative vertical clinical attachment level
HA: Hydroxyapatite	RSV: Rosuvastatin
HI: Healing index	TGF: Transforming Growth Factor
IBD: Intra-bony defects	VEGF: Vascular endothelial growth factor
INR: International normalized ratio	VRD: Vertical gingival recession depth
ISQ: Implant stability quotients	WKT: Width of the keratinized tissue

Table 3: All the data collected was organized by field of dentistry. The numbers in bracket [] match with references included in the article. *Indications: (BR/A) Bone regeneration and augmentation, (PC) pain control (PRA) periodontal attachment regeneration, (R/A) revascularization/apexification of non-vital tooth, (SD) surgical depigmentation, (SL) sinus lift, (SP) socket preservation and (WH) wound healing.

Level	Type of scientific evidence
Ia	Scientific evidence obtained from meta-analyses of randomized clinical trials
Ib	Scientific evidence obtained from at least one randomized clinical trial
IIa	Scientific evidence obtained from at least one well-designed, non-randomized controlled prospect study
IIb	Scientific evidence obtained from at least one well-designed, quasi-experimental study
III	Scientific evidence obtained from well-designed observational studies, such as comparative studies, correlation studies or case control studies.
IV	Scientific evidence obtained from documents or opinions of experts committees and/or clinical experiences of renowned opinion leaders.



Autologous graft/Biomaterial	Comparison/ Ratio of mixture	Reference
Freeze-dried bone allograft	Not reported	[19]
Deproteinized bovine bone mineral	Not reported	[35]
Bovine bone graft mixture	1:2	[38]
40% beta-tricalcium phosphate (beta-TCP) and 60%hydroxyapatite (HA) beta-tricalcium phosphate (beta-TCP)	Not reported	[41]
Deproteinized bovine bone	Not reported	[50]
Allogeneous freeze dried corticocancellous bone chips	Not reported	[62]
Cortico-Cancellous Particulate Allograft and Deproteinized bovine bone (3:2)	Not reported	[69]
1% Metformin	1:1	[70]
Porous hydroxyapatite + Rosuvastatin 1.2 mg	1:1	[71]
Freeze-dried bone allograft	Not reported	[72]
Decalcified freeze-dried bone allograft (DFDBA)	1:1	[75]
Autogenous bone graft	Compared	[76]
Demineralized freeze dried bone allograft	No reported	[77]
Gengigel - hyaluronic acid	Not reported	[78]
Amnion allograft membrane	Not reported	[84]
Bioactive glass	Not reported	[86]
Bioactive glass	Not mixed	[88]
Anorganic bovine bone mineral	Not reported	[89]
Demineralized bone matrix	Not reported	[90]
1% Alendronate Gel	1:1	[91]
Deproteinized bovine bone	Not reported	[95]
Enamel matrix derivative	Compared	[97]
Bovine porous bone mineral	1:1	[98]
hydroxyapatite graft	Not reported	[99]



[Click here to access/download](#)

Cover Letter

BMQYP4DB-1EB3-52C2-7C65-CEFE-48E9.pdf





[Click here to access/download](#)

Copyright Form

[JOI_Copyright_Form_041317s - Flattened.pdf](#)

