

CORRECTION

Correction: Access to Interdental Brushing in Periodontal Healthy Young Adults: A Cross-Sectional Study

The PLOS ONE staff

The values in Tables $\underline{2}$ and $\underline{4}$ are incorrectly duplicated. In <u>Table 2</u>, lines 1–15 are the same as lines 16–31. Lines 16–31 feature the correct formatting. In <u>Table 4</u>, lines 1–11 are the same as lines 12–22. Lines 12–22 feature the correct formatting. Please see the corrected Tables $\underline{2}$ and $\underline{4}$ here. The publisher apologizes for the error.



G OPEN ACCESS

Citation: The *PLOS ONE* staff (2016) Correction: Access to Interdental Brushing in Periodontal Healthy Young Adults: A Cross-Sectional Study. PLoS ONE 11(6): e0158252. doi:10.1371/journal.pone.0158252

Published: June 21, 2016

Copyright: © 2016 The PLOS ONE staff. This is an open access article distributed under the terms of the Creative Commons Attribution License, which permits unrestricted use, distribution, and reproduction in any medium, provided the original author and source are credited.



Table 2. Distribution (mean ± sd) of the number of interproximal sites according to the diameter of the interproximal brush in 99 adults.

	n	IDB diameter (mm.)						
		0.6	0.7	0.8	0.9	1.1		
All	99	8.78 ± 5.18	10.83 ± 4.56	3.39 ± 3.47	0.83 ± 1.42	0.49 ± 1.66		
Sex								
Male	55	8.27 ± 5.07	10.80 ± 4.50	3.74 ± 3.52	74 ± 3.52 0.98 ± 1.51			
Female	44	9.41 ± 5.31	10.86 ± 4.68	2.95 ± 3.38	0.64 ± 1.30	0.36 ± 1.06		
p-value		0.280	0.944	0.260	0.229	0.510		
Smoker								
No	75	9.03 ± 5.29	10.83 ± 4.35	3.11 ± 3.15	0.79 ± 1.33	0.37 ± 0.98		
Yes	24	8.00 ± 4.86	10.83 ± 5.26	4.29 ± 4.27	0.96 ± 1.71	0.87 ± 2.89		
p-value		0.594	0.991	0.142	0.615	0.195		
Subject's Baseline Bleeding								
Low (< 30% bleeding sites)	39	7.85 ± 4.74	10.74 ± 4.87	4.18 ± 3.82	1.10 ± 1.82	0.67 ± 1.32		
High (≥ 30% bleeding sites)	60	9.38 ± 5.40	10.88 ± 4.39	2.88 ± 3.14	0.65 ± 1.07	0.38 ± 1.84		
p-value		0.147	0.877	0.065	0.118	0.587		

http://dx.doi.org/10.1371/journal.pone.0155467.t002

doi:10.1371/journal.pone.0158252.t001

Table 4. Distribution of interproximal sites (n = 2408a) from 99 adults according to adult's variables.

	n	0.6	0.7	0.8	0.9	1.1	p-value ^b
Sex							0.161
Female	1066	38.8	44.8	12.2	2.6	1.5	
Male	1342	33.9	44.3	15.4	4.0	2.5	
Smoker							0.205
Yes	599	32.1	43.4	17.2	3.8	3.5	
No	1809	37.4	44.9	12.9	3.3	1.5	
Patient's periodontal risk							0.061
Low (< 30% bleeding sites)	957	32.0	43.8	17.0	4.5	2.7	
High (\geq 30% bleeding sites)	1451	38.8	45.0	11.9	2.7	1.6	

^a: Interproximal sites with enough space to introduce the IDB (interdental brush).

http://dx.doi.org/10.1371/journal.pone.0155467.t004

doi:10.1371/journal.pone.0158252.t002

Reference

 Carrouel F, Llodra JC, Viennot S, Santamaria J, Bravo M, Bourgeois D (2016) Access to Interdental Brushing in Periodontal Healthy Young Adults: A Cross-Sectional Study. PLoS ONE 11(5): e0155467. doi: 10.1371/journal.pone.0155467 PMID: 27192409

b: DESCRIPT procedure in SUDAAN 7.0 used to adjust for clustering (multiple sites within the patient).