

UNSW SCHOOL OF PHYSICS  
**PHYS2111 – Quantum Mechanics**  
**Tutorial 7**

**Question 1**

Prove the identity  $[\hat{A}\hat{B}, \hat{C}] = \hat{A}[\hat{B}, \hat{C}] + [\hat{A}, \hat{C}]\hat{B}$ .

**Question 2**

Show that  $\delta(ax) = \delta(x)/|a|$ .

*Hint:* Consider  $\int \delta(ax) d(ax)$  and remember that  $\delta(x) = \delta(-x)$ .

**Question 3**

Show that the Hamiltonian operator

$$\hat{H} = -\frac{\hbar^2}{2m} \frac{d^2}{dx^2} + V(x)$$

is Hermitian.

**Question 4**

An anti-hermitian operator is equal to minus its Hermitian conjugate  $\hat{L}^\dagger = -\hat{L}$ .

- (a) Show that the expectation value of an anti-hermitian operator is imaginary.
- (b) Show that the commutator of two Hermitian operators is anti-hermitian. How about the commutator of two anti-hermitian operators?

**Question 5**

Is the ground state of the infinite square potential well an eigenfunction of momentum? If so, what is its momentum? If not, why not?

**Question 6**

(Liboff 3.4) The displacement operator  $\hat{\mathcal{D}}$  is defined by the equation

$$\hat{\mathcal{D}}f(x) = f(x + \zeta).$$

Show that the eigenfunctions of  $\hat{\mathcal{D}}$  are of the form

$$\phi_\beta = e^{\beta x} g(x)$$

where

$$g(x + \zeta) = g(x)$$

and  $\beta$  is any complex number. What is the eigenvalue corresponding to  $\phi_\beta$ ?

**Question 7**

(Liboff 3.17) Show that for an arbitrary “well-behaved” function  $f(x)$ ,

$$\exp\left(\frac{i\zeta\hat{p}}{\hbar}\right) f(x) = f(x + \zeta)$$

where  $\hat{p}$  is the momentum operator and the constant  $\zeta$  represents a small displacement. In this problem, you should demonstrate that the left-hand side of the equation is the Taylor series expansion of the right-hand side about  $\zeta = 0$ .

**Question 8**

- (a) Substitute the Fourier transform

$$f(k) = \frac{1}{\sqrt{2\pi}} \int_{-\infty}^{\infty} e^{-ikx} f(x) dx$$

into its own inverse to show that

$$\frac{1}{2\pi} \int_{-\infty}^{\infty} e^{-ik(x'-x)} dk = \delta(x' - x) \quad (1)$$

- (b) Alternatively, starting with the basis definition (
- $p = \hbar k$
- )

$$\langle k|k' \rangle = \delta(k - k')$$

and writing  $|k\rangle = Ae^{ikx}$ , insert the continuity formula for  $x$

$$\int_{-\infty}^{\infty} |x\rangle \langle x| dx = 1$$

to obtain the same equation (1). What value of the normalisation constant  $A$  should be used?

**Question 9**

(Liboff 3.7) Show that the following are valid representation of  $\delta(y)$ :

- (a)

$$2\pi\delta(y) = \int_{-\infty}^{\infty} e^{iky} dk$$

- (b)

$$\pi\delta(y) = \lim_{\eta \rightarrow \infty} \frac{\sin \eta y}{y}$$

*Note:* In mathematics an object such as  $\delta(y)$ , which is defined in terms of its integral properties, is called a *distribution*. Consider all functions  $\chi(y)$  defined on the interval  $(-\infty, \infty)$  for which

$$\int_{-\infty}^{\infty} |\chi(y)|^2 dy < \infty.$$

Then two distributions  $\delta_1$  and  $\delta_2$  are *equivalent* if for all such  $\chi(y)$ ,

$$\int_{-\infty}^{\infty} \chi(y) \delta_1 dy = \int_{-\infty}^{\infty} \chi(y) \delta_2 dy.$$

When one establishes that a mathematical form such as those shown in parts (a) and (b) above are a representation of  $\delta(y)$ , one is in effect demonstrating that these two object are equivalent as distributions.

**Question 10**

(past exam question) A particle with an energy  $E$  is tunneling through a potential barrier (see figure below). Note that such a tunneling process takes place for example at a metal-insulator-semiconductor thin film structure used in semiconducting industry.

- (a) Sketch the form of the wave function expected in each of the 3 regions and give the functional forms of these wave functions. Explain the choice of the wave function for each region in a few words.




Figure 1:

- (b) What are the mathematical boundary conditions for the waves when the particle enters and leaves the barrier?
- (c) Derive the equations that must be solved to obtain the constants appearing in the wave functions
- (d) If the potential energy in Region 3 is exactly  $E/2$ , calculate the ratio of the wavelength in Regions 1 and 3.

#### Question 11

Complete the analysis of the finite quantum well scattering states (i.e.  $E > 0$  and  $V = V_0$ ) from the notes (or otherwise) to derive an analytical expression for the reflectance ( $R$ ) and transmittance ( $T$ ). Find an expression for the energies at which the transmission is 100%.

#### Question 12

(adapted Griffiths 2.35) A particle of mass  $m$  and kinetic energy  $E > 0$  approaches an abrupt potential drop  $V_0$ .

- (a) What is the probability that it will “reflect” back, if  $E = V_0/3$ ?
- (b) When a free neutron enters a nucleus, it experiences a sudden drop in potential energy, from  $V = 0$  to  $V = -12 \text{ MeV}$  inside. Suppose a neutron, emitted with kinetic energy  $V = 4 \text{ MeV}$  by a fission event, strikes such a nucleus. What is the probability it will be absorbed, initiating another fission? *Hint: You calculated the probability of reflection in part (a): use  $T = 1 - R$  to get the probability of transmission through the surface.*