Arrival Control in Quasi-Reversible Queueing Systems: Optimization and Reinforcement Learning

Céline Comte ⊠ 😭 📵

LAAS-CNRS, Université de Toulouse, CNRS, Toulouse, France

Pascal Moval ☑��

IECL, Université de Lorraine / Inria PASTA, Nancy, France

Abstract

In this paper, we introduce a versatile scheme for optimizing the arrival rates of quasi-reversible queueing systems. We first propose an alternative definition of quasi-reversibility that encompasses reversibility and highlights the importance of the definition of customer classes. In a second time, we introduce balanced arrival control policies, which generalize the notion of balanced arrival rates introduced in the context of Whittle networks, to the much broader class of quasi-reversible queueing systems. We prove that supplementing a quasi-reversible queueing system with a balanced arrival-control policy preserves the quasi-reversibility, and we specify the form of the stationary measures. We revisit two canonical examples of quasi-reversible queueing systems, Whittle networks and order-independent queues. Lastly, we focus on the problem of admission control and leverage our results in the frameworks of optimization and reinforcement learning.

2012 ACM Subject Classification Mathematics of computing \rightarrow Markov processes; Applied computing \rightarrow Operations research

Keywords and phrases Admission control, quasi-reversibility, reversibility, linear programming, policy-gradient reinforcement learning

Acknowledgements Thank you to Fabien Mathieu for insightful discussions that helped construct Section 5.3.

1 Introduction

In this paper, we introduce a versatile optimization scheme for the control of arrival rates into quasi-reversible queueing systems, a broad class of multi-class queueing systems that has been studied extensively in the literature [18, 19, 8]. Our definition of quasi-reversibility differs slightly from the definitions given in these references. In a nutshell, under our definition, the quasi-reversibility of a queueing system implies the existence of an aggregation of the states of the natural Markov chain of the system into macrostates, such that the macrostate process thereby obtained satisfies a reversibility property with respect to the stationary measures of the Markov chain. Note that the macrostate process is not Markov in general, yet it coincides in distribution with a reversible Markov chain, see Section 2.2.3 below. This aggregation is obtained through a so-called counting function, which maps the microstate to the macrostate by only retaining the number of customers of each class in the system.

This alternative definition of quasi-reversibility is instrumental to our first main contribution, namely, the definition of balanced arrival-control policies. These generalize the balanced arrival rates introduced in the context of Whittle networks without internal routing [5, 16], to the much broader class of quasi-reversible queueing systems. As our first main result (Theorem 1), we show that the quasi-reversibility of the queueing system under consideration is preserved by any balanced arrival-control policy, and we specify the form of the stationary measures in this case. Roughly speaking, we show that supplementing a quasi-reversible

2 Arrival Control in Quasi-Reversible Queueing Systems

queueing system with a balanced arrival control boils down to multiplying the stationary measures by the balance function associated with the arrival control.

To illustrate the definition of balanced policies, we focus on the special case of admission control, where arrival control is interpreted as admitting or rejecting incoming customers. For a quasi-reversible queueing system with a finite state space, we formulate the problem of finding the best balanced admission-control policy as a linear optimization problem. This allows us to show that there exists a policy optimal among balanced policies that is deterministic (Theorem 8), thus generalizing a result shown in [16] in the context of Whittle networks. We conclude with a digression, to compare the performance and structure of the best balanced policy with that of an optimal policy. To do so, we leverage both insights from the literature on (quasi-)reversible queueing systems and numerical results in a toy example.

Lastly, we show the crucial interest of quasi-reversibility and balanced arrival control for implementing efficient reinforcement learning (RL) algorithms that optimize the access control. Given a family of parameterized balanced policies, we show that, thanks to quasi-reversibility, the stationary distribution of the parameterized system satisfies a simple gradient equation with respect to the system parameter. This allows us to construct a gradient estimator that we use in a gradient-ascent type algorithm common in policy-gradient reinforcement learning (PGRL). The exploitation of product-form like stationary measures to implement this class of RL algorithms, is reminiscent of the approach recently developed in [12].

To illustrate these results, we discuss two canonical examples of quasi-reversible systems, which gather a wide class of queueing systems: multi-type Whittle networks as defined in [23], and order-independent (OI) queues introduced in [3, 4]. After checking that these two queueing systems (already known to be quasi-reversible, at least under some assumptions) satisfy our definition of quasi-reversibility, we show that the PGRL procedure defined above can be fruitfully applied to optimize admission control into a variant, with customer abandonment, of the redundancy model with customer-server compatibility introduced in [15] which, as observed in [6], is an example of OI queues.

1.1 Overview

This paper is organized as follows. In Section 2, we introduce quasi-reversible systems, and we review the literature on reversibility as well as the various variants of quasi-reversibility. Second, in Section 3, we introduce the notion of balanced arrival control in quasi-reversible systems and their interest in the context of RL. In Section 4, we show how this approach can be applied to two well-known general examples of quasi-reversible systems, namely, multi-class Whittle systems in Section 4.1; and order-independent queues in Section 4.2. Then, in Section 5 we set our optimization problem for admission control. Optimality of deterministic policies among balanced policies is shown in Section 5.2 for quasi-reversible queueing systems with a finite state space. The cost of implementing balanced policies is investigated in Section 5.3. In Section 6, we present a case study, and we implement an RL algorithm to optimize the admission control in a redundancy model subject to server-customer compatibility. Section 7 concludes this work.

1.2 Notation

We let \mathbb{N} , $\mathbb{N}_{>0}$, \mathbb{R} , and $\mathbb{R}_{>0}$ denote the sets of non-negative integers, positive integers, reals, and positive reals, respectively. For $a \leq b \in \mathbb{N}$, we use [a, b] to denote the integer interval $\{a, a+1, \ldots, b\}$, and define [a, b[, a, b[, a, b]]] accordingly.

Fix $m, n \in \mathbb{N}_{>0}$. We let $\mathcal{M}_{n,m}(\mathbb{N})$ denote the set of matrices with n rows and m columns of integer entries. With some abuse of notation, the null vectors of \mathbb{R}^n and the null matrix of $\mathcal{M}_{n,m}(\mathbb{N})$ are both denoted by $\mathbf{0}$. For any $i \in [\![1,n]\!]$, we let e_i be the vector of \mathbb{R}^n with one in component i and zero elsewhere. Likewise, for all $i \in [\![1,n]\!]$ and $k \in [\![1,m]\!]$, we let e_{ik} be the matrix of $\mathcal{M}_{n,m}(\mathbb{N})$ with one in component ik, and zero elsewhere. The vector space \mathbb{R}^n is equipped with the coordinate-wise semi-ordering, namely, for all $x = (x_1, \ldots, x_n), y = (y_1, \ldots, y_n) \in \mathbb{R}^n, x \leq y$ if and only if $x_i \leq y_i$ for all $i \in [\![1,n]\!]$. We also let $\mathbb{R}^n_{>0}$ be the subset of vectors of \mathbb{R}^n having positive entries.

A subset $\mathcal{X} \subseteq \mathbb{N}^n$ is called a *Ferrers set* if $0 \in \mathcal{X}$ and \mathcal{X} is coordinate convex, in the sense that, for $x \in \mathbb{N}^n$, $x \in \mathcal{X}$ implies $y \in \mathcal{X}$ for each $y \in \mathbb{N}^n$ such that $y \leq x$.

Let \mathcal{A} be a finite set. We denote by $\operatorname{card}(\mathcal{A})$, the cardinality of \mathcal{A} . We let \star denote the Kleene-star operator of \mathcal{A} , namely, \mathcal{A}^{\star} is the set of words on the alphabet \mathcal{A} . A word $w \in \mathcal{A}^{\star}$ is then written as $w = w_1 \cdots w_{\ell}$, where $\ell \in \mathbb{N}_{>0}$ is the *length* of w. The empty word of \mathcal{A}^{\star} is denoted by \emptyset , is supposed to be of null length.

The identity function of any set into itself is denoted by Id. Given a real-valued function f, supp(f) denotes the support of f, i.e., the preimage of $\mathbb{R} \setminus \{0\}$ by f.

Quasi-reversible queueing systems

In this section, we introduce quasi-reversible queueing systems, a broad class of multi-class queueing systems described for instance in [18, Section 3.2]. In Section 2.1, we define what we intend by a queueing system: It is formally defined as a (discrete-state-space) continuous-time Markov chain (CTMC) whose state contains enough information to count the number of customers of each class. Section 2.2 gives our definition of quasi-reversibility. To prepare the ground for our first main result (Theorem 1 in Section 3.3), we pay particular attention to assumptions regarding the CTMC's transition diagram. In case the reader would like to make these definitions more concrete, we encourage them to take a look at Section 4 right away, as it describes two canonical examples of quasi-reversible systems, and applies our results to these examples.

2.1 Queueing systems

In this paper, a queueing system is defined as a triple $Q = (\mathcal{S}, |\cdot|, q)$, where \mathcal{S} is called the (micro-)state space, $|\cdot|: \mathcal{S} \to \mathbb{N}^n$ the counting function, and $q: \mathcal{S} \times \mathcal{S} \to \mathbb{R}_{\geq 0}$ the transition kernel. Let us specify each of these elements.

2.1.1 State space, counting function, and microstate-independence

Consider a queueing system with a finite number $n \in \mathbb{N}_{>0}$ of customer classes. Intuitively, a class is a label that is attached to a customer and that may carry information about this customer, such as its service requirement. As we will see later, we assume that the class of a customer is fixed throughout its stay in the queueing system. Let \mathcal{S} denote the state space of the system, assumed to be countable.

The system state is assumed to contain enough information to recover the number of customers of each class via a mapping $|\cdot|$ called the *counting* function: for each $s \in \mathcal{S}$, $|s| = (|s|_1, |s|_2, \ldots, |s|_n) \in \mathbb{N}^n$ gives the number of customers of each class when the system is in state s. We let $|\mathcal{S}| = \{|s|; s \in \mathcal{S}\}$ denote the image of \mathcal{S} by $|\cdot|$, and we assume that $|\mathcal{S}|$ is a Ferrers set. Also, for each $x \in \mathbb{N}^n$ we let $\mathcal{S}_x = \{s \in \mathcal{S}; |s| = x\}$ (which is empty if $x \notin |\mathcal{S}|$

4 Arrival Control in Quasi-Reversible Queueing Systems

by definition). An element of S is called a *microstate* and an element of |S| a *macrostate*¹. We assume that there exists exactly one state in S, called the *empty state* and denoted by \varnothing , such that $|\varnothing| = 0$.

We will often consider functions $f: \mathcal{S} \to \mathbb{R}$ that are microstate-independent in the sense that f(s) = f(t) for each $s, t \in \mathcal{S}$ such that |s| = |t|. This is equivalent to the existence of a function $\tilde{f}: |\mathcal{S}| \to \mathbb{R}$ such that $f = \tilde{f} \circ |\cdot|$, that is, $f(s) = \tilde{f}(|s|)$ for each $s \in \mathcal{S}$. With a slight abuse of notation, we will neglect the distinction between f and \tilde{f} and see f as a function defined on both \mathcal{S} and $|\mathcal{S}|$, possibly writing f(x) for f(s) when x = |s|, depending on what is convenient from a notation point of view; whether the input is a microstate or a macrostate will be clear from the notation. In the same spirit, given a function $f: |\mathcal{S}| \to \mathbb{R}$, we will often write f(s) for f(x) when x = |s|.

2.1.2 Transition kernel

We assume that the evolution of the microstate over time defines a CTMC with transition kernel $q: \mathcal{S} \times \mathcal{S} \to \mathbb{R}_{\geq 0}$. This CTMC will be called the *microstate chain* and will often be identified with the kernel q or the triple $Q = (\mathcal{S}, |\cdot|, q)$. For each $s, t \in \mathcal{S}$, q(s, t) gives the rate at which the system transitions to state t when it is in state s. In particular, q(s, s) gives the transition rate from a state $s \in \mathcal{S}$ to itself (so that s is not an infinitesimal generator). We assume that transitions out of each state $s \in \mathcal{S}$ can be partitioned as follows:

- Arrivals of class i, that is, transitions to a state $t \in \mathcal{S}_{|s|+e_i}$, for each $i \in [1, n]$;
- Departures of class i, i.e., transitions to a state $t \in S_{|s|-e_i}$, for each $i \in [1, n]$;
- Internal transitions, that is, transitions to a state $t \in S_{|s|}$.

In particular, we assume that q(s,t) = 0 if $t \notin \mathcal{S}_{|s|} \cup \bigcup_{i=1}^n (\mathcal{S}_{|s|+e_i} \cup \mathcal{S}_{|s|-e_i})$. Intuitively, customers arrive and leave one at a time, and a customer's class is constant over time.

To avoid pathological cases (e.g., to guarantee that the empty state is reachable) and simplify the definitions in Section 3.1, we make the following technical assumption.

▶ Assumption 1. There exists $S_{rec} \subseteq S$ such that:

- 1. For each $s \in \mathcal{S}_{rec}$, there is a sequence $\emptyset = t_1, t_2, \dots, t_\ell = s$ in \mathcal{S} , with $\ell \geq 1$, such that $|t_k| \leq |t_{k+1}|$ and $q(t_k, t_{k+1}) > 0$ for each $k \in [\![1, \ell-1]\!]$.
- 2. For each $s \in \mathcal{S} \setminus \mathcal{S}_{rec}$, there is no sequence $\emptyset = t_1, t_2, \ldots, t_\ell = s$ in \mathcal{S} , with $\ell \geq 1$, such that $q(t_k, t_{k+1}) > 0$ for each $k \in [1, \ell 1]$.
- 3. For each $s \in \mathcal{S}$, there is a sequence $s = t_1, t_2, \ldots, t_{\ell} = \emptyset$ in \mathcal{S} , with $\ell \geq 1$, such that $|t_k| \geq |t_{k+1}|$ and $q(t_k, t_{k+1}) > 0$ for each $k \in [1, \ell 1]$.
- **4.** The set $|\mathcal{S}_{rec}| = \{|s|; s \in \mathcal{S}_{rec}\}$ is a Ferrers set.

Assumption 1 implies in particular that Q is unichain: the microstate chain admits \mathcal{S}_{rec} as a single recurrent class, plus a possibly empty set $\mathcal{S} \setminus \mathcal{S}_{rec}$ of transient states. More specifically, under Assumption 1, the empty state \varnothing is recurrent, and other states in \mathcal{S}_{rec} communicate with \varnothing via paths that are monotonic with respect to $|\cdot|$. We will see in Section 4 that, despite these restrictions, this structure is sufficiently versatile to capture broad families

The state spaces S and |S| may not be disjoint in general (for instance, if $|\cdot|$ is the identify function). In this case, the formal distinction between microstates and macrostates is mostly intuitive and related to the counting function: a microstate is an input of this function, while a macrostate is an output. As we will see in Sections 2.1.2 and 2.2.3, the stochastic process describing the evolution of the microstate over time satisfies the Markov property, while the stochastic process describing the evolution of the macrostate does not in general.

of systems, such as Whittle networks and order-independent (OI) queues, equipped with sophisticated routing mechanisms, such as those described by the method of stages [17, 1].

Unless stated otherwise, a queueing system will be understood as a triple $Q = (\mathcal{S}, |\cdot|, q)$ whose associated microstate chain satisfies Assumption 1 and is in particular unichain.

2.2 Stationary measures and quasi-reversibility

2.2.1 General definition

Consider a queueing system $Q = (S, |\cdot|, q)$ as defined in Section 2.1. Due to Assumption 1, the microstate chain has a stationary measure $\Pi : S \to \mathbb{R}_{\geq 0}$ that is unique up to a positive multiplicative constant, and characterized by the balance equations

$$\Pi(s) \sum_{t \in \mathcal{S}} q(s, t) = \sum_{t \in \mathcal{S}} \Pi(t) q(t, s), \text{ for each } s \in \mathcal{S},$$

along with the normalizing condition $\sum_{s\in\mathcal{S}}\Pi(s)=1$. As we assumed q(s,t)=0 for each $s,t\in\mathcal{S}$ with $t\notin\mathcal{S}_{|s|}\cup\bigcup_{i=1}^n(\mathcal{S}_{|s|+e_i}\cup\mathcal{S}_{|s|-e_i})$, these balance equations rewrite as follows:

$$\Pi(s) \left(\sum_{i=1}^{n} \sum_{t \in \mathcal{S}_{|s|+e_i}} q(s,t) + \sum_{i=1}^{n} \sum_{t \in \mathcal{S}_{|s|-e_i}} q(s,t) + \sum_{t \in \mathcal{S}_{|s|}} q(s,t) \right)$$

$$= \left(\sum_{i=1}^{n} \sum_{t \in \mathcal{S}_{|s|+e_i}} \Pi(t)q(t,s) + \sum_{i=1}^{n} \sum_{t \in \mathcal{S}_{|s|-e_i}} \Pi(t)q(t,s) + \sum_{t \in \mathcal{S}_{|s|}} \Pi(t)q(t,s) \right), \quad \text{for each } s \in \mathcal{S}.$$
(1)

▶ **Definition 1.** We say that a queueing system $Q = (S, |\cdot|, q)$ is quasi-reversible if its stationary measures Π also satisfy the following set of equations:

$$\Pi(s) \sum_{t \in \mathcal{S}_{|s|+e_i}} q(s,t) = \sum_{t \in \mathcal{S}_{|s|+e_i}} \Pi(t)q(t,s), \quad \text{for each } s \in \mathcal{S} \text{ and } i \in [1,n].$$

Equation (2) in Definition 1 states that the probability flow out of state s due to a class-i arrival is equal to the probability flow into state s due to a class-i departure, for each $s \in \mathcal{S}$ and $i \in [1, n]$, which is equivalent to [18, Equation (3.11)]. Subtracting (2) from (1) yields as a by-product

$$\Pi(s) \left(\sum_{i=1}^{n} \sum_{t \in \mathcal{S}_{|s|-e_i}} q(s,t) + \sum_{t \in \mathcal{S}_{|s|}} q(s,t) \right)$$

$$= \sum_{i=1}^{n} \sum_{t \in \mathcal{S}_{|s|-e_i}} \Pi(t)q(t,s) + \sum_{t \in \mathcal{S}_{|s|}} \Pi(t)q(t,s), \text{ for each } s \in \mathcal{S}. \quad (3)$$

In fact, satisfying (1)–(2) is equivalent to satisfying (2)–(3), hence quasi-reversibility is stronger than stationarity. Lastly, observe that quasi-reversibility does not require positive recurrence, as it is stated as a property of the stationary measures.

2.2.2 Other definitions of quasi-reversibility

For clarity, let us compare Definition 1 with two other classical definitions of quasi-reversibility, namely, [18, Section 3.2] and [8, Definition 1]. All three definitions have many consequences in common, but they also have differences:

1. Definition 1 is more general than [18, Section 3.2] and [8, Definition 1] in that it does not require the arrival rate of each class to be independent of the current state, i.e., we do not impose the existence, for each $i \in [1, n]$, of a $\lambda_i > 0$ such that

$$\sum_{t \in \mathcal{S}_{|s|+e_i}} q(s,t) = \lambda_i \mathbb{1}\left[|s| + e_i \in \mathcal{X}\right], \quad \text{for each } s \in \mathcal{S}.$$
(4)

2. Definition 1 is more restrictive than [8, Definition 1] in that it restricts the sets of states towards which each type of transitions (arrival, departure, and internal) can lead. This restriction imposed by Definition 1 is necessary to prove Theorem 1. It rules out the negative customers considered in [8], but is still general enough to capture complex queueing systems such as those of Section 4.

All in all, both Definition 1 and [8, Definition 1] are strict extensions of [18, Section 3.2], but neither definition can be seen as a special case of the other. Point 1 above has the following consequences:

- Under Definition 1, quasi-reversibility cannot be interpreted as a condition of independence between the system state and the arrival and departure processes, as it was the case in [18, Section 3.2] and [8, Definition 2]. However, other consequences of quasi-reversibility continue to hold; in particular, a network of quasi-reversible queueing systems is still quasi-reversible (see [18, Theorem 3.7(iv)] and Section 4.1).
- Relaxing assumption (4) allows Definition 1 to capture queueing systems that are (trully) reversible, even when the arrival rates are state dependent. On this question, see Section 2.2.3 below, as well as the discussion in [18, Equation (3.11)].

2.2.3 Quasi-reversibility and (true) reversibility

We make a small digression to clarify the relationship between quasi-reversibility and reversibility. The first-time reader can skip this section if necessary, and possibly come back to it after reading the equivalence result of Proposition 7, which is a special case.

Reversibility as a special case of quasi-reversibility

Consider a queueing system $Q^* = (\mathcal{S}^*, |\cdot|^*, q^*)$ where $|\cdot|^* = \text{Id}$ is the identity function: microstates and macrostates are identical, $\mathcal{S}^* \subseteq \mathbb{N}^n$, and $\mathcal{S}^*_{|x|} = \{x\}$ for each $x \in \mathcal{S}^*$. Then the definition of quasi-reversibility (Definition 1) simplifies as follows:

$$\Pi^{\star}(x)q^{\star}(x,x+e_i) = \Pi^{\star}(x+e_i)q^{\star}(x+e_i,x), \text{ for each } x \in \mathcal{S}^{\star} \text{ and } i \in [1,n],$$

where Π^* is the stationary distribution of Q^* . This is exactly the definition of reversibility. In this way, under the assumption that $|\cdot|^* = \mathrm{Id}$, quasi-reversibility is equivalent to reversibility.

Reversibility as an aggregation of quasi-reversibility

Now consider an arbitrary quasi-reversible queueing system $Q = (\mathcal{S}, |\cdot|, q)$. To simplify the discussion, let us assume that $\mathcal{S} = \mathcal{S}_{rec}$, so that the stationary distribution Π of the microstate chain of Q is positive on \mathcal{S} . In that case, quasi-reversibility of the microstate chain implies reversibility of the macrostate process, even if this process does not have the Markov property in general. To see this, let $x \in |\mathcal{S}|$ and $i \in [1, n]$ such that $x + e_i \in |\mathcal{S}|$. Summing (2) over all $s \in \mathcal{S}_x$ yields

$$\sum_{s \in \mathcal{S}_x} \Pi(s) \sum_{t \in \mathcal{S}_{x+e_i}} q(s,t) = \sum_{s \in \mathcal{S}_x} \sum_{t \in \mathcal{S}_{x+e_i}} \Pi(t) q(t,s).$$

Given an arbitrary distribution Π^* with support $|\mathcal{S}|$, we can rewrite this equation as

$$\Pi^{\star}(x) \underbrace{\sum_{s \in \mathcal{S}_x} \sum_{t \in \mathcal{S}_{x+e_i}} \frac{\Pi(s)}{\Pi^{\star}(x)} q(s, t)}_{q^{\star}(x, x+e_i)} = \Pi^{\star}(x + e_i) \underbrace{\sum_{s \in \mathcal{S}_x} \sum_{t \in \mathcal{S}_{x+e_i}} \frac{\Pi(t)}{\Pi^{\star}(x + e_i)} q(t, s)}_{q^{\star}(x + e_i, x)}.$$
(5)

Using (5), there are two ways of interpreting q^* and Π^* :

- If we abstract ourselves away from the original queueing system Q, we can simply see $Q^* = (|\mathcal{S}|, \operatorname{Id}, q^*)$ as a reversible queueing system. In particular, q^* is the transition kernel of a reversible CTMC with stationary distribution Π^* .
- Focusing on the distribution Π^* given by $\Pi^*(x) = \sum_{s \in \mathcal{S}_x} \Pi(s)$ for each $x \in |\mathcal{S}|$, (5) shows that, even if the macrostate process does not have the Markov property in general, this process is reversible, and under stationarity it has the same transition rates as a reversible CTMC with transition kernel q^* .

3 Balanced arrival control

In this section, we consider balanced arrival control, a way of modifying the arrival rates in a quasi-reversible queueing system that preserves the quasi-reversibility property. Balanced arrival control was already considered in the special case of Whittle networks, see [23, 5, 16]. We recall the definition in Section 3.1 and provide examples in Section 3.2. Our main contribution, in Section 3.3, consists of showing that balanced arrival control is also compatible with quasi-reversibility, in the sense that supplementing a quasi-reversible queueing system with a balanced arrival control again yields a quasi-reversible queueing system. Lastly, Section 3.4 shows a consequence of this result that is useful for optimizing arrival rates via gradient descent.

3.1 Definition

Consider a quasi-reversible queueing system $Q = (\mathcal{S}, |\cdot|, q)$. An arrival-control policy, or policy for short, is a function $\gamma : \mathcal{S} \to \mathbb{R}^n_{\geq 0}$. The queueing system under policy γ is $Q_{\gamma} = (\mathcal{S}, |\cdot|, q_{\gamma})$, where $q_{\gamma} : \mathcal{S} \times \mathcal{S} \to \mathbb{R}_{\geq 0}$ is given as follows: for each $s, t \in \mathcal{S}$,

$$q_{\gamma}(s,t) = \begin{cases} q(s,t)\gamma_{i}(s) & \text{if } t \in \mathcal{S}_{|s|+e_{i}} \text{ for some } i \in [1,n], \\ q(s,t) & \text{otherwise.} \end{cases}$$
(6)

If $\gamma_i(s) \in [0,1]$ for each $s \in \mathcal{S}$ and $i \in [1,n]$, then γ can be interpreted as an admission-control policy, i.e., Q_{γ} is a variant of Q where some incoming customers are rejected. If $\sum_{i=1}^{n} \gamma_i(s) = 1$ for each $s \in \mathcal{S}$, then γ can be interpreted as a load-balancing policy. Examples of such policies in the context of Whittle networks without internal transitions were considered in [5, 16].

Let us briefly discuss the assumptions of Section 2.1. Q_{γ} again satisfies $q_{\gamma}(s,t)=0$ if $t \notin \mathcal{S}_{|s|} \cup \bigcup_{i=1}^{n} (\mathcal{S}_{|s|+e_{i}} \cup \mathcal{S}_{|s|-e_{i}})$, it is unichain (which follows from (6) and the fact that Q satisfies Assumption 1), and it admits \varnothing as a recurrent state. Therefore, Q_{γ} has a stationary measure Π_{γ} that is unique up to a positive multiplicative constant, and this measure satisfies $\Pi_{\gamma}(\varnothing) > 0$. However, in general, Q_{γ} does not satisfy Assumption 1 and is not quasi-reversible.

This motivates us to introduce balanced policies in Definition 2 below. As we will see in Section 3.3, if the policy γ is balanced, then Q_{γ} does satisfy Assumption 1 and is quasi-reversible. This result will be exploited in the remainder of the paper to provide efficient ways of searching balanced policies that optimize certain performance criteria.

- ▶ **Definition 2** (Balanced policy). Consider a queueing system $Q = (S, |\cdot|, q)$. A policy $\gamma: \mathcal{S} \to [0,1]^n$ is said to be balanced if it satisfies the following two assumptions:
- 1. Microstate-independence: $\gamma(s) = \gamma(t)$ for each $s, t \in \mathcal{S}$ such that |s| = |t|.
- **2.** Balance condition: There exists a function $\Gamma: |\mathcal{S}| \to \mathbb{R}_{\geq 0}$ such that $\Gamma(0) = 1$, and

$$\Gamma(x)\gamma_i(x) = \Gamma(x+e_i), \quad \text{for each } x \in |\mathcal{S}| \text{ and } i \in [1,n] \text{ so that } x+e_i \in |\mathcal{S}|.$$
 (7)

In this case, we say that the policy γ is Γ -balanced, or equivalently that Γ is the balance function of the policy γ , and we write $\gamma = \partial \Gamma$.

$$\begin{array}{c} \Gamma(0,2) \xrightarrow{\gamma_1(0,2)} \Gamma(1,2) \xrightarrow{\gamma_1(1,2)} \Gamma(2,2) \xrightarrow{\gamma_1(2,2)} \Gamma(3,2) \xrightarrow{\gamma_1(3,2)} \Gamma(4,2) \\ \hline \\ \widehat{\mathbb{C}} \\ \widehat{\mathbb{C}} \\ \hline \\ \Gamma(0,1) \xrightarrow{\gamma_1(0,1)} \Gamma(1,1) \xrightarrow{\gamma_1(1,1)} \Gamma(2,1) \xrightarrow{\gamma_1(2,1)} \Gamma(3,1) \xrightarrow{\gamma_1(3,1)} \Gamma(4,1) \\ \hline \\ \widehat{\mathbb{C}} \\ \widehat{\mathbb{C}} \\ \hline \\ \Gamma(0,0) \xrightarrow{\gamma_1(0,0)} \Gamma(1,0) \xrightarrow{\gamma_1(1,0)} \Gamma(2,0) \xrightarrow{\gamma_1(2,0)} \Gamma(3,0) \xrightarrow{\gamma_1(3,0)} \Gamma(4,0) \\ \hline \end{array}$$

Figure 1 An intuitive description of the balance condition in a toy queueing system of dimension n=2 with macrostate space $\mathcal{X}=\{x\in\mathbb{N}^2:x_1\leq 4,x_2\leq 2\}$. For each $x\in\mathcal{X},\,\Gamma(x)$ is the product of admission probabilities along an arbitrary increasing path going from the origin to x. The balance condition guarantees that $\Gamma(x)$ is well-defined in the sense that the product is independent of the path.

Before giving examples, let us discuss several points in Definition 2. Definition 2 consists of two conditions. Microstate-independence is tied to the notions of classes and counting function introduced in Section 2.1.1. In general, there may be several ways of defining classes and the counting function in a queueing system, and these may lead to different definitions of balance. If a policy γ is microstate-independent, it can be seen either as a (microstate-independent) function on \mathcal{S} , or as a function on $\mathcal{X} = |\mathcal{S}|$. In the remainder, we will often take the later point of view, in which case we will ignore S and say that γ is a policy on \mathcal{X} . The balance condition, which is the second part of Definition 2, is illustrated in Figure 1. Intuitively, for any x, Equation (7) states that $\Gamma(x)$ is the product of the policy's transition rates over any increasing path from the origin to x, so that this product is independent of the chosen path. In fact, one can verify that there exists a balance function Γ that satisfies (7) if and only if

$$\gamma_{i}(x)\gamma_{j}(x+e_{i}) = \gamma_{i}(x+e_{j})\gamma_{j}(x), \quad \begin{cases} \text{for each } x \in |\mathcal{S}|, i, j \in [1, n] \text{ such that} \\ x+e_{i}, x+e_{j} \in \text{supp}(\Gamma), x+e_{i}+e_{j} \in |\mathcal{S}|. \end{cases}$$
(8)

Observe that (7) would still be satisfied if we replaced Γ with $\frac{1}{Z}\Gamma$ for any Z>0. The original queueing system Q is retrieved by taking $\Gamma(x) = 1$ for each $x \in |\mathcal{S}|$.

Given a Ferrers set $\mathcal{X} \subseteq \mathbb{N}^n$ interpreted as a queueing system's macrostate space, we will see that there is a bijection between the set of balanced policies on \mathcal{X} and the set of balance functions on \mathcal{X} . The set of balance functions on \mathcal{X} is given by

$$\mathcal{B}_{\mathcal{X}} = \{ \Gamma : \mathcal{X} \to \mathbb{R}_{\geq 0} \, ; \, \Gamma(0) = 1 \text{ and } \operatorname{supp}(\Gamma) \text{ is a Ferrers set} \} \,. \tag{9}$$

Indeed, if Γ satisfies the balance condition (7) for a certain policy γ , then $\Gamma(x) = 0$ for some $x \in \mathcal{X}$ implies that $\Gamma(y) = 0$ for each $y \in \mathcal{X}$ such that $x \leq y$, which means that supp(Γ) is a Ferrers set. Conversely, for each $\Gamma \in \mathcal{B}_{\mathcal{X}}$, we can construct a policy γ on \mathcal{X} that satisfies (7) by letting

$$\gamma_i(x) = \frac{\Gamma(x+e_i)}{\Gamma(x)}, \text{ for each } x \in \text{supp}(\Gamma) \text{ and } i \in [1,n] \text{ such that } x+e_i \in \mathcal{X},$$
(10)

and $\gamma(x) = 0$ for each $x \in \mathcal{X} \setminus \text{supp}(\Gamma)$. Now, if we let γ and γ' denote two Γ -balanced policies, (10) implies that $\gamma_i(x) = \gamma_i'(x)$ for each $x \in \text{supp}(\Gamma)$ and $i \in [\![1,n]\!]$ such that $x + e_i \in \mathcal{X}$. In particular, the restrictions of Q_{γ} and $Q_{\gamma'}$ to $\text{supp}(\Gamma)$ are identical, i.e., they have the same transition rates. Since one can also verify that the recurrent sets of Q_{γ} and $Q_{\gamma'}$ are included into $\text{supp}(\Gamma)$, it follows that Q_{γ} and $Q_{\gamma'}$ have the same stationary distribution. Therefore, by assimilating two policies if they lead to the same stationary distribution if necessary, it follows that (7) defines a bijection between the set $\mathcal{B}_{\mathcal{X}}$ of balance functions and the set of balanced policies. In general, when we write $\gamma = \partial \Gamma$, we mean that γ satisfies (10) and takes an arbitrary value on $\mathcal{X} \setminus \text{supp}(\Gamma)$, say $\gamma(x) = 0$ for each $x \in \mathcal{X} \setminus \text{supp}(\Gamma)$.

Balanced arrival control has been studied extensively in the context of Whittle networks without internal transitions [5, 16]. The key novelty of Definition 2 is to apply balanced policies in the much more general context of quasi-reversible queueing systems, via the definitions of the counting function $|\cdot|$ and of microstate-independence.

3.2 Examples

Consider a queueing system $Q = (\mathcal{S}, |\cdot|, q)$. To make the definition of balanced policies more concrete, let us focus on the special case of admission control, where $\gamma_i(x) \in [0, 1]$ is the admission probability of class-i customers when the macrostate is x, for each $x \in |\mathcal{S}|$ and $i \in [1, n]$. Examples of balanced policies are shown in Table 1.

| Name | Admission probability (with $\gamma_i(x) = 0$ if $\Gamma(x) = 0$ or $x + e_i \notin \mathcal{X}$) | Balance function (with $\Gamma(x) = 0$ if $x \notin \mathcal{X}$) | Parameter | |
|-----------------------------------|----------------------------------------------------------------------------------------------------------|---------------------------------------------------------------------|---------------------------------------------------------------------------------------------------------------------------|--|
| Static | $\gamma_i(x) = \alpha_i$ | $\Gamma(x) = \prod_{i=1}^{n} \alpha_i^{x_i}$ | Reals $\alpha_i \in [0, 1]$ | |
| Decentralized | $\gamma_i(x) = \psi_i(x_i)$ | $\Gamma(x) = \prod_{i=1}^{n} \prod_{\ell=0}^{x_i - 1} \psi_i(\ell)$ | Functions $\psi_i : \overline{\mathcal{X}_i} \to [0, 1],$ with $\overline{\mathcal{X}_i} = \{x_i; x \in \mathcal{X}\}$ | |
| Size-based | $\gamma_i(x) = \psi(x _1)$ | $\Gamma(x) = \prod_{\ell=0}^{i=1} \psi(\ell)$ | Function $\psi : \overline{\mathcal{X}} \to [0, 1],$ with $\overline{\mathcal{X}} = \{ \ x\ _1; x \in \mathcal{X} \}$ | |
| Deterministic | $\gamma_i(x) = \mathbb{1}\left[x + e_i \in \mathcal{A}\right]$ | $\Gamma(x) = \mathbb{1}\left[x \in \mathcal{A}\right]$ | A Ferrers set $A \subseteq \mathcal{X}$ called the policy's mask | |
| Cum-prod (for cumulative-product) | $\gamma_i(x) = \prod_{\substack{y \in \mathcal{X}: y \le x + e_i, \\ y_i = x_i + 1}} \psi(y)$ | $\Gamma(x) = \prod_{y \in \mathcal{X}: y \le x} \psi(y)$ | Function $\psi: \mathcal{X} \to [0,1]$ | |

Table 1 Examples of balanced policies over an arbitrary Ferrers set $\mathcal{X} \subseteq \mathbb{N}^n$. Here, $x \in \mathcal{X}$ and $i \in [1, n]$.

The first row shows a simple but common family of balanced policies: *static policies*, where the admission decision does not depend on the current state. The next two rows show that balanced policies include two important families of admission policies: *decentralized policies*, where the decision to admit a customer of a given class only depends on the number

of customers of this class in the system, and size-based policies, where the decision to admit a customer only depends on the total number of customers in the system (irrespective of their classes). It follows that, if we know a priori that the optimal policy belongs to one of these two classes, then we can search the optimal policy among balanced policies. Deterministic policies, on the fourth row of Table 1, will play a fundamental role in Section 5, as we will see that the best balanced policy (in a sense that we will define in Section 5.2) can be chosen of this form, at least when the state space is finite. Observe that deterministic balanced policies form a subset of deterministic possibly-imbalanced policies. Indeed, under a deterministic balanced policy, the only states $x \in \mathcal{X}$ and classes $i, j \in [1, n]$ such that $\gamma_i(x) \neq \gamma_j(x)$ are such that $x + e_i \in \mathcal{A}$ and $x + e_j \notin \mathcal{A}$ or $x + e_i \notin \mathcal{A}$ and $x + e_j \in \mathcal{A}$, where $\mathcal{A} \subseteq |\mathcal{S}|$ is called the policy's mask. Cum-prod policies, on the fifth row of Table 1, is a family of balanced policies that includes deterministic policies but admit a continuous parameterization.

The simple examples of Table 1 can be generalized in various ways depending on the problem at hand. For instance, decentralized policies can be augmented by factors involving multiple variables, for instance,

$$\Pi(x) = \left(\prod_{i=1}^{n} \prod_{\ell=0}^{x_{i}-1} \psi_{i}(\ell) \right) \left(\prod_{\{i,j\} \subseteq [\![1,n]\!]} \prod_{\ell=0}^{x_{i}+x_{j}-1} \psi_{i,j}(\ell) \right), \quad x \in \mathcal{X},$$

where ψ_i is as defined for decentralized policies, and $\psi_{i,j}: \overline{\mathcal{X}}_{i,j} \to [0,1]$ with $\overline{\mathcal{X}}_{i,j} = \{x_i + x_j; x \in \mathcal{X}\}$ for each $\{i,j\} \subseteq [1,n]$. One may restrict the latter product to only some pairs of classes, using for instance information on the structure of the system.

One may also consider balance functions Γ of the form $\Gamma(x) = f(x)$ for each $x \in \mathcal{X}$, where $f: [0, +\infty)^n \to [0, +\infty)$ is a differentiable function with non-positive first partial derivatives (in particular, f is componentwise non-increasing). One broad class of functions that satisfy this requirement are complementary cumulative distribution functions (CCDFs) of multivariate distributions, such as multivariate Gaussian distributions.

3.3 Balanced arrival rates preserve quasi-reversibility

Theorem 1 hereafter shows that supplementing a quasi-reversible queueing system with a $(\Gamma$ -)balanced policy preserves quasi-reversibility and boils down to modulating the stationary distribution by Γ , in the sense of (11). Theorem 1 was inspired by two observations: (i) [23, Proposition 1.23] and [5, Equation (10)], two special cases of Theorem 1 that were stated in a family of reversible queueing systems, namely Whittle networks without internal transitions (see Section 4.1), and (ii) the reversibility of the macrostate process of a quasi-reversible queueing system discussed in Section 2.2.3.

▶ Theorem 1. Consider a quasi-reversible queueing system $Q = (S, |\cdot|, q)$, and let Π denote one of its stationary measures. For each balance function $\Gamma \in \mathcal{B}_{|S|}$, the triplet $Q_{\partial\Gamma} = (S, |\cdot|, q_{\partial\Gamma})$ is a quasi-reversible queueing system, and its stationary measures are obtained by taking $\frac{1}{Z}\Pi_{\partial\Gamma}$ for any Z > 0, where

$$\Pi_{\partial\Gamma}(s) = \Pi(s)\Gamma(s), \quad \text{for each } s \in \mathcal{S}.$$
 (11)

Proof. We proceed in two steps: first we prove that $Q_{\partial\Gamma}$ satisfies Assumption 1, and then we prove (at once) that it is quasi-reversible and that its stationary measures are proportional to (11).

Step 1: $Q_{\partial\Gamma}$ satisfies Assumption 1. Let us first show that $Q_{\partial\Gamma}$ satisfies Assumption 1 and that its recurrent class is $S_{\partial\Gamma,\text{rec}} = S_{\text{rec}} \cap S_{\Gamma>0}$, where S_{rec} is the recurrent class of Q and $S_{\Gamma>0} = \{s \in S; \Gamma(s) > 0\}$. We verify each point of Assumption 1 one after the other:

- 1. Let $s \in \mathcal{S}_{\partial\Gamma,\text{rec}}$. Since $s \in \mathcal{S}_{\text{rec}}$, there is a sequence $\emptyset = t_1, t_2, \dots, t_\ell = s$ in \mathcal{S} such that $|t_k| \leq |t_{k+1}|$ and $q(t_k, t_{k+1}) > 0$ for each $k \in [1, \ell 1]$. Combining $\Gamma(s) > 0$, (6), (10), and the microstate-independence of $\partial\Gamma$, we obtain that $q_{\partial\Gamma}(t_k, t_{k+1}) > 0$ for $k \in [1, \ell 1]$.
- **2.** Let $s \in \mathcal{S} \setminus \mathcal{S}_{\partial\Gamma, \text{rec}}$: either $s \notin \mathcal{S}_{\text{rec}}$, or $s \notin \mathcal{S}_{\Gamma > 0}$, or both:
 - If $s \notin \mathcal{S}_{rec}$, we know that there is no sequence $\emptyset = t_1, t_2, \ldots, t_{\ell} = s$ in \mathcal{S} such that $q(t_k, t_{k+1}) > 0$ for each $k \in [1, \ell 1]$, and the same follows if we replace q with $q_{\partial \Gamma}$ by (6).
 - If $s \notin \mathcal{S}_{\Gamma>0}$, we have $\Gamma(t) = 0$ for each $t \in \mathcal{S}$ such that $|t| \geq |s|$ because $\mathcal{S}_{\Gamma>0}$ is a Ferrers set. Combining this observation with (6) and (10), we conclude that for each sequence $\emptyset = t_1, t_2, \ldots, t_{\ell} = s$, there must exit an index $k \in [1, \ell-1]$ such that $q_{\partial\Gamma}(t_k, t_{k+1}) = 0$.
- 3. Let $s \in \mathcal{S}$. There is a sequence $s = t_1, t_2, \dots, t_\ell = \emptyset$ in \mathcal{S} such that $|t_k| \geq |t_{k+1}|$ and $q(t_k, t_{k+1}) > 0$ for each $k \in [1, \ell 1]$. To conclude, it suffices to observe that $q_{\partial \Gamma}(t_k, t_{k+1}) = q(t_k, t_{k+1})$ for each $k \in [1, \ell 1]$.
- 4. We have $|\mathcal{S}_{\partial\Gamma,\text{rec}}| = |\mathcal{S}_{\text{rec}}| \cap \text{supp}(\Gamma)$, where $\text{supp}(\Gamma) = \{x \in |\mathcal{S}|; \Gamma(x) > 0\}$. The set $|\mathcal{S}_{\text{rec}}|$ is a Ferrers set because Q satisfies Assumption 1. The set $\text{supp}(\Gamma)$ also is also a Ferrers set because $\Gamma \in \mathcal{B}_{|\mathcal{S}|}$. Hence, $|\mathcal{S}_{\partial\Gamma,\text{rec}}|$ also satisfies these two properties.

Step 2: Stationary distribution and quasi-reversibility. Now, to show that $Q_{\partial\Gamma}$ is quasi-reversible and admits $\Pi_{\partial\Gamma}$ given by (11) as a stationary measure, it is enough to show that $\Pi_{\partial\Gamma}$ satisfies the following partial balance equations:

$$\Pi_{\partial\Gamma}(s) \sum_{t \in \mathcal{S}_{|s|+e_i}} q(s,t) \partial\Gamma_i(s) = \sum_{t \in \mathcal{S}_{|s|+e_i}} \Pi_{\partial\Gamma}(t) q(t,s), \text{ for each } s \in \mathcal{S}, i \in [1, n],$$
(12)

and

$$\Pi_{\partial\Gamma}(s) \left(\sum_{i=1}^{n} \sum_{t \in \mathcal{S}_{|s|-e_i}} q(s,t) + \sum_{t \in \mathcal{S}_{|s|}} q(s,t) \right) \\
= \sum_{t \in \mathcal{S}_{|s|-e_i}} \Pi_{\partial\Gamma}(t) q(t,s) \partial\Gamma_i(s) + \sum_{t \in \mathcal{S}_{|s|}} \Pi_{\partial\Gamma}(t) q(t,s), \quad \text{for each } s \in \mathcal{S}. \quad (13)$$

Indeed, (12)–(13) is the counterpart of (2)–(3) in $Q_{\partial\Gamma}$, and we observed below Definition 1 that (2)–(3) was equivalent to stationarity and quasi-reversibility. It will be useful to observe that, if we see Γ as a function on the microstate space \mathcal{S} , (7) can be rewritten as

$$\partial \Gamma_i(s)\Gamma(s) = \Gamma(t), \quad \text{for each } s \in \mathcal{S}, i \in [1, n], \text{ and } t \in \mathcal{S}_{|s|+e_i}.$$
 (14)

Let us first prove (12). For each $s \in \mathcal{S}$ and $i \in [1, n]$, we have successively, starting from the left-hand side of (12),

$$\begin{split} \Pi_{\partial\Gamma}(s) \sum_{t \in \mathcal{S}_{|s|+e_i}} q(s,t) \partial\Gamma_i(s) &= \Gamma(s) \partial\Gamma_i(s) \Pi(s) \sum_{t \in \mathcal{S}_{|s|+e_i}} q(s,t), & \text{by (11)}, \\ &= \Gamma(s) \partial\Gamma_i(s) \sum_{t \in \mathcal{S}_{|s|+e_i}} \Pi(t) q(t,s), & \text{by (2)}, \\ &= \sum_{t \in \mathcal{S}_{|s|+e_i}} \Pi(t) \Gamma(t) q(t,s), & \text{by (14)}, \\ &= \sum_{t \in \mathcal{S}_{|s|+e_i}} \Pi_{\partial\Gamma}(t) q(t,s), & \text{by (11)}. \end{split}$$

Now let us prove (13). For each $s \in \mathcal{S}$, we have successively, starting from the left-hand side of (13),

$$\begin{split} \Pi_{\partial\Gamma}(s) \left(\sum_{i=1}^n \sum_{t \in \mathcal{S}_{|s|-e_i}} q(s,t) + \sum_{t \in \mathcal{S}_{|s|}} q(s,t) \right) \\ &= \Gamma(s) \Pi(s) \left(\sum_{i=1}^n \sum_{t \in \mathcal{S}_{|s|-e_i}} q(s,t) + \sum_{t \in \mathcal{S}_{|s|}} q(s,t) \right), \qquad \text{by (11)}, \\ &= \Gamma(s) \left(\sum_{i=1}^n \sum_{t \in \mathcal{S}_{|s|-e_i}} \Pi(t) q(t,s) + \sum_{t \in \mathcal{S}_{|s|}} \Pi(t) q(t,s) \right), \qquad \text{by (3)}, \\ &= \sum_{i=1}^n \sum_{t \in \mathcal{S}_{|s|-e_i}} \Pi(t) \Gamma(t) q(t,s) \partial \Gamma_i(t) + \sum_{t \in \mathcal{S}_{|s|}} \Pi(t) \Gamma(t) q(t,s), \quad \text{by (14)}, \\ &= \sum_{i=1}^n \sum_{t \in \mathcal{S}_{|s|-e_i}} \Pi_{\partial\Gamma}(t) q(t,s) \partial \Gamma_i(t) + \sum_{t \in \mathcal{S}_{|s|}} \Pi_{\partial\Gamma}(t) q(t,s), \quad \text{by (11)}. \end{split}$$

This completes the proof of Theorem 1.

3.4 Gradient of parameterized systems

In the literature, results like Theorem 1 have been applied mainly to evaluate performance, i.e., to compute an average performance criterion that can be written as an expectation under the stationary distribution, when this expectation exists; see for instance [5, 16]. Here, we instead show a consequence of Theorem 1 in case of parameterized balance functions, as explained below. This results is useful for sensitivity analysis and in the context of policy-gradient reinforcement learning (PGRL), as we will develop in Section 6.

Consider a quasi-reversible queueing system $Q = (S, |\cdot|, q)$ and let $d \in \mathbb{N}_{>0}$. We call parameterization of the balance function a mapping $\theta \in \mathbb{R}^d \mapsto \Gamma_\theta \in \mathcal{B}_{|S|}$ such that:

- 1. For each $s \in \mathcal{S}$, the function $\theta \in \mathbb{R}^d \to \Gamma_{\theta}(s)$ is differentiable and positive, so that in particular $\nabla_{\theta} \log \Gamma_{\theta}(s)$ is well defined for each $\theta \in \mathbb{R}^d$.
- 2. For each $\theta \in \mathbb{R}^d$, the queueing system $Q_{\theta} \triangleq Q_{\partial \Gamma_{\theta}}$, which is quasi-reversible by Theorem 1, is also positive recurrent.

A simple example of parameterization is given by $\Pi_{\theta}(s) = \prod_{i=1}^{n} \sigma(\theta_i)^{|s|_i}$, where d = n and $\sigma : x \in \mathbb{R} \mapsto 1/(1 + \mathrm{e}^{-x}) \in \mathbb{R}$ is a sigmoid function. One can then verify that the set of balanced policies generated by letting θ vary in \mathbb{R}^n is the set of static policies mentioned in Section 3.2. In general, in practice, the parameterization should be such that we can compute $\Gamma_{\theta}(s)$ and $\nabla_{\theta} \log \Gamma_{\theta}(s)$ efficiently, for each $\theta \in \mathbb{R}^d$ and $s \in \mathcal{S}_{rec}$ (e.g., in closed form or by automatic differentiation).

For each $\theta \in \mathbb{R}^d$, let $\Pi_{\theta} \triangleq \Pi_{\partial \Gamma_{\theta}}$ denote the stationary distribution of Q_{θ} , so that

$$\log \Pi_{\theta}(s) = \log \Pi(s) + \log \Gamma_{\theta}(s) - \log Z_{\theta}, \quad \text{for each } s \in \mathcal{S}_{rec}, \tag{15}$$

$$Z_{\theta} = \sum_{s \in \mathcal{S}_{rec}} \Pi(s) \Gamma_{\theta}(s), \tag{16}$$

where (15) follows from (11), and (16) from the normalizing condition $\sum_{s \in \mathcal{S}} \Pi_{\theta}(s) = 1$. Corollary 2 below gives a closed-form expression for the gradient $\nabla_{\theta} \log \Pi_{\theta}(s)$ for each $s \in \mathcal{S}_{rec}$, when this gradient is properly defined. This result can be seen as an extension of [12, Equation (12) in Theorem 1], and it remains valid if we replace \mathbb{R}^d with any open set $\Omega \subseteq \mathbb{R}^d$. The key take-away of (17) is that $\nabla_{\theta} \log \Pi_{\theta}(s)$ can be written as an expectation under the distribution Π_{θ} .

▶ Corollary 2. Consider a quasi-reversible queueing system $Q = (S, |\cdot|, q)$ and a parameterization $\theta \in \mathbb{R}^d \mapsto \Gamma_\theta \in \mathcal{B}_{|S|}$ of the balance function. Assume that, for each $\theta \in \mathbb{R}^d$, Q_θ is positive recurrent and the random vector $\nabla_\theta \log \Pi_\theta(S_\theta)$, where $S_\theta \sim \Pi_\theta$, is integrable. Then, for each $\theta \in \mathbb{R}^d$, we have

$$\nabla_{\theta} \log \Pi_{\theta}(s) = \nabla_{\theta} \log \Gamma_{\theta}(s) - \mathbb{E} \left(\nabla_{\theta} \log \Gamma_{\theta}(S_{\theta}) \right), \quad \text{for each } s \in \mathcal{S}_{rec}. \tag{17}$$

Proof. Let $\theta \in \mathbb{R}^d$. Let us first compute $\nabla_{\theta} \log Z_{\theta}$, and then we will inject this result to compute $\nabla_{\theta} \log \Pi_{\theta}(s)$ for each $s \in \mathcal{S}$. We have successively:

$$\nabla_{\theta} \log Z_{\theta} \stackrel{\text{(a)}}{=} \frac{1}{Z_{\theta}} \sum_{s \in \mathcal{S}_{\text{rec}}} \Pi(s) \Gamma_{\theta}(s) \nabla_{\theta} \log \Gamma_{\theta}(s)$$

$$\stackrel{\text{(b)}}{=} \sum_{s \in \mathcal{S}_{\text{rec}}} \Pi_{\theta}(s) \nabla_{\theta} \log \Gamma_{\theta}(s)$$

$$\stackrel{\text{(c)}}{=} \mathbb{E} \left(\nabla_{\theta} \log \Gamma_{\theta}(S_{\theta}) \right), \tag{18}$$

where (a) follows by applying the rule $\nabla_{\theta} f(\theta) = f(\theta) \nabla \log f(\theta)$ to $f \in \{\Gamma, Z\}$ and applying (16), (b) follows by definition of the stationary distribution Π_{θ} , and (c) follows by recognizing an expectation under this distribution. Now, for each $s \in \mathcal{S}$, we have

$$\nabla_{\theta} \log \Pi_{\theta}(s) \stackrel{(d)}{=} \nabla_{\theta} \log \Gamma_{\theta}(s) - \nabla_{\theta} \log Z_{\theta} \stackrel{(e)}{=} \nabla_{\theta} \log \Gamma_{\theta}(s) - \mathbb{E} \left(\nabla_{\theta} \log \Gamma_{\theta}(S_{\theta}) \right),$$

where (d) follows by differentiating (15) and (e), by injecting (18).

4 Two canonical examples

We consider two well-known queueing systems that are quasi-reversible in the sense of Section 2.1, and provide different interpretations of the distinction between microstates and macrostates introduced in Section 2.1.1. We then briefly explain how the concepts of Section 3 can be applied in these two examples. Notation that was already introduced in Section 2.1 is reused in a consistent way.

4.1 Multi-class Whittle process

We first address a particular case of multi-class Whittle processes [23, Section 3.1]. Let n and m be two positive integers, and let $\mathcal{L} = [\![1,n]\!] \times [\![1,m]\!]$. We consider a Markovian discrete-event system in which elements, which are labelled in \mathcal{L} , come, change labels, and eventually leave. We suppose that the events are of one of the three following types:

- \blacksquare Arrival of an element of label $ik \in \mathcal{L}$;
- Internal transition of an element from a label $ik \in \mathcal{L}$ to a label $i\ell \in \mathcal{L}$,
- Departure of an element of label $ik \in \mathcal{L}$.

At any time, the state of the system is represented by the matrix $s \in \mathcal{M}_{n,m}(\mathbb{N})$, where for all $ik \in \mathcal{L}$, s_{ik} is the number of elements having label ik in the system. We let $\mathcal{S} \subseteq \mathcal{M}_{n,m}(\mathbb{N})$ be the state space of the process, to be be entailed by the transitions below. We denote

by $0 \in \mathcal{S}$, the null matrix. We assume that the only non-zero transitions of the underlying CTMC are the following ones,

$$\begin{cases}
q(s, s + e_{i\ell}) = P_{0,i\ell}\phi_0, & s \in \mathcal{S}, i\ell \in \mathcal{L}, \\
q(s, s - e_{ik} + e_{i\ell}) = P_{ik,i\ell}\phi_{ik}(s), & s \in \mathcal{S}, k, \ell \in [1, m]^2, i \in [1, n] \text{ s.t. } s_{ik} > 0, \\
q(s, s - e_{ik}) = P_{ik,0}\phi_{ik}(s), & s \in \mathcal{S}, ik \in \mathcal{L} \text{ s.t. } s_{ik} > 0,
\end{cases}$$
(19)

where $\phi_0 > 0$, $\phi_{ik}(s) > 0$ for each $s \in \mathcal{S}$ and $ik \in \mathcal{L}$ such that $s_{ik} > 0$, and $P = (P_{x,y})_{(x,y)\in(\{0\}\cup(\mathcal{L}))^2}$ is a stochastic matrix on $(\{0\}\cup(\mathcal{L}))^2$ that is irreducible, in the sense that for any $ik \in \mathcal{L}$, there exists $p \geq 1$ and a p-uple $(k_1, \dots, k_p) \in [1, m]^p$ that includes k and is such that

$$P_{0,ik_1} \left(\prod_{l=1}^{p-1} P_{ik_l,ik_{l+1}} \right) P_{ik_p,0} > 0, \tag{20}$$

where, throughout, a product over an empty set of indices is understood as 1. It follows that the set of equations

$$\lambda_{ik} = \lambda_0 P_{0,ik} + \sum_{\ell=1}^{m} \lambda_{i\ell} P_{i\ell,ik}, \quad ik \in \mathcal{L},$$

admits a unique solution $(\lambda_x)_{x\in\{0\}\cup\mathcal{L}}$ in $(\mathbb{R}_+)^{1+nm}$ up to a positive multiplicative constant. Therefore, setting $\lambda_0 = 1$, the system

$$\lambda_{ik} = P_{0,ik} + \sum_{\ell=1}^{m} \lambda_{i\ell} P_{i\ell,ik}, \quad ik \in \mathcal{L},$$
(21)

admits a unique solution $(\lambda_{ik})_{ik\in\mathcal{L}}$ in $(\mathbb{R}_+)^{nm}$.

As we now check, the model can be cast in the framework of Section 2.1, by viewing *elements* as *customers*. It follows that the counting function $|\cdot|$ can be rewritten as

$$|\cdot|: \begin{cases} \mathcal{S} & \longrightarrow \mathbb{N}^n; \\ s & \longmapsto |s| \text{ s.t. } |s|_i = \sum_{k=1}^m s_{ik}, \quad i \in \llbracket 1, n \rrbracket, \end{cases}$$

and we have the following.

▶ Lemma 3. The triple $Q = (S, |\cdot|, q)$ is a queueing system in the sense of Section 2.1. Proof. See Appendix C.

▶ **Definition 3.** The service rate functions $\phi := \{\phi_x ; x \in \mathcal{L}\}$ are said to be Φ -balanced if, for some function $\Phi : \mathcal{S} \to \mathbb{R}_{>0}$,

$$\Phi(s)\phi_0 = \Phi(s + e_{i\ell})\phi_{i\ell}(s + e_{i\ell}), \quad s \in \mathcal{S}, \ i\ell \in \mathcal{L} \ s.t. \ s + e_{i\ell} \in \mathcal{S}.$$
(22)

It is a well-known fact, that Whittle networks without internal routing (namely, no internal transitions in the definition above, or in other words, $P_{ik,i\ell} = 0$ for all i,k,ℓ) are reversible when the service rates are balanced, see e.g. Chapter 1 of [23]. Given the form of the counting function above, it is therefore natural that the same model becomes quasi-reversible in the sense of Definition 1, if internal routing are allowed. This fact is rigourously proven in Proposition 4 below. In particular, the results of Section 3 will then imply that balanced arrival rates can be applied to this network, to obtain a quasi-reversible queueing system. Note that the application of balanced arrival control to Whittle networks without internal routing was already considered in [5, 16].

▶ **Proposition 4.** Assume that the mappings ϕ are Φ -balanced, and consider the positive measure Π on S defined by

$$\Pi(s) = \Phi(s) \prod_{ik \in \mathcal{L}} \lambda_{ik}^{s_{ik}}, \quad s \in \mathcal{S}.$$
(23)

Then, the queueing system $Q := (S, |\cdot|, q)$ has stationary measures proportional to Π . Moreover Q is quasi-reversible in the sense of Definition 1.

Proof. Let us first verify that the assumptions of [23, Theorem 3.1] are satisfied, which will allow us to conclude directly that Π defined by (23) is a stationary measure of the CTMC of the model. First observe that for all $s \in \mathcal{S}$ and all $(ik, i\ell) \in \mathcal{L}^2$ such that $s_{ik} > 0$,

$$\Phi(s)\phi_{ik}(s) = \Phi(s - e_{ik} + e_{i\ell})\phi_{i\ell}(s - e_{ik} + e_{i\ell}).$$

This equality follows by applying (22) to $s := s - e_{ik}$, and successively to $i\ell$ and to ik, to obtain that

$$\Phi(s - e_{ik} + e_{i\ell})\phi_{i\ell}(s - e_{ik} + e_{i\ell}) = \Phi(s - e_{ik})\phi_0,$$

= $\Phi(s - e_{ik} + e_{ik})\phi_{ik}(s - e_{ik} + e_{ik}),$
= $\Phi(s)\phi_{ik}(s).$

Moreover, summing (21) over $i \in [1, n]$ for all fixed $k \in [1, m]$, yields to

$$\sum_{i=1}^{n} \lambda_{ik} P_{ik,0} + \sum_{i=1}^{n} \sum_{\ell=1}^{m} \lambda_{ik} P_{ik,i\ell} = \sum_{i=1}^{n} P_{0,ik} + \sum_{\ell=1}^{m} \sum_{i=1}^{n} \lambda_{i\ell} P_{i\ell,ik}, \quad k \in [1, m],$$

which is in turn equivalent to

$$\sum_{i=1}^{n} P_{0,ik} = \sum_{i=1}^{n} \lambda_{ik} P_{ik,0}, \quad k \in [1, m].$$
(24)

In particular, this readily implies that

$$\sum_{ik\in\mathcal{L}} P_{0,ik} = \sum_{ik\in\mathcal{L}} \lambda_{ik} P_{ik,0}.$$

Therefore, all the assumptions of [23, Theorem 3.1] are satisfied, showing that Π defined by (23) is a stationary measure of the CTMC of the model.

We finally check that the system Q is quasi-reversible in the sense of Definition 1. For this, take $s \in \mathcal{S}$ and $i \in [1, n]$. We have that

$$\sum_{t \in \mathcal{S}_{|s|+e_i}} \Pi(t)q(t,s) = \sum_{\ell=1}^m \Pi(s+e_{i\ell})P_{i\ell,0}\phi_{i\ell}(s+e_{i\ell}),$$

$$= \sum_{\ell=1}^m P_{i\ell,0}\Phi(s+e_{i\ell})\phi_{i\ell}(s+e_{i\ell}) \left(\prod_{ik \in \mathcal{L}} \lambda_{ik}^{(s+e_{i\ell})_{ik}}\right),$$

$$= \Phi(s)\phi_0 \sum_{\ell=1}^m P_{i\ell,0}\lambda_{i\ell} \left(\prod_{ik \in \mathcal{L}} \lambda_{ik}^{s_{ik}}\right),$$

$$= \Pi(s)\phi_0 \sum_{\ell=1}^m P_{i\ell,0}\lambda_{i\ell},$$

16

$$\begin{split} &= \Pi(s)\phi_0 \sum_{\ell=1}^m P_{0,i\ell}, \\ &= \Pi(s) \sum_{\ell=1}^m \phi_0 P_{0,i\ell}, \\ &= \Pi(s) \sum_{t \in \mathcal{S}_{\text{baller}}} q(s,t), \end{split}$$

applying (22) in the third equality and (24) in the fifth one. But as was noted above, Π is a stationary measure for the considered CTMC. This concludes the proof.

- ▶ Example 1. This model is very versatile. The key assumption is that the first index of the label is the class of the customer and remains constant over time. The second index, on the other hand, can change and convey other information about the customer, such as the queue at which is currently in service, the sequence of queues it visits in the network, or its service requirement. Here is a (non-exhaustive) list of particular cases:
- (Multiclass) Jackson networks, as pointed out in [23, Section 3.1], when $\phi_{ik}(s)$ depends only on s_{ik} , for all s, i, k;
- \blacksquare (Simple) Whittle networks, when m = 1;
- As was observed above, Whittle networks without internal routing, whenever $P_{ik,i\ell} = 0$ for $i \in [1, n]$ and $k, \ell \in [1, m]$ such that $k \neq \ell$. In this case, the queueing system is not only quasi-reversible, but also reversible.
- The method of stages [17, 1] can also be applied to capture more general service-time distributions.

4.2 Order-independent (OI) queue

Consider a queue with n customer classes, where the queue state is described by the word $s = s_1 s_2 \cdots s_\ell \in \mathcal{S} \subseteq [\![1,n]\!]^*$, where $\ell \in \mathbb{N}$ is the number of customers in the queue and, if $\ell \neq 0$, s_p is the class of the p-th oldest customer, for each $p \in [\![1,\ell]\!]$. We assume that the microstate space \mathcal{S} is:

- Stable by prefix, in the sense that for any $s \in \mathcal{S}$, $\breve{s} \in \mathcal{S}$ for any prefix \breve{s} of s;
- Stable by permutation, namely, if $s = s_1 \cdots s_\ell \in \mathcal{S}$, then any permutation $s_{\sigma(1)} \cdots s_{\sigma(\ell)}$ of s is also an element of \mathcal{S} .

To cast this model in the framework of Section 2.1, the counting function $|\cdot|$ takes the form

$$|\cdot|: \begin{cases} \mathcal{S} & \longrightarrow \mathbb{N}^n; \\ s = s_1 \cdots s_\ell & \longmapsto |s|, \text{ s.t. } |s|_i = \sum_{k=1}^{\ell} \mathbf{1}_{s_k = i}, \quad i \in [1, n]. \end{cases}$$

For each $i \in [\![1,n]\!]$, we assume that class-i customers arrive according to a Poisson process with rate $\nu_i > 0$, and that there exists a function $\mu : |\mathcal{S}| \to \mathbb{R}_{\geq 0}$, called the *rate* function, such that for each microstate $s \in \mathcal{S}$, $\mu(s)$ gives the total departure rate of customers in state s. We assume that the function μ is such that $\mu(\mathbf{0}) = 0$, $\mu(s) > 0$ for each $s \in \mathcal{S} \setminus \{\emptyset\}$, and μ is component-wise non-decreasing, in the sense that $\mu(|s|) \leq \mu(|s| + e_i)$ for each $s \in \mathcal{S}$ and $i \in [\![1,n]\!]$ so that $|s| + e_i \in |\mathcal{S}|$. For each $s = s_1 s_2 \cdots s_\ell \in \mathcal{S} \setminus \{\emptyset\}$, the departure rate of the customer in position $p \in [\![1,\ell]\!]$, of class s_p , is given by

$$\Delta\mu(s_1s_2\cdots s_p) := \mu(s_1s_2\cdots s_p) - \mu(s_1s_2\cdots s_{p-1}).$$

In particular, although the *total* departure rate only depends on the macrostate, how this rate is distributed among customers also depends on the microstate. As a consequence, with the notation of Section 2.1, the microstate CTMC's transition kernel is given by

$$q(s,t) = \begin{cases} \nu_i & \text{if } t = s_1 \cdots s_{\ell}i, \text{ for each } i \text{ s.t. } s_1 \cdots s_{\ell}i \in \mathcal{S}, \\ \Delta \mu(s_1 s_2 \cdots s_p) & \text{if } t = s_1 \cdots s_{p-1} s_{p+1} \cdots s_{\ell}, \text{ for each } p \in [1, \ell], \end{cases}$$
 (25)

and q(s,t) = 0 otherwise. More details about OI queues can be found in [19]. We have the following results.

▶ Lemma 5. The triple $Q = (S, |\cdot|, q)$ is a queueing system in the sense of Section 2.1.

Proof. See Appendix D.

As for Whittle networks, Proposition 6 below recalls that the OI queue defined above is quasi-reversible, implying again that, thanks to the results of Section 3, we can apply balanced arrival rates to OI queues and still obtain a quasi-reversible queueing system. While it was already known that OI queues were quasi-reversible, we believe that the application of state-dependent arrival rates to OI queues was only considered in [9] in the special case of redundancy systems. Besides this, to the best of our knowledge, the only state-dependent arrival rates that were considered so far in the context of OI queues only depended on the total number of customers in the system [3, 4, 19].

ightharpoonup Proposition 6. The triple Q is quasi-reversible in the sense of Definition 1, and its stationary measures are given by

$$\Pi(s) \propto \prod_{p=1}^{\ell} \frac{\nu_{s_p}}{\mu(s_1 s_2 \cdots s_p)}, \quad s_1 s_2 \cdots s_{\ell} \in \mathcal{S}.$$

$$(26)$$

Proof. Equation (26) is shown in [3, Theorem 1], and the quasi-reversibility property is exactly (4) in [3].

▶ Example 2. As shown in [6], the redundancy model introduced in [15] can be seen as an OI queue when the servers apply the first-come-first-served (FCFS) scheduling policy. An extension of this model with customer abandonments will be considered in the case study of Section 6. As shown in [11], the *stochastic matching model* introduced in [20] is also an OI queue when the matching policy is first-come-first-matched (FCFM). In [21], the product-form result specializing (26) to the FCFM matching model, is shown using a dynamic reversibility argument for an extended version of the state space \mathcal{S} . See also [2] for the corresponding model supplemented with self-loops in the compatibility graph.

4.3 Equivalence between Whittle networks and OI queues

Proposition 7 below shows that multiclass Whittle networks and OI queues supplemented with balanced arrival control are macroscopically equivalent in steady-state. This result extends [6, Theorem 2] and [10, Theorem 3.2]. In the special case where the Whittle network has no internal routing (and is therefore reversible - see Example 1), this result can be seen as a special case of the digression of Section 2.2.3.

▶ Proposition 7. Consider two integers $n, m \in \mathbb{N}_{>0}$, a Ferrers set $\mathcal{X} \subseteq \mathcal{M}_{n,m}(\mathbb{N})$, and balance function $\Gamma \in \mathcal{B}_{\mathcal{X}}$. Consider the following two systems:

- A Whittle network with state space \mathcal{X} and Φ -balanced service rates ϕ , as defined in Section 4.1;
- An OI queue as defined in Section 4.2, with nm classes, state space S such that |S| = Xwhen replacing n with nm in the definition of $|\cdot|$, and with arrival rates $\nu_{ik} := P_{0,ik}\phi_0$, $ik \in$ $[1,n] \times [1,m]$, and departure rate function μ such that

$$\mu(x) = \sum_{i=1}^{n} \sum_{k=1}^{m} \frac{\Phi(x - e_{ik})}{\Phi(x)}, \quad x \in \mathcal{X}, \quad ik \in [1, n] \times [1, m] : x_{ik} > 0.$$

Suppose that the two models are also supplemented with the same Γ -balanced admission control. Then, the macrostate of the controlled OI queue has the same distribution as the controlled Whittle network at equilibrium, that is, if we denote by $\Pi_{W,\Gamma}$ and $\Pi_{OL\Gamma}$, the stationary measures of these systems, and assume that $\Pi_{W,\Gamma}(\mathbf{0}) = \Pi_{OI,\Gamma}(\varnothing)$, we have

$$\Pi_{W,\Gamma}(x) = \sum_{s \in \mathcal{S}: |s| = x} \Pi_{OI,\Gamma}(s), \quad x \in \mathcal{X}.$$

Proof. With these assumptions on the two uncontrolled models, we fall into the settings of Chapter 3 of [10]. Indeed, in the Whittle network, in any state $x \in \mathcal{X}$ the arrival rates to the various classes are precisely $P_{0,ik}\phi_0 = \nu_{ik}, ik \in [1,n] \times [1,m]$, whereas the departure rates are given by $\phi_{ik}(x)$, $ik \in [1, n] \times [1, m]$. Therefore, we can apply [10, Theorem 3.2], entailing that the result holds for the uncontrolled systems ($\Gamma \equiv 1$), that is, with obvious notation,

$$\Pi_{\mathrm{W,Id}}(x) = \sum_{s \in \mathcal{S}: |s| = x} \Pi_{\mathrm{OI,Id}}(s), \quad x \in \mathcal{X},$$

since by our very assumption, we have $\Pi_{W,Id}(0) = \Pi_{OI,Id}(\emptyset)$. But from Theorem 1, Proposition 4, and Proposition 6, we also have $\Pi_{W,\Gamma}(x) = \Pi_{W,\mathrm{Id}}(x)\Gamma(x)$ and $\Pi_{\mathrm{OI},\Gamma}(s) =$ $\Pi_{\text{OLId}}(s)\Gamma(x)$ for each $x \in \mathcal{X}$ and $s \in \mathcal{S}$ such that |s| = x, which concludes the proof.

▶ Remark 1. This result is reminiscent of the discussion of Section 2.2.3. It shows that we in fact have $\Pi_{OI,\Gamma}^{\star} = \Pi_{W,\Gamma}$, that is, the (truly reversible, as an immediate consequence of Proposition 4 in the case m=1) CTMC representing the controlled simple Whittle network has the same steady state distribution as the (non-Markov) macrostate process of the controlled OI queue.

5 Admission control

To illustrate balanced policies, we focus on the admission-control problem, which we define formally in Section 5.1. Section 5.2 characterizes the set of balanced admission-control policies and proves that there exists a deterministic policy that is optimal among balanced policies. In Section 5.3, we give insights into the optimality gap of balanced policies, i.e., the difference in performance between the best balanced policy and the optimal policy. To simplify the discussion, we focus throughout this section on queueing systems with a finite state space. We believe that the methodology developed in this section, and in particular Theorem 8, can be adapted to other arrival-control problems.

5.1 Admission-control problem and policies

We define an admission-control problem as a 3-tuple $\mathcal{P} = (Q, r_{\text{cont}}, r_{\text{disc}})$, where $Q = (\mathcal{S}, |\cdot|, q)$ is a quasi-reversible queueing system, $r_{\text{cont}}: \mathcal{S} \to \mathbb{R}$ is the occupation-reward(-rate) function, and $r_{\rm disc}: \mathcal{S} \times \mathcal{S} \to \mathbb{R}$ is the transition-reward function. For example, one can take $r_{\rm cont}(s) = -\sum_{i=1}^n \eta_i |s|_i$ and $r_{\rm disc}(s,t) = \sum_{i=1}^n r_i \mathbb{1}[|t| = |s| + e_i]$ for each $s,t \in \mathcal{S}$, where η_i is the holding cost rate of class i and r_i the admission reward of class i. An admission-control policy is a function $\gamma: \mathcal{S} \to [0,1]^n$, such that $\gamma_i(s)$ is interpreted as the probability that a class-i customer is admitted into the system while the microstate is s, for each $s \in \mathcal{S}$ and $i \in [\![1,n]\!]$. The queueing system under policy s0 is denoted by s1 and is as defined in Section 3.1. We are interested in solving the following optimization problem:

Maximize
$$\gamma: \mathcal{S} \to [0,1]^n, \\
\Pi_{\gamma}: \mathcal{S} \to [0,1]$$

$$G_{\gamma} \triangleq \sum_{s \in \mathcal{S}} \Pi_{\gamma}(s) r_{\text{cont}}(s) + \sum_{s,t \in \mathcal{S}} \Pi_{\gamma}(s) q_{\gamma}(s,t) r_{\text{disc}}(s,t), \tag{27a}$$

Subject to
$$\Pi_{\gamma}(s) \sum_{t \in \mathcal{S}} q_{\gamma}(s, t) = \sum_{t \in \mathcal{S}} \Pi_{\gamma}(t) q_{\gamma}(t, s)$$
, for each $s \in \mathcal{S}$, (27b)

$$\sum_{s \in S} \Pi_{\gamma}(s) = 1. \tag{27c}$$

Here, G_{γ} is called the *gain* of the policy γ , $r_{\text{cont}}(s)$ is the reward per time unit in microstate s, and $r_{\text{disc}}(s,t)$ is a reward received upon transition from state s to state t. Equations (27b)–(27c) say that Π_{γ} is the stationary distribution of Q_{γ} . Existence and unicity of Π_{γ} follow from the fact that Q_{γ} is unichain, which follows from our assumption that Q satisfies Assumption 1. Following the standard approach in Markov decision processs (MDPs) solving as described in [22, Section 8.8.1], (27) can be rewritten as a linear program; more details are given in Appendix A for the special case of reversible queueing systems. It follows in particular that the optimal policy is deterministic, i.e., it satisfies $\gamma_i(s) \in \{0,1\}$ for each $s \in \mathcal{S}$.

5.2 Balanced policies

Our goal in this section is to characterize the set of balanced admission-control policies and to identify a solution of (27) that is optimal among all balanced policies. Consider an admission-control problem $\mathcal{P} = (Q, r_{\text{cont}}, r_{\text{disc}})$, where $Q = (\mathcal{S}, |\cdot|, q)$ is a quasi-reversible queueing system with a finite microstate space \mathcal{S} , and let $\mathcal{X} = |\mathcal{S}|$.

Since there is a one-to-one mapping between the set of balanced policies and the set $\mathcal{B}_{\mathcal{X}}$ of balance functions, looking for the best balanced admission-control policy is equivalent to looking for the best balance function. By combining Theorem 1 and (7), we can rewrite the restriction of (27) to balanced policies as an optimization problem that is linear with respect to the balance function:

Maximize
$$G^{\partial\Gamma} = \sum_{x \in \mathcal{X}} \sum_{s \in \mathcal{S}} \Pi(s)\Gamma(x)r_{\text{cont}}(s)$$
 (28a)

$$+ \sum_{x \in \mathcal{X}} \sum_{i=1}^{n} \sum_{s \in \mathcal{S}_{x}} \sum_{t \in \mathcal{S}_{x+e_{i}}} \Pi(s)\Gamma(x+e_{i})q(s,t)r_{\text{disc}}(s,t)$$

$$+ \sum_{x \in \mathcal{X}} \sum_{s \in \mathcal{S}_{x}} \sum_{t \in \mathcal{S} \setminus \bigcup_{i=1}^{n} \mathcal{S}_{x+e_{i}}} \Pi(s)\Gamma(x)q(s,t)r_{\text{disc}}(s,t),$$

Subject to
$$\Gamma(x) \ge 0$$
 for each $x \in \mathcal{X}$, (28b)

$$\Gamma(x) \ge \Gamma(x + e_i)$$
 for each $x \in \mathcal{X}$ and $i \in [1, n]$, (28c)

$$\sum_{s \in S} \Pi(s)\Gamma(s) = 1,\tag{28d}$$

where Π is a stationary measure of Q, given by (1). One can verify that Γ^* is an optimal solution of (28) if and only if $\gamma^* = \partial \Gamma^*$ is the best solution of (27) among balanced

policies. Indeed, Equation (28a) is a specialization of (27a), simplified using (7) and (11); Equations (28b), (28c), and (28d) are satisfied if and only if (i) there exists Z>0 such that $\frac{1}{Z}\Gamma\in\mathcal{B}_{\mathcal{X}}$ and (ii) $\partial\Gamma_i(x)\in[0,1]$ for each $x\in\operatorname{supp}(\Gamma)$. Equation (28d) guarantees in particular that Z is such that $s\in\mathcal{S}\mapsto\Pi(s)\Gamma(s)$ is the stationary distribution of $Q_{\partial\Gamma}$, and not an arbitrary stationary measure.

Theorem 8 below gives the form of an optimal solution of (28) and, correspondingly, of an admission-control policy with maximal gain among all balanced policies. It shows in particular that, when we consider the restriction of problem (27) to balanced policies, the solution is a (balanced) deterministic policy.

▶ **Theorem 8.** Consider an admission-control problem $\mathcal{P} = (Q, r_{cont}, r_{disc})$, where $Q = (\mathcal{S}, |\cdot|, q)$ is a quasi-reversible queueing system with a finite microstate space \mathcal{S} . There exists a Ferrers set $\mathcal{A} \subseteq |\mathcal{S}|$ such that the function $\Gamma_{\mathcal{A}} : x \in \mathcal{X} \mapsto \frac{1}{Z}\mathbb{1}[x \in \mathcal{A}]$ is an optimal solution of (28), where $Z = \sum_{y \in \mathcal{A}} \Pi(y)$. Correspondingly, the following policy is an optimal solution of (27) among all balanced admission policies:

$$\gamma_i(x) = \mathbb{1}[x + e_i \in A], \text{ for each } x \in |S| \text{ and } i \in [1, n].$$

Proof. See Appendix B.

Though the detail of the proof of Theorem 8 can be found in Appendix B, let us briefly discuss the intuition behind this result. Problem (28) is a linear optimization problem, hence it has an optimal solution that is a vertex of its polytope $\tilde{\mathcal{B}}_{\mathcal{X}}^{\mathrm{ac}}$ of feasible solutions:

$$\tilde{\mathcal{B}}^{\mathrm{ac}}_{\mathcal{X}} = \left\{ \Gamma: \mathcal{X} \to \mathbb{R}_{\geq 0}; \Gamma \text{ is componentwise non-increasing and } \sum_{x \in \mathcal{X}} \Pi(x) \Gamma(x) = 1 \right\}.$$

To prove Theorem 8, it suffices to show that $\tilde{\mathcal{B}}_{\mathcal{X}}^{\mathrm{ac}}$ is the convex hull of $\{\Gamma_{\mathcal{A}}; \mathcal{A} \in \mathcal{F}_{\mathcal{X}}\}$, where $\mathcal{F}_{\mathcal{X}}$ is the family of Ferrers subsets of \mathcal{X} . Characterizing the vertices of $\tilde{\mathcal{B}}_{\mathcal{X}}^{\mathrm{ac}}$ is made complicated by the normalizing condition $\sum_{s \in \mathcal{S}} \Pi(s)\Gamma(s) = 1$ so, as an intermediary step, we focus on the state $\mathcal{B}_{\mathcal{X}}^{\mathrm{ac}} \subseteq \mathcal{B}_{\mathcal{X}}$ defined as follows:

$$\mathcal{B}^{ac}_{\mathcal{X}} = \left\{\Gamma: \mathcal{X} \to \mathbb{R}_{\geq 0}; \Gamma \text{ is componentwise non-increasing and } \Gamma(0) = 1\right\}.$$

The only difference between $\mathcal{B}_{\mathcal{X}}^{\mathrm{ac}}$ and $\tilde{\mathcal{B}}_{\mathcal{X}}^{\mathrm{ac}}$ is the normalizing condition, given by $\Gamma(0) = 1$ for $\mathcal{B}_{\mathcal{X}}^{\mathrm{ac}}$ and $\sum_{x \in \mathcal{X}} \Pi(x) \Gamma(x) = 1$ for $\tilde{\mathcal{B}}_{\mathcal{X}}^{\mathrm{ac}}$. Proposition 9 below characterizes the vertices of $\mathcal{B}_{\mathcal{X}}^{\mathrm{ac}}$. It is used as an intermediary step in the proof of Proposition 10, which characterizes the vertices of $\tilde{\mathcal{B}}_{\mathcal{X}}^{\mathrm{ac}}$.

- ▶ Proposition 9 (Theorem 4.2 in [16]). Let $n \in \mathbb{N}$ and consider a Ferrers set $\mathcal{X} \subseteq \mathbb{N}^n$. The set $\mathcal{B}_{\mathcal{X}}$ is the convex hull of $\{\mathbb{1}_{\mathcal{A}}; \mathcal{A} \in \mathcal{F}_{\mathcal{X}}\}$, where $\mathcal{F}_{\mathcal{X}}$ is the family of Ferrers subsets of \mathcal{X} , and $\mathbb{1}_{\mathcal{A}}: x \in \mathcal{X} \mapsto \mathbb{1} [x \in \mathcal{A}]$.
- ▶ Proposition 10. Let $n \in \mathbb{N}$ and consider a Ferrers set $\mathcal{X} \subseteq \mathbb{N}^n$. The set $\tilde{\mathcal{B}}_{\mathcal{X}}$ is the convex hull of $\{\Gamma_{\mathcal{A}}; \mathcal{A} \in \mathcal{F}_{\mathcal{X}}\}$, where

$$\Gamma_{\mathcal{A}}: x \in \mathcal{X} \mapsto \frac{1}{\sum_{y \in \mathcal{A}} \Pi(y)} \mathbb{1} [x \in \mathcal{A}], \text{ for each } \mathcal{A} \in \mathcal{F}_{\mathcal{X}}.$$

Proofs of Propositions 9 and 10. See Appendix B.

Observe that Proposition 9 is a rephrasing of [16, Theorem 4.2], which was stated in the special case of Whittle networks without internal transitions. We propose an alternative proof of this result that is more constructive. Our proof shows in particular that, even if $\mathcal{B}_{\mathcal{X}}$ has as many vertices as there are Ferrers subsets of \mathcal{X} , a particular $\Gamma \in \mathcal{B}_{\mathcal{X}}$ can be written as a convex combination of only $\operatorname{card}(\mathcal{X})$ of these functions.

5.3 The cost of balance

Section 6 will illustrate one benefit of balanced policies when it comes to optimizing performance in systems with an infinite state space. Before that, we make a digression to give insight into the performance loss of balanced policies with respect to an optimal policy, still focusing on admission control and systems with a finite state space for simplicity. We combine insights from the literature on (quasi-)reversibility with numerical results in toy admission-control examples solved by linear programming.

5.3.1 Dimension reduction

Consider an admission-control problem $\mathcal{P}=(Q,r_{\mathrm{cont}},r_{\mathrm{disc}})$, where $Q=(\mathcal{S},|\cdot|,q)$ has a finite state space \mathcal{S} , and let $\mathcal{X}=|\mathcal{S}|$. An obvious but insightful observation is that the balance condition reduces the degree of freedom in the choice of the admission probabilities, as the admission policy $\gamma=\delta\Gamma$ is fully specified once the balance function Γ is given. The number of values necessary to specify these, is therefore reduced from $\Theta(nm)$ to $\Theta(m)$, where n is the number of classes and $m=\mathrm{card}(\mathcal{X})$ is the number of states.

Monotonicity

One way of illustrating this reduction of dimensionality consists of describing how it restricts the set of feasible admission probabilities. This is the point of Proposition 11 below, which is a direct consequence of the balance condition.

▶ Proposition 11. Let $n \in \mathbb{N}_{>0}$ and $i, j \in [1, n]$. Consider $\gamma = \delta \Gamma$, where $\Gamma \in \mathcal{B}_{\mathcal{X}}$ and $\mathcal{X} \subseteq \mathbb{N}^n$ is a Ferrers set. For each $\diamond \in \{<, \leq, \geq, >\}$ and $x \in \mathcal{X}$ such that $x + e_i, x + e_j \in supp(\Gamma)$ and $x + e_i + e_j \in |\mathcal{S}|$, we have $\gamma_i(x) \diamond \gamma_i(x + e_j)$ if and only if $\gamma_j(x) \diamond \gamma_j(x + e_i)$. In particular, the class-i admission probability γ_i is increasing (resp. non-decreasing, non-increasing, decreasing) with respect to the number of class j customers if and only if the class-j admission probability γ_j is increasing (resp. non-decreasing, non-increasing, decreasing) with respect to the number of class-i customers.

Proof. Let $\diamond \in \{\leq, <, \geq, >\}$ and $x \in \mathcal{X}$ such that $x + e_i, x + e_j \in \operatorname{supp}(\Gamma)$ and $x + e_i + e_j \in |\mathcal{S}|$. The equivalence of $\gamma_i(x) \diamond \gamma_i(x + e_j)$ and $\gamma_j(x) \diamond \gamma_j(x + e_i)$ follows by rewriting (8) as

$$\frac{\gamma_i(x+e_j)}{\gamma_i(x)} = \frac{\gamma_j(x+e_i)}{\gamma_j(x)},$$

after recalling that $x + e_i$, $x + e_j \in \text{supp}(\Gamma)$ implies that $\gamma_i(x) > 0$ and $\gamma_j(x) > 0$. Alternately, this equivalence follows by observing that, according to (7), $\gamma_i(x) \diamond \gamma_i(x + e_j)$ and $\gamma_j(x) \diamond \gamma_j(x + e_i)$ are both equivalent to $\Gamma(x + e_i)\Gamma(x + e_j) \diamond \Gamma(x + e_i + e_j)\Gamma(x)$.

Alternative parameterization

To understand more concretely how the balance condition restricts the choice of the admission probabilities, let us consider an alternative way of specifying the admission probabilities. Consider a Ferrers set $\mathcal{X} \subseteq \mathbb{N}^n$, a balance function $\Gamma \in \mathcal{B}^{\mathrm{ac}}_{\mathcal{X}}$, and a Γ -balanced admission policy γ . By replacing \mathcal{X} with $\mathrm{supp}(\Gamma)$ if necessary, we can assume that Γ is positive. We can show that, under the balance condition (7)–(8), all admission probabilities under γ are specified once we set the values of the following admission probabilities: $\gamma_i(x_1, x_2, \ldots, x_i, 0, \ldots, 0)$, for each $i \in [1, n]$ and $(x_1, x_2, \ldots, x_{i-1}, x_i + 1, 0, \ldots, 0) \in \mathcal{X}$. Indeed, unfolding recursion (7) shows that, for each $x \in \mathcal{X}$, $\Gamma(x)$ can be computed by taking the product of admission

Figure 2 A complete and consistent definition of the balanced admission probabilities in a system n=2 classes and macrostate space $\mathcal{X}=[\![0,4]\!]\times[\![0,2]\!]$. The admission probabilities along the dashed arrows are fixed once we fix the admission probabilities along the solid edges.

probabilities along the increasing path going from the origin to x by going parallel first to the x_1 -axis, then to the x_2 -axis, and so on:

$$\Gamma(x) = \prod_{i=1}^{n} \prod_{y=0}^{x_i-1} \gamma_i(x_1, \dots, x_{i-1}, y, 0, \dots, 0), \text{ for each } x \in \mathcal{X}.$$

An example is shown in Figure 2 for a queueing system with n=2 classes and macrostate space $\mathcal{X}=\llbracket 0,4\rrbracket \times \llbracket 0,2\rrbracket$. We only need to specify $\gamma_1(x_1,0)$ for $x_1\in \llbracket 0,4\llbracket$ and $\gamma_2(x_1,x_2)$ for $x_1\in \llbracket 0,4\rrbracket$ and $x_2\in \llbracket 0,2\llbracket$. The value of $\gamma_1(x_1,x_2)$ for $x_1\in \llbracket 0,4\llbracket$ and $x_2\in \llbracket 0,2\rrbracket$ follows by writing

$$\gamma_1(x_1, x_2) = \frac{\Gamma(x_1 + 1, x_2)}{\Gamma(x_1, x_2)} = \gamma_1(x_1, 0) \prod_{i=0}^{x_2 - 1} \frac{\gamma_2(x_1 + 1, j)}{\gamma_2(x_1, j)}.$$

Intuitively, the class-1 admission probability in state (x_1, x_2) is "small" if the class-1 admission probability is already "small" in the absence of class 2 and/or if the class-2 admission probability is "significantly reduced" by adding a class-1 customer in a state (x_1, y) for $y \in [0, x_2]$.

Toy example

Let us illustrate this concept on a toy example. Consider a 2-class processor-sharing queue. Customers of each class arrive according to an independent Poisson process with rate 0.1. The queue can contain at most 5 customers of each class, so that an incoming customer is rejected if the queue already contains 5 customers of the corresponding class. The service requirement of each customer is exponentially distributed with mean 1, and the service capacity of the server is also 1. In this way, we consider a queueing system $Q_{\gamma} = (\mathcal{S}, \mathrm{Id}, q_{\gamma})$, where $\mathcal{S} = [0, 5]^2$,

$$\left. \begin{array}{l} q_{\gamma}(s,s+e_i) = 0.1 \gamma_i(s) \mathbb{1} \left[s_i < 5 \right], \\ q_{\gamma}(s,s-e_i) = \frac{s_i}{s_1 + s_2}, \end{array} \right\} \quad \text{for each } s \in \mathcal{S} \text{ and } i \in \llbracket 1,2 \rrbracket,$$

and other entries of q_{γ} are zero. The reward functions are given by $r_{\text{cont}}(s) = 0$ and $r_{\text{disc}}(s,t) = \mathbb{1}[s \in \mathcal{L}, t \in \mathcal{L}, \text{ and } |s| \leq |t|]$, for each $s,t \in \mathcal{S}$, where

$$\mathcal{L} = \Big\{ (0,0), (1,0), (2,0), (2,1), (3,1), (3,2), (3,3), (3,4), (4,4), (4,5), (5,5) \Big\}.$$




Figure 3 Comparison of the optimal policy and the best balanced policy in the toy example of Section 5.3.1. Blue edges represent admissions that are authorized (with probability 1) by the policy. The value inside each state gives the logarithm in base 10 of the value of the stationary distribution in this state under the depicted policy.

In other words, a reward is received each time an admission transition occurs along the path \mathcal{L} , and no reward is received otherwise. Note that this reward structure is chosen, not to be realistic, but to illustrate the impact of the dimension reduction in balanced policies. Recall that, by Sections 5.1 and 5.2, the optimal policy and the best balanced policy can both be chosen to be deterministic.

Figure 3 shows the optimal policy and the best balanced policy. States are depicted as circles on a grid. Thick blue edges represent admissions that are authorized by the corresponding policy, and the value inside each state shows the logarithm of the stationary probability of being in this state under the corresponding policy. In Figure 3a, we see that the optimal policy allows transitions along the path \mathcal{P} , plus a few additional transitions. On




Figure 4 Percentage loss of the best balanced policy with respect to the optimal policy, as a function of the per-class arrival rates in the toy example of Section 5.3.1.

the contrary, the best balanced policy shown in Figure 3b admits only arrivals of class 1, and only in states x = (0,0) and x = (0,1). With our choice of arrival rates, such a significant difference in policy definition has a small impact on the policy's gain, as large states have a small stationary probability anyway. More precisely, the suboptimality of the best balanced policy in this toy example is given by

$$\label{eq:loss} \text{Loss} = \frac{G^{\gamma^*} - G^{\delta \Gamma *}}{G^{\gamma^*} - G^{\gamma_*}} \simeq 0.82\%,$$

where γ^* is an optimal policy, γ_* a worst policy, and $\partial \Gamma^*$ the best balanced policy. In this particular example, γ_* is the reject-all policy and $G^{\gamma_*} = 0$.

Figure 4 shows the evolution of the loss of the best balanced policy with respect to the optimal policy, as a function of the arrival rates of the two classes. The loss of approximately 0.82% mentioned above appears on the lower left square, corresponding to both classes having an arrival rate of 0.1. In general, the structure of the optimal and best balanced policies remain similar to Figure 3, but the loss increases when larger states become more likely. The suboptimality of balanced policies is maximal when the arrival rate of class 1 is 0.5 and that of class 2 is 1. Detailed results (not shown here) reveal that the optimal policy tends to accept class-1 customers in more states than in Figure 3a, while the best balanced policy is the same as in Figure 3b.

5.3.2 No net circulation in state space

Classical works on reversibility also help shed light on the balance condition. According to [25], "a reversible Markov process shows no net circulation in state-space". This means that, in a reversible stationary CTMC, a cycle in the transition diagram is traversed equally frequently, whether we orient this cycle one way or the other, as formalized by Kolmogorov's criterion [19, Section 1.5]. We observed in Section 2.2.3 that the macrostate-process of a




Figure 5 Comparison of the optimal policy and the best balanced policy in the toy example of Section 5.3.2. Blue edges represent admissions that are authorized (with probability 1) by the policy. The value inside each state gives the logarithm in base 10 of the value of the stationary distribution in this state under the depicted policy. We see that state x = (5,0) is more likely under the optimal policy (log $\Pi^{\gamma^*}(5,0) = -5.01$) than under the best balanced policy (log $\Pi^{\delta\Gamma^*}(5,0) = -5.05$).




Figure 6 Percentage loss of the best balanced policy with respect to the optimal policy, as a function of the per-class arrival rates in the toy example of Section 5.3.2.

quasi-reversible queueing system has the same stationary distribution and transition rates as a reversible CTMC. Therefore, Theorem 1 implies that a quasi-reversible queueing system subject to a balanced policy also shows no net circulation in state space, when looked at the macrostate level.

Toy example

Let us illustrate this concept with the same toy example as in Section 5.3.1, namely, a processor-sharing queue with two customer classes, each arriving as a Poisson process with rate 0.1, and with state space $\mathcal{S} = [\![0,5]\!]^2$. Compared to Section 5.3.1, we change the reward functions, which are now given by $r_{\text{cont}}(s) = \mathbbm{1}[s_1 = 5, s_2 = 0]$ and $r_{\text{disc}}(s,t) = 0$, for each $s,t \in \mathcal{S}$. In other words, the goal is to maximize the probability of being in the state where the system is full of class-1 customers and empty of class-2 customers. Again, recall that this example has been built to provide insights on the impact of the balance condition, and not to be realistic.

The results are shown in Figure 5. Figure 5a shows the optimal policy, and Figure 5b shows the best balanced policy. States are depicted as circles on a grid. Thick edges represent admissions that are authorized by the corresponding policy, and the value inside each state shows the logarithm of the stationary probability of being in this state under the corresponding policy. The main observation is that, compared to $\delta\Gamma^*$, the optimal policy γ^* leads to a higher probability of being in state (5,0) by admitting class-2 customers in states with 0 or 1 class-1 customers. Intuitively, since the arrival rates are small compared to the service rates, it is advantageous to exploit circulation along cycles that visit states with a positive number of class-2 customers. To give an order of magnitude of the suboptimality of balanced policies in this toy example, we have

$$\operatorname{Loss} = \frac{G^{\gamma^*} - G^{\delta \Gamma^*}}{G^{\gamma^*} - G^{\gamma_*}} \simeq 7.96\%.$$

Figure 6 shows the evolution of the loss as a function of the arrival rates of both classes. In this example, the loss of the best balanced policy with respect to the optimal policy decreases with the arrival rate of class 1 and increases with that of class 2. Intuitively, as the arrival

rate of class 1 increases, the optimal policies shifts from the policy of Figure 5a (leading to a maximal loss) to the best balanced policy shown in Figure 5b (leading to a zero loss).

5.3.3 A more realistic scenario

Let us consider one last variant of the toy example of Section 5.3.1, corresponding to a more realistic application. We still have a processor-sharing queue with two customer classes, each arriving as a Poisson process with rate 0.1, and with state space $S = [0, 5]^2$. We consider the following cost structure: for each $s, t \in S$,

$$r_{\text{cont}}(s) = -s_1 - s_2,$$
 $r_{\text{disc}}(s, t) = 31 [|t| = |s| + e_1] + 61 [|t| = |s| + e_2].$

In this way, r_{cont} represents a holding cost of 1 per job per time unit, and r_{disc} represents an admission reward of 3 for class-1 jobs and 6 for class-2 jobs.

The optimal policy and the best balanced policy are shown in Figure 7. The optimal policy and the best balanced policy have approximately the same set of recurrent states, but the admission probabilities inside this set are different. The optimal balanced policy is constrained by the admission space, i.e., there is a single deterministic balanced policy with a particular set of recurrent states. On the contrary, without the balance condition, there are many deterministic policies with a particular set of recurrent states. In this example, having approximately the same set of recurrent states seems to be enough to yield the good performance, as we have

$$\operatorname{Loss} = \frac{G^{\gamma^*} - G^{\delta \Gamma^*}}{G^{\gamma^*} - G^{\gamma_*}} \simeq 0.26\%.$$

Figure 8 shows the evolution of the loss when the arrival rates of the two classes vary. The loss is (approximately) zero when the arrival rate of class 2 is 1. Looking closer at the case where the arrival rate of class 1 is 0.5, we see that both the optimal policy and the best




Figure 7 Comparison of the optimal policy and the best balanced policy in the toy example of Section 5.3.3. Blue edges represent admissions that are authorized (with probability 1) by the policy. The value inside each state gives the logarithm in base 10 of the value of the stationary distribution in this state under the depicted policy.




Figure 8 Percentage loss of the best balanced policy with respect to the optimal policy, as a function of the per-class arrival rates in the toy example of Section 5.3.3.

balanced policy always reject class 1 customers and only accept class 2. The admission thresholds of class 2 differ slightly: it is one value larger for the optimal policy than for the best balanced policy; we attribute this to rounding errors, as the difference in performance is of order 10^{-8} . The worst case on Figure 8 is when both classes have arrival rate 0.5. In this case, the discussion is similar to that we had above: the optimal policy and the best balanced policy have approximately the same set of recurrent states, but within this set they sometimes make different decisions.

6 Case study

To illustrate the results of this paper, we consider a multi-class multi-server redundancy system with customer abandonment. We will see that this queueing system can be described as an OI queue and is therefore quasi-reversible. After describing the queueing system and the admission-control problem, we propose a gradient-based method to search numerically for the best policy within a parametric family of balanced policies. This method is inspired by [12] and leverages the gradient result of Corollary 2. The numerical results we obtain in this case study illustrate two aspects of our work: (i) we compare numerically the performance of different families of balance functions, going from static to dynamic, and (ii) we illustrate how to use the gradient expression of Corollary 2 to design a stochastic gradient ascent algorithm that finds a (locally) optimal policy within each family.

6.1 Admission-control problem

Consider a queueing system with n customer classes and m servers, where $n, m \in \mathbb{N}_{>0}$. For each $i \in [\![1,n]\!]$, class-i customers enter the system according to a Poisson process with rate $\nu_i > 0$, and each class-i customer in the system abandons at rate $\theta_i > 0$. Furthermore, for each $j \in [\![1,m]\!]$, the completion time of a customer at server j is exponentially distributed with rate $\mu_j > 0$. We let $\nu = (\nu_1, \nu_2, \dots, \nu_n)$, $\theta = (\theta_1, \theta_2, \dots, \theta_n)$, and $\mu = (\mu_1, \mu_2, \dots, \mu_m)$. Compatibility between customers and servers is described by a bipartite graph with adjacency matrix $B = (b_{i,j})_{i \in [\![1,n]\!] \times j \in [\![1,m]\!]}$, such that $b_{i,j} = 1$ if class-i customers can be processed by server j, and $b_{i,j} = 0$ otherwise. An example compatibility graph is shown in Figure 9.




Figure 9 A compatibility graph and its adjacency matrix.

We assume that the scheduling discipline is redundancy cancel-on-complete, and that each server processes its compatible jobs in first-come-first-served order. In this way, at each time, each customer has a replica in service on each server that is compatible with this customer class but not with other customers that arrived earlier. We assume that the service times of the replicas of a job are independent of each other and of the service times of other jobs' replicas.

Under the appropriate independence assumptions, one can verify that this system is an OI queue, as described in Section 4.2, with rate function given by

$$\mu(s_1, s_2, \dots, s_n) = \sum_{\substack{j=1\\(|s|B)_i > 0}}^m \mu_j + \sum_{i=1}^n \theta_i |s|_i, \quad \text{for each } s_1 s_2 \dots s_\ell \in \mathcal{S},$$

where $S = [1, n]^*$ and, for each $j \in [1, m]$, $(|s|B)_j = \sum_{i=1}^n |s|_i b_{i,j}$ is the number of customers in the system that are compatible with server j. In particular, the stationary distribution of the CTMC defined by the system's microstate is given by (6). In the special case where $\theta = 0$ (i.e., no abandonment), we obtain the queueing system introduced in [15] and described as an OI queue in [6]. Under appropriate stability conditions, closed-form expressions for the normalizing constant as well as various performance metrics have been derived in [14, 13, 7]. In the remainder, we focus instead on the case where $\theta_i > 0$ for each $i \in [1, n]$. Stability is guaranteed by the abandonments, irrespective of the relative values of λ and μ , but deriving a closed-form expression for the normalizing constant is an open question that seems challenging.

Assume a reward $r_i > 0$ is received every time a class-i customer completes service, for each $i \in [\![1,n]\!]$. If a customer is rejected or abandons before service completion, no reward is received. Our goal is to maximize the long-run average reward. This problem can written as a special case of the admission-control problem (27) (albeit with an infinite state space) as follows. For each $s = s_1 s_2 \cdots s_\ell \in \mathcal{S}$ and $i \in [\![1,n]\!]$ such that $|s|_i > 0$, we define, with $p = \arg\min\{q \in [\![1,\ell]\!]; s_q = i\}$, and $t = s_1 s_2 \cdots s_{p-1} s_{p+1} \cdots s_\ell$,

$$\begin{split} r_{\mathrm{disc}}(s,t) &= r_i \gamma, \quad \text{where } \gamma = \frac{\sum_{j \in \mathcal{J}} \mu_j}{\sum_{j \in \mathcal{J}} \mu_j + \theta_i (d-p+1)}, \\ \mathcal{J} &= \{j \in [\![1,m]\!]; b_{i,j} = 1 \text{ and } b_{s_q,j} = 0 \text{ for each } q \in [\![1,p-1]\!]\}, \text{ and } \\ d &= \max \left\{q \in [\![p,\ell]\!]; c_r = i \text{ for each } r \in [\![p,q]\!]\right\}. \end{split}$$

Here, γ is the probability that the departure of the class-i customer in the transition from state s to state t is due to a service completion, \mathcal{J} is the set of servers that are processing a class-i customer in state s, and d is the latest position in the sequence of class-i customers that follow the oldest one. Other rewards are zero.

Finding an optimal policy for this problem is made complicated by the fact that the state space is infinite. More specifically, computing the optimal policy is unfeasible (e.g., using the linear programming approach mentioned in Section 5.1, or dynamic programming), because

the state space is infinite, and because we do not have a closed-form expression for long-run indicators such as a the normalizing constant of the stationary distribution and the state value function.

Also due to the fact that the state space is infinite, the results of Section 5.2 cannot be applied directly, though we conjecture that the best balanced policy is again deterministic. In the remainder of this section, we explain how to search for the best balanced policy, by adapting a PGRL algorithm introduced in [12]. While classical PGRL algorithms without function approximation again fall short in this problem, the PGRL algorithm we describe below exploits the form of the stationary distribution to provide an estimator of the gradient that works without function approximation, even when the state space is infinite.

6.2 Policy-gradient reinforcement learning

Let us cast the problem in the framework of policy-gradient reinforcement learning (PGRL). Though this paragraph was written to be self-contained, the interested reader can refer to [24, Chapter 13] for a brief introduction on this branch of reinforcement learning (RL). Since the only decision points are customer arrival times, which are distributed as a Poisson process with rate $\sum_{i=1}^{n} \nu_i$, we can reformulate the problem as a (discrete-time) MDP by focusing on the state observed at arrival times:

- State space $\mathcal{S} \times [\![1,n]\!]$: The MDP is in state $(s,i) \in \mathcal{S} \times [\![1,n]\!]$ if the state observed by the incoming customer is s and if this customer belongs to class i.
- Action space $A = \{0,1\}$: Action 0 means that the incoming customer is rejected and action 1, that the customer is admitted.
- Reward space $\mathcal{R} = \{\sum_{i=1}^n x_i r_i : x \in \mathbb{N}^n\}$: The reward is $\sum_{i=1}^n x_i r_i$ if, for each $j \in [1, n]$, x_j is the number of class-j customers that leave *due to service completion* between the (class-i) arrival for which we make a decision and the next arrival.

Now consider a balance function parameterization $\theta \in \mathbb{R}^d \mapsto \Gamma_{\theta} \in \mathcal{B}^{\mathrm{ac}}_{|\mathcal{S}|}$ as described in Section 3.4, where $d \in \mathbb{N}_{>0}$. In the numerical results, we will consider in particular the following three balanced function parameterizations:

- 1. Static: $\Gamma_{\theta}(x) = \prod_{i=1}^{n} \sigma(\theta_{i})^{x_{i}}$, where $\sigma : \mathbb{R} \to (0,1)$ is the logistic function, given by $\sigma(\phi) = 1/(1 + e^{-\phi})$ for each $\phi \in \mathbb{R}$, and $\theta \in \mathbb{R}^{n}$.
- **2.** Semi-static: $\Gamma_{\theta}(x) = \prod_{i=1}^{n} \sigma(\theta_i)^{x_i} \prod_{j=1}^{n} \sigma(\theta_{i,j})^{x_i x_j}$, where $\theta \in \mathbb{R}^{n+n^2}$.
- 3. Dynamic: We consider a simplified variant of the cum-prod policies described in Section 3.2, where large queue lengths are collapsed: $\Gamma_{\theta}(x) = \prod_{y \in |\mathcal{S}|: y \leq x} \psi_{\theta}(y)$ with $\psi_{\theta}(y) = \sigma(\theta_{\xi(y)})$, where $\theta \in \mathbb{R}^{15^n}$ and the feature function $\xi : \mathbb{N}^n \to [0, 15]^n$ is defined in Table 2. This parameterization can be used only for small values of n.

Intuitively, there is a trade-off between the dimension d of the parameter (equal to n, $n + n^2$, and 15^n , respectively, in the examples above) and the versatility of the family of policies. Under all three parameterizations, deterministic policies can be obtained asymptotically, when the components of the parameter θ tend to $\pm \infty$. This is common in the context of PGRL, see for instance [24, Section 13.1].

| Value range of y_i | $[\![0,9]\!]$ | [10, 19] | $[\![20,29]\!]$ | [30, 39] | [40, 49] | $\llbracket 50, +\infty \rrbracket$ |
|-----------------------|---------------|----------|-----------------|----------|----------|-------------------------------------|
| Value of $\xi_i(y_i)$ | y_i | 10 | 11 | 12 | 13 | 14 |

Table 2 Definition of the feature function ξ used for dynamic policies.

According to the PASTA property [26], the stationary distribution of the discrete-time Markov chain (DTMC) of the system state observed at arrival times of customers is again

given by Π_{θ} . Therefore, the gain function can be rewritten as

$$G_{\theta} = \sum_{s,i,a,r,s'} \Pi_{\theta}(s) \frac{\nu_i}{\sum_j \nu_j} \pi_{\theta}(s,i,a) p(r,s'|s,i,a) r, \quad \text{for each } \theta \in \mathbb{R}^d,$$
 (29)

where the sum is over $s \in \mathcal{S}$, $i \in [1, n]$, $a \in \mathcal{A}$, $r \in \mathcal{R}$, and $s' \in \mathcal{S}$. Here, we let π_{θ} denote the policy in the sense of PGRL, that is,

$$\pi_{\theta}(s, i, a) = (\partial \Gamma_{\theta})_{i}(s) \mathbb{1} [a = 1] + (1 - (\partial \Gamma_{\theta})_{i}(s)) \mathbb{1} [a = 0]$$

is the probability of taking action a in state (s, i), and we let p denote the transition kernel of the MDP. We do not explicitate p since its form will not play a role in the remainder (we will only need to sample from this kernel, which is done by simulating the process).

6.3 Gradient estimator

Roughly speaking, the idea of PGRL is to run a (stochastic) gradient ascent algorithm with updates of the form

$$\theta \leftarrow \theta + \widehat{\nabla_{\theta} G_{\theta}},$$

30

where $\widehat{\nabla_{\theta}G_{\theta}}$ is a stochastic estimator for $\nabla_{\theta}G_{\theta}$, as $\nabla_{\theta}G_{\theta}$ cannot be computed explicitly in general. Proposition 12 below builds on Corollary 2 to derive an expression for $\nabla_{\theta}G_{\theta}$, which we will use to construct an estimator for $\nabla_{\theta}G_{\theta}$. This result can be seen as an extension of [12, Equation (13) in Theorem 1]. For each $\theta \in \mathbb{R}^d$, we let $(S_{\theta}, I_{\theta}, A_{\theta}, R_{\theta})$ denote a stationary 4-tuple under policy π_{θ} , that is, having the following distribution: for each $s \in \mathcal{S}$, $i \in [\![1, n]\!]$, $a \in \mathcal{A}$, and $r \in \mathcal{R}$,

$$\mathbb{P}_{\theta}(s, i, a, r) = \mathbb{P}(S_{\theta} = s, I_{\theta} = i, A_{\theta} = a, R_{\theta} = r)$$

$$= \Pi_{\theta}(s) \frac{\nu_{i}}{\sum_{j} \nu_{j}} \pi_{\theta}(s, i, a) \sum_{s' \in \mathcal{S}} p(r, s'|s, i, a). \tag{30}$$

For each $k \in [1, d]$, we let ∂_k denote the partial derivative with respect to θ_k .

▶ Proposition 12. For each $\theta \in \mathbb{R}^d$, we have

$$\partial_k G_{\theta} = Cov(R_{\theta}, \partial_k \log \Gamma_{\theta}(S_{\theta})) + \mathbb{E}\left(R \,\partial_k \log \pi_{\theta}(S_{\theta}, I_{\theta}, A_{\theta})\right). \tag{31}$$

Proof. Let $k \in [1, d]$ and $\theta \in \mathbb{R}^d$. Observe that we have for all s, i, a, d

$$\partial_k (\Pi_{\theta}(s)\pi_{\theta}(s,i,a)) = \Pi_{\theta}(s)\pi_{\theta}(s,i,a)\partial_k \log (\Pi_{\theta}(s)\pi_{\theta}(s,i,a)),$$

= $\Pi_{\theta}(s)\pi_{\theta}(s,i,a) (\partial_k \log \Pi_{\theta}(s) + \partial_k \log \pi_{\theta}(s,i,a))$

Thus, taking the partial derivative of (29) yields

$$\partial_k G_{\theta} = \sum_{s,i,a,r,s'} \Pi_{\theta}(s) \frac{\nu_i}{\sum_j \nu_j} \pi_{\theta}(s,i,a) p(r,s'|s,i,a) r \left[\partial_k \log \Pi_{\theta}(s) + \partial_k \log \pi_{\theta}(s,i,a) \right].$$

Therefore, by injecting Corollary 2 and recalling (30), we obtain

$$\partial_k G_{\theta} = \sum_{s,i,a,r,s'} r \left[\partial_k \log \Gamma_{\theta}(s) - \mathbb{E} \left(\partial_k \log \Gamma_{\theta}(S_{\theta}) \right) + \partial_k \log \pi_{\theta}(s,i,a) \right] \mathbb{P}_{\theta}(s,i,a,r),$$

which is equivalent to (31).


We use Proposition 12 to construct an estimator for $\nabla_{\theta}G_{\theta}$ and train a PGRL algorithm in a similar way as in [12]. See in particular [12, Sections 3.4 and 4.3] for a description of the algorithm and [12, Section 6.1] for a toy admission-control example. The algorithm's hyperparameters are also the same as in [12].

6.4 Numerical results

We show numerical results for the compatibility graph of Figure 9, with arrival rate vector $\nu = (0.5, 0.5, 0.5)$, abandonment rate vector $\theta = (0.1, 0.2, 0.5)$, service rate vector $\mu = (0.5, 0.5, 0.5)$, and reward completion rate vector r = (1, 2, 16). We ran the PGRL algorithm described in Sections 6.2 and 6.3 for $2 \cdot 10^6$ steps, with the three policy parameterizations described in Section 6.2.

Figure 10a shows the running average reward, and Figures 10b–10d show the running average of the admission probabilities of the classes. All three policy parameterizations are shown on the same plot to facilitate comparison. Under all policy parameterizations, the parameter vector is initialized so that the initial admission probability is approximately $\frac{1}{2}$ in all states.

A first glance at Figure 10 shows the following trend. During the first 10^4 time steps, the running averages are relatively noisy, and the admission probabilities do not seem to increase or decrease significantly. Beyond 10^4 steps, the running averages are less noisy, and



rigule 10 Empirical average reward

the admission probabilities move away from $\frac{1}{2}$. This suggests that, under the initial policy, 10^4 is the order of magnitude of the number of takes necessary for the system to mix and the ergodic theorem to apply approximately.

Figure 10a shows that the running average reward initially increases faster under the static and semi-static policies, but that the dynamic policy eventually catches up and slightly outperforms the static and semi-static policies. More precisely, as we can see on Figure 10a, while the static and semi-static policies stabilize below 2.55 during the second half of the simulation (between 10^6 and $2 \cdot 10^6$ steps), that under the dynamic policy keeps increasing between 2.55 and 2.57 during that period. Our intuition is that the dynamic parameterization eventually yields larger reward because the family of dynamic policies is larger than the families of static and semi-static policies, with the side-effect is that convergence is also

slower.

Now focusing on the admission probabilities in Figures 10b–10d, our main observation is that the difference in performance between the static and semi-static policies on the one hand and the dynamic policy on the other hand seems to be related to the admission probability of class 2. While this probability is almost monotonically increasing under the static and semi-static policies, it increases and then *decreases* again under the dynamic policy. We conjecture that this is again due to the fact that the family of dynamic policies is larger than the families of static and semi-static policies.

7 Conclusion

In this paper, we have introduced a general definition of *quasi-reversibility* for queueing systems, that slightly differs from existing definitions in the literature. We have shown that the class of quasi-reversible systems embraces a wide range of systems of particular interest in the recent queueing literature: multi-class Whittle networks and Order-Independent queues, which encompass in turn, among others, redundancy systems and stochastic matching models.

For this class of systems, we have introduced the notion of balanced arrival control policies, which generalizes the existing notion of balanced arrival rates for Whittle networks, to any quasi-reversible queueing system. We have shown that supplementing a quasi-reversible queueing system with a balanced arrival control policy, preserves the quasi-reversibility. This has allowed us, first, to obtain the stationary measures of the controlled system in an explicit form, and second, to set the problem of optimization of the admission control to the queueing system, as a linear programming problem. To illustrate the cost of balance, we have conducted a comparison between the best balanced admission control and the optimal (unconstrained) control, on various simple examples.

Last, we have shown the crucial interest of balanced admission control in quasi-reversible systems, to deploy a policy-gradient reinforcement learning (PGRL) algorithm in order to optimize the admission control. We have conducted a first, simple, case study, by implementing a PGRL algorithm to a redundancy model with compatibility constraints, and analyzed a set of numerical results on this model.

This paper thus provides the theoretical basis for the optimization of quasi-reversible systems with RL algorithms, and consists of a first step in that direction. It naturally opens the way for future work. First, for a better understanding of the optimality gap of balanced policies (which are crucial in the present framework) and optimal, unconstrained, policies. This raises the following complementary open question: is there a broader class of policies (beyond balanced policies) that also preserve quasi-reversibility, and is still convenient to parameterize?

Third, the last part of this work suggests that PGRL algorithms can be fruitfully applied to a wide range of multi-class Whittle networks and OI queues, among which, open (multi-class) Jackson networks, redundancy models and matching queues. The optimization of admission control for the last two classes of models is the topic of our current investigations on this class of problems.

Last, many *closed* multi-class systems also enjoy the quasi-reversibility property. A balanced control would then be interpreted as an *internal* load balancing rather than an admission control. However, it appears that the procedure introduced in this work is also amenable to optimize this internal control. This is left for future work.

References

- 1 Andrew D. Barbour. Networks of queues and the method of stages. *Advances in Applied Probability*, 8(3):584–591. doi:10.2307/1426145.
- 2 Jocelyn Begeot, Irène Marcovici, Pascal Moyal, and Youssef Rahme. A general stochastic matching model on multigraphs. (18):1325-1351, 2021.
- 3 S. A. Berezner, C. F. Kriel, and A. E. Krzesinski. Quasi-reversible multiclass queues with order independent departure rates. *Queueing Systems*, 19(4):345–359. doi:10.1007/BF01151928.
- 4 S. A. Berezner and A. E. Krzesinski. Order independent loss queues. *Queueing Systems*, 23(1):331–335. doi:10.1007/BF01206565.
- T. Bonald, M. Jonckheere, and A. Proutière. Insensitive load balancing. In Proceedings of the Joint International Conference on Measurement and Modeling of Computer Systems, SIGMETRICS '04/Performance '04, pages 367–377. ACM. doi:10.1145/1005686.1005729.
- 6 Thomas Bonald and Céline Comte. Balanced fair resource sharing in computer clusters. Performance Evaluation, 116:70-83. doi:10.1016/j.peva.2017.08.006.
- 7 Thomas Bonald, Céline Comte, and Fabien Mathieu. Performance of balanced fairness in resource pools: A recursive approach. *Proceedings of the ACM on Measurement and Analysis of Computing Systems*, 1(2):41:1–41:25. doi:10.1145/3154500.
- 8 Xiuli Chao. Networks with customers, signals, and product form solution. In Richard J. Boucherie and Nico M. van Dijk, editors, *Queueing Networks: A Fundamental Approach*, International Series in Operations Research & Management Science, pages 217–267. Springer US. doi:10.1007/978-1-4419-6472-4_5.
- 9 Céline Comte. Dynamic load balancing with tokens. In 17th International IFIP TC6 Networking Conference Networking 2018, pages 343-351. IFIP Open Digital Library. doi:10.23919/IFIPNetworking.2018.8697018.
- 10 Céline Comte. Resource management in computer clusters: Algorithm design and performance analysis. URL: https://pastel.archives-ouvertes.fr/tel-02413496.
- Céline Comte. Stochastic non-bipartite matching models and order-independent loss queues. 38(1):1–36. Publisher: Taylor & Francis. doi:10.1080/15326349.2021.1962352.
- Céline Comte, Matthieu Jonckheere, Jaron Sanders, and Albert Senen-Cerda. Score-aware policy-gradient methods and performance guarantees using local lyapunov conditions: Applications to product-form stochastic networks and queueing systems. arXiv:2312.02804[cs,math], doi:10.48550/arXiv.2312.02804.
- Kristen Gardner, Mor Harchol-Balter, Esa Hyytiä, and Rhonda Righter. Scheduling for efficiency and fairness in systems with redundancy. 116:1-25. URL: http://www.sciencedirect.com/science/article/pii/S0166531617300457, doi:10.1016/j.peva.2017.07.001.
- 14 Kristen Gardner, Mor Harchol-Balter, Alan Scheller-Wolf, Mark Velednitsky, and Samuel Zbarsky. Redundancy-d: The power of d choices for redundancy. 65(4):1078–1094. doi: 10.1287/opre.2016.1582.
- Kristen Gardner, Samuel Zbarsky, Sherwin Doroudi, Mor Harchol-Balter, Esa Hyytiä, and Alan Scheller-Wolf. Queueing with redundant requests: exact analysis. *Queueing Systems*, 83(3):227–259. doi:10.1007/s11134-016-9485-y.
- Matthieu Jonckheere and Jean Mairesse. Towards an Erlang formula for multiclass networks. Queueing Systems, 66(1):53–78. doi:10.1007/s11134-010-9185-y.
- 17 F. P. Kelly. Networks of queues. Advances in Applied Probability, 8(2):416-432. doi: 10.2307/1425912.
- 18 F. P. Kelly. Reversibility and Stochastic Networks. Cambridge University Press.
- A. E. Krzesinski. Order independent queues. In Richard J. Boucherie and Nico M. van Dijk, editors, *Queueing networks: A fundamental approach*, number 154 in International Series in Operations Research & Management Science, pages 85–120. Springer US. doi: 10.1007/978-1-4419-6472-4_2.
- 20 Jean Mairesse and Pascal Moyal. Stability of the stochastic matching model. 53(4):1064–1077. Publisher: Cambridge University Press. doi:10.1017/jpr.2016.65.

- Pascal Moyal, Ana Bušić, and Jean Mairesse. A product form for the general stochastic matching model. 58(2):449–468. Publisher: Cambridge University Press. doi:10.1017/jpr. 2020.100.
- 22 Martin L. Puterman. Markov Decision Processes: Discrete Stochastic Dynamic Programming. John Wiley & Sons, Ltd.
- 23 Richard Serfozo. Introduction to Stochastic Networks. Springer. URL: http://link.springer.com/10.1007/978-1-4612-1482-3.
- 24 Richard S. Sutton and Andrew G. Barto. *Reinforcement Learning: An Introduction*. Adaptive Computation and Machine Learning series. A Bradford Book, 2 edition.
- P. Whittle. Reversibility and acyclicity. Journal of Applied Probability, 12:217–224. Publisher: Cambridge University Press. doi:10.1017/S0021900200047677.
- Ronald W. Wolff. Poisson arrivals see time averages. *Operations Research*, 30(2):223-231. Publisher: INFORMS. doi:10.1287/opre.30.2.223.

A Linear programs for reversible queueing systems

Consider a quasi-reversible queueing system $Q = (S, |\cdot|, q)$ with $|\cdot| = \text{Id}$, so that $S = \mathcal{X}$, $S_x = \{x\}$ for each $x \in \mathcal{X}$, and q is reversible (see Section 2.2.3). In this case, all policies trivially satisfy the microstate-independence condition of Definition 2, and the only difference between an arbitrary admission policy and balanced policies is the balance condition. Problem (A1) below is a rewriting of (27), accounting for the fact that $S = \mathcal{X}$:

$$\underset{\gamma:\mathcal{X}\to[0,1]^n,\\\Pi:\mathcal{X}\to[0,1]}{\text{Maximize}} \quad G_{\gamma} \triangleq \sum_{x\in\mathcal{X}} \Pi(x) r_{\text{cont}}(x) + \sum_{x,y\in\mathcal{S}} \Pi(x) q_{\gamma}(x,y) r_{\text{disc}}(x,y), \tag{A1a}$$

Subject to
$$\Pi(x) \sum_{y \in \mathcal{S}} q_{\gamma}(x, y) = \sum_{y \in \mathcal{S}} \Pi(y) q_{\gamma}(y, x)$$
, for each $x \in \mathcal{X}$, (A1b)

$$\sum_{x \in \mathcal{S}} \Pi(x) = 1. \tag{A1c}$$

This, in turn, is equivalent to the following linear program (LP):

$$\underset{\eta: \mathcal{X} \to [0,1]^n, \\ \Pi: \mathcal{X} \to [0,1]}{\text{Maximize}} \quad G_{\gamma} \triangleq \sum_{x \in \mathcal{X}} \Pi(x) r_{\text{cont}}(x) + \sum_{x,y \in \mathcal{X}} \eta(x,y) r_{\text{disc}}(x,y), \tag{A2a}$$

Subject to
$$\sum_{y \in \mathcal{X}} \eta(x, y) = \sum_{y \in \mathcal{X}} \eta(y, x)$$
, for each $x \in \mathcal{X}$, (A2b)

$$\sum_{x \in \mathcal{X}} \Pi(x) = 1,\tag{A2c}$$

$$\eta(x, x + e_i) \le \Pi(x)q(x, x + e_i), \text{ for each } x \in \mathcal{X} \text{ and } i \in [1, n],$$
 (A2d)

$$\eta(x,y) = \Pi(x)q(x,y), \text{ for each } x \in \mathcal{X} \text{ and } y \in \mathcal{X} \setminus \{x + e_i; i \in [1, n]\}.$$
(A2e)

Equivalence between Problems (A1) and (A2) is established by injecting the definition (6) of q_{γ} , and letting

$$\eta(x,y) = \begin{cases} \Pi(x)q(x,x+e_i)\gamma_i(x) & \text{if } y = x+e_i \text{ for some } i \in [1,n], \\ \Pi(x)q(x,y) & \text{otherwise.} \end{cases}$$

Problem (A2) can be further simplified by recalling that q(x,y) = 0 for each $x, y \in \mathcal{X}$ such that $y \notin \{x - e_i; i \in [1, n]\} \cup \{x\} \cup \{x + e_i; i \in [1, n]\}$. Lastly, one can verify that in this

special case, restricting the problem to balanced policies boils down to replacing the balance equations (A2b) with local balance equations, so as to obtain:

$$\underset{\substack{\eta: \mathcal{X} \to [0,1]^n, \\ \Pi: \mathcal{X} \to [0,1]}}{\text{Maximize}} \quad G_{\gamma} \triangleq \sum_{x \in \mathcal{X}} \Pi(x) r_{\text{cont}}(x) + \sum_{x,y \in \mathcal{X}} \eta(x,y) r_{\text{disc}}(x,y), \tag{A3a}$$

Subject to
$$\eta(x,y) = \eta(y,x)$$
, for each $x,y \in \mathcal{X}$, (A3b)

$$\sum_{x \in \mathcal{X}} \Pi(x) = 1,\tag{A3c}$$

$$\eta(x, x + e_i) \le \Pi(x)q(x, x + e_i), \text{ for each } x \in \mathcal{X} \text{ and } i \in [1, n],$$
 (A3d)

$$\eta(x,y) = \Pi(x)q(x,y), \text{ for each } x \in \mathcal{X} \text{ and } y \in \mathcal{X} \setminus \{x + e_i; i \in [1, n]\}.$$
(A3e)

B Proofs of Theorem 8, Proposition 9, and Proposition 10

We prove the results in the following order. First, we prove Proposition 9, which gives the vertices of the set $\mathcal{B}_{\mathcal{X}}^{\mathrm{ac}}$. Then, we build on this proof to show Proposition 10, which characterizes the vertices of the set $\tilde{\mathcal{B}}_{\mathcal{X}}^{\mathrm{ac}}$. Lastly, we use Proposition 10 to prove our final result, Theorem 8.

Proof of Proposition 9. Consider a finite Ferrers set $\mathcal{X} \subseteq \mathbb{N}^n$ and let $m = \operatorname{card}(\mathcal{X})$. We proceed in three steps.

 $\mathcal{B}_{\mathcal{X}}$ is a convex polytope. It follows from the fact that $\mathcal{B}_{\mathcal{X}}$ can be seen as a set of vectors (with coordinates in \mathcal{X} and values in $\mathbb{R}_{\geq 0}$) defined by linear constraints, both equality ($\Gamma(0) = 1$) and inequality (Γ is non-negative and component-wise non-increasing).

Deterministic function are extreme points of $\mathcal{B}_{\mathcal{X}}$. Let $\mathcal{A} \in \mathcal{F}_{\mathcal{X}}$. Assume for the sake of contradiction that there exist $\alpha \in (0,1)$ and two distinct functions $\Gamma, \Delta \in \mathcal{B}_{\mathcal{X}}$ such that $\mathbb{1}_{\mathcal{A}} = \alpha\Gamma + (1-\alpha)\Delta$. Since $\Gamma \neq \Delta$, there exists $x \in \mathcal{X}$ such that $\Gamma(x) \neq \Delta(x)$. By exchanging symbols if necessary, we can assume without loss of generality that $\Gamma(x) > \Delta(x)$. We now distinguish two cases:

- If $x \notin \mathcal{A}$, then $0 = \mathbb{1}_{\mathcal{A}}(x) = \alpha\Gamma(x) + (1 \alpha)\Delta(x) > \Delta(x)$. It follows that $\Delta(x) < 0$, which contradicts our assumption that $\Delta \in \mathcal{B}_{\mathcal{X}}$.
- If $x \in \mathcal{A}$, then $1 = \mathbb{1}_{\mathcal{A}}(x) = \alpha\Gamma(x) + (1 \alpha)\Delta(x) < \Gamma(x)$. As Γ is non-increasing, it follows that $\Gamma(0) > 1$, which contradicts our assumption that $\Gamma \in \mathcal{B}_{\mathcal{X}}$.

Both cases lead to a contradiction, hence $\mathbb{1}_{\mathcal{A}}$ is an extreme point of $\mathcal{B}_{\mathcal{X}}$.

Every element of $\mathcal{B}_{\mathcal{X}}$ is a convex combination of deterministic balance functions. Let $\Gamma \in \mathcal{B}_{\mathcal{X}}$. We first describe our approach to write Γ as a convex combination of deterministic balance functions, and then we prove the validity of this approach (e.g., that the coefficients of the combination are indeed non-negative and sum up to one).

Starting with $V_0 = \mathcal{X}$, let us define recursively, for each $k \in \{1, 2, \dots, m\}$,

$$y_{k-1} \in \underset{x \in \partial \mathcal{V}_{k-1}}{\arg \min} \Gamma(x),$$
 (B4)

$$\mathcal{V}_k = \mathcal{V}_{k-1} \setminus \{y_{k-1}\},\tag{B5}$$

where $\partial \mathcal{V} = \{x \in \mathcal{V}; x \not\leq y \text{ for each } y \in \mathcal{V}\}$ is the set of maximal (or non-dominated) points of \mathcal{V} , for each $\mathcal{V} \in \mathcal{F}_{\mathcal{X}}$. By construction, $y_{m-1} = 0$ and $\mathcal{V}_m = \emptyset$, as we remove maximal elements one by one starting from $\mathcal{V}_0 = \mathcal{X}$. Furthermore, $\mathcal{V}_0, \mathcal{V}_1, \ldots, \mathcal{V}_{m-1}$ are Ferrers

subsets of \mathcal{X} , hence $\mathbb{1}_{\mathcal{V}_k} \in \mathcal{B}_{\mathcal{X}}$ for each $k \in \{0, 1, \dots, m-1\}$. We will verify that we can write Γ as follows:

$$\Gamma(x) = \sum_{k=0}^{m-1} \alpha_k \mathbb{1}_{\mathcal{V}_k}(x) \quad \text{for each } x \in \mathcal{X},$$
(B6)

where $\alpha_0 \triangleq \Gamma(y_0)$, and $\alpha_k \triangleq \Gamma(y_k) - \Gamma(y_{k-1})$ for each $k \in \{1, 2, ..., m-1\}$. We will also verify that $\alpha_k \geq 0$ for each $k \in \{0, 1, ..., m-1\}$ and that $\sum_{k=0}^{m-1} \alpha_k = 1$, which will conclude the proof. Let us prove the required points:

■ Proof of (B6). Let us first verify (B6) for each $x \in \mathcal{X}$. Since the sequence $y_0, y_1, \ldots, y_{m-1}$ visits every element of \mathcal{X} , there exists (a unique) $\ell \in \{0, 1, \ldots, m-1\}$ such that $x = y_{\ell}$. By (B5), if follows that $x \in \mathcal{V}_k$ for each $k \in \{\ell, 1, \ldots, \ell\}$ and $x \notin \mathcal{V}_k$ for each $k \in \{\ell+1, \ell+2, \ldots, m-1\}$. The conclusion follows by simplifying the right-hand side of (B6):

$$\sum_{k=0}^{m-1} \alpha_k \mathbb{1}_{\mathcal{V}_k}(x) = \sum_{k=0}^{\ell} \alpha_k \mathbb{1}_{\mathcal{V}_k}(y_\ell) = \Gamma(y_0) + \sum_{k=0}^{\ell} (\Gamma(y_k) - \Gamma(y_{k-1})) = \Gamma(y_\ell) = \Gamma(x).$$

- Proof that $\alpha_k \geq 0$ for each $k \in [0, m-1]$. We have directly $\alpha_0 = \Gamma(y_0) \geq 0$ since Γ takes values in $\mathbb{R}_{\geq 0}$. Now, given $k \in [1, m-1]$, we want to prove that $\alpha_k \geq 0$, i.e., that $\Gamma(y_k) \geq \Gamma(y_{k-1})$. We know that $y_k \in \mathcal{V}_{k-1}$ (because $y_k \in \mathcal{V}_k = \mathcal{V}_{k-1} \setminus \{y_{k-1}\}$), hence it suffices to prove that $\Gamma(x) \geq \Gamma(y_{k-1})$ for each $x \in \mathcal{V}_{k-1}$. For each $x \in \mathcal{V}_{k-1}$, we can verify that $\Gamma(x) \geq \Gamma(y) \geq \Gamma(y_{k-1})$, where $y \in \delta \mathcal{V}_{k-1}$ is such that $x \leq y$; the first inequality follows because Γ is component-wise non-increasing, and the second follows from the definition of y_{k-1} .
- definition of y_{k-1} .

 Proof that $\sum_{k=0}^{m-1} \alpha_k = 1$. This follows by observing that this is a telescoping sum and recalling that $\Gamma(y_{m-1}) = \Gamma(0) = 1$.

Proof of Proposition 10. Consider a finite Ferrers set $\mathcal{X} \subseteq \mathbb{N}^n$ and let $m = \operatorname{card}(\mathcal{X})$. Building on the proof of Proposition 9, we again proceed in three steps.

 $\tilde{\mathcal{B}}_{\mathcal{X}}$ is a convex polytope. Like for $\mathcal{B}_{\mathcal{X}}$, it follows from the fact that $\tilde{\mathcal{B}}_{\mathcal{X}}$ can be seen as a set of vectors (with coordinates in \mathcal{X} and values in $\mathbb{R}_{\geq 0}$) defined by linear constraints, both equality $(\sum_{x \in \mathcal{X}} \Pi(x)\Gamma(x) = 1)$ and inequality (Γ is non-negative and component-wise non-increasing).

Deterministic function are extreme points of $\tilde{\mathcal{B}}_{\mathcal{X}}$. Let $\mathcal{A} \in \mathcal{F}_{\mathcal{X}}$ and assume for the sake of contradiction that there exist $\alpha \in (0,1)$ and two distinct functions $\Gamma, \Delta \in \tilde{\mathcal{B}}_{\mathcal{X}}$ such that $\Gamma_{\mathcal{A}} = \alpha\Gamma + (1-\alpha)\Delta$. Let $\mu = 1/(\sum_{y \in \mathcal{A}} \Pi(y))$.

Let us first verify that $\Gamma(x) = \tilde{\Delta}(x) = 0$ for each $x \in \mathcal{X} \setminus \mathcal{A}$. Assume for the sake of contradiction that there exists $x \in \mathcal{X} \setminus \mathcal{A}$ such that $\Gamma(x) \neq \Delta(x)$. By exchanging symbols if necessary, we can assume without loss of generality that $\Gamma(x) > \Delta(x)$. It follows that $0 = \Gamma_{\mathcal{A}}(x) = \alpha \Gamma(x) + (1 - \alpha)\Delta(x) > \Delta(x)$, which contradicts our assumption that $\Delta \in \tilde{\mathcal{B}}_{\mathcal{X}}$. Therefore, for each $x \in \mathcal{X} \setminus \mathcal{A}$, we have $\Gamma(x) = \Delta(x)$ which, combined with the fact that $0 = \Gamma_{\mathcal{A}}(x) = \alpha \Gamma(x) + (1 - \alpha)\Delta(x)$, implies that $\Gamma(x) = \Delta(x) = 0$.

Since $\Gamma \neq \Delta$, there exists $x \in \mathcal{X}$ such that $\Gamma(x) \neq \Delta(x)$. The previous paragraph implies that $x \in \mathcal{A}$. By exchanging symbols if necessary, we can assume without loss of generality that $\Gamma(x) > \Delta(x)$ which, combined with $\mu = \Gamma_{\mathcal{A}}(x) = \alpha\Gamma(x) + (1 - \alpha)\Delta(x)$, implies $\Gamma(x) > \mu > \Delta(x)$. Since Γ is non-increasing, it follows that $\Gamma(0) > \mu$. Since we also have $\mu = \Gamma_{\mathcal{A}}(0) = \alpha\Gamma(0) + (1 - \alpha)\Delta(0)$, we obtain $\Delta(0) < \mu$. Since Δ is also non-increasing, we obtain $\Delta(x) < \mu$ for each $x \in \mathcal{A}$. All in all, we obtain

$$\sum_{x \in \mathcal{X}} \Pi(x) \Delta(x) \stackrel{\text{(a)}}{=} \sum_{x \in \mathcal{A}} \Pi(x) \Delta(x) \stackrel{\text{(b)}}{<} \sum_{x \in \mathcal{A}} \Pi(x) \mu \stackrel{\text{(c)}}{=} 1,$$

where (a) follows from the fact that $\Delta(x) = 0$ for each $x \in \mathcal{X} \setminus \mathcal{A}$, (b) from the fact that $\Delta(x) < \mu$ for each $x \in \mathcal{A}$, and (c) from the definition of μ . This contradicts our assumption that $\Delta \in \tilde{\mathcal{B}}_{\mathcal{X}}$.

Every element of $\tilde{\mathcal{B}}_{\mathcal{X}}$ is a convex combination of deterministic balance functions. Let $\Gamma \in \tilde{\mathcal{B}}_{\mathcal{X}}$. Start off from the decomposition derived in the proof of Proposition 9, we have

$$\Gamma(x) = \sum_{k=0}^{m-1} \alpha_k \mathbb{1}[x \in \mathcal{V}_k], \text{ for each } x \in \mathcal{X},$$

where $y_0, y_1, \ldots, y_{m-1}$ and $\mathcal{V}_0, \mathcal{V}_1, \ldots, \mathcal{V}_{m-1}$ are built according to (B4)–(B5), $\alpha_0 = \Gamma(y_0)$, and $\alpha_k = \Gamma(y_k) - \Gamma(y_{k-1})$ for each $k \in [1, m-1]$. We also showed in the proof of Proposition 9 that $\alpha_k \geq 0$ for each $k \in [0, m-1]$. We can rewrite this equality as

$$\Gamma(x) = \sum_{k=0}^{m-1} \tilde{\alpha}_k \Gamma_{\mathcal{V}_k}(x), \quad \text{for each } x \in \mathcal{X},$$

where $\tilde{\alpha}_k = \alpha_k \sum_{y \in \mathcal{V}_k} \Pi(y)$ for each $k \in [0, m-1]$. We have $\tilde{\alpha}_k \geq 0$ for each $k \in [0, m-1]$. All that remains is to check that $\sum_{k=0}^{m-1} \tilde{\alpha}_k = 1$:

$$\begin{split} \sum_{k=0}^{m-1} \tilde{\alpha}_k &\stackrel{\text{(a)}}{=} \sum_{k=0}^{m-1} \alpha_k \sum_{y \in \mathcal{V}_k} \Pi(y), \\ &\stackrel{\text{(b)}}{=} \Gamma(y_0) \sum_{y \in \mathcal{V}_0} \Pi(y) + \sum_{k=1}^{m-1} (\Gamma(y_k) - \Gamma(y_{k-1})) \sum_{y \in \mathcal{V}_k} \Pi(y), \\ &\stackrel{\text{(c)}}{=} \Gamma(y_0) \sum_{y \in \mathcal{V}_0} \Pi(y) + \sum_{k=1}^{m-1} \Gamma(y_k) \sum_{y \in \mathcal{V}_k} \Pi(y) - \sum_{k=0}^{m-2} \Gamma(y_k) \sum_{y \in \mathcal{V}_{k+1}} \Pi(y), \\ &\stackrel{\text{(d)}}{=} \sum_{k=0}^{m-2} \Gamma(y_k) \left(\sum_{y \in \mathcal{V}_k} \Pi(y) - \sum_{y \in \mathcal{V}_{k+1}} \Pi(y) \right) + \Gamma(y_{m-1}) \sum_{y \in \mathcal{V}_{m-1}} \Pi(y), \\ &\stackrel{\text{(e)}}{=} \sum_{k=0}^{m-2} \Gamma(y_k) \Pi(y_k) + \Gamma(0) \Pi(0) \stackrel{\text{(f)}}{=} 1, \end{split}$$

where (a) follows from the definition of $\tilde{\alpha}_k$ for $k \in [0, m-1]$, (b) follows from the definition of α_k for $k \in [0, m-1]$, (c) follows by applying the distributivity property and then making a change of variable in the second sum, (d) follows by rearranging the terms, (e) follows by applying (B4)–(B5), and in particular by recalling that $\mathcal{V}_{k+1} = \mathcal{V}_k \setminus \{y_k\}$ for each $k \in [0, m-2]$, $y_{m-1} = 0$, and $\mathcal{V}_{m-1} = \{0\}$, and (f) follows by recalling that $y_0, y_1, \ldots, y_{m-1}$ is a permutation of \mathcal{X} and from our assumption that $\Gamma \in \tilde{\mathcal{B}}_{\mathcal{X}}$.

Proof of Theorem 8. Consider an admission control problem $\mathcal{P} = (Q, r_{\text{cont}}, r_{\text{disc}})$, where $Q = (\mathcal{S}, |\cdot|, q)$. Let Π be a stationary measure of Q. Since Problem (28) is a linear optimization problem whose set of feasible solutions is $\tilde{\mathcal{B}}_{|\mathcal{S}|}$, the conclusion follows from the fundamental theorem of linear programming and by applying Proposition 10 to $|\mathcal{S}|$.

C Proof of Lemma 3

First observe that (19) implies that, from any state $s \in \mathcal{S}$, the only possible transitions are to a state t of $\mathcal{S}_{|s|}$ (if t is of the form $t = s - e_{ik} + e_{i\ell}$), of $\bigcup_{i=1}^{n} \mathcal{S}_{|s|+e_i}$ if t is of the form

- ▶ Lemma 13. Let $s \in S \setminus \{0\}$. Then, for any $i \in [1, m]$ such that $|s|_i > 0$,
- 1. There exist a positive integer r, and $u_0, u_1, \dots, u_{r-1} \in \mathcal{S}$ such that $|u_0| + e_i = |u_1| = \dots = |u_{r-1}| = |s|$, and such that

$$\left(\prod_{j=0}^{r-2} q(u_j, u_{j+1})\right) q(u_{r-1}, s) > 0.$$

2. There exist a positive interger m, and $v_1, \dots, v_m \in \mathcal{S}$ such that $|v_m| + e_i = |v_{m-1}| = \dots = |v_1| = |s|$, and such that

$$q(s, v_1) \left(\prod_{j=1}^{m-1} q(v_j, v_{j+1}) \right) > 0.$$

Proof. Fix $s \in \mathcal{S}$ and $i \in [1, n]$ such that $|s|_i > 0$. Let $k \in [1, n]$ be such that $s_{ik} > 0$, and take $p \equiv p_{ik}$ and k_1, \dots, k_p be such that (20) holds. Let $r \in [1, p]$ be such that $k_r = k$. It then readily follows that, by defining the descending sequence

$$\begin{cases} u_r = s; \\ u_j = u_{j+1} + e_{ik_j} - e_{ik_{j+1}}, & j \in [1, r-1], \text{ if } r \ge 2; \\ u_0 = u_1 - e_{ik_1}, \end{cases}$$

we have

$$|u_0| + e_i = |u_1| = \dots = |u_{r-1}| = |s|.$$

Moreover, it follows from (20) that

$$\prod_{j=0}^{r-1} q(u_j, u_{j+1}) = P_{0,ik_1} \phi_0(u_1 - e_{ik_1}) \left(\prod_{j=1}^{r-1} P_{ik_j, ik_{j+1}} \phi_{ik_j}(u_j) \right) > 0,$$

proving the first assertion. To show the second one we reason similarly, by letting

$$\begin{cases} v_0 &= s; \\ v_{j+1} &= v_j + e_{ik_{r+j+1}} - e_{ik_{r+j}}, \quad j = [0, p-r-1], \text{ if } r \leq p-1; \\ v_{p-r+1} &= v_{p-r} - e_{ik_p}, \end{cases}$$

in a way that

$$|v_{p-r+1}| + e_i = |v_{p-r}| = \dots = |v_1| = |s|,$$

and

$$\prod_{j=0}^{p-r} q(v_j, v_{j+1}) = \left(\prod_{j=0}^{p-r-1} P_{ik_{r+j}, ik_{r+j+1}} \phi_{ik_{r+j}}(v_j)\right) P_{ik_p, 0} \phi_{ik_p}(v_{p-r}) > 0,$$

concluding the proof by setting m = p - r + 1.

We are now ready to show that multi-class Whittle networks are indeed a queueing system, in the sense of Section 2.1.

Proof of Lemma 3. We have to check that the system satisfies Assumptions 1, and we show that it is so for $S_{\text{rec}} \equiv S$. For any s it suffices to apply Assertion 1 of Lemma 13, a number $||s|| := \sum_{i \in [\![1,n]\!], k \in [\![1,m]\!]} s_{ik}$ of times, to obtain, by induction, the existence of ||s|| positive integers $r^1, \cdots, r^{||s||}$ and of ||s|| array of elements $\left(\left(u_0^i, ..., u_{r^i-1}^i\right)\right)_{i=1, \cdots, ||s||}$, such that $u_0^1 = \mathbf{0}$ and such that, by setting $u_0^{||s||+1} := s$ we have

$$|u_0^i|<|u_1^i|=|u_2^i|=\cdots=|u_{r^i-1}^i|=|u_0^{i+1}|, \ \ \text{for all} \ i\in [\![1,\|s\|]\!]$$

and

$$\prod_{i=1}^{\|s\|} \left(\prod_{j=0}^{r^i-2} q(u^i_j, u^i_{j+1}) \right) q(u^i_{r^i-1}, u^{i+1}_0) > 0,$$

implying Assertion 1. Likewise, Assertion 3 follows by applying Assertion 2 of Lemma 13, a number $\|s\| := \sum_{i \in [\![1,n]\!], k \in [\![1,m]\!]} s_{ik}$ of times, to obtain, by induction, the existence of $\|s\|$ positive integers $m^1, \cdots, m^{\|s\|}$ and of $\|s\|$ array of elements $\left(\left(v_1^i, ..., v_{m^i}^i\right)\right)_{i=1, \cdots, \|s\|}$, such that $v_{m^{\|s\|}}^{\|s\|} = \mathbf{0}$ and such that, by setting $v_{m^0}^0 := s$ we have

$$|v_{m^{i-1}}^{i-1}| = |v_1^i| = \dots = |v_{m^{i-1}}^i| > |v_{m^i}^i|, \text{ for all } i \in [1, ||s|]$$

and

$$\prod_{i=1}^{\|s\|} q(v_{m^{i-1}}^{i-1}, v_1^i) \left(\prod_{j=1}^{m^i-1} q(v_j^i, v_{j+1}^i) \right) > 0,$$

in view of Assertion 3. To finish, to check Assertion 4 of Assumptions 1, take two elements $u, s \in \mathcal{S}$ such that $|u| \leq |s|$. As above, we can apply Assertion 2 of Lemma 13 a number ||u|| of times, to construct a path from $\mathbf{0}$ to u having positive probability, whence the fact that \mathcal{S} is coordinate-convex. This concludes the proof of Assumption 1.

D Proof of Lemma 5

In view of (19), from any state $s = s_1 \cdots c_\ell \in \mathcal{S}$, the only possible transitions are to a state of the form $s_1 \cdots s_\ell i$, i.e., an element of $\bigcup_{i=1}^n \mathcal{S}_{|s|+e_i}$, or to a state of the form $s_1 \cdots s_{p-1} s_{p+1} \cdots s_\ell$, that is, an element of $\bigcup_{i=1}^n \mathcal{S}_{|s|-e_i}$.

First, fix $s = s_1 \cdots s_\ell \in \mathcal{S} \setminus \{\varnothing\}$. Set $t_1 = \varnothing$ and, for any $k \in [\![2,\ell+1]\!]$, $t_k = s_1 \cdots s_{k-1}$. Then, for all $k \in [\![1,\ell]\!]$, as \mathcal{S} is prefix-stable, t_k is an element of \mathcal{S} . Moreover we have $|t_k| = |t_{k+1}| - e_{s_k} \leq |t_{k+1}|$, and also $q(t_k,t_{k+1}) = \nu_{s_k} > 0$. This shows Assumption 1. Then, to show Assumption 3, take again $s = s_1 \cdots s_\ell \in \mathcal{S} \setminus \{\varnothing\}$. Observe that, by setting $s_1 \cdots s_0 \equiv \varnothing$,

$$\sum_{p=1}^{\ell} \Delta \mu(s_1 \cdots s_p) = \sum_{p=1}^{\ell} (\mu(s_1 \cdots s_p) - \mu(s_1 \cdots s_{p-1})) = \mu(s) > 0,$$

implying that $\Delta \mu(s_1 \cdots s_p) > 0$ for some $p \in [1, \ell]$. Therefore, setting $t_1 = s$, for such p there exists $t_2 = s_1 \cdots s_{p-1} s_{p+1} \cdots s_{\ell}$, such that $|t_2| = |t_1| - e_{s_p} \le |t_1|$ and $q(t_1, t_2) = |t_1| - e_{s_p} \le |t_1|$

40 Arrival Control in Quasi-Reversible Queueing Systems

 $\Delta\mu(s_1\cdots s_p)>0$. We can then iterate this argument by induction, to prove the existence of a sequence of elements $s=t_1,t_2,\cdots,t_{\ell+1}=\emptyset$, such that $q(t_k,t_{k+1})>0$ and $|t_k|\geq |t_{k+1}|$ for all $k\in [1,\ell]$, completing the proof of Assertion 3.

It remains to check Assertion 4. Let $s = s_1 \cdots s_\ell \in \mathcal{S} \setminus \{\emptyset\}$, and let $y \in \mathbb{N}^n$ be such that $y \leq |s|$. This means exactly that for all $i \in [1, n]$, $|s|_i - y_i \geq 0$. Then, set $\mathcal{I} = \{i \in [1, n]; t \mid s \mid_i - y_i > 0\}$, and for any $i \in \mathcal{I}$, $\alpha_i = |s|_i - y_i > 0$. Then, if we denote by i_1, \dots, i_q , the elements of \mathcal{I} , let \tilde{s} be a permuted version of s ending with the suffix

$$\underline{s} := \underbrace{i_1 \cdots i_1}_{\alpha_{i_1} \text{ times}} \underbrace{i_2 \cdots i_2}_{\alpha_{i_2} \text{ times}} \cdots \underbrace{i_q \cdots i_q}_{\alpha_{i_q} \text{ times}} \,.$$

Then, by permutation stability, \tilde{s} is an element of S. Moreover, the prefix \tilde{s} obtained from \tilde{s} by erasing the suffix \underline{s} , is such that $|\tilde{s}| = y$ by construction, and it is an element of S by prefix-stability. This concludes the proof of Assumptions 1.