SHARP CONCENTRATION OF UNIFORM GENERALIZATION ERRORS IN BINARY LINEAR CLASSIFICATION

SHOGO NAKAKITA

ABSTRACT. We examine the concentration of uniform generalization errors around their expectation in binary linear classification problems via an isoperimetric argument. In particular, we establish Poincaré and log-Sobolev inequalities for the joint distribution of the output labels and the label-weighted input vectors, which we apply to derive concentration bounds. The derived concentration bounds are sharp up to moderate multiplicative constants by those under well-balanced labels. In asymptotic analysis, we also show that almost sure convergence of uniform generalization errors to their expectation occurs in very broad settings, such as proportionally high-dimensional regimes. Using this convergence, we establish uniform laws of large numbers under dimension-free conditions.

1. Introduction

Binary classification is one of the most fundamental tasks in machine learning and statistics. It concerns estimating a high-performing model for predicting a $\{\pm 1\}$ -valued random variable Y (output label) from its corresponding d-dimensional random vector X (input vector) from a sequence of n independent and identically distributed random variables $\{(X_i, Y_i)\}_{i=1}^n$ whose distributions are the same as (X, Y). Binary linear classifiers are fundamental models in this task; we estimate a reasonable $\hat{w} \in \mathbb{R}^d$ by the training data and use $\text{sgn}(\langle X, \hat{w} \rangle)$ as a prediction for the corresponding value of Y. There are multiple approaches to estimating \hat{w} from training data; in this paper, we consider an empirical risk minimization problem such that

minimize
$$\mathcal{R}_n(\mathbf{w}, b) \left(= \frac{1}{n} \sum_{i=1}^n \ell\left(Y_i\left(\langle \mathbf{X}_i, \mathbf{w} \rangle + b\right)\right) \right)$$
 such that $\|\mathbf{w}\| \le R_{\mathbf{w}}, \ |b| \le R_b,$ (1)

where $\ell: \mathbb{R} \to [0, +\infty)$ is an L-Lipschitz continuous function (for some L > 0), $\|\cdot\|$ is the Euclidean norm, $R_w > 0$ is the radius of the space of w, and $R_b \ge 0$ is the radius of the space of b. We use the notation $\mathcal{T} := \{w \in \mathbb{R}^d: \|w\| \le R_w\} \times [-R_b, R_b] \subset \mathbb{R}^{d+1}$. We refer to \mathcal{R}_n as the empirical risk function and ℓ as the loss function. The logistic loss function $\ell_L(t) = \log(1 + \exp(-t))$ and hinge loss function $\ell_H = \max\{0, 1-t\}$ serve as prime examples of Lipschitz loss functions. A basic idea of these loss functions is that their subderivatives satisfy $\ell'(t) \le 0$, and thus they penalize the difference in the signs of Y_i and $\langle X_i, w \rangle + b$. Hence, we can interpret the problem (1) as the selection of (w, b) so that the signs of Y_i and $\langle X_i, w \rangle + b$ are the same as possible.

A motivation for the empirical risk minimization (1) is that we expect that the problem should be close to the following population risk minimization problem:

minimize
$$\mathcal{R}(\mathbf{w}, b) = \mathbb{E}\left[\ell\left(Y(\langle X, \mathbf{w} \rangle + b)\right)\right]$$
 such that $\|\mathbf{w}\| \le R_{\mathbf{w}}, |b| \le R_{b},$ (2)

where $\mathcal{R}(w,b)$ is referred to as a population risk function. Under a mild condition, we know that $\mathcal{R}_n(w,b) \to \mathcal{R}(w,b)$ almost surely for every $(w,b) \in \mathbb{R}^{d+1}$, which is a *pointwise* law of large numbers. However, we need a *uniform* law on \mathcal{T} to verify that the optimization problem (1) is indeed close to (2). In particular, the following type of concentration inequalities for given $\delta \in (0,1]$ has been widely used to characterize the discrepancy between them in studies of machine learning and statistics:

$$\mathbb{P}\left(\sup_{(\boldsymbol{w},b)\in\mathcal{T}}\left(\mathcal{R}(\boldsymbol{w},b)-\mathcal{R}_n(\boldsymbol{w},b)\right)\geq \text{(residual dependent on }(\delta,n,R_{\boldsymbol{w}},R_b))\right)\leq \delta.$$

1

The term $\sup_{(\mathbf{w},b)\in\mathcal{T}}(\mathcal{R}(\mathbf{w},b)-\mathcal{R}_n(\mathbf{w},b))$ is referred to as a *uniform generalization error* or *worst-case generalization error*. If the residual dependent on $(\delta,n,R_{\mathbf{w}},R_b)$ takes small values even for small $\delta>0$, we conclude that the problem (1) is close to (2).

In this work, we analyze the discrepancy between the problems (1) and (2) by showing the following concentration of a uniform generalization error around its expectation:

$$\mathbb{P}\left(\sup_{(\boldsymbol{w},b)\in\mathcal{T}}\left(\mathcal{R}(\boldsymbol{w},b)-\mathcal{R}_{n}(\boldsymbol{w},b)\right)\geq\mathbb{E}\left[\sup_{(\boldsymbol{w},b)\in\mathcal{T}}\left(\mathcal{R}(\boldsymbol{w},b)-\mathcal{R}_{n}(\boldsymbol{w},b)\right)\right]+\left(\operatorname{residual}\right)_{\delta,n,R_{\boldsymbol{w}},R_{b}}\right)\leq\delta. \quad (3)$$

Given a large n, we show that uniform generalization errors concentrate around their expectation in many settings, and the residual terms tend to zero. Hence, our strategy for analysing a uniform generalization error is a divide-and-conquer one; we characterize the behaviour of a uniform generalization error in terms of its expectation, which may be small or large. If both the expectation and residual converge to zero, then we conclude that the empirical risk minimization problem (1) is a good alternative to the population risk minimization problem (2). On the other hand, if the residual converges to zero, but the expectation does not, then the empirical risk minimization problem does not serve as a good approximation of the population risk minimization, and we can identify that the cause of the difficulty is not the random behaviour of uniform generalization errors but the essential difficulty represented by expected uniform generalization errors.

Such a strategy is reasonable and even classical since the evaluation of expected uniform generalization errors is a classical problem in statistical learning theory. For example, the Rademacher complexity argument is a very established strategy to give estimates for the expectation (Ledoux and Talagrand, 1991; Bartlett and Mendelson, 2002; Wainwright, 2019; Bach, 2024), and it is straightforward to derive such a bound in our settings (see Proposition 2). Regarding the residual term, McDiarmid's inequality is a well-known and convenient approach to evaluating it. If the loss function ℓ is bounded, then McDiarmid's inequality yields an explicit and simple residual $\|\ell\|_{\infty}\sqrt{\log(1/\delta)/2n}$ (e.g., see Bach, 2024). The order of this residual is much smaller than that of the Rademacher complexity bound, and thus we can characterize the behaviour of a uniform generalization error by its expectation as the main term.

However, McDiarmid's inequality estimates the residual term much greater than the expectation term obtained by the Rademacher complexity argument when ℓ is Lipschitz but unbounded. For example, unbounded losses (logistic or hinge) combined with unbounded X_i are realistic and often of interest, but McDiarmid's inequality is not directly applicable. Even if X_i is bounded, the residual is of order $O(L(R_w || ||X_i|| ||_{L^\infty} + R_b) \sqrt{\log(1/\delta)/n})$. Since the expected uniform generalization error is of order $O(L(R_w || ||X_i|| ||_{L^2} + R_b)/\sqrt{n})$ by the Rademacher complexity argument, the residual dominates the bound in Eq. (3). Hence, if ℓ is unbounded, McDiarmid's inequality does not capture the concentration of uniform generalization errors around their expectation well.

1.1. Main idea: Isoperimetry in binary linear classification. This study establishes that uniform generalization errors in binary linear classification problems concentrate sharply around their expectation, even if loss functions are unbounded. To obtain residuals significantly smaller than the expectation of uniform generalization errors, we employ isoperimetric properties derived from functional inequalities (Poincaré and log-Sobolev) for the joint distributions of $(Y_iX_i,Y_i)=:(Z_i,Y_i)$. Let us introduce an intuitive example. Assume that $X_i \sim \mathcal{N}(\mathbf{0}, \mathbf{\Sigma})$ with $\|\mathbf{\Sigma}\|_{op} = 1$ and $Y_i \sim \text{Unif}(\{\pm 1\})$ are independent random variables. Then, the joint distribution of (Z_i,Y_i) coincides with that of (X_i,Y_i) . Owing to the independence and tensorization argument (Bakry et al., 2014), we have the following logarithmic Sobolev inequality: for any $f: (\mathbb{R}^d \times \{\pm 1\})^n$ being smooth in the Euclidean arguments,

$$\operatorname{Ent}[f^2((\mathbf{Z}_i, Y_i)_{i=1}^n)] \le 2\mathbb{E}\left[\Gamma(f)((\mathbf{Z}_i, Y_i)_{i=1}^n)\right],$$

where $\mathrm{Ent}[f^2]=\mathbb{E}[f^2\log f^2]-\mathbb{E}[f^2]\log \mathbb{E}[f^2]$ is the entropy operator,

$$\Gamma(f)((z_i, y_i)_{i=1}^n) = \sum_{i=1}^n \left(\|\nabla_{z_i} f((z_i, y_i)_{i=1}^n)\|^2 + |\nabla_{y_i} f((z_i, y_i)_{i=1}^n)|^2 \right)$$

is a carré du champ operator, ∇_{z_i} is the gradient operator with respect to z_i , and ∇_{y_i} is a discrete gradient operator with respect to y_i such that $\nabla_{y_i} f((z_i, y_i)_{i=1}^n) = (f((z_i, y_i)_{i=1}^n) - f((z_j, y_j)_{j=1}^{i-1}, (z_i, -y_i), (z_j, y_j)_{j=i+1}^n))/2$. Then, we have

$$\left\|\Gamma\left(\sup_{(\boldsymbol{w},b)\in\mathcal{T}}\left(\mathcal{R}(\boldsymbol{w},b)\right)-\frac{1}{n}\sum_{i=1}^{n}\ell(\langle z_{i},\boldsymbol{w}\rangle+y_{i}b)\right)\right\|_{\infty}\leq\frac{L^{2}(R_{\boldsymbol{w}}^{2}+R_{b}^{2})}{n};$$

that is, the empirical risk function is $L\sqrt{(R_w^2 + R_b^2)/n}$ -Lipschitz (with respect to the carré du champ operator Γ). Noting that $\ell(Y_i(\langle X_i, w \rangle + b)) = \ell(\langle Z_i, w \rangle + bY_i)$ and employing Herbst's argument (Ledoux, 1999; Bakry et al., 2014), we obtain the following concentration: for any $\delta \in (0, 1]$,

$$\mathbb{P}\left(\sup_{(\boldsymbol{w},b)\in\mathcal{T}}(\mathcal{R}(\boldsymbol{w},b)-\mathcal{R}_n(\boldsymbol{w},b))\geq \mathbb{E}\left[\sup_{(\boldsymbol{w},b)\in\mathcal{T}}(\mathcal{R}(\boldsymbol{w},b)-\mathcal{R}_n(\boldsymbol{w},b))\right]+\sqrt{\frac{2L^2(R_w^2+R_b^2)\log(1/\delta)}{n}}\right)\leq \delta.$$

Although Z_i and Y_i are not independent in general, we expect that suitable isoperimetric properties of the joint distributions of (Z_i, Y_i) will enable the extension of this argument.

Therefore, the problem is when the joint distributions of Z_i and Y_i are isoperimetric in the sense of Poincaré or log-Sobolev inequalities. To develop such functional inequalities, we extend functional inequalities for mixture distributions, which have gathered interest recently (Bardet et al., 2018; Schlichting, 2019; Chen et al., 2021). Let us restrict ourselves to the case with $R_b = 0$ to obtain an intuition; we need functional inequalities for the marginal distribution of Z_i in such a setting. The marginal distribution is a weighted sum of the conditional distributions of Z_i given $Y_i = y \in \{\pm 1\}$, which is a mixture distribution. In general, if the joint state space of two random variables has a good geometric structure (such as an Euclidean space or Riemannian manifold), we can establish functional inequalities for the joint distribution of the variables via standard arguments like the Bakry–Émery condition, and its marginalization is trivial. Hence, the fact that marginal distributions are mixture distributions is not problematic under good geometric structures. However, we need functional inequalities for a marginal distribution of a measure on hybrid continuous–discrete state spaces, which are not manifolds; alternative geometric arguments on such hybrid spaces are not readily available. As a direct strategy to solve this problem, we employ recent results of functional inequalities for mixture distributions. More precisely, considering $R_b > 0$, we extend them to functional inequalities for the joint distribution of continuous and discrete random variables.

1.2. **Literature review.** Asymptotic/non-asymptotic behaviours of high-dimensional binary linear classification problems have received much attention in this decade. Studies on this topic can be mainly classified into two categories: (i) relatively low-dimensional settings $d/n \to 0$ or $d^*/n \to 0$ for some intrinsic dimension d^* (e.g., Kuchelmeister and van de Geer, 2024; Hsu and Mazumdar, 2024; Nakakita, 2024, and references therein), and (ii) proportional high-dimensional settings $d/n \to \kappa \in (0, \infty)$ (e.g., Sur and Candès, 2019; Sur et al., 2019; Salehi et al., 2019; Candès and Sur, 2020). Typical examples of d^* are the number of non-zero signals in sparse settings and an effective rank $\operatorname{tr}(\mathbb{E}[X_iX_i^{\mathsf{T}}])/\|\mathbb{E}[X_iX_i^{\mathsf{T}}]\|_{\operatorname{op}}$ in non-sparse settings. In particular, d^* can be much smaller than d, the dimension of the ambient space, and they characterize the problem better than d itself. Our results in the setting $d/n \to 0$ or $d^*/n \to 0$ yield powerful conclusions, uniform laws of large numbers; hence, this study is well-contextualized in a series of studies with $d/n \to 0$ or $d^*/n \to 0$ (see Section 5.1). Moreover, our results can also be applied to asymptotic studies in proportionally high-dimensional settings; the deviation of the empirical risk minimization problem from the population one under $d/n \to \kappa \in (0, \infty)$ can be partially characterized by expected uniform generalization errors (see Section 5.2).

Let us review studies for relatively low-dimensional settings in detail. Liang and Du (2012) guarantee the asymptotic normality of maximum likelihood estimation in logistic regression under $d/n \to 0$. Kuchelmeister and van de Geer (2024) identify two distinct rates of convergence of maximum likelihood estimation in logistic regression in small and large noise schemes, the probit model as a data generating model, and $d/n \to 0$. Hsu and Mazumdar (2024) show the phase transition of the sample complexities for

logistic regression in three temperature settings and $d/n \to 0$. van de Geer (2008) gives non-asymptotic guarantees for generalized linear models including logistic regression, and the bounds converge to zero under $d^*/n \to 0$ (ignoring logarithmic factors for simplicity), where d^* is the number of non-zero elements of the true value of the parameter. Nakakita (2024) gives a dimension-free concentration bound for uniform generalized errors around zero in logistic regression, which converges under $d^*/n \to 0$, where d^* is the effective rank of the covariance matrix of X_i . Our results resemble the result of Nakakita (2024) but have some essential differences; we compare our results with it in Section 5.1.

It is remarkable that a number of studies in high-dimensional binary linear classification problems do not address the bias of Y_i (that is to say, they restrict themselves to the settings $R_b = 0$ and $\mathbb{P}(Y_i = +1) = \mathbb{P}(Y_i = -1) = 1/2$). Bias is the most fundamental parameter in the prediction of Y given a new input vector X, and large prediction errors can occur if we ignore it. If the distribution of X is sign-invariant, then for any \hat{w} independent of X, $\mathbb{P}(\text{sgn}(\langle X, \hat{w} \rangle) = +1) = \mathbb{P}(\text{sgn}(\langle X, \hat{w} \rangle) = -1) = 1/2$; then, the prediction results in large errors if $\mathbb{P}(Y = +1) \neq \mathbb{P}(Y = -1)$. This study is not the first one enabling us to characterize such problems in the presence of both bias and high dimensionality (e.g., Nakakita, 2024, can address it by considering non-centred input vectors), but is one of the few studies to estimate the effect of biases under high-dimensional settings explicitly.

1.3. **Paper organization.** Section 2 provides notation and definitions used in this paper. Section 3 exhibits functional inequalities for distributions on $\mathbb{R}^d \times \{\pm 1\}$, which are the main technical contributions of this study. Section 4 gives concentration inequalities for uniform generalization errors in binary linear classification problems. Section 5 is for applications of our non-asymptotic results to asymptotic studies in two settings: a dimension-free regime and proportionally high-dimensional regime. Section 6 gives some concluding remarks and possible future directions of research. All the technical proofs are deferred to the Appendix.

2. Notation and definitions

In this section, we introduce notation and definitions used throughout this paper.

For any matrix $\mathbf{A} \in \mathbb{R}^{p \times q}$, $\|\mathbf{A}\|_{\text{op}} := \sup_{\mathbf{x} \in \mathbb{R}^q : \|\mathbf{x}\| \neq 0} \|\mathbf{A}\mathbf{x}\| / \|\mathbf{x}\|$. $\sharp(\cdot)$ denotes the counting measure on $\{\pm 1\}$. For $f: \mathbb{R}^d \times \{\pm 1\} \to \mathbb{R}$, we define a discrete gradient $\nabla_y f(z,y) := (f(z,y) - f(z,-y))/2$. For $f: (\mathbb{R}^d \times \{\pm 1\})^n \to \mathbb{R}$, we also define a discrete gradient $\nabla_{y_i} f((z_i,y_i)_{i=1}^n) := (f((z_i,y_i)_{i=1}^n) - f((z_j,y_j)_{j=1}^{i-1},(z_i,-y_i),(z_j,y_j)_{j=i+1}^n)))/2$. \mathcal{A}_1 denotes the family of all functions $f: \mathbb{R}^d \times \{\pm 1\} \to \mathbb{R}$ (resp. $f: (\mathbb{R}^d \times \{\pm 1\})^n \to \mathbb{R}$) such that for both $y \in \{\pm 1\}$, f(z,y) is locally Lipschitz continuous with respect to $z \in \mathbb{R}^d$. Similarly, \mathcal{A}_n is the family of all functions $f: (\mathbb{R}^d \times \{\pm 1\})^n \to \mathbb{R}$ such that for any $(y_i)_{i=1}^n \in \{\pm 1\}^n$, $f((z_i,y_i)_{i=1}^n)$ is locally Lipschitz with respect to $(z_i)_{i=1}^n \in \mathbb{R}^{dn}$.

For any nonnegative random variable ξ , we let $\operatorname{Ent}_{\rho}(\xi)$ denote the entropy of ξ with a probability measure ρ defined as follows:

$$\operatorname{Ent}_{\rho}(\xi) := \mathbb{E}_{\rho}[\xi \log \xi] - \mathbb{E}_{\rho}[\xi] \log \mathbb{E}_{\rho}[\xi].$$

We use subscripts for the expectation \mathbb{E} , variance Var, and entropy Ent to indicate the underlying measure. In particular, when these measures are equipped with subscripts (e.g., P_X), we adopt the same subscripts for the corresponding expectation, variance, and entropy to specify the measure they are taken with respect to (e.g., \mathbb{E}_X , Var_X, and Ent_X).

We call a symmetric positive bilinear operator Γ on $\mathcal{A}_1 \times \mathcal{A}_1$ (resp. $\mathcal{A}_n \times \mathcal{A}_n$) such that (i) $\Gamma(f_1, f_2)$: $\mathbb{R}^d \times \{\pm 1\} \to \mathbb{R}$ for $f_1, f_2 \in \mathcal{A}_1$ (res. $\Gamma(f_1, f_2)$: $(\mathbb{R}^d \times \{\pm 1\})^n \to \mathbb{R}$ for $f_1, f_2 \in \mathcal{A}_n$) is measurable, (ii) $\Gamma(f_1, f_1) \geq 0$, (iii) $\Gamma(f_1, f_2) = \Gamma(f_2, f_1)$, and (iv) $\Gamma(1, 1) = 0$ as a *carré du champ operator*. A typical example is $\Gamma(f_1, f_2) = \langle \nabla_z f_1, \nabla_z f_2 \rangle + \langle \nabla_y f_1, \nabla_y f_2 \rangle$ for $f_1, f_2 \in \mathcal{A}_1$. For notational convenience, we write $\Gamma(f) = \Gamma(f, f)$.

We provide the formal definitions of the evenness of functions and m-convexity.

Definition 1. $f : \mathbb{R}^d \to \mathbb{R}$ is an *even* function if $f(\mathbf{x}) = f(-\mathbf{x})$ for all $\mathbf{x} \in \mathbb{R}^d$.

Definition 2. A convex function $f: \mathbb{R}^d \to \mathbb{R}$ is m-convex with $m \ge 0$ if for any $x, y \in \mathbb{R}^d$ and any $g \in \mathbb{R}^d$ being a subgradient of f at $x \in \mathbb{R}^d$,

$$f(\mathbf{y}) - f(\mathbf{x}) - \langle \mathbf{g}, \mathbf{y} - \mathbf{x} \rangle \ge \frac{m}{2} \|\mathbf{y} - \mathbf{x}\|^2.$$

All convex functions are 0-convex by definition.

3. Fundamental results for functional inequalities

In this section, we discuss functional inequalities for the joint distribution of general random variables (\mathbf{Z}, Y) on $\mathbb{R}^d \times \{\pm 1\}$. We suppose that (\mathbf{Z}, Y) are absolutely continuous with respect to the product measure of the d-dimensional Lebesgue measure and the counting one \sharp .

3.1. **Settings.** We introduce the notation of the joint, marginal, and conditional measures of Y and Z. We let $P_{ZY}(dzdy)$ denote the joint measure of (Z,Y) on $\mathbb{R}^d \times \{\pm 1\}$.

We use the following notation for marginal/conditional measures and densities:

$$P_{\mathbf{Z}}(\mathrm{d}z) := P_{\mathbf{Z}Y}(\mathrm{d}z \times \{\pm 1\}), \qquad P_{Y}(\mathrm{d}y) := P_{\mathbf{Z}Y}(\mathbb{R}^{d} \times \mathrm{d}y) =: p_{Y}(y)\sharp(\mathrm{d}y),$$

$$P_{\mathbf{Z}|Y}(\mathrm{d}z|y) := \frac{1}{p_{Y}(y)}P_{\mathbf{Z}Y}(\mathrm{d}z \times \{y\}), \qquad P_{Y|\mathbf{Z}}(\mathrm{d}y|z) := \frac{P_{\mathbf{Z}Y}(\mathrm{d}z\mathrm{d}y)}{P_{\mathbf{Z}}(\mathrm{d}z)} =: p_{Y|\mathbf{Z}}(y|z)\sharp(\mathrm{d}y).$$

We also set the following carré du champ operators on $\mathcal{A}_1 \times \mathcal{A}_1$:

$$\Gamma_{\mathbf{Z}}(f_1, f_2)(z, y) := \langle \nabla_{\mathbf{z}} f_1(z, y), \nabla_{\mathbf{z}} f_2(z, y) \rangle, \ \Gamma_{\mathbf{Y}}(f_1, f_2)(z, y) := (\nabla_{\mathbf{y}} f_1(z, y) \nabla_{\mathbf{y}} f_2(z, y)).$$

We define the following constants:

$$K_{P} := \max_{y \in \{\pm 1\}} C_{P}(P_{\mathbf{Z}|Y}(\cdot|y)), K_{LS} = \max_{y \in \{\pm 1\}} C_{LS}(P_{\mathbf{Z}|Y}(\cdot|y)),$$

where $C_P(P(\cdot))$ and $C_{LS}(P(\cdot))$ with $P(\cdot)$, an arbitrary probability measure on \mathbb{R}^d , are defined as the minimal constants satisfying a Poincaré inequality and log-Sobolev inequality such that for any locally Lipschitz $f: \mathbb{R}^d \to \mathbb{R}$,

$$\operatorname{Var}_{P}(f) \leq C_{P}(P(\cdot))\mathbb{E}_{P}\left[\|\nabla f\|^{2}\right], \operatorname{Ent}_{P}(f) \leq 2C_{\operatorname{LS}}(P(\cdot))\mathbb{E}_{P}\left[\|\nabla f\|^{2}\right].$$

We also set the following notation:

$$K_{\chi^2} := \max_{y \in \{\pm 1\}} \chi^2 \left(P_{\mathbf{Z}}(\cdot) \| P_{\mathbf{Z}|Y}(\cdot|y) \right), \ K_{\mathbf{V}} := 4p_Y(+1)p_Y(-1), \ K_{\mathbf{U}} := \max \left\{ \frac{1}{p_Y(+1)}, \frac{1}{p_Y(-1)} \right\}.$$

 K_{χ^2} quantifies the dependence between **Z** and *Y*. If **Z** \perp *Y*, then $K_{\chi^2} = 0$. K_V is the variance of *Y*. The constant K_U represents the degree of deviation from the uniform distribution on $\{\pm 1\}$.

3.2. **Statement.** We present Poincaré and log-Sobolev inequalities for weighted sums of the carré du champ operators $\Gamma_{\mathbf{Z}}$ and $\Gamma_{\mathbf{Y}}$. Although the inequalities formally appear with constant 1, the associated carré du champ operators may carry large scaling factors. We adopt the convention that $\infty \times 0 = 0$.

Theorem 1. The following Poincaré and log-Sobolev inequalities hold: for any $f \in \mathcal{A}_1$,

$$Var(f(\mathbf{Z}, Y)) \leq \mathbb{E}\left[\Gamma_{P}(f)(\mathbf{Z}, Y)\right],$$

$$Ent(f^{2}(\mathbf{Z}, Y)) \leq 2\mathbb{E}\left[\Gamma_{LS}(f)(\mathbf{Z}, Y)\right],$$

where Γ_P and Γ_{LS} are the following weighted carré du champ operators for any conjugate pair $c, c^* \in [1, \infty]$ such that $1/c + 1/c^* = 1$: for any $f_1, f_2 \in \mathcal{A}_1$,

$$\begin{split} &\Gamma_{\mathrm{P}}(f_{1},f_{2}) := K_{\mathrm{P}}\left(1 + cK_{\chi^{2}}\right)\Gamma_{\mathbf{Z}}(f_{1},f_{2}) + c^{*}K_{\mathrm{V}}\Gamma_{Y}(f_{1},f_{2}), \\ &\Gamma_{\mathrm{LS}}(f_{1},f_{2}) := \left(1 + \frac{1}{2}\log K_{\mathrm{U}}\right)\Gamma_{\mathrm{P}}(f_{1},f_{2}) + 2K_{\mathrm{LS}}\Gamma_{\mathbf{Z}}(f_{1},f_{2}). \end{split}$$

If $\mathbf{Z} \perp Y$ and thus $K_{Y^2} = 0$, then we can select $c = \infty$ and $c^* = 1$ and obtain

$$\Gamma_{P}(f_{1}, f_{2}) = C_{P}(P_{Z}(\cdot))\Gamma_{Z}(f_{1}, f_{2}) + \Gamma_{Y}(f_{1}, f_{2}),$$

which exactly recovers the result by tensorization. Γ_{LS} can also be recovered up to a constant $1+(1/2)\log K_{\rm U}$, which should be multiplied with $\Gamma_{\rm Y}$ due to the well-known logarithmic Sobolev constant for binomial distributions (see Boucheron et al., 2013); the multiplicative constant of $\Gamma_{\rm Z}$ in the logarithmic Sobolev inequality above is optimal up to a numerical constant and $\log K_{\rm U}$.

4. Concentration inequalities for uniform generalization errors

We now apply Theorem 1 to concentration bounds on uniform generalization errors for binary linear classification problems.

We first introduce a basic estimate for the expectation of uniform generalization errors, which should be the main term of the bound (3).

Proposition 2. We have

$$\mathbb{E}\left[\sup_{(\boldsymbol{w},b)\in\mathcal{T}}(\mathcal{R}(\boldsymbol{w},b)-\mathcal{R}_n(\boldsymbol{w},b))\right] \leq \frac{2L\left(R_{\boldsymbol{w}}\sqrt{\operatorname{tr}(\mathbb{E}[\boldsymbol{X}_i\boldsymbol{X}_i^{\top}])}+R_b\right)}{\sqrt{n}}.$$

Note that this result is dimension-independent, but depends on a possibly large quantity $\operatorname{tr}(\mathbb{E}[X_iX_i^{\top}])$. The purpose of this section is to show that the residual term in Equation (3) is much smaller than this standard estimate for expected uniform generalization errors.

4.1. **Basic settings.** We set the probability measure of X_i as

$$P_X(\mathrm{d}x) = Z^{-1} \exp\left(-U(x)\right) \mathrm{d}x,$$

where $U: \mathbb{R}^d \to [0, \infty)$ is the potential function, and $\mathcal{Z} = \int_{\mathbb{R}^d} \exp(-U(x)) dx$ is the normalizing constant assumed to be finite. Although theorems below assume the evenness of U, we can relax them to the settings with non-central input vectors; see Section 4.4 for details. We also set the conditional probability mass function of Y_i on $\{\pm 1\}$ given $X_i = x_i \in \mathbb{R}^d$ as

$$p_{Y|X}(y|x) = g(y(\langle x, \theta_1 \rangle + \theta_0)),$$

where $g: \mathbb{R} \to [0, 1]$ is a function called a link function, $\theta_1 \in \mathbb{R}^d$ is a fixed vector, and $\theta_0 \in \mathbb{R}$ is a fixed number. By definition, we have g(t) + g(-t) = 1 for all $t \in \mathbb{R}$.

Under these settings, we derive the following joint probability measure of (\mathbf{Z}_i, Y_i) :

$$P_{ZY}(\mathrm{d}z\mathrm{d}y) = \mathcal{Z}^{-1}g(\langle z, \theta_1 \rangle + y\theta_0) \exp(-U(yz))\mathrm{d}z\sharp(\mathrm{d}y).$$

The marginal density/mass functions and conditional density/mass functions are given as

$$p_{\mathbf{Z}}(z) := \frac{P_{\mathbf{Z}}(\mathrm{d}z)}{\mathrm{d}z} = \sum_{y \in \{\pm 1\}} \mathcal{Z}^{-1} g(\langle z, \theta_1 \rangle + y\theta_0) \exp(-U(yz)),$$

$$p_{Y}(y) := \frac{P_{y}(\mathrm{d}y)}{\sharp(\mathrm{d}y)} = \frac{1}{\mathcal{Z}} \int_{\mathbb{R}^d} g(\langle z, \theta_1 \rangle + y\theta_0) \exp(-U(yz)) \mathrm{d}z =: \frac{\mathcal{Z}_y}{\mathcal{Z}},$$

$$p_{\mathbf{Z}|Y}(z|y) = \mathcal{Z}_y^{-1} g(\langle z, \theta_1 \rangle + y\theta_0) \exp(-U(yz)),$$

$$p_{Y|\mathbf{Z}}(y|z) = \frac{g(\langle z, \theta_1 \rangle + y\theta_0) \exp(-U(yz))}{\sum_{y' \in \{\pm 1\}} g(\langle z, \theta_1 \rangle + y'\theta_0) \exp(-U(y'z))}.$$

Here, we employ the fact $\mathcal{Z} = \int_{\mathbb{R}^d} \exp(-U(x)) dx = \sum_{y \in \{\pm 1\}} \int_{\mathbb{R}^d} g(y(\langle x, \theta_1 \rangle + \theta_0)) \exp(-U(x)) dx = \sum_{y \in \{\pm 1\}} \int_{\mathbb{R}^d} g(\langle z, \theta_1 \rangle + y\theta_0) \exp(-U(yz)) dz$. Our loss function $\ell(\cdot)$ does not necessarily coincide with $-\log(g(\cdot))$.

We set the assumptions on functional inequalities of the conditional distributions $P_{\mathbf{Z}|Y}(\cdot|y)$ for $y \in \{\pm 1\}$.

Assumption 1 (Poincaré inequality). For both $y \in \{\pm 1\}$, the conditional distribution $P_{\mathbf{Z}|Y}(\cdot|y)$ satisfies a Poincare inequality (with respect to $\Gamma_{\mathbf{Z}}$) for some constant $K_{\mathbf{P}} > 0$.

Assumption 2 (log-Sobolev inequality). For both $y \in \{\pm 1\}$, the conditional distribution $P_{Z|Y}(\cdot|y)$ satisfies a log-Sobolev inequality (with respect to Γ_Z) for some constant $K_{LS} > 0$.

Note that a Poincaré inequality yields exponential concentration of arbitrary Lipschitz maps, while a log-Sobolev inequality yields Gaussian concentration of arbitrary Lipschitz maps. Since $K_P \le K_{LS}$ in general, sufficient conditions for a log-Sobolev inequality are stronger than those for a Poincaré inequality.

Let us introduce some natural sufficient conditions for Assumption 2. For example, if U is K^{-1} -convex, $\log \circ g$ is concave, and they are of C^2 , then the potential of the conditional distribution is K^{-1} -convex, and the conditional distribution satisfies a log-Sobolev inequality with constant K by the Bakry-Émery theory. Note that $U(x) = (1/2)\langle x, \Sigma^{-1}x \rangle$, the potential function of $\mathcal{N}_d(\mathbf{0}_d, \Sigma)$ for positive definite Σ , is $\|\Sigma\|_{\text{op}}$ -convex. Moreover, if U is even, $\log \circ g$ is G-Lipschitz, and a log-Sobolev inequality for the probability measure $(\int_{\mathbb{R}^d} g(\langle z, \theta_1 \rangle) e^{-U(z)} \mathrm{d}z)^{-1} g(\langle z, \theta_1 \rangle) e^{-U(z)} \mathrm{d}z$ holds with constant $Ke^{-2G|\theta_0|}$, then the perturbation theory (Bakry et al., 2014; Cattiaux and Guillin, 2022) gives Assumption 2. It is because $g(t + |\theta_0|)/g(t) \leq e^{G|\theta_0|}$ and $g(t - |\theta_0|)/g(t) \geq e^{-G|\theta_0|}$ for all $t \in \mathbb{R}$. For example, the 1-Lipschitz continuity of $\log \circ \sigma$ holds, where $\sigma(t) := 1/(1 + \exp(-t))$ is the logistic link function.

We also introduce sufficient conditions for Assumption 1. Since $K_P \leq K_{LS}$, sufficient conditions for Assumption 2 are also sufficient for Assumption 1. A log-Sobolev inequality in Theorem 1 includes the multiplicative constant $(1 + (1/2) \log K_U)$, which can be large in some situations (Theorems 6 and 7); hence, considering exponential concentration is sometimes beneficial even if we know that Assumption 2 holds. In addition, if both U and $(-1) \log \circ g$ are convex (here, we do not require the m-convexity of U for some positive m > 0), we can give estimates for the Poincaré constants by applying recent results on the Kannan– Lovász–Simonovits (KLS) conjecture (Kannan et al., 1995), which is a topical problem (Lee and Vempala, 2018; Chen, 2021; Klartag and Lehec, 2022; Jambulapati et al., 2022; Klartag, 2023; Lee and Vempala, 2024). Let us introduce a constant $C_{\text{KLS}}(d) = \sup_{\mu: \text{isotropic and log-concave on } \mathbb{R}^d} C_P(\mu(\cdot))$, where $C_P(\mu(\cdot))$ is the minimal constant in a Poincaré inequality of μ such that for any locally Lipschitz $f: \mathbb{R}^d \to \mathbb{R}$, $\operatorname{Var}_{\mu}(f) \leq C_{P}(\mu(\cdot))\mathbb{E}_{\mu}[\|\nabla f\|^{2}].$ We notice that $C_{P}(P_{\mathbf{Z}_{i}|Y}(\cdot|y)) \leq \|\mathbb{E}[\mathbf{Z}_{i}\mathbf{Z}_{i}^{\top}] - \mathbb{E}[\mathbf{Z}_{i}]\mathbb{E}[\mathbf{Z}_{i}]^{\top}\|_{op}C_{KLS}(d) \leq C_{P}(\mu(\cdot))\mathbb{E}_{\mu}[\|\nabla f\|^{2}].$ $\|\mathbb{E}[X_iX_i^{\mathsf{T}}]\|_{\mathrm{op}}C_{\mathrm{KLS}}(d)$. The statement of the KLS conjecture is " $C_{\mathrm{KLS}} := \sup_{d \in \mathbb{N}} C_{\mathrm{KLS}}(d) < \infty$ "; to our best knowledge, the most recent estimate is $C_{KLS}(d) \le c\sqrt{\log d}$ for some absolute constant c > 0 by Klartag (2023). Hence, we can let $K_P = c\sqrt{\log d} \|\mathbb{E}[X_i X_i^{\mathsf{T}}]\|_{\text{op}}$ under the (weak) convexity of U and $(-1) \log \circ g$. Moreover, we can apply an upper bound given by Lee and Vempala (2024) in the place of $\|\mathbb{E}[X_i X_i^{\mathsf{T}}]\|_{\text{op}} C_{\text{KLS}}(d)$; they yield $C_P(P_{\mathbf{Z}|Y}(\cdot|y)) \leq c' \text{tr}(\mathbb{E}[X_i X_i^{\mathsf{T}}]^2)^{1/2}$ for an absolute constant c' > 0. Hence, we can let $K_P = c' \operatorname{tr}(\mathbb{E}[X_i X_i^{\top}]^2)^{1/2}$. The residual derived by this result is smaller than that by McDiarmid's inequality and the Rademacher complexity bound for the expected uniform generalization error given by Proposition 2 up to $\log(3/\delta)$ and a numerical constant.

4.2. Concentration under no bias via tensorization. First, we consider the case under $\theta_0 = 0$ as the baseline analysis; to our best knowledge, this sharp result itself is novel as dimension-free concentration inequalities for uniform generalization errors in binary linear classification problems without boundedness. The setting $\theta_0 = 0$ results in the independence of \mathbf{Z}_i and Y_i .

Proposition 3. Assume that the potential function U is even and $\theta_0 = 0$. Then, \mathbf{Z}_i and Y_i are independent.

Using Proposition 3, we derive the following sharp concentration inequality via the classical tensorization argument without our Theorem 1. Hence, we take this result as a baseline for comparison, and we will give concentration inequalities similar to this inequality in more general settings.

Proposition 4 (concentration under no bias). Suppose that the same assumption as Proposition 3 holds true.

(i) Under Assumption 1, for all $\delta \in (0, 1]$, with probability at least $1 - \delta$,

$$\sup_{(\boldsymbol{w},b)\in\mathcal{T}} (\mathcal{R}(\boldsymbol{w},b) - \mathcal{R}_n(\boldsymbol{w},b)) - \mathbb{E}\left[\sup_{(\boldsymbol{w},b)\in\mathcal{T}} (\mathcal{R}(\boldsymbol{w},b) - \mathcal{R}_n(\boldsymbol{w},b))\right] \leq \sqrt{\frac{L^2(K_PR_w^2 + R_b^2)(\log(3/\delta))^2}{n}}.$$

(ii) Under Assumption 2, for all $\delta \in (0, 1]$, with probability at least $1 - \delta$,

$$\sup_{(\boldsymbol{w},b)\in\mathcal{T}}(\mathcal{R}(\boldsymbol{w},b)-\mathcal{R}_n(\boldsymbol{w},b))-\mathbb{E}\left[\sup_{(\boldsymbol{w},b)\in\mathcal{T}}(\mathcal{R}(\boldsymbol{w},b)-\mathcal{R}_n(\boldsymbol{w},b))\right]\leq \sqrt{\frac{2L^2(K_{\mathrm{LS}}R_{\boldsymbol{w}}^2+R_b^2)\log(1/\delta)}{n}}.$$

Remark 1. The typical examples of log-concave g is the logistic link function σ and probit link function Φ defined as for all $t \in \mathbb{R}$,

$$\sigma(t) := \frac{1}{1 + \exp(-t)}, \ \Phi(t) := \int_{-\infty}^{t} \frac{\exp(-s^2/2)}{\sqrt{2\pi}} ds.$$
 (4)

4.3. Concentration in general settings via Theorem 1. Theorem 4 yields a sharp bound, but it needs the assumption $\theta_0 = 0$. We use our Theorem 1 to investigate the concentration of uniform generalization errors in more general settings. We exhibit four concentration inequalities, which are sharp under the following four edge cases respectively (see Figure 1): small bias $|\theta_0| = O(1)$; large bias $|\theta_0| \gg 1$; weak signal $||\theta_1|| = O(1)$; and strong signal $||\theta_1|| \gg 1$. In particular, the cases $|\theta_0| = O(1)$ (Proposition 5) and $||\theta_1|| \gg 1$ (Proposition 8) yield upper bounds matching that of Theorem 4 up to mild multiplicative constants.




FIGURE 1. A schematic representation of the regimes covered in Propositions 5–8. The cyan line represents the zero bias region (studied by Proposition 4). Situations in the region close to the left edge (that is, small $|\theta_0|$) can be evaluated by Proposition 5; the same applies to other cases. If both $|\theta_0|$ and $||\theta_1||$ are moderately large (the light orange region in the middle), then we can choose an arbitrary edge minimizing the tail.

4.3.1. Concentration under small bias. In the next place, we examine the case where $|\theta_0|$ is non-zero but not very large. If $\theta_0 \neq 0$, then \mathbf{Z}_i and Y_i are dependent in general. Therefore, we apply Theorem 1 instead of tensorization and derive the following result.

Proposition 5 (concentration under small bias). Assume that the potential function U is even, g is non-decreasing, and $\log \circ g$ is G-Lipschitz.

(i) Suppose that Assumption 1 holds true. Then, for all $\delta \in (0,1]$, with probability at least $1-\delta$,

$$\sup_{(\boldsymbol{w},b)\in\mathcal{T}} (\mathcal{R}(\boldsymbol{w},b) - \mathcal{R}_n(\boldsymbol{w},b)) - \mathbb{E}\left[\sup_{(\boldsymbol{w},b)\in\mathcal{T}} (\mathcal{R}(\boldsymbol{w},b) - \mathcal{R}_n(\boldsymbol{w},b))\right]$$

$$\leq \sqrt{\frac{L^2 \left[K_P \left(e^{2G|\theta_0|} + \sqrt{e^{2G|\theta_0|} - 1}\right) R_{\boldsymbol{w}}^2 + (1 + \sqrt{e^{2G|\theta_0|} - 1}) R_b^2\right] (\log(3/\delta))^2}{n}}.$$

(ii) Suppose that Assumption 2 holds true. Then, for all $\delta \in (0,1]$, with probability at least $1-\delta$,

$$\sup_{(\boldsymbol{w},b)\in\mathcal{T}} (\mathcal{R}(\boldsymbol{w},b) - \mathcal{R}_n(\boldsymbol{w},b)) - \mathbb{E}\left[\sup_{(\boldsymbol{w},b)\in\mathcal{T}} (\mathcal{R}(\boldsymbol{w},b) - \mathcal{R}_n(\boldsymbol{w},b))\right]$$

$$\leq \sqrt{\frac{2L^2 (3 + 2G|\theta_0|) \left[K_{LS} \left(\frac{1}{2} + e^{2G|\theta_0|}\right) R_{\boldsymbol{w}}^2 + R_b^2\right] \log(1/\delta)}{n}}.$$

The upper bound above is non-vacuous compared to that of Proposition 4 as long as $|\theta_0|$ is not very large. In particular, if $|\theta_0| = 0$, then the right-hand side in (i) coincides with that of Proposition 4-(i), and the right-hand side in (ii) coincides with that of Proposition 4 up to a constant factor of 3.

Remark 2. It is easy to see that $\log \circ \sigma$, the composition of the logarithmic function and the logistic link function (4), is 1-Lipschitz. On the other hand, $\log \circ \Phi$, the composition of the logarithmic function and the probit link function (4), is not Lipschitz. Hence, we can assume logistic models as the true model, but cannot set probit models to apply Proposition 5 under $\theta_0 \neq 0$.

4.3.2. Concentration under large bias. If $|\theta_0|$ is large, then the upper bound given by Proposition 5 can be vacuous; however, we should guess that Y_i is almost constant if $|\theta_0|$ is very large, and then Z_i and Y_i are again approximately independent. In this section, we restrict ourselves to discussing the logistic link function $g = \sigma$ and see that the guess is certainly correct in a specified setting.

We define the following moment-generating function:

$$M_X(t) := \mathbb{E}\left[\exp\left(\langle X, t \rangle\right)\right], \ t \in \mathbb{R}^d.$$

If U is even, then M_X is also an even function.

Proposition 6 (concentration under large bias and logistic models). Assume that the potential function U is even and $g(t) = \sigma(t) = 1/(1 + \exp(-t))$.

(i) Suppose that Assumption 1 holds. Then, for all $\delta \in (0,1]$, with probability at least $1-\delta$,

$$\begin{split} \sup_{(\boldsymbol{w},b)\in\mathcal{T}} & (\mathcal{R}(\boldsymbol{w},b) - \mathcal{R}_n(\boldsymbol{w},b)) - \mathbb{E}\left[\sup_{(\boldsymbol{w},b)\in\mathcal{T}} (\mathcal{R}(\boldsymbol{w},b) - \mathcal{R}_n(\boldsymbol{w},b))\right] \\ \leq & \sqrt{\frac{L^2\left[K_{\mathrm{P}}\left(8M_X^2(\boldsymbol{\theta}_1) - 1\right)R_{\boldsymbol{w}}^2 + 8e^{-|\boldsymbol{\theta}_0|}M_X(\boldsymbol{\theta}_1)R_b^2\right]\left(\log(3/\delta)\right)^2}{n}}. \end{split}$$

(ii) Suppose that Assumption 2 holds. Then, for all $\delta \in (0,1]$, with probability at least $1-\delta$,

$$\begin{split} &\sup_{(\boldsymbol{w},b)\in\mathcal{T}}(\mathcal{R}(\boldsymbol{w},b)-\mathcal{R}_n(\boldsymbol{w},b)) - \mathbb{E}\left[\sup_{(\boldsymbol{w},b)\in\mathcal{T}}(\mathcal{R}(\boldsymbol{w},b)-\mathcal{R}_n(\boldsymbol{w},b))\right] \\ &\leq \sqrt{\frac{L^2\left(e+|\theta_0|\right)\left[K_{\mathrm{LS}}\left(1/2+8M_X^2(\theta_1)\right)R_{\boldsymbol{w}}^2+8e^{-|\theta_0|}M_X(\theta_1)R_b^2\right]\log(1/\delta)}{n}}. \end{split}$$

The right-hand side in Proposition 6-(i) is decreasing in $|\theta_0|$; therefore, large $|\theta_0|$ is beneficial to yield a sharp bound from this proposition. Proposition 6-(ii) gives a residual with dependence on $\sqrt{|\theta_0|}$ (ignoring other parameters), which is much milder than $\exp(G|\theta_0|)$ of Proposition 5-(ii). For instance, if $|\theta_0| = o(n)$

in statistical experiments, then Proposition 6-(ii) concludes that a uniform generalization error sharply concentrates around its expectation. Removing the dependence on $|\theta_0|$ via the improvement of Theorem 1 is an important direction of further investigation.

4.3.3. Concentration under weak signal. We can guess that the residual in Equation (3) should be close to that of Proposition 4 if $\|\theta_1\|$ is very small; it is because the dependence between X_i and Y_i or that between Z_i and Y_i should be weak.

Let us define the following functions resembling the moment-generating function of X:

$$M_X^{\text{num}}(t) := \mathbb{E}\left[\exp\left(\max\{0,\langle X,t\rangle\}\right)\right], \ M_X^{\text{den}}(t) := \mathbb{E}\left[\exp\left(\min\{0,\langle X,t\rangle\}\right)\right], \ t \in \mathbb{R}^d.$$

We know that $M_X^{\mathrm{den}}(t) \geq \exp(\mathbb{E}[\min\{0,\langle X,t\rangle\}]) = \exp(-\mathbb{E}[|\langle X,t\rangle|]/2) > 0$ by Jensen's inequality if U is even. For instance, if $X \sim \mathcal{N}_d(\mathbf{0}_d, \Sigma)$ for some positive definite $\Sigma \in \mathbb{R}^{d \times d}$, $M_X^{\mathrm{den}}(t) \geq \exp(-\langle t, \Sigma t \rangle / \sqrt{2\pi})$. If U is even, $M_X^{\mathrm{num}}(t) \leq \mathbb{E}[\exp(|\langle X,t\rangle|)] \leq \mathbb{E}[\exp(\langle X,t\rangle) + \exp(\langle -X,t\rangle)] \leq 2M_X(t)$.

Proposition 7 (concentration under weak signal). Assume that the potential function U is even, g is non-decreasing, and $\log \circ g$ is G-Lipschitz.

(i) Suppose that Assumption 1 holds true. We have for all $\delta \in (0,1]$, with probability at least $1-\delta$,

$$\sup_{(\boldsymbol{w},b)\in\mathcal{T}} (\mathcal{R}(\boldsymbol{w},b) - \mathcal{R}_n(\boldsymbol{w},b)) - \mathbb{E}\left[\sup_{(\boldsymbol{w},b)\in\mathcal{T}} (\mathcal{R}(\boldsymbol{w},b) - \mathcal{R}_n(\boldsymbol{w},b))\right]$$

$$\leq \sqrt{\frac{L^2\left[K_P\left(4M_X^{\text{num}}(G\boldsymbol{\theta}_1)M_X^{\text{num}}(2G\boldsymbol{\theta}_1) - 1\right)R_{\boldsymbol{w}}^2 + 8g(+\theta_0)g(-\theta_0)M_X^{\text{num}}(G\boldsymbol{\theta}_1)R_b^2\right](\log(3/\delta))^2}{n}}.$$

(ii) Suppose that Assumption 2 holds true. We have for all $\delta \in (0,1]$, with probability at least $1-\delta$,

$$\sup_{(\boldsymbol{w},b)\in\mathcal{T}} (\mathcal{R}(\boldsymbol{w},b) - \mathcal{R}_n(\boldsymbol{w},b)) - \mathbb{E}\left[\sup_{(\boldsymbol{w},b)\in\mathcal{T}} (\mathcal{R}(\boldsymbol{w},b) - \mathcal{R}_n(\boldsymbol{w},b))\right]$$

$$\leq \sqrt{\frac{2L^2\log(1/\delta)}{n}} \sqrt{1 + \frac{1}{2} \left(\log\frac{1}{g(-|\theta_0|)} + \log\frac{1}{M_X^{\text{den}}(G\theta_1)}\right)}$$

$$\times \sqrt{5K_{\text{LS}}M_X^{\text{num}}(G\theta_1)M_X^{\text{num}}(2G\theta_1)R_{\boldsymbol{w}}^2 + 8g(+\theta_0)g(-\theta_0)M_X^{\text{num}}(G\theta_1)R_b^2}.$$

Lebesgue's dominated convergence theorem gives that $\lim_{\|t\|\to 0} M_X^{\text{num}}(t) = 1$ and $\lim_{\|t\|\to 0} M_X^{\text{den}}(t) = 1$. Therefore, θ_1 around zero achieves the same upper bounds for the residuals as the tensorization argument (Proposition 4) up to a numerical constant factor of 2 and a factor of $\sqrt{5(1+(1/2)\log(1/g(-|\theta_0|)))}$. The multiplicative numerical constants of Proposition 7 are better than those of Proposition 6 in the limit $\|\theta_1\|\to 0$ (here, for simplicity, we use rough upper bounds for both cases; however, the tightest bounds derived in our proofs yields the same conclusion). On the other hand, if $\|\theta_1\|$ grows, then $M_X^{\text{num}}(\theta_1)M_X^{\text{num}}(2\theta_1) = \omega(M_X^2(\theta_1))$ in general (we let G=1 for the logistic link); hence, Proposition 6 gives a better bound. Note that the multiplicative factor $\log(1/g(-|\theta_0|))$ for $g(+\theta_0)g(-\theta_0)$ must appear for a log-Sobolev inequality for Bernoulli distributions corresponding to Γ_Y ; see Boucheron et al. (2013).

4.3.4. Concentration under strong signal. We now examine the case with a strong signal such that $\|\theta_1\| \gg 1$. If $\|\theta_1\|$ is large, then the input of g inflate; $\langle X_i, \theta_1 \rangle$ takes very large or small values, and θ_0 should not affect the residual as long as they are of order $o(\|\theta_1\|)$. It is well-known that the empirical minimization problem (1) performs well (for example, see Kuchelmeister and van de Geer, 2024; Hsu and Mazumdar, 2024). Our result also shows that such a setting makes uniform generalization errors concentrate around their expectation sharply.

We define the following function:

$$\tilde{M}_{X}(t) := \frac{1}{\mathbb{E}\left[\exp\left(-|\langle X, t \rangle|\right)\right]}, \ t \in \mathbb{R}^{d}.$$

Note that $1/\tilde{M}_X(t) \to 0$ as $||t|| \to \infty$ in general (we notice that $\lim_{t\to\infty}(1/\tilde{M}_X(t))=0$ is equivalent to the statement that for any sequence $\{t_n\}_{n\in\mathbb{N}}$ with $||t_n|| \to \infty$, there exists a subsequence $\{t_n'\}_{n\in\mathbb{N}}\subset \{t_n\}_{n\in\mathbb{N}}$ such that $\lim_{n\to\infty}(1/\tilde{M}_X(t_n'))=0$; by the dominated convergence theorem, $\lim_{n\to\infty}(1/\tilde{M}_X(t_n'))=\mathbb{E}[\lim_{n\to\infty}\exp(-|\langle X,t_n'\rangle|)]$ as long as the limits exist; therefore, it is sufficient to show $1/|\langle X,t_n'\rangle|\to 0$ almost surely for some $\{t_n'\}\subset \{t_n\}$ for any $\{t_n\}$; since P_X is absolutely continuous with respect to the d-dimensional standard Gaussian measure, it is sufficient to consider $X\sim \mathcal{N}_d(\mathbf{0}_d,\mathbf{I}_d)$; $\langle X,t_n\rangle\sim \mathcal{N}_1(0,||t_n||^2)$, and the Borel–Cantelli lemma and condition $||t_n||\to\infty$ yield the conclusion). If $X\sim \mathcal{N}_d(\mathbf{0}_d,\Sigma)$ for some positive definite $\Sigma\in\mathbb{R}^{d\times d}$, then we obtain a concrete bound $1/\tilde{M}_X(t)\leq 1/\sqrt{2\pi}\langle t,\Sigma t\rangle$ (we use $1/\tilde{M}_X(t)=2\int_0^\infty\exp(-t)\exp(-t^2/(2\langle t,\Sigma t\rangle))/(2\pi\langle t,\Sigma t\rangle)^{d/2}\mathrm{d}t=2\exp(\langle t,\Sigma t\rangle/2)(1-\Phi(\sqrt{\langle t,\Sigma t\rangle}))$ and $1-\Phi(t)\leq (1/t)(1/\sqrt{2\pi})\exp(-t^2/2)$; see Proposition 2.1.2 of Vershynin, 2018). In addition, we use the following notation:

$$p_{\theta_1,\theta_0} := \mathbb{P}\left(\langle X, \theta_1 \rangle \geq |\theta_0|\right).$$

For example, if we consider $X \sim \mathcal{N}_d(\mathbf{0}_d, \Sigma)$, then $\mathbb{P}(\langle X, \theta_1 \rangle \geq |\theta_0|) = 1 - \Phi(|\theta_0|/\sqrt{\langle \theta_1, \Sigma \theta_1 \rangle})$.

Proposition 8 (concentration under strong signal and logistic models). Assume that the potential function U is even and $g(t) = 1/(1 + \exp(-t))$.

(i) Suppose that Assumption 1 holds. Then, for all $\delta \in (0,1]$, with probability at least $1-\delta$,

$$\sup_{(\boldsymbol{w},b)\in\mathcal{T}} (\mathcal{R}(\boldsymbol{w},b) - \mathcal{R}_n(\boldsymbol{w},b)) - \mathbb{E}\left[\sup_{(\boldsymbol{w},b)\in\mathcal{T}} (\mathcal{R}(\boldsymbol{w},b) - \mathcal{R}_n(\boldsymbol{w},b))\right]$$

$$\leq \sqrt{\frac{L^2\left[K_P\left(1 + e^{5|\theta_0|}/\tilde{M}_X(\theta_1) + \sqrt{e^{5|\theta_0|}/\tilde{M}_X(\theta_1)}\right)R_w^2 + \left(1 + \sqrt{e^{5|\theta_0|}/\tilde{M}_X(\theta_1)}\right)R_b^2\right] (\log(3/\delta))^2}{n}}.$$

(ii) Suppose that Assumption 2 holds. Then, for all $\delta \in (0,1]$, with probability at least $1-\delta$,

$$\begin{split} \sup_{(\boldsymbol{w},b)\in\mathcal{T}} & \left(\mathcal{R}(\boldsymbol{w},b) - \mathcal{R}_n(\boldsymbol{w},b)\right) - \mathbb{E}\left[\sup_{(\boldsymbol{w},b)\in\mathcal{T}} \left(\mathcal{R}(\boldsymbol{w},b) - \mathcal{R}_n(\boldsymbol{w},b)\right)\right] \\ \leq & \sqrt{\frac{2L^2\left(e + \log\left(1/p_{\theta_1,\theta_0}\right)\right)\left(K_{\mathrm{LS}}\left(5/4 + e^{5|\theta_0|}/\tilde{M}_X(\theta_1)\right)R_{\boldsymbol{w}}^2 + R_b^2\right)\log(1/\delta)}{n}}. \end{split}$$

The right-hand side in Proposition 8-(i) coincides that in Proposition 4-(i) in the limit $\|\theta_1\| \to \infty$. We also observe that the upper bound in Proposition 8-(ii) coincides with that of Proposition 4-(ii) up to a moderate numerical constant $(\sqrt{(e+\log(2))5/4} \le 2.07)$ if $p_{\theta_1,\theta_0} \approx 1/2$ and $1/\tilde{M}_X(\theta_1) \ll 1$. For example, if we assume $X_i \sim \mathcal{N}_d(\mathbf{0}_d, \Sigma)$, then $\langle \theta_1, \Sigma \theta_1 \rangle \gg |\theta_0|^2$ is a sufficient condition for them.

4.4. **Non-central input vectors.** We have discussed concentration inequalities based on the evenness of the potential function U. However, one can be interested in situations where the input vectors have the nonzero mean vector $\mu \neq 0$. To address this concern, let us consider the shifted dataset $\{(X_i + \mu, Y_i)\}_{i=1}^n$ for $\mu \in \mathbb{R}^d$. We can redefine the empirical risk function and conditional mass function of Y_i given $X_i + \mu$ as follows:

$$\begin{split} \mathcal{R}_{n}^{\mu}(\boldsymbol{w},b) &:= \frac{1}{n} \sum_{i=1}^{n} \ell \left(Y_{i} \left(\langle \boldsymbol{X}_{i} + \boldsymbol{\mu}, \boldsymbol{w} \rangle + b \right) \right) = \frac{1}{n} \sum_{i=1}^{n} \ell \left(\langle Y_{i} \boldsymbol{X}_{i}, \boldsymbol{w} \rangle + Y_{i} (b + \langle \boldsymbol{\mu}, \boldsymbol{w} \rangle) \right), \\ p_{\boldsymbol{Y} \mid \boldsymbol{X}}^{\mu}(\boldsymbol{y} | \boldsymbol{x}) &= g(\langle \boldsymbol{x} + \boldsymbol{\mu}, \boldsymbol{\theta}_{1} \rangle + \theta_{0}) = g(\langle \boldsymbol{x}, \boldsymbol{\theta}_{1} \rangle + \theta_{0} + \langle \boldsymbol{\mu}, \boldsymbol{\theta}_{1} \rangle). \end{split}$$

We can immediately obtain non-central versions of the results above using this reparameterization. The following corollary is an example using Proposition 5-(ii); other examples using other propositions can be shown immediately.

Corollary 9. Assume that the potential function U is even, g is non-decreasing, and $\log \circ g$ is G-Lipschitz. Furthermore, suppose that Assumption 1 holds true. Then, for any $\delta \in (0, 1]$, with probability $1 - \delta$,

$$\begin{split} \sup_{(w,b)\in\mathcal{T}} & (\mathcal{R}^{\mu}(w,b) - \mathcal{R}^{\mu}_{n}(w,b)) - \mathbb{E}\left[\sup_{(w,b)\in\mathcal{T}} (\mathcal{R}^{\mu}(w,b) - \mathcal{R}^{\mu}_{n}(w,b))\right] \\ \leq & \sqrt{\frac{L^{2}\left[K_{P}\left(e^{2G|\tilde{\theta}_{0}|} + \sqrt{e^{2G|\tilde{\theta}_{0}|} - 1}\right)R_{w}^{2} + (1 + \sqrt{e^{2G|\tilde{\theta}_{0}|} - 1})\left(\|\mu\|^{2}R_{w}^{2} + R_{b}^{2}\right)\right]\left(\log(3/\delta)\right)^{2}}{n}, \end{split}$$

where $\mathcal{R}^{\mu}(\mathbf{w},b) = \mathbb{E}[\mathcal{R}_{n}^{\mu}(\mathbf{w},b)]$ and $\tilde{\theta}_{0} = \theta_{0} + \langle \mu, \theta \rangle$

Note that a trivial modification of Proposition 2 derives

$$\mathbb{E}\left[\sup_{(\boldsymbol{w},b)\in\mathcal{T}}(\mathcal{R}^{\mu}(\boldsymbol{w},b)-\mathcal{R}^{\mu}_{n}(\boldsymbol{w},b))\right]\leq \frac{2L\left(R_{\boldsymbol{w}}\sqrt{\operatorname{tr}(\mathbb{E}[(\boldsymbol{X}_{i}+\boldsymbol{\mu})(\boldsymbol{X}_{i}+\boldsymbol{\mu})^{\top}])}+R_{b}\right)}{\sqrt{n}}.$$

The order of the upper bound in Corollary 9 is much milder than the order of this bound.

5. Applications to asymptotic analysis

In this section, we apply the concentration results of Section 4 to asymptotic regimes. While we consider asymptotics and a sequence of statistical experiments indexed by n, we omit the explicit dependence of L, R_w, R_b, K_P, K_{LS} , and other quantities and random variables on n for notational simplicity.

We now observe that the following almost sure convergences hold under some concise and reasonable conditions (for example, $|\theta_0| = O(1)$, L = O(1), and $(K_{LS}R_w^2 + R_b^2) \log n/n \to 0$ when we use Proposition 5-(ii)):

$$\lim_{n \to \infty} \left| \sup_{(\mathbf{w}, b) \in \mathcal{T}} (\mathcal{R}(\mathbf{w}, b) - \mathcal{R}_n(\mathbf{w}, b)) - \mathbb{E} \left[\sup_{(\mathbf{w}, b) \in \mathcal{T}} (\mathcal{R}(\mathbf{w}, b) - \mathcal{R}_n(\mathbf{w}, b)) \right] \right| = 0,$$

$$\lim_{n \to \infty} \left| \sup_{(\mathbf{w}, b) \in \mathcal{T}} (\mathcal{R}_n(\mathbf{w}, b) - \mathcal{R}(\mathbf{w}, b)) - \mathbb{E} \left[\sup_{(\mathbf{w}, b) \in \mathcal{T}} (\mathcal{R}_n(\mathbf{w}, b) - \mathcal{R}(\mathbf{w}, b)) \right] \right| = 0,$$
(6)

$$\lim_{n \to \infty} \left| \sup_{(\mathbf{w}, b) \in \mathcal{T}} (\mathcal{R}_n(\mathbf{w}, b) - \mathcal{R}(\mathbf{w}, b)) - \mathbb{E} \left[\sup_{(\mathbf{w}, b) \in \mathcal{T}} (\mathcal{R}_n(\mathbf{w}, b) - \mathcal{R}(\mathbf{w}, b)) \right] \right| = 0, \tag{6}$$

almost surely. This follows immediately from the Borel-Cantelli lemma and the fact that sign changes preserve Lipschitz continuity. The difference in signs does not matter because $-\ell$ is also L-Lipschitz if ℓ is.

5.1. Effective rank and uniform law of large numbers. If the expectation terms in Eqs. (5) and (6) tend to zero as $n \to \infty$, the next uniform law of large numbers follows as a direct corollary of the almost sure convergences (5) and (6):

$$\lim_{n \to \infty} \sup_{(\mathbf{w}, b) \in \mathcal{T}} |\mathcal{R}_n(\mathbf{w}, b) - \mathcal{R}(\mathbf{w}, b)| = 0.$$
 (7)

Proposition 2 derives a sufficient condition for convergence of the expectation terms by flipping the sign of ℓ ; if $L^2 R_w^2 \operatorname{tr}(\mathbb{E}[X_i X_i^{\top}])/n \to 0$ and $L^2 R_h^2/n \to 0$, then the uniform law (7) holds true. In particular, consider the case where L, R_w, R_b, K_{LS} , and $\|\mathbb{E}[X_i X_i^\top]\|_{op}$ are fixed in n and $d^* = \text{tr}(\mathbb{E}[X_i X_i^\top]) / \|\mathbb{E}[X_i X_i^\top]\|_{op}$ varies according to n. Then, the sufficient condition for the uniform law is $d^*/n \to 0$. This generalizes the classical condition $d/n \to 0$ by replacing the ambient dimension with the effective rank.

Remark 3. We explain the differences between our asymptotic and non-asymptotic results and those of Nakakita (2024) for clarity. Nakakita (2024) considers a uniform law only for the logistic loss function under the isoperimetry of the marginal distribution of X_i . Our results imply uniform laws for arbitrary Lipschitz loss functions under the isoperimetry of the joint distribution of (\mathbf{Z}_i, Y_i) . Hence, our results and that of Nakakita (2024) are complementary in the regime $d^*/n \to 0$, but differ in the form of concentration and the distributions they assume isoperimetry over. Moreover, our results can show the almost sure convergences (5) and (6) even if $d^*/n = \Theta(1)$, while Nakakita (2024) only yields the tightness of the uniform generalization error. This is because we consider concentration around means, and Nakakita (2024) deals with concentration around zero.

5.2. **Biased convergence in proportionally high-dimensional settings.** We also consider proportionally high-dimensional settings following the setup of prior studies such as Sur and Candès (2019) and Salehi et al. (2019). Let us set $X_i \sim^{\text{i.i.d.}} \mathcal{N}_d(\mathbf{0}_d, (1/d)\mathbf{I}_d)$ and $g = \sigma$ (logistic link); then $U(x) = (d/2)||x||^2$ and $K_{\text{LS}} = 1/d$ since U is d-convex. We set $R_w = \sqrt{d}R_1$ for some fixed $R_1 > 0$. This is our original setting, but it reflects the typical assumption that each element of the true value of the parameter is independent and identically distributed.

A Rademacher complexity bound for the expected uniform generalization error is of order $O(LR_1)$ (Proposition 2), and thus does not vanish asymptotically when $R_1 = \Omega(1)$. This result can be predicted by the result of Sur and Candès (2019); they show that the maximum likelihood estimator for logistic regression converges with bias in proportional high-dimensional settings. Since our setting for ℓ includes the logistic regression loss, the expected uniform generalization error should not converge to zero.

On the other hand, the residuals in Propositions 4–8 under Assumption 2 are of order $O(L(R_1 + R_b)\sqrt{\log(1/\delta)/n})$ (ignoring θ_0 and θ_1 for a unified treatment). Therefore, uniform generalization errors rapidly converge to their expectation in proportionally high-dimensional settings. This does not contradict existing results, which primarily characterize convergence of the maximum likelihood estimator to a biased limit in high dimensions. The point is that maximum likelihood estimation converges anyway. Our result is similar to it that the generalization error concentrates around its (possibly biased) expectation, even if the expectation itself does not vanish asymptotically.

Note that we can examine and characterize arbitrary Lipschitz functions of (\mathbf{Z}_i, Y_i) in proportionally high-dimensional settings because our proof strategy can be extended to other Lipschitz functions of them.

6. Concluding remarks

We have developed functional inequalities tailored to binary linear classification, and derived sharp concentration bounds under a variety of bias and signal regimes. The small numerical constants in our results allow for tight evaluation of uncertainty in binary linear classification tasks. Moreover, our proof strategy can be adapted to different problem setups. In particular, the idea to evaluate the isoperimetry of some problem-specific random vectors should be helpful in other supervised and unsupervised learning problems.

We refer to potential future research directions. As Johnson (2017) considers the Bakry-Émery condition for a discrete space \mathbb{Z}_+ , more tractable criteria for functional inequalities on hybrid continuous-discrete spaces such as $\mathbb{R}^d \times \{\pm 1\}$ and derivation of sufficient conditions generalizing Theorem 1 should be of interest. Extension to nonlinear classification problems (in particular, classification with neural networks) is also an important and topical problem.

ACKNOWLEDGEMENTS

This work is supported by JSPS KAKENHI Grant Number JP24K02904.

REFERENCES

Bach, F. (2024). Learning Theory from First Principles. MIT Press.

Bakry, D., Gentil, I., and Ledoux, M. (2014). *Analysis and Geometry of Markov Diffusion Operators*. Springer Science & Business Media.

Bardet, J.-B., Gozlan, N., Malrieu, F., and Zitt, P.-A. (2018). Functional inequalities for Gaussian convolutions of compactly supported measures: Explicit bounds and dimension dependence. *Bernoulli*, 24(1):333–353.

Bartlett, P. L. and Mendelson, S. (2002). Rademacher and Gaussian complexities: Risk bounds and structural results. *J. Mach. Learn. Res.*, 3(Nov):463–482.

- Bobkov, S. and Ledoux, M. (1997). Poincaré's inequalities and talagrand's concentration phenomenon for the exponential distribution. *Probab. Theory Related Fields*, 107:383–400.
- Boucheron, S., Lugosi, G., and Massart, P. (2013). *Concentration Inequalities: A Nonasymptotic Theory of Independence*. Oxford University Press.
- Candès, E. J. and Sur, P. (2020). The phase transition for the existence of the maximum likelihood estimate in high-dimensional logistic regression. *Ann. Statist.*, 48(1):27–42.
- Cattiaux, P. and Guillin, A. (2022). Functional inequalities for perturbed measures with applications to log-concave measures and to some Bayesian problems. *Bernoulli*, 28(4):2294–2321.
- Chen, H.-B., Chewi, S., and Niles-Weed, J. (2021). Dimension-free log-Sobolev inequalities for mixture distributions. *J. Funct. Anal.*, 281(11):109236.
- Chen, Y. (2021). An almost constant lower bound of the isoperimetric coefficient in the KLS conjecture. *Geom. Funct. Anal.*, 31:34–61.
- Hsu, D. and Mazumdar, A. (2024). On the sample complexity of parameter estimation in logistic regression with normal design. In *The Thirty Seventh Annual Conference on Learning Theory*, pages 2418–2437. PMLR.
- Jambulapati, A., Lee, Y. T., and Vempala, S. S. (2022). A slightly improved bound for the KLS constant. *arXiv* preprint *arXiv*:2208.11644.
- Johnson, O. (2017). A discrete log-Sobolev inequality under a Bakry-Émery type condition. *Ann. Inst. Henri Poincaré Probab. Stat.*, 53(4):1952–1970.
- Kannan, R., Lovász, L., and Simonovits, M. (1995). Isoperimetric problems for convex bodies and a localization lemma. *Discrete Comput. Geom.*, 13:541–559.
- Klartag, B. (2023). Logarithmic bounds for isoperimetry and slices of convex sets. Ars Inven. Anal.
- Klartag, B. and Lehec, J. (2022). Bourgain's slicing problem and KLS isoperimetry up to polylog. *Geom. Funct. Anal.*, 32(5):1134–1159.
- Kuchelmeister, F. and van de Geer, S. (2024). Finite sample rates for logistic regression with small noise or few samples. *Sankhya A*. Advance online publication.
- Ledoux, M. (1999). Concentration of measure and logarithmic Sobolev inequalities. *Séminaire de probabilités de Strasbourg*, 33:120–216.
- Ledoux, M. and Talagrand, M. (1991). *Probability in Banach Spaces: Isoperimetry and Processes*, volume 23. Springer Science & Business Media.
- Lee, Y. T. and Vempala, S. S. (2018). The Kannan-Lovász-Simonovits Conjecture. *arXiv preprint* arXiv:1807.03465.
- Lee, Y. T. and Vempala, S. S. (2024). Eldan's stochastic localization and the KLS conjecture: Isoperimetry, concentration and mixing. *Ann. of Math.* (2), 199(3):1043–1092.
- Liang, H. and Du, P. (2012). Maximum likelihood estimation in logistic regression models with a diverging number of covariates. *Electron. J. Stat.*, 6:1838–1846.
- Nakakita, S. (2024). Dimension-free uniform concentration bound for logistic regression. *arXiv preprint* arXiv:2405.18055.
- Salehi, F., Abbasi, E., and Hassibi, B. (2019). The impact of regularization on high-dimensional logistic regression. *Advances in Neural Information Processing Systems*, 32.
- Schlichting, A. (2019). Poincaré and log-Sobolev inequalities for mixtures. *Entropy*, 21(1):89.
- Sur, P. and Candès, E. J. (2019). A modern maximum-likelihood theory for high-dimensional logistic regression. *Proc. Natl. Acad. Sci. USA*, 116(29):14516–14525.
- Sur, P., Chen, Y., and Candès, E. J. (2019). The likelihood ratio test in high-dimensional logistic regression is asymptotically a rescaled chi-square. *Probab. Theory Related Fields*, 175:487–558.
- van de Geer, S. A. (2008). High-dimensional generalized linear models and the lasso. *Ann. Statist.*, 36(1):614–645.
- Vershynin, R. (2018). *High-Dimensional Probability: An Introduction with Applications in Data Science*. Cambridge University Press.

Wainwright, M. J. (2019). *High-Dimensional Statistics: A Non-Asymptotic Viewpoint*. Cambridge University Press.

APPENDIX A. PROOF OF THEOREM 1

We cite a technical result used in the proof of Theorem 1.

Lemma 10 (Chen et al., 2021). Let π and ρ be two probability measures. For any non-negative function f with $\mathbb{E}_{\pi}[f] < \infty$, We have

$$\mathbb{E}_{\pi} \left[f \log \left(\frac{\mathbb{E}_{\pi}[f]}{\mathbb{E}_{\rho}[f]} \right) \right] \leq \operatorname{Ent}_{\pi}(f) + \mathbb{E}_{\pi}[f] \log \left(1 + \chi^{2}(\pi \| \rho) \right).$$

Proof of Theorem 1. As the basis of our proof strategy, we employ the proof of Chen et al. (2021) partially and introduce a decoupling-like argument.

(Step 1: Poincaré inequality) We set a random variable Y', satisfying the following two conditions:

$$Y' \stackrel{\mathcal{L}}{=} Y, \ (\mathbf{Z}, Y) \perp Y'.$$

Note the following decomposition:

$$Var(f(\mathbf{Z}, Y)) = \mathbb{E}[Var(f(\mathbf{Z}, Y)|Y)] + Var(\mathbb{E}[f(\mathbf{Z}, Y)|Y]).$$

The first term on the right-hand side is bounded as

$$\mathbb{E}[\operatorname{Var}(f(\mathbf{Z},Y)|Y)] \leq \mathbb{E}[K_{\mathbf{P}}\Gamma_{\mathbf{Z}}(f)(\mathbf{Z},Y)].$$

Regarding the second term, by definition, we have

$$\operatorname{Var}(\mathbb{E}[f(\mathbf{Z}, Y)|Y]) = \mathbb{E}\left[\left|\mathbb{E}[f(\mathbf{Z}, Y)|Y] - \mathbb{E}[f(\mathbf{Z}, Y)]\right|^{2}\right].$$

We have for each $y \in \{\pm 1\}$,

$$\mathbb{E}[f(\mathbf{Z},Y)|Y=y] - \mathbb{E}[f(\mathbf{Z},Y)] = (\mathbb{E}_{\mathbf{Z}|Y}[f(\mathbf{Z},Y)|Y=y] - \mathbb{E}_{\mathbf{Z}}[f(\mathbf{Z},y)]) + \mathbb{E}_{\mathbf{Z}Y}[f(\mathbf{Z},y) - f(\mathbf{Z},Y)].$$

Therefore, for any $c, c^* \in [1, \infty]$ with $1/c + 1/c^* = 1$,

$$\operatorname{Var}(\mathbb{E}[f(\boldsymbol{Z},Y)|Y]) \leq c \, \mathbb{E}_{Y'} \left[\left| \mathbb{E}_{\boldsymbol{Z}|Y}[f(\boldsymbol{Z},Y)|Y=Y'] - \mathbb{E}_{\boldsymbol{Z}}[f(\boldsymbol{Z},Y')] \right|^2 \right]$$

$$+ c^* \, \mathbb{E}_{Y'} \left[\left| \mathbb{E}_{\boldsymbol{Z}Y}[f(\boldsymbol{Z},Y) - f(\boldsymbol{Z},Y')] \right|^2 \right].$$
Term (ii): How \boldsymbol{Z} is informative to \boldsymbol{Y} .

For the term (i), since for each $y \in \{\pm 1\}$ and $c \in \mathbb{R}$,

$$\mathbb{E}_{\mathbf{Z}|Y}[f(\mathbf{Z},Y)|Y=y] - \mathbb{E}_{\mathbf{Z}}[f(\mathbf{Z},y)] = \int_{\mathbb{R}^d} f(z,y) \left(1 - \frac{P_{\mathbf{Z}}(\mathrm{d}z)}{P_{\mathbf{Z}|Y}(\mathrm{d}z|y)}\right) P_{\mathbf{Z}|Y}(\mathrm{d}z|y)$$

$$= \int_{\mathbb{R}^d} (f(z,y) - c) \left(1 - \frac{P_{\mathbf{Z}}(\mathrm{d}z)}{P_{\mathbf{Z}|Y}(\mathrm{d}z|y)}\right) P_{\mathbf{Z}|Y}(\mathrm{d}z|y),$$

the Cauchy-Schwarz inequality yields

$$\begin{split} \mathbb{E}_{Y'}\left[\left|\mathbb{E}_{\mathbf{Z}|Y}[f(\mathbf{Z},Y)|Y=Y'] - \mathbb{E}_{\mathbf{Z}}[f(\mathbf{Z},Y')]\right|^2\right] &\leq \mathbb{E}_{Y'}\left[\operatorname{Var}(f(\mathbf{Z},Y)|Y=Y')\chi^2(P_{\mathbf{Z}}(\cdot)||P_{\mathbf{Z}|Y}(\cdot|Y'))\right] \\ &\leq K_{\chi^2}\mathbb{E}\left[\operatorname{Var}(f(\mathbf{Z},Y)|Y)\right] \\ &\leq K_{\chi^2}K_{\mathbf{P}}\mathbb{E}\left[\Gamma_{\mathbf{Z}}(f)(\mathbf{Z},Y)\right]. \end{split}$$

For the term (ii), we have

$$\begin{split} &\mathbb{E}_{Y'}\left[\left|\mathbb{E}_{ZY}[f(Z,Y)-f(Z,Y')]\right|^{2}\right] \\ &= \sum_{y'\in\{\pm 1\}} p_{Y}(y') \left(\sum_{y\in\{\pm 1\}} \int_{\mathbb{R}^{d}} (f(z,y)-f(z,y')) P_{ZY}(\mathrm{d}z\times\{y\})\right)^{2} \\ &= \sum_{y'\in\{\pm 1\}} p_{Y}(y') \left(\int_{\mathbb{R}^{d}} (f(z,-y')-f(z,y')) P_{ZY}(\mathrm{d}z\times\{-y'\})\right)^{2} \\ &= \sum_{y'\in\{\pm 1\}} p_{Y}(y') \left(\int_{\mathbb{R}^{d}} 2\nabla_{y} f(z,y') P_{ZY}(\mathrm{d}z\times\{-y'\})\right)^{2} \\ &= \sum_{y'\in\{\pm 1\}} p_{Y}(y') p_{Y}(-y')^{2} \left(\frac{1}{p_{Y}(-y')} \int_{\mathbb{R}^{d}} 2\nabla_{y} f(z,y') P_{ZY}(\mathrm{d}z\times\{-y'\})\right)^{2} \\ &= p_{Y}(+1) p_{Y}(-1) \sum_{y'\in\{\pm 1\}} p_{Y}(-y') \left(\frac{1}{p_{Y}(-y')} \int_{\mathbb{R}^{d}} 2\nabla_{y} f(z,y') P_{ZY}(\mathrm{d}z\times\{-y'\})\right)^{2} \\ &\leq p_{Y}(+1) p_{Y}(-1) \sum_{y'\in\{\pm 1\}} p_{Y}(-y') \frac{1}{p_{Y}(-y')} \int_{\mathbb{R}^{d}} 4(\nabla_{y} f(z,y'))^{2} P_{ZY}(\mathrm{d}z\times\{-y'\}) \\ &= 4 p_{Y}(+1) p_{Y}(-1) \mathbb{E}\left[(\nabla_{y} f(Z,Y))^{2}\right], \end{split}$$

where the Cauchy-Schwarz inequality derives the inequality above. Therefore, we obtain that

$$Var(f(\mathbf{Z}, Y)) = \mathbb{E}[Var(f(\mathbf{Z}, Y)|Y)] + Var(\mathbb{E}[f(\mathbf{Z}, Y)|Y])$$

$$\leq K_{P} \left(1 + cK_{\chi^{2}}\right) \mathbb{E}\left[\Gamma_{\mathbf{Z}}(f)(\mathbf{Z}, Y)\right] + c^{*}K_{V}\mathbb{E}\left[\left(\nabla_{y} f(\mathbf{Z}, Y)\right)^{2}\right]$$

$$\leq \mathbb{E}\left[\Gamma_{P}(f)(\mathbf{Z}, Y)\right].$$

(Step 2: log-Sobolev inequality) Note the following decomposition:

$$\operatorname{Ent}(f^2(\mathbf{Z},Y)) = \mathbb{E}\left[\operatorname{Ent}_{\mathbf{Z}|Y}(f^2(\mathbf{Z},Y))\right] + \operatorname{Ent}_Y\left(\mathbb{E}_{\mathbf{Z}|Y}[f^2(\mathbf{Z},Y)|Y]\right).$$

The first term can be bounded as the proof of the Poincaré inequality above. We analyse the second term on the right-hand side in detail. We have

$$\begin{split} &\operatorname{Ent}_{Y}\left(\mathbb{E}_{\mathbf{Z}|Y}[f^{2}(\mathbf{Z},Y)|Y]\right) \\ &= \mathbb{E}_{Y}\left[\mathbb{E}_{\mathbf{Z}|Y}[f^{2}(\mathbf{Z},Y)|Y]\log\frac{\mathbb{E}_{\mathbf{Z}|Y}[f^{2}(\mathbf{Z},Y)|Y]}{\mathbb{E}_{\mathbf{Z}Y}[f^{2}(\mathbf{Z},Y)]}\right] \\ &\leq \mathbb{E}_{Y}\left[\operatorname{Ent}_{\mathbf{Z}|Y}(f^{2}(\mathbf{Z},Y))\right] + \mathbb{E}_{Y}\left[\mathbb{E}_{\mathbf{Z}|Y}[f^{2}(\mathbf{Z},Y)|Y]\log\left(1 + \chi^{2}\left(P_{\mathbf{Z}|Y}(\cdot|y)\delta_{y}(\cdot)\|P_{\mathbf{Z}Y}(\cdot)\right)\right|_{y=Y}\right)\right] \\ &= \mathbb{E}_{Y}\left[\operatorname{Ent}_{\mathbf{Z}|Y}(f^{2}(\mathbf{Z},Y))\right] + \mathbb{E}_{Y}\left[\mathbb{E}_{\mathbf{Z}|Y}[f^{2}(\mathbf{Z},Y)|Y]\log\frac{1}{p_{Y}(Y)}\right] \\ &\leq \mathbb{E}_{Y}\left[\operatorname{Ent}_{\mathbf{Z}|Y}(f^{2}(\mathbf{Z},Y))\right] + \log K_{\mathbf{U}}\mathbb{E}[f^{2}], \end{split}$$

where the first inequality holds via Lemma 10 with letting $\pi = P_{\mathbf{Z}|Y}(\cdot|y)\delta_y(\cdot)$ and $\rho = P_{\mathbf{Z}Y}(\cdot)$, and the second identity is given as follows: for each $y \in \{\pm 1\}$,

$$1 + \chi^{2} \left(P_{\mathbf{Z}|Y}(\cdot|y) \delta_{y}(\cdot) || P_{\mathbf{Z}Y}(\cdot) \right) = \int_{\mathbb{R}^{d}} \frac{P_{\mathbf{Z}|Y}(\mathrm{d}z|y)}{P_{\mathbf{Z}Y}(\mathrm{d}z \times \{y\})} P(\mathrm{d}z|y)$$

$$= \frac{1}{p_{Y}(y)^{2}} \int_{\mathbb{R}^{d}} \frac{P_{\mathbf{Z}Y}(\mathrm{d}z \times \{y\})}{P_{\mathbf{Z}Y}(\mathrm{d}z \times \{y\})} P_{\mathbf{Z}Y}(\mathrm{d}z \times \{y\})$$

$$= \frac{1}{p_{Y}(y)}.$$

In total, we have

$$\operatorname{Ent}(f^{2}(\mathbf{Z}, Y)) \leq 2\mathbb{E}\left[\operatorname{Ent}_{\mathbf{Z}|Y}(f^{2}(\mathbf{Z}, Y))\right] + \log K_{\mathbf{U}}\mathbb{E}[f^{2}].$$

By letting $\hat{f} = f - \mathbb{E}f$, we adapt the argument of Section 5.1.2 of Bakry et al. (2014) for deficit log-Sobolev inequalities and yield

$$\operatorname{Ent}(f^{2}) \leq \operatorname{Ent}(\hat{f}^{2}) + 2\operatorname{Var}(f) \leq 2\mathbb{E}\left[\operatorname{Ent}_{\mathbf{Z}|Y}(\hat{f}^{2})\right] + \log K_{\mathbf{U}}\mathbb{E}[\hat{f}^{2}] + 2\operatorname{Var}(f)$$

$$\leq 2\left(K_{\mathrm{LS}}\mathbb{E}\left[\Gamma_{\mathbf{Z}}(f)\right] + \left(1 + \log K_{\mathbf{U}}\right)\operatorname{Var}(f)\right)$$

$$\leq 2\left(2K_{\mathrm{LS}}\mathbb{E}\left[\Gamma_{\mathbf{Z}}(f)\right] + \left(1 + \frac{\log K_{\mathbf{U}}}{2}\right)\mathbb{E}\left[\Gamma_{\mathbf{P}}(f)\right]\right).$$

This is the desired conclusion.

Appendix B. Proofs of the results in Section 4

We introduce exponential concentration and Gaussian concentration for Lipschitz maps under a Poincaré inequality and log-Sobolev inequality respectively; we restrict our arguments to the state space $(\mathbb{R}^d \times \{\pm 1\})^n$, but this argument is used in much broader settings (see Bakry et al., 2014).

Lemma 11 (exponential concentration, Bobkov and Ledoux, 1997; Bakry et al., 2014). Let P be a probability measure on a state space $(\mathbb{R}^d \times \{\pm 1\})^n$ such that the following Poincaé inequality holds true: for some carré du champ operator Γ and a positive constant C > 0 (Poincaé constant of P), for any $f \in \mathcal{A}_n$,

$$\operatorname{Var}_{P}(f) \leq C\mathbb{E}_{P}\left[\Gamma(f)\right].$$

Then, for any $f \in \mathcal{A}_n$ with $\|\Gamma(f)\|_{L^{\infty}} < \infty$ and $t \ge 0$, we have

$$P(f - \mathbb{E}_P[f] \ge t) \le 3 \exp\left(-\frac{t}{\sqrt{C} \|\Gamma(f)\|_{L^{\infty}}}\right).$$

Lemma 12 (Gaussian concentration via Herbst's argument, Ledoux, 1999; Bakry et al., 2014). Let P be a probability measure on a state space $(\mathbb{R}^d \times \{\pm 1\})^n$ such that the following log-Sobolev inequality holds true: for some carré du champ operator Γ and a positive constant C > 0 (log-Sobolev constant of P), for any $f \in \mathcal{A}_n$,

$$\operatorname{Ent}_P(f^2) \leq 2C\mathbb{E}_P\left[\Gamma(f)\right].$$

Then, for any $f \in \mathcal{A}_n$ with $\|\Gamma(f)\|_{L^{\infty}} < \infty$ and $t \geq 0$, we have

$$P(f - \mathbb{E}_P[f] \ge t) \le \exp\left(-\frac{t^2}{2C\|\Gamma(f)\|_{L^{\infty}}^2}\right).$$

We also introduce tensorization under independence.

Lemma 13 (tensorization, Propositions 4.3.1 and 5.2.7 of Bakry et al., 2014). Let P_i (i = 1, ..., n) be a probability measure on a state space $\mathbb{R}^d \times \{\pm 1\}$ and suppose that every P_i satisfies a Poincaré inequality (resp. log-Sobolev inequality) for constant C > 0 with a carré du champ operator Γ_i . Then, $\bigotimes_{i=1}^n P_i$ on $(\mathbb{R}^d \times \{\pm 1\})^n$ satisfies a Poincaré inequality (resp. log-Sobolev inequality) for constant C with the carré du champ operator $\Gamma = \bigoplus_{i=1}^n \Gamma_i$.

B.1. **Proof of the result in Section 4.1.** Though the proof of Proposition 2 is a very standard one, we present it for readers' convenience.

Proof of Proposition 2. Proposition 4.2 of Bach (2024) (symmetrization) gives

$$\mathbb{E}\left[\sup_{(\boldsymbol{w},b)\in\mathcal{T}}(\mathcal{R}(\boldsymbol{w},b)-\mathcal{R}_n(\boldsymbol{w},b))\right]\leq 2\mathbb{E}\left[\sup_{(\boldsymbol{w},b)\in\mathcal{T}}\frac{1}{n}\sum_{i=1}^n\varepsilon_i\ell(Y_i(\langle X_i,\boldsymbol{w}\rangle+b))\right],$$

where $\varepsilon_i \sim^{\text{i.i.d.}} \text{Unif}(\{\pm 1\})$ independent of $\{X_i, Y_i\}_{i=1}^n$. Proposition 4.3 of Bach (2024) (contraction) yields

$$\mathbb{E}\left[\sup_{(\boldsymbol{w},b)\in\mathcal{T}}\frac{1}{n}\sum_{i=1}^{n}\varepsilon_{i}\ell(Y_{i}(\langle\boldsymbol{X}_{i},\boldsymbol{w}\rangle+b))\right]\leq L\mathbb{E}\left[\sup_{(\boldsymbol{w},b)\in\mathcal{T}}\frac{1}{n}\sum_{i=1}^{n}\varepsilon_{i}(Y_{i}(\langle\boldsymbol{X}_{i},\boldsymbol{w}\rangle+b))\right].$$

Since

$$\mathbb{E}\left[\sup_{(\boldsymbol{w},b)\in\mathcal{T}}\frac{1}{n}\sum_{i=1}^{n}\varepsilon_{i}(Y_{i}(\langle\boldsymbol{X}_{i},\boldsymbol{w}\rangle+b))\right] = \frac{1}{n}\mathbb{E}\left[R_{\boldsymbol{w}}\left\|\sum_{i=1}^{n}\varepsilon_{i}Y_{i}\boldsymbol{X}_{i}\right\| + R_{b}\left|\sum_{i=1}^{n}\varepsilon_{i}Y_{i}\right|\right]$$

$$= \frac{1}{n}\left(R_{\boldsymbol{w}}\mathbb{E}\left[\left\|\sum_{i=1}^{n}\varepsilon_{i}Y_{i}\boldsymbol{X}_{i}\right\|\right] + R_{b}\mathbb{E}\left[\left|\sum_{i=1}^{n}\varepsilon_{i}Y_{i}\right|\right]\right)$$

$$\leq \frac{1}{n}\left(R_{\boldsymbol{w}}\mathbb{E}\left[\left\|\sum_{i=1}^{n}\varepsilon_{i}Y_{i}\boldsymbol{X}_{i}\right\|^{2}\right]^{1/2} + R_{b}\mathbb{E}\left[\left(\sum_{i=1}^{n}\varepsilon_{i}Y_{i}\right)^{2}\right]^{1/2}\right)$$

$$= \frac{1}{n}\left(R_{\boldsymbol{w}}\mathbb{E}\left[\sum_{i=1}^{n}\|Y_{i}\boldsymbol{X}_{i}\|^{2}\right]^{1/2} + R_{b}\mathbb{E}\left[\sum_{i=1}^{n}Y_{i}^{2}\right]^{1/2}\right)$$

$$= \frac{1}{n}\left(R_{\boldsymbol{w}}\sqrt{n}\mathbb{E}\left[\|\boldsymbol{X}_{i}\|^{2}\right]^{1/2} + R_{b}\sqrt{n}\right),$$

we obtain the desired result.

B.2. Proofs of the results in Section 4.2.

Proof of Proposition 3. Let $P_{X|Y}(dx|y)$ be the conditional measure of X_i given $Y_i = y \in \{\pm 1\}$. For each $y \in \{\pm 1\}$,

$$P_{X|Y}(\mathrm{d}x|y) = \frac{\exp\left(-U\left(x\right) - \left(-\log(g(y\langle x, \theta_1\rangle)\right)\mathrm{d}x}{\int_{\mathbb{R}^d} \exp\left(-U\left(x\right) - \left(-\log(g(y\langle x, \theta_1\rangle)\right)\mathrm{d}x}.$$

We consider $P_{Z|Y}$, the conditional measure of Z_i given $Y_i = y \in \{\pm 1\}$. We have

$$P_{Z|Y}(\mathrm{d}z|y) = \frac{\exp\left(-U\left(yz\right) - \left(-\log\left(g\left(\langle z, \theta_1 \rangle\right)\right) \mathrm{d}z\right)}{\int_{\mathbb{R}^d} \exp\left(-U\left(yz\right) - \left(-\log\left(g\left(\langle z, \theta_1 \rangle\right)\right) \mathrm{d}z\right)} = \frac{\exp\left(-U\left(z\right) - \left(-\log\left(g\left(\langle z, \theta_1 \rangle\right)\right) \mathrm{d}z\right)}{\int_{\mathbb{R}^d} \exp\left(-U\left(z\right) - \left(-\log\left(g\left(\langle z, \theta_1 \rangle\right)\right) \mathrm{d}z\right)}$$

where the evenness of U gives the second identity. Hence, \mathbf{Z}_i is independent of Y_i .

Proof of Proposition 4. We only prove the statement (ii) since (i) can be shown by Lemma 11 and a parallel manner.

The conditional distribution of Z_i given Y_i is equal to the marginal distribution of Z_i due to Proposition 3, and $Y_i \sim \text{Unif}(\{\pm 1\})$ holds. Tensorization of log-Sobolev inequalities (Lemma 13) and Assumption 2 yield the following log-Sobolev inequality: for all $f \in \mathcal{A}_n$,

$$\operatorname{Ent}(f^2) \le 2\mathbb{E}\left[\sum_{i=1}^n \left(K_{\operatorname{LS}}\Gamma_{\mathbf{Z}_i}(f) + \Gamma_{Y_i}(f)\right)\right] =: 2\mathbb{E}[\Gamma_{\operatorname{i}}(f)].$$

What we need is an almost sure upper bound on Γ_i , owing to Lemma 12. We easily see that for arbitrary $z_i, z_i' \in \mathbb{R}^d$ and $y, y' \in \{\pm 1\}$,

$$\sup_{(\boldsymbol{w},b)\in\mathcal{T}} (\mathcal{R}(\boldsymbol{w},b) - \ell(\langle z_i,\boldsymbol{w}\rangle + y_ib)) - \sup_{(\boldsymbol{w},b)\in\mathcal{T}} (\mathcal{R}(\boldsymbol{w},b) - \ell(\langle z_i',\boldsymbol{w}\rangle + y_ib)) \leq LR_{\boldsymbol{w}} \|\boldsymbol{z} - \boldsymbol{z}'\|,$$

$$\sup_{(\boldsymbol{w},b)\in\mathcal{T}} (\mathcal{R}(\boldsymbol{w},b) - \ell(\langle z_i,\boldsymbol{w}\rangle + y_ib)) - \sup_{(\boldsymbol{w},b)\in\mathcal{T}} (\mathcal{R}(\boldsymbol{w},b) - \ell(\langle z_i,\boldsymbol{w}\rangle + y_i'b)) \leq 2LR_{\boldsymbol{w}},$$

and

$$\Gamma_{\mathbf{Z}_{i}}\left(\sup_{(\mathbf{w},b)\in\mathcal{T}}\left(\mathcal{R}(\mathbf{w},b)-\frac{1}{n}\sum_{i=1}^{n}\ell(\langle z_{i},\mathbf{w}\rangle+y_{i}b)\right)\right)\leq\frac{L^{2}R_{\mathbf{w}}^{2}}{n^{2}},\tag{8}$$

$$\Gamma_{Y_i} \left(\sup_{(\boldsymbol{w}, b) \in \mathcal{T}} \left(\mathcal{R}(\boldsymbol{w}, b) - \frac{1}{n} \sum_{i=1}^n \ell(\langle \boldsymbol{z}_i, \boldsymbol{w} \rangle + y_i b) \right) \right) \le \frac{L^2 R_b^2}{n^2}. \tag{9}$$

Therefore,

$$\Gamma_{i}\left(\sup_{(\boldsymbol{w},b)\in\mathcal{T}}(\mathcal{R}(\boldsymbol{w},b))-\frac{1}{n}\sum_{i=1}^{n}\ell(\langle z_{i},\boldsymbol{w}\rangle+y_{i}b)\right)\leq\frac{L^{2}}{n}\left(K_{LS}R_{\boldsymbol{w}}^{2}+R_{b}^{2}\right).$$

Hence, we obtain the desired conclusion by Lemma 12.

B.3. **Proofs of the results in Section 4.3.** Let us define the normalizing constants of conditional measures as follows:

$$\mathcal{Z}_{+\theta_{0}} := \int_{\mathbb{R}^{d}} g(\langle z, \theta_{1} \rangle + \theta_{0}) \exp(-U(z)) dz, \ \mathcal{Z}_{-\theta_{0}} := \int_{\mathbb{R}^{d}} g(\langle z, \theta_{1} \rangle - \theta_{0}) \exp(-U(z)) dz.$$

Note that $\mathcal{Z}_{+\theta_0} + \mathcal{Z}_{-\theta_0} = \mathcal{Z}$ under the evenness of the potential function U.

Proof of Proposition 5. We only prove (ii) since the proof of (i) is analogous and more straightforward (we use Theorem 1 with $c = 1 + 1/\sqrt{e^{2G|\theta_0|} - 1}$ and $c^* = 1 + \sqrt{e^{2G|\theta_0|} - 1}$).

We use that $K_P \le K_{LS}$. Therefore, we need estimates for the following constants:

$$\begin{split} K_{\chi^2} &:= \max_{y \in \{\pm 1\}} \chi^2 \left(P_{\mathbf{Z}}(\cdot) \| P_{\mathbf{Z}|Y}(\cdot|y) \right), \\ K_{V} &:= 4p_Y(+1)p_Y(-1), \\ K_{U} &:= \max \left\{ \frac{1}{p_Y(+1)}, \frac{1}{p_Y(-1)} \right\}. \end{split}$$

The assumption yields that for all $\theta_0 \in \mathbb{R}$ and $t \in \mathbb{R}$,

$$|\log(g(t+\theta_0)) - \log(g(t-\theta_0))| \le 2G|\theta_0|.$$
 (10)

Let us assume $\theta_0 \ge 0$ without loss of generality. Equation (10) gives $g(t + \theta_0)/g(t - \theta_0) \le e^{2G\theta_0}$, which is frequently used in the subsequent discussions. Note that $\mathcal{Z}_{+\theta_0}/\mathcal{Z}_{-\theta_0} \in [1, e^{2G\theta_0}]$ for any $\theta_0 \ge 0$ since the inequality (10) yields

$$\mathcal{Z}_{+\theta_0} = \int_{\mathbb{R}^d} \exp\left(-U(z) + \log(g(\langle z, \theta_1 \rangle + \theta_0))\right) dz$$

$$= \int_{\mathbb{R}^d} \exp\left(-U(z) + \log(g(\langle z, \theta_1 \rangle + \theta_0)) - \log(g(\langle z, \theta_1 \rangle - \theta_0)) + \log(g(\langle z, \theta_1 \rangle - \theta_0))\right) dz$$

$$\leq \int_{\mathbb{R}^d} \exp\left(-U(z) + 2G\theta_0 + \log(g(\langle z, \theta_1 \rangle - \theta_0))\right) dz$$

$$= e^{2G\theta_0} \mathcal{Z}_{-\theta_0},$$

and the monotonicty of g gives $\mathcal{Z}_{+\theta_0} \geq \mathcal{Z}_{-\theta_0}$. Similarly, $\mathcal{Z}_{-\theta_0}/\mathcal{Z}_{+\theta_0} \in [e^{-2G\theta_0}, 1]$ holds.

Since $g(t - \theta_0)/g(t + \theta_0) \le 1$, we obtain

$$\chi^{2} \left(P_{\mathbf{Z}}(\cdot) \| P_{\mathbf{Z}|Y}(\cdot | + 1) \right) = \int_{\mathbb{R}^{d}} \frac{Z_{+\theta_{0}} \sum_{y' \in \{\pm 1\}} g(\langle z, \theta_{1} \rangle + y' \theta_{0})}{Zg(\langle z, \theta_{1} \rangle + \theta_{0})} P_{\mathbf{Z}} (dz) - 1$$

$$\leq 2 \frac{Z_{+\theta_{0}}}{Z} - 1 = 2 \frac{Z_{+\theta_{0}}}{Z_{+\theta_{0}} + Z_{-\theta_{0}}} - 1 = 2 \frac{Z_{+\theta_{0}}/Z_{-\theta_{0}}}{1 + Z_{+\theta_{0}}/Z_{-\theta_{0}}} - 1$$

$$\leq Z_{+\theta_{0}}/Z_{-\theta_{0}} - 1 \leq e^{2G\theta_{0}} - 1.$$

In addition, the fact that $Z_{-\theta}/Z \leq 1/2$ along with $g(t+\theta_0)/g(t-\theta_0) \leq e^{2G\theta_0}$ gives

$$\chi^{2}\left(P_{\mathbf{Z}}(\cdot)||P_{\mathbf{Z}|Y}(\cdot|-1)\right) = \int_{\mathbb{R}^{d}} \frac{\mathcal{Z}_{-\theta_{0}}\sum_{y'\in\{\pm 1\}}g(\langle z,\theta_{1}\rangle + y'\theta_{0})}{\mathcal{Z}g(\langle z,\theta_{1}\rangle - \theta_{0})} P_{\mathbf{Z}}(dz) - 1$$

$$\leq \frac{\mathcal{Z}_{-\theta_{0}}}{\mathcal{Z}} \int_{\mathbb{R}^{d}} \left(1 + \frac{g(\langle z,\theta_{1}\rangle + \theta_{0})}{g(\langle z,\theta_{1}\rangle - \theta_{0})}\right) P_{\mathbf{Z}}(dz) - 1$$

$$= \frac{1}{2}(1 + e^{2G\theta_{0}}) - 1 \leq e^{2G\theta_{0}} - 1.$$

Therefore, we obtain

$$K_{\chi^2} = \max_{y \in \{\pm 1\}} \chi^2 \left(P_{\mathbf{Z}}(\cdot) || P_{\mathbf{Z}|Y}(\cdot|y) \right) \le e^{2G\theta_0} - 1.$$

The remaining constants can be evaluated as follows:

$$K_{V} = \frac{4Z_{+\theta_{0}}Z_{-\theta_{0}}}{(Z_{+\theta_{0}} + Z_{-\theta_{0}})^{2}} \le 1, K_{U} = \frac{Z_{+\theta_{0}} + Z_{-\theta_{0}}}{Z_{-\theta_{0}}} \le 1 + e^{2G\theta_{0}}.$$

Hence, we obtain the following log-Sobolev inequality by Theorem 1 ($c = c^* = 2$) and Lemma 13: for all $f \in \mathcal{A}_n$,

$$\operatorname{Ent}(f^2((\mathbf{Z}_i, Y_i)_{i=1}^n)) \le 2\mathbb{E}[\Gamma_{ii}(f)((\mathbf{Z}_i, Y_i)_{i=1}^n)],$$

where

$$\begin{split} \Gamma_{ii}(f) &:= \left(\left(1 + \frac{1}{2} \log \left(1 + e^{2G\theta_0} \right) \right) K_{LS} \left(1 + 2 \left(e^{2G\theta_0} - 1 \right) \right) + 2K_{LS} \right) \sum_{i=1}^{n} \Gamma_{\mathbf{Z}_i}(f) \\ &+ 2 \left(1 + \frac{1}{2} \log \left(1 + e^{2G\theta_0} \right) \right) \sum_{i=1}^{n} \Gamma_{\mathbf{Y}_i}(f) \\ &\leq K_{LS} \left(3 + 2G\theta_0 \right) \left(\frac{1}{2} + e^{2G\theta_0} \right) \sum_{i=1}^{n} \Gamma_{\mathbf{Z}_i}(f) + (3 + 2G\theta_0) \sum_{i=1}^{n} \Gamma_{\mathbf{Y}_i}(f); \end{split}$$

here, we use that $2 + \log(1 + e^{2G\theta_0}) \le 3 + 2G\theta_0$, and

$$\left(1+\frac{1}{2}\log\left(1+e^{2G\theta_0}\right)\right)\left(1+2\left(e^{2G\theta_0}-1\right)\right)+2\leq \left(1+\frac{1}{2}\log\left(1+e^{2G\theta_0}\right)\right)\left(1+2e^{2G\theta_0}\right).$$

Therefore, using Equations (8) and (9) and Lemma 12, we obtain that for all $\delta \in (0, 1]$, with probability at least $1 - \delta$,

$$\sup_{(\boldsymbol{w},b)\in\mathcal{T}} (\mathcal{R}(\boldsymbol{w},b) - \mathcal{R}_n(\boldsymbol{w},b)) - \mathbb{E}\left[\sup_{(\boldsymbol{w},b)\in\mathcal{T}} (\mathcal{R}(\boldsymbol{w},b) - \mathcal{R}_n(\boldsymbol{w},b))\right]$$

$$\leq \sqrt{\frac{2L^2(3 + 2G\theta_0)\left[K_{LS}\left(\frac{1}{2} + e^{2G\theta_0}\right)R_w^2 + R_b^2\right]\log(1/\delta)}{n}}.$$

Hence, we obtain the desired conclusion.

Proof of Proposition 6. We only prove (ii) since we can prove (i) in a similar and simpler manner.

Again, we consider $\theta_0 > 0$ and focus on estimate for K_{χ^2} , K_V , K_U . We obtain the following bound for $\mathcal{Z}_{-\theta_0}$:

$$\mathcal{Z}_{-\theta_0} \leq \int_{\mathbb{R}^d} \exp(\langle z, \theta_1 \rangle - \theta_0) \exp(-U(z)) dz = e^{-\theta_0} \mathcal{Z} M_X(\theta_1),$$

$$\mathcal{Z}_{-\theta_0} = \int_{\mathbb{R}^d} \frac{1}{1 + \exp(-\langle z, \theta_1 \rangle + \theta_0)} \exp(-U(z)) dz$$

$$\geq \int_{\mathbb{R}^d} \frac{1}{e^{\theta_0} + \exp(-\langle z, \theta_1 \rangle + \theta_0)} \exp(-U(z)) dz$$

$$\geq e^{-\theta_0} \int_{\mathbb{R}^d} \frac{1}{1 + \exp(-\langle z, \theta_1 \rangle)} \exp(-U(z)) dz = e^{-\theta_0} \frac{\mathcal{Z}}{2},$$

where we use $1/(1 + \exp(-t)) \le \exp(t)$ for the upper bound. Since $\mathcal{Z}_{+\theta_0} \le \mathcal{Z}$,

$$\chi^{2}\left(P_{\mathbf{Z}}(\cdot)||P_{\mathbf{Z}|Y}(\cdot|+1)\right) = \int_{\mathbb{R}^{d}} \frac{\mathcal{Z}_{+\theta_{0}}\sum_{y'\in\{\pm 1\}}\sigma(\langle z,\theta_{1}\rangle + y'\theta_{0})}{\mathcal{Z}\sigma(\langle z,\theta_{1}\rangle + \theta_{0})} P_{\mathbf{Z}}\left(\mathrm{d}z\right) - 1$$

$$\leq 2\frac{\mathcal{Z}_{+\theta_{0}}}{\mathcal{Z}} - 1 \leq 1.$$

In addition, the inequality $\sigma(t + \theta_0) + \sigma(t - \theta_0) \le 2$ gives

$$\chi^{2}\left(P_{\mathbf{Z}}(\cdot)||P_{\mathbf{Z}|Y}(\cdot|-1)\right) = \int_{\mathbb{R}^{d}} \frac{\mathcal{Z}_{-\theta_{0}} \sum_{y' \in \{\pm 1\}} \sigma(\langle z, \theta_{1} \rangle + y'\theta_{0})}{\mathcal{Z}\sigma(\langle z, \theta_{1} \rangle - \theta_{0})} P_{\mathbf{Z}}\left(dz\right) - 1$$

$$\leq 2e^{-\theta_{0}} M_{X}(\theta_{1}) \int_{\mathbb{R}^{d}} \frac{1}{\sigma(\langle z, \theta_{1} \rangle - \theta_{0})} P_{\mathbf{Z}}\left(dz\right) - 1$$

$$\leq 2e^{-\theta_{0}} M_{X}(\theta_{1}) \int_{\mathbb{R}^{d}} \exp\left(-\langle z, \theta_{1} \rangle + \theta_{0}\right) P_{\mathbf{Z}}\left(dz\right) - 1$$

$$= 2e^{-\theta_{0}} M_{X}(\theta_{1}) \int_{\mathbb{R}^{d}} \exp\left(-\langle z, \theta_{1} \rangle + \theta_{0}\right) \sum_{y \in \{\pm 1\}} \sigma(\langle z, \theta_{1} \rangle + y\theta_{0}) \mathcal{Z}^{-1} e^{-U(z)} dz - 1$$

$$\leq 4e^{-\theta_{0}} M_{X}(\theta_{1}) \int_{\mathbb{R}^{d}} \exp\left(-\langle z, \theta_{1} \rangle + \theta_{0}\right) \mathcal{Z}^{-1} e^{-U(z)} dz - 1$$

$$= 4M_{X}^{2}(\theta_{1}) - 1.$$

Therefore, we obtain

$$K_{\chi^2} = \max_{y \in \{\pm 1\}} \chi^2 \left(P_{\mathbf{Z}}(\cdot) || P_{\mathbf{Z}|Y}(\cdot|y) \right) \le 4M_X^2(\theta_1) - 1;$$

here, we use $M_X(\theta_1) = \mathbb{E}[\exp(\langle \theta_1, X \rangle)] \ge \exp(\mathbb{E}[\langle \theta_1, X \rangle]) = \exp(0) = 1$ by the evenness of U. For the other constants, we have

$$K_{\rm V} \le 4e^{-\theta_0} M_X(\theta_1), \ K_U = \frac{\mathcal{Z}}{\mathcal{Z}_{-\theta_0}} \le 2e^{\theta_0}$$

Hence, we obtain the following log-Sobolev inequality by Theorem 1 ($c = c^* = 2$) and Lemma 13: for any $f \in \mathcal{A}_n$,

$$\operatorname{Ent}(f^{2}((\mathbf{Z}_{i}, Y_{i})_{i=1}^{n})) \leq 2\mathbb{E}[\Gamma_{\operatorname{iii}}(f)((\mathbf{Z}_{i}, Y_{i})_{i=1}^{n})],$$

where

$$\begin{split} \Gamma_{\text{iii}}(f) &:= \left(\left(1 + \frac{1}{2} \log \left(2e^{\theta_0} \right) \right) K_{\text{LS}} \left(1 + 2(4M_X^2(\theta_1) - 1) \right) + 2K_{\text{LS}} \right) \sum_{i=1}^n \Gamma_{\mathbf{Z}_i}(f) \\ &+ 8e^{-\theta_0} M_X(\theta_1) \left(1 + \frac{1}{2} \log \left(2e^{\theta_0} \right) \right) \sum_{i=1}^n \Gamma_{Y_i}(f) \\ &\leq K_{\text{LS}} \frac{e + \theta_0}{2} \left(\frac{1}{2} + 8M_X^2(\theta_1) \right) \sum_{i=1}^n \Gamma_{\mathbf{Z}_i}(f) + 4e^{-\theta_0} M_X(\theta_1) \left(e + \theta_0 \right) \sum_{i=1}^n \Gamma_{Y_i}(f); \end{split}$$

here, we use $2 + \log 2 \le e$ and

$$\left(1 + \frac{1}{2}\log\left(2e^{\theta_0}\right)\right) \left(8M_X^2(\theta_1) - 1\right) + 2 \le \left(\frac{e}{2} + \frac{\theta_0}{2}\right) \left(8M_X^2(\theta_1) - 1\right) + 2
\le \left(\frac{e}{2} + \frac{\theta_0}{2}\right) \left(8M_X^2(\theta_1) - 1 + \frac{4}{e}\right) \le \left(\frac{e}{2} + \frac{\theta_0}{2}\right) \left(8M_X^2(\theta_1) + \frac{1}{2}\right).$$

Therefore, using Equations (8) and (9) and Lemma 12, we obtain that for all $\delta \in (0, 1]$, with probability at least $1 - \delta$,

$$\sup_{(\boldsymbol{w},b)\in\mathcal{T}} (\mathcal{R}(\boldsymbol{w},b) - \mathcal{R}_n(\boldsymbol{w},b)) - \mathbb{E}\left[\sup_{(\boldsymbol{w},b)\in\mathcal{T}} (\mathcal{R}(\boldsymbol{w},b) - \mathcal{R}_n(\boldsymbol{w},b))\right]$$

$$\leq \sqrt{\frac{L^2\left(e + \theta_0\right)\left[K_{\text{LS}}\left(1/2 + 8M_X^2(\boldsymbol{\theta}_1)\right)R_{\boldsymbol{w}}^2 + 8e^{-\theta_0}M_X(\boldsymbol{\theta}_1)R_b^2\right]\log(1/\delta)}{n}}.$$

This is the desired conclusion.

Proof of Proposition 7. We only prove (ii) as the proof of (i) is analogous and simpler.

We assume that $\theta_0 \ge 0$ without loss of generality. Using Equation 10, we obtain

$$\mathcal{Z}_{+\theta_0} = \int_{\mathbb{R}^d} g(\langle z, \theta_1 \rangle + \theta_0) \exp(-U(z)) dz = g(+\theta_0) \int_{\mathbb{R}^d} \frac{g(\langle z, \theta_1 \rangle + \theta_0)}{g(+\theta_0)} \exp(-U(z)) dz
\leq g(+\theta_0) \int_{\mathbb{R}^d} \exp(\max\{0, \langle z, G\theta_1 \rangle\}) \exp(-U(z)) dz = g(+\theta_0) M_X^{\text{num}}(G\theta_1) \mathcal{Z}.$$

Similarly, we have

$$\mathcal{Z}_{+\theta_0} \geq g(+\theta_0) M_{\boldsymbol{X}}^{\text{den}}(G\boldsymbol{\theta}_1) \mathcal{Z}, \ \mathcal{Z}_{-\theta_0} \leq g(-\theta_0) M_{\boldsymbol{X}}^{\text{num}}(G\boldsymbol{\theta}_1) \mathcal{Z}, \ \mathcal{Z}_{-\theta_0} \geq g(-\theta_0) M_{\boldsymbol{X}}^{\text{den}}(G\boldsymbol{\theta}_1) \mathcal{Z}.$$

In total, $\mathcal{Z}_{+\theta_0}/(g(+\theta_0)\mathcal{Z})$, $\mathcal{Z}_{-\theta_0}/(g(-\theta_0)\mathcal{Z}) \in [M_X^{\text{den}}(G\theta_1), M_X^{\text{den}}(G\theta_1)]$. Then, since $g(t-\theta_0)/g(t+\theta_0) \le 1$ and $g(t+\theta_0)+g(t-\theta_0) \le 2$ for any $t \in \mathbb{R}$,

$$\begin{split} \chi^{2}(P_{\mathbf{Z}}(\cdot) || P_{\mathbf{Z}|Y}(\cdot|+1)) \\ &= \int_{\mathbb{R}^{d}} \frac{\mathcal{Z}_{+\theta_{0}} \sum_{y' \in \{\pm 1\}} g(\langle z, \theta_{1} \rangle + y'\theta_{0})}{\mathcal{Z}g(\langle z, \theta_{1} \rangle + \theta_{0})} P_{\mathbf{Z}}(\mathrm{d}z) - 1 \\ &\leq g(+\theta_{0}) M_{X}^{\mathrm{num}}(G\theta_{1}) \int_{\mathbb{R}^{d}} \left(1 + \frac{g(\langle z, \theta_{1} \rangle - \theta_{0})}{g(\langle z, \theta_{1} \rangle + \theta_{0})} \right) P_{\mathbf{Z}}(\mathrm{d}z) - 1 \\ &\leq g(+\theta_{0}) M_{X}^{\mathrm{num}}(G\theta_{1}) \int_{\mathbb{R}^{d}} \left(1 + \frac{g(-\theta_{0})}{g(+\theta_{0})} \frac{g(\langle z, \theta_{1} \rangle - \theta_{0})/g(-\theta_{0})}{g(\langle z, \theta_{1} \rangle + \theta_{0})/g(+\theta_{0})} \right) P_{\mathbf{Z}}(\mathrm{d}z) - 1 \\ &\leq M_{X}^{\mathrm{num}}(G\theta_{1}) \int_{\mathbb{R}^{d}} \left(g(+\theta_{0}) + g(-\theta_{0}) \exp\left(2\langle z, G\theta_{1} \rangle \right) \right) P_{\mathbf{Z}}(\mathrm{d}z) - 1 \\ &= M_{X}^{\mathrm{num}}(G\theta_{1}) \int_{\mathbb{R}^{d}} \left(g(+\theta_{0}) + g(-\theta_{0}) \exp\left(2\langle z, G\theta_{1} \rangle \right) \right) \sum_{y \in \{\pm 1\}} g(\langle z, \theta_{1} \rangle + y\theta_{0}) \mathcal{Z}^{-1} e^{-U(z)} \mathrm{d}z - 1 \end{split}$$

$$\leq 2M_X^{\text{num}}(G\boldsymbol{\theta}_1)\left(g(+\boldsymbol{\theta}_0)+g(-\boldsymbol{\theta}_0)M_X^{\text{num}}(2G\boldsymbol{\theta}_1)\right)-1.$$

Moreover,

$$\chi^{2}(P_{\mathbf{Z}}(\cdot)||P_{\mathbf{Z}|Y}(\cdot|-1)) = \int_{\mathbb{R}^{d}} \frac{\mathcal{Z}_{-\theta_{0}} \sum_{y' \in \{\pm 1\}} g(\langle z, \theta_{1} \rangle + y'\theta_{0})}{\mathcal{Z}g(\langle z, \theta_{1} \rangle - \theta_{0})} P_{\mathbf{Z}}(\mathrm{d}z) - 1$$

$$\leq g(-\theta_{0}) M_{X}^{\mathrm{num}}(G\theta_{1}) \int_{\mathbb{R}^{d}} \left(1 + \frac{g(\langle z, \theta_{1} \rangle + \theta_{0})}{g(\langle z, \theta_{1} \rangle - \theta_{0})} \right) P_{\mathbf{Z}}(\mathrm{d}z) - 1$$

$$\leq g(-\theta_{0}) M_{X}^{\mathrm{num}}(G\theta_{1}) \int_{\mathbb{R}^{d}} \left(1 + \frac{g(+\theta_{0})}{g(-\theta_{0})} \frac{g(\langle z, \theta_{1} \rangle + \theta_{0})/g(+\theta_{0})}{g(\langle z, \theta_{1} \rangle - \theta_{0})/g(-\theta_{0})} \right) P_{\mathbf{Z}}(\mathrm{d}z) - 1$$

$$\leq M_{X}^{\mathrm{num}}(G\theta_{1}) \int_{\mathbb{R}^{d}} \left(g(-\theta_{0}) + g(+\theta_{0}) \exp\left(2\langle z, G\theta_{1} \rangle\right) \right) P_{\mathbf{Z}}(\mathrm{d}z) - 1$$

$$\leq 2M_{X}^{\mathrm{num}}(G\theta_{1}) \left(g(-\theta_{0}) + g(+\theta_{0}) M_{X}^{\mathrm{num}}(2G\theta_{1}) \right) - 1.$$

Therefore, since $g(+\theta_0) \ge g(-\theta_0)$ and $M_X^{\text{num}}(t) \ge 1$ for $t \in \mathbb{R}^d$,

$$K_{\chi^2} \leq 2M_{X}^{\text{num}}(G\theta_1) \left(g(-\theta_0) + g(+\theta_0) M_{X}^{\text{num}}(2G\theta_1) \right) - 1 \leq 2M_{X}^{\text{num}}(G\theta_1) M_{X}^{\text{num}}(2G\theta_1) - 1.$$

We also have

$$K_{V} = 4g(+\theta_{0})g(-\theta_{0})M_{X}^{\text{num}}(G\theta_{1}), K_{U} = \frac{1}{g(-\theta_{0})M_{X}^{\text{den}}(G\theta_{1})}.$$

Hence, Theorem 1 ($c = c^* = 2$) along with Lemma 13 gives the following log-Sobolev inequality: for arbitrary $f \in \mathcal{A}_n$,

$$\operatorname{Ent}(f^2((\mathbf{Z}_i, Y_i)_{i=1}^n)) \le 2\mathbb{E}[\Gamma_{iv}(f)((\mathbf{Z}_i, Y_i)_{i=1}^n)],$$

where

$$\begin{split} \Gamma_{\text{iv}}(f) &:= \left(\left(1 - \frac{1}{2} \log \left(g(-\theta_0) M_X^{\text{den}}(G\theta_1) \right) \right) K_{\text{LS}} \left(4 M_X^{\text{num}}(G\theta_1) M_X^{\text{num}}(2G\theta_1) - 1 \right) + 2 K_{\text{LS}} \right) \sum_{i=1}^n \Gamma_{\mathbf{Z}_i}(f) \\ &+ 8 g(+\theta_0) g(-\theta_0) M_X^{\text{num}}(G\theta_1) \left(1 - \frac{1}{2} \log \left(g(-\theta_0) M_X^{\text{den}}(G\theta_1) \right) \right) \sum_{i=1}^n \Gamma_{Y_i}(f) \\ &\leq K_{\text{LS}} \left(1 - \frac{1}{2} \log \left(g(-\theta_0) M_X^{\text{den}}(G\theta_1) \right) \right) \left(5 M_X^{\text{num}}(G\theta_1) M_X^{\text{num}}(2G\theta_1) \right) \sum_{i=1}^n \Gamma_{\mathbf{Z}_i}(f) \\ &+ 8 g(+\theta_0) g(-\theta_0) M_X^{\text{num}}(G\theta_1) \left(1 - \frac{1}{2} \log \left(g(-\theta_0) M_X^{\text{den}}(G\theta_1) \right) \right) \sum_{i=1}^n \Gamma_{Y_i}(f). \end{split}$$

Therefore, Equations (8) and (9) and Lemma 12 give that for all $\delta \in (0, 1]$, with probability at least $1 - \delta$,

$$\sup_{(\boldsymbol{w},b)\in\mathcal{T}} (\mathcal{R}(\boldsymbol{w},b) - \mathcal{R}_n(\boldsymbol{w},b)) - \mathbb{E}\left[\sup_{(\boldsymbol{w},b)\in\mathcal{T}} (\mathcal{R}(\boldsymbol{w},b) - \mathcal{R}_n(\boldsymbol{w},b))\right]$$

$$\leq \sqrt{\frac{2L^2 \log(1/\delta)}{n}} \sqrt{1 - \frac{1}{2} \log\left(g(-\theta_0) M_X^{\text{den}}(G\theta_1)\right)}$$

$$\times \sqrt{5K_{\text{LS}} M_X^{\text{num}}(G\theta_1) M_X^{\text{num}}(2G\theta_1) R_{\boldsymbol{w}}^2 + 8g(+\theta_0)g(-\theta_0) M_X^{\text{num}}(G\theta_1) R_b^2}$$

Hence, the statement holds true.

Proof of Proposition 8. We give the proof of (ii) and omit that of (i), as that of (i) can be given with fewer steps by applying Theorem 1 with $c = 1 + 1/\sqrt{e^{5|\theta_0|}/\tilde{M}_X(\theta_1)}$ and $c^* = 1 + \sqrt{e^{5|\theta_0|}/\tilde{M}_X(\theta_1)}$.

Let $\theta_0 \ge 0$ without loss of generality. Since $\sigma(t) \le \exp(t)$ for any $t \in \mathbb{R}$,

$$\begin{split} \mathcal{Z}_{+\theta_0} &= \int_{\mathbb{R}^d} \sigma(\langle z, \theta_1 \rangle + \theta_0) \exp(-U(z)) \mathrm{d}z \\ &\leq \int_{\langle z, \theta_1 \rangle \geq 0} \exp(-U(z)) \mathrm{d}z + \int_{\langle z, \theta_1 \rangle < 0} \sigma(\langle z, \theta_1 \rangle + \theta_0) \exp(-U(z)) \mathrm{d}z \\ &= \int_{\langle z, \theta_1 \rangle \geq 0} \exp(-U(z)) \mathrm{d}z + \exp(+\theta_0) \int_{\langle z, \theta_1 \rangle < 0} \exp(\langle z, \theta_1 \rangle) \exp(-U(z)) \mathrm{d}z \\ &\leq \frac{\mathcal{Z}}{2} \left(1 + \exp(+\theta_0) \mathbb{E} \left[\exp\left(- |\langle X, \theta_1 \rangle| \right) \right] \right), \\ \mathcal{Z}_{-\theta_0} &\leq \frac{\mathcal{Z}}{2} \left(1 + \exp(-\theta_0) \mathbb{E} \left[\exp\left(- |\langle X, \theta_1 \rangle| \right) \right] \right), \end{split}$$

and by $\sigma(t) \ge 1/2$ for $t \ge 0$,

$$\mathcal{Z}_{-\theta_0} = \int_{\mathbb{R}^d} \sigma(\langle z, \theta_1 \rangle - \theta_0) \exp(-U(z)) dz \ge \int_{\langle z, \theta_1 \rangle - \theta_0 \ge 0} \sigma(\langle z, \theta_1 \rangle - \theta_0) \exp(-U(z)) dz \\
\ge \frac{\mathcal{Z}}{2} \mathbb{P} \left(\langle X, \theta_1 \rangle \ge \theta_0\right).$$

We obtain

$$\chi^{2}(P_{\mathbf{Z}}(\cdot)||P_{\mathbf{Z}|Y}(\cdot|+1)) = \int_{\mathbb{R}^{d}} \frac{\mathcal{Z}_{+\theta_{0}} \sum_{y' \in \{\pm 1\}} \sigma(\langle z, \theta_{1} \rangle + y'\theta_{0})}{\mathcal{Z}\sigma(\langle z, \theta_{1} \rangle + \theta_{0})} P_{\mathbf{Z}}(dz) - 1$$

$$\leq \frac{1}{2} \left(1 + \exp(+\theta_{0}) / \tilde{M}_{X}(\theta_{1}) \right) \int_{\mathbb{R}^{d}} \left(1 + \frac{\sigma(\langle z, \theta_{1} \rangle - \theta_{0})}{\sigma(\langle z, \theta_{1} \rangle + \theta_{0})} \right) P_{\mathbf{Z}}(dz) - 1$$

$$\leq 1 + \exp(+\theta_{0}) / \tilde{M}_{X}(\theta_{1}) - 1$$

$$\leq \exp(+\theta_{0}) / \tilde{M}_{X}(\theta_{1})$$

by the monotonicity of σ , and

$$\chi^{2}(P_{\mathbf{Z}}(\cdot)||P_{\mathbf{Z}|Y}(\cdot|-1)) = \int_{\mathbb{R}^{d}} \frac{Z_{-\theta_{0}} \sum_{y' \in \{\pm 1\}} \sigma(\langle z, \theta_{1} \rangle + y'\theta_{0})}{Z \sigma(\langle z, \theta_{1} \rangle - \theta_{0})} P_{\mathbf{Z}}(dz) - 1$$

$$\leq \frac{1}{2} \left(1 + \exp(-\theta_{0}) / \tilde{M}_{X}(\theta_{1}) \right) \int_{\mathbb{R}^{d}} \left(1 + \frac{\sigma(\langle z, \theta_{1} \rangle + \theta_{0})}{\sigma(\langle z, \theta_{1} \rangle - \theta_{0})} \right) P_{\mathbf{Z}}(dz) - 1$$

$$\leq \left(1 + \exp(-\theta_{0}) / \tilde{M}_{X}(\theta_{1}) \right) - 1$$

$$+ \frac{1}{2} e^{+\theta_{0}} (e^{+4\theta_{0}} - 1) / \tilde{M}_{X}(\theta_{1}) \left(1 + \exp(-\theta_{0}) / \tilde{M}_{X}(\theta_{1}) \right)$$

$$= \left(1 / \tilde{M}_{X}(\theta_{1}) \right) \left(e^{-\theta_{0}} + \frac{1}{2} e^{+\theta_{0}} (e^{+4\theta_{0}} - 1) \left(1 + e^{-\theta_{0}} / \tilde{M}_{X}(\theta_{1}) \right) \right)$$

because

$$\frac{\sigma(t+\theta_0)}{\sigma(t-\theta_0)} - 1 = \frac{\sigma(t+\theta_0)}{\sigma(t-\theta_0)} - 1 = \frac{1+e^{-t}e^{+\theta_0}}{1+e^{-t}e^{-\theta_0}} - 1 = \frac{e^{-t}(e^{+\theta_0}-e^{-\theta_0})}{1+e^{-t}e^{-\theta_0}} = (e^{+2\theta_0}-1)\frac{e^{-t-\theta_0}}{1+e^{-t-\theta_0}}$$
$$= (e^{+2\theta_0}-1)\frac{1}{1+e^{t+\theta_0}} = (e^{+2\theta_0}-1)\sigma(-t-\theta_0),$$

and thus the facts $\sigma(t)\sigma(-t) \le \sigma(-|t|)$ and $\sigma(t) \le \exp(t)$ give

$$\begin{split} \int_{\mathbb{R}^{d}} \frac{\sigma(t+\theta_{0})}{\sigma(t-\theta_{0})} P_{\mathbf{Z}}(\mathrm{d}z) - 1 &= (e^{+2\theta_{0}}-1) \int_{\mathbb{R}^{d}} \sigma(-\langle z,\theta_{1}\rangle - \theta_{0}) P_{\mathbf{Z}}(\mathrm{d}z) \\ &= (e^{+2\theta_{0}}-1) \int_{\mathbb{R}^{d}} \sigma(-\langle x,\theta_{1}\rangle - \theta_{0}) \left(\sigma(\langle x,\theta_{1}\rangle + \theta_{0}) + \sigma(\langle x,\theta_{1}\rangle - \theta_{0})\right) P_{\mathbf{X}}(\mathrm{d}x) \\ &\leq (e^{+2\theta_{0}}-1) \int_{\mathbb{R}^{d}} \sigma(-\langle x,\theta_{1}\rangle - \theta_{0}) (1+e^{+2\theta_{0}}) \sigma(\langle x,\theta_{1}\rangle + \theta_{0}) P_{\mathbf{X}}(\mathrm{d}x) \\ &\leq (e^{+4\theta_{0}}-1) \int_{\mathbb{R}^{d}} \frac{1}{1+\exp(-\langle x,\theta_{1}\rangle - \theta_{0})} \frac{1}{1+\exp(+\langle x,\theta_{1}\rangle + \theta_{0})} P_{\mathbf{X}}(\mathrm{d}x) \\ &\leq (e^{+4\theta_{0}}-1) \int_{\mathbb{R}^{d}} \sigma(-|\langle x,\theta_{1}\rangle + \theta_{0}|) P_{\mathbf{X}}(\mathrm{d}x) \\ &\leq (e^{+4\theta_{0}}-1) \mathbb{E} \left[\exp(-|\langle X,\theta_{1}\rangle + \theta_{0}|)\right] \\ &\leq e^{+\theta_{0}} (e^{+4\theta_{0}}-1) \mathbb{E} \left[\exp(-|\langle X,\theta_{1}\rangle + \theta_{0}|)\right]. \end{split}$$

In total, since both $e^{-\theta_0} \le 1$ and $1/\tilde{M}_X(\theta_1) \le 1$ by definition, we have

$$\begin{split} K_{\chi^2} & \leq e^{+\theta_0}/\tilde{M}_X(\theta_1) \left(1 + \frac{1}{2}(e^{+4\theta_0} - 1) \left(1 + e^{-\theta_0}/\tilde{M}_X(\theta_1)\right)\right) \\ & \leq e^{+\theta_0}/\tilde{M}_X(\theta_1) \left(1 + (e^{+4\theta_0} - 1)\right) \leq e^{+5\theta_0}/\tilde{M}_X(\theta_1), \end{split}$$

and

$$K_{\mathrm{U}} \leq 1, \ K_{\mathrm{V}} \leq \frac{2}{p_{\theta_{1},\theta_{0}}}.$$

Therefore, the following log-Sobolev inequality holds by Theorem 1 ($c = c^* = 2$) and Lemma 13: for any $f \in \mathcal{A}_n$,

$$\operatorname{Ent}(f^{2}((\mathbf{Z}_{i}, Y_{i})_{i-1}^{n})) \leq 2\mathbb{E}[\Gamma_{V}(f)((\mathbf{Z}_{i}, Y_{i})_{i-1}^{n})],$$

where

$$\begin{split} \Gamma_{\mathbf{v}}(f) &:= K_{\mathrm{LS}}\left(\left(1 + \frac{1}{2}\log\left(\frac{2}{p_{\theta_{1},\theta_{0}}}\right)\right)\left(1 + 2e^{+5\theta_{0}}/\tilde{M}_{X}(\theta_{1})\right) + 2\right) \sum_{i=1}^{n} \Gamma_{\mathbf{Z}_{i}}(f) \\ &+ 2\left(1 + \frac{1}{2}\log\left(\frac{2}{p_{\theta_{1},\theta_{0}}}\right)\right) \sum_{i=1}^{n} \Gamma_{Y_{i}}(f) \\ &\leq K_{\mathrm{LS}}\left(\frac{e}{2} + \frac{1}{2}\log\left(\frac{1}{p_{\theta_{1},\theta_{0}}}\right)\right) \left(\frac{5}{2} + 2e^{+5\theta_{0}}/\tilde{M}_{X}(\theta_{1})\right) \sum_{i=1}^{n} \Gamma_{\mathbf{Z}_{i}}(f) \\ &+ \left(e + \log\left(\frac{1}{p_{\theta_{1},\theta_{0}}}\right)\right) \sum_{i=1}^{n} \Gamma_{Y_{i}}(f), \end{split}$$

where we use $1+(1/2)\log(2) \le e/2$ and $1+4/e \le 5/2$. Therefore, as the proofs of the previous propositions above, for all $\delta \in (0,1]$, with probability at least $1-\delta$,

$$\sup_{(\boldsymbol{w},b)\in\mathcal{T}} (\mathcal{R}(\boldsymbol{w},b) - \mathcal{R}_n(\boldsymbol{w},b)) - \mathbb{E}\left[\sup_{(\boldsymbol{w},b)\in\mathcal{T}} (\mathcal{R}(\boldsymbol{w},b) - \mathcal{R}_n(\boldsymbol{w},b))\right] \\ \leq \sqrt{\frac{2L^2\left(e + \log\left(1/p_{\theta_1},\theta_0\right)\right)\left(K_{\text{LS}}\left(5/4 + e^{+5\theta_0}/\tilde{M}_X(\theta_1)\right)R_{\boldsymbol{w}}^2 + R_b^2\right)\log(1/\delta)}{n}}.$$

Hence, the statement holds true.

 $\mathsf{K}\mathsf{o}\mathsf{m}\mathsf{a}\mathsf{b}\mathsf{a}$ Institute for Science, University of Tokyo, 3-8-1 $\mathsf{K}\mathsf{o}\mathsf{m}\mathsf{a}\mathsf{b}\mathsf{a}$, Meguro-ku, Tokyo 153-8902, Japan