

Body size trends in Neogene tortoises

Did tortoises evolve towards a smaller body size?

Julia Joos

Humboldt-Universität zu Berlin

20.07.2016

Terrestrial Tortoises

- 2 clades: Meiolaniidae **P**
and Testudinidae **P**

Terrestrial Tortoises

- 2 clades: Meiolaniidae **P** and Testudinidae **P**
- Meiolaniidae: extinct, used to be present in Australia + South America

Terrestrial Tortoises

- 2 clades: Meiolaniidae **P** and Testudinidae **P**
- Meiolaniidae: extinct, used to be present in Australia + South America
- Testudinidae: comprise all extant terrestrial tortoises (America, Europe, Africa, Asia)

Terrestrial Tortoises

- 2 clades: Meiolaniidae **P** and Testudinidae **P**
- Meiolaniidae: extinct, used to be present in Australia + South America
- Testudinidae: comprise all extant terrestrial tortoises (America, Europe, Africa, Asia)
- Testudinidae: probably Asian origin, oldest fossils: North America + Europe

Terrestrial Tortoises

- 2 clades: Meiolaniidae **P** and Testudinidae **P**
- Meiolaniidae: extinct, used to be present in Australia + South America
- Testudinidae: comprise all extant terrestrial tortoises (America, Europe, Africa, Asia)
- Testudinidae: probably Asian origin, oldest fossils: North America + Europe
- today's most famous

Terrestrial Tortoises

- 2 clades: Meiolaniidae **P** and Testudinidae **P**
- Meiolaniidae: extinct, used to be present in Australia + South America
- Testudinidae: comprise all extant terrestrial tortoises (America, Europe, Africa, Asia)
- Testudinidae: probably Asian origin, oldest fossils: North America + Europe
- today's most famous

Megafauna

- animals with body mass > 44 kg = megafauna

Megafauna

- animals with body mass $> 44 \text{ kg}$ = megafauna
- mammalian megafauna:
popular, famous, well
investgated

Megafauna

- animals with body mass $> 44 \text{ kg}$ = megafauna
- mammalian megafauna: popular, famous, well investigated
- megafauna extinctions! – humans or climate change

Megafauna

- animals with body mass $> 44 \text{ kg}$ = megafauna
- mammalian megafauna: popular, famous, well investigated
- megafauna extinctions! – humans or climate change
- ...

Pleistocene Extinctions

- human influence

Pleistocene Extinctions

- human influence
- climate change

Pleistocene Extinctions

- human influence
- climate change
- ...

Pleistocene Extinctions


- human influence
- climate change
- ...
- ...

① Body size distribution of Testudinidae?

- ① Body size distribution of Testudinidae?
- ② Body size differences on spatial/temporal scale?


- ① Body size distribution of Testudinidae?
- ② Body size differences on spatial/temporal scale?
- ③ General body size trends?

Data set




- occurrences (black): 796 individuals, 647 localities (Eocene – recent)

Data set



- occurrences (black): 796 individuals, 647 localities (Eocene – recent)
- body size data (blue): 376 individuals, 193 localities (Miocene – recent)

Data set



- occurrences (black): 796 individuals, 647 localities (Eocene – recent)
- body size data (blue): 376 individuals, 193 localities (Miocene – recent)

Carapace length measurements/estimations

- exact measurements or estimations by original authors (n=...)

Carapace length measurements/estimations

- exact measurements or estimations by original authors (n=...)
- estimations from CL/PL ratio (– extant measurements)

Carapace length measurements/estimations

- exact measurements or estimations by original authors (n=...)
- estimations from CL/PL ratio (– extant measurements)
- estimations from humeri/femora length, others (claw phalanges, verbal description etc.)

Methods

- SACs to check if sampling was sufficient – ζ generic level!
add figure!!

Methods

- SACs to check if sampling was sufficient – $\hat{\epsilon}$ generic level!
add figure!!
- histograms/density plots and boxplots (plus kruskal-wallis test, wilcoxon rank test)

Methods

- SACs to check if sampling was sufficient – ζ generic level!
add figure!!
- histograms/density plots and boxplots (plus kruskal-wallis test, wilcoxon rank test)
- paleoTS: fit different models to evolutionary trajectory of testudinid body size

- ① Body size distribution of Testudinidae?
- ② Body size differences on spatial/temporal scale?
- ③ General body size trends?

Inhalt...

- ① Body size distribution of Testudinidae?
- ② Body size differences on spatial/temporal scale?
- ③ General body size trends?

- 1 Body size distribution of Testudinidae?
- 2 Body size differences on spatial/temporal scale?
- 3 General body size trends?

① Body size distribution of Testudinidae?

- 1 Body size distribution of Testudinidae?
→ Bimodal distribution (left-skewed on islands!)

- ① Body size distribution of Testudinidae?
→ Bimodal distribution (left-skewed on islands!)
- ② Body size differences on spatial/temporal scale?

- ① Body size distribution of Testudinidae?
→ Bimodal distribution (left-skewed on islands!)
- ② Body size differences on spatial/temporal scale?
→ modern < fossil

- ① Body size distribution of Testudinidae?
 - Bimodal distribution (left-skewed on islands!)
- ② Body size differences on spatial/temporal scale?
 - modern < fossil
 - continental < insular

- ① Body size distribution of Testudinidae?
 - Bimodal distribution (left-skewed on islands!)
- ② Body size differences on spatial/temporal scale?
 - modern < fossil
 - continental < insular
- ③ General body size trends?

- ① Body size distribution of Testudinidae?
 - Bimodal distribution (left-skewed on islands!)
- ② Body size differences on spatial/temporal scale?
 - modern < fossil
 - continental < insular
- ③ General body size trends?
 - stasis (generally and on islands)

- ❶ Body size distribution of Testudinidae?
 - Bimodal distribution (left-skewed on islands!)
- ❷ Body size differences on spatial/temporal scale?
 - modern < fossil
 - continental < insular
- ❸ General body size trends?
 - stasis (generally and on islands)
 - URW (continental tortoises)

Image sources

①

②