

MAthesis

Julia Joos

14 August 2017

Table 1: Time bins with age range, epoch name, mean age and corresponding sample sizes (on individual, species and genus level)

| bin | EpochBins | MeanBins | nIndividuals | nSpecies | nGenera |
|----------------|--------------------|-----------|--------------|----------|---------|
| (0,1e-06] | Modern | 0.0000005 | 240 | 58 | 17 |
| (1e-06,0.0117] | Holocene | 0.0058500 | 12 | 6 | 4 |
| (0.0117,0.126] | Upper Pleistocene | 0.0688500 | 46 | 15 | 7 |
| (0.126,0.781] | Middle Pleistocene | 0.4535000 | 46 | 9 | 6 |
| (0.781,2.59] | Lower Pleistocene | 1.6845000 | 68 | 24 | 11 |
| (2.59,3.6] | Upper Pliocene | 3.0940000 | 21 | 14 | 9 |
| (3.6,5.33] | Lower Pliocene | 4.4660000 | 27 | 16 | 8 |
| (5.33,11.6] | Upper Miocene | 8.4700000 | 41 | 21 | 9 |




Figure 1: Map displaying all fossil occurrences of testudinids, with color indicating whether relevant literature was available (black if not) and if it was, whether body size data was available or not (yes and no, respectively).







Figure 2: Map displaying all localities for which body size data for testudinids was available in the literature. Size of points denotes sample size, color denotes approximate age.






CL


Fossil species, CL, per Locality




Fossil species, CL, per Reference



Fossil genera, CL, per Locality




Fossil genera, CL, per Reference



Warning: Removed 16 rows containing missing values (geom_pointrange).

Warning: Removed 16 rows containing missing values (geom_pointrange).






Figure 4: Mean body size and standard deviation per genus in each time bin

1 including Island species (n=2215)

Table 2: paleoTS object (mm= mean CL, nn = sample size, vv = variance (CL), tt = Age)

| mm | nn | vv | tt |
|-----------|------|--------------|-----------|
| 246.8335 | 1968 | 2.126636e+09 | 0.0000005 |
| 688.5455 | 11 | 1.245041e+05 | 0.0058500 |
| 447.6480 | 45 | 8.098707e+04 | 0.0688500 |
| 333.8707 | 45 | 3.704545e+04 | 0.4535000 |
| 415.0939 | 66 | 1.833202e+05 | 1.6845000 |
| 642.0167 | 18 | 2.812598e+05 | 3.0940000 |
| 1004.9909 | 22 | 5.319102e+05 | 4.4660000 |
| 582.7750 | 40 | 3.097159e+05 | 8.4700000 |

##

Comparing 3 models [n = 7, method = AD]

##

| ## | | logL | K | AICc | Akaike.wt |
|-----------|--|-----------|---|----------|-----------|
| ## GRW | | -50.30596 | 2 | 107.6119 | 0.038 |
| ## URW | | -50.78070 | 1 | 104.3614 | 0.191 |
| ## Stasis | | -47.28232 | 2 | 101.5646 | 0.772 |

Table 3: Model-fitting results for testudinidae, individuals, including island species

| | logL | K | AICc | Akaike.wt |
|--------|-----------|---|----------|-----------|
| GRW | -50.30596 | 2 | 107.6119 | 0.038 |
| URW | -50.78070 | 1 | 104.3614 | 0.191 |
| Stasis | -47.28232 | 2 | 101.5646 | 0.772 |




Figure 5: individuals, including island species

| EpochBins | meanSpeciesCL | nSpecies | MeanBins |
|--------------------|---------------|----------|----------|
| Holocene | 671.4667 | 5 | 0.00585 |
| Upper Pleistocene | 521.4533 | 17 | 0.06885 |
| Middle Pleistocene | 384.8626 | 10 | 0.45350 |
| Lower Pleistocene | 581.2039 | 28 | 1.68450 |
| Upper Pliocene | 610.4591 | 11 | 3.09400 |
| Lower Pliocene | 1009.4738 | 14 | 4.46600 |
| Upper Miocene | 680.7708 | 24 | 8.47000 |




Figure 6: paleoTS plot with species mean, including island species

| tt | mm | vv | nn |
|-----------|----------|-----------|----|
| 0.0000005 | 400.5972 | 104321.19 | 50 |
| 0.0058500 | 671.4667 | 195810.92 | 5 |
| 0.0688500 | 521.4533 | 67149.55 | 17 |

| | tt | mm | vv | nn |
|--|-----------|-----------|-----------|----|
| | 0.4535000 | 384.8626 | 99603.12 | 10 |
| | 1.6845000 | 581.2039 | 319998.46 | 28 |
| | 3.0940000 | 610.4591 | 260640.34 | 11 |
| | 4.4660000 | 1009.4738 | 437400.60 | 14 |
| | 8.4700000 | 680.7708 | 349806.99 | 24 |

##

Comparing 3 models [n = 7, method = AD]

##

| ## | | logL | K | AICc | Akaike.wt |
|-----------|-----------|------|-----------|------|-----------|
| ## GRW | -47.86862 | 2 | 102.73724 | | 0.059 |
| ## URW | -47.99319 | 1 | 98.78638 | | 0.422 |
| ## Stasis | -45.68754 | 2 | 98.37508 | | 0.519 |

| | logL | K | AICc | Akaike.wt |
|--------|-----------|---|-----------|-----------|
| GRW | -47.86862 | 2 | 102.73724 | 0.059 |
| URW | -47.99319 | 1 | 98.78638 | 0.422 |
| Stasis | -45.68754 | 2 | 98.37508 | 0.519 |




Figure 7: paleoTS plot with species mean, including island species