



## LEBENSWISSENSCHAFTLICHE FAKULTÄT INSTITUT FÜR BIOLOGIE

#### **MASTERARBEIT**

# ZUM ERWERB DES AKADEMISCHEN GRADES MASTER OF SCIENCE

"Körpergrößentrends in fossilen Landschildkröten aus dem Neogen"

"Body size trends in Neogene testudinid tortoises"

vorgelegt von

Julia Joos

geb. am 18.05.1991 in Freudenstadt

angefertigt in der Arbeitsgruppe Paläozoologie am Institut für Biologie/Museum für Naturkunde

Berlin, im Septenber 2017

## Contents

Lis	st of Figures	3
Lis	st of Tables	4
1.	Introduction	5
2.	Material & Methods	6
	2.1. Data collection	6
	2.2. Body size estimation	7
	2.3. Analyses	7
	2.3.1. Data strucure and statistics	8
	2.3.2. Body size trends over time	8
3.	Results	10
4.	paleoTS analysis	15
	4.1. all (continental and insular)	15
	4.2. continental (excluding insular species)	16
	4.3. insular (excluding continental)	18
	4.4. per continent	20
	4.4.1. Europe, genera	20
	4.4.2. Eurasia, genera	23
	4.4.3. Eurasia, genera, continental	24
	4.4.4. Eurasia, genera, insular	26
5.	References	28
Αŗ	opendix A. Sampling Accumulation Curves	29
Αp	ppendix B. Histograms	30
Αŗ	opendix C. Boxplots	31
Αp	opendix D. Random Sampling	33
Αp	ppendix E. Tables	33

## **List of Figures**

١.	map: fossil occurences	Ь
2.	Carapace length over time	8
3.	Map: body size localities	12
4.	Sampling Accumulation Curve on generic level	13
5.	CL distribution	13
6.	Boxplots of mean CL per time bin, including mean and sd CL for each genus (as	
	pointrange)	14
7.	paleoTS plot with genus mean, all	16
8.	paleoTS plot with genus mean, continental	18
9.	paleoTS plot with genus mean, insular	19
10.	Genera, Europe	21
11.	paleoTS, genera, Europe, continental	21
12.	paleoTS, genera, Europe, insular	23
13.	paleoTS, genera, Eurasia	24
14.	paleoTS, genera, Eurasia, continental	26
15.	paleoTS, genera, Eurasia, insular	27
16.	Additional Sampling Accumulation Curves	29
17.	Testing normal distribution	30
18.	Additional histograms	30
19.	Boxplots fossil vs. modern, continental vs. insular species	31
20.	Boxplot: body size on different continents, genera summarised	32
21.	Random sampling	33
22	Bandom sampling continents	33

## **List of Tables**

1.	Sample sizes per time bins	11
2.	paleoTS object, all data	15
3.	Model-fitting results for testudinidae, genera, all	15
4.	paleoTS object, continental	16
5.	Model-fitting results for testudinidae, genera, continental	18
6.	paleoTS object, insular	18
7.	Model-fitting results for testudinidae, genera, insular	19
8.	paleoTS object, Europe	20
9.	Model-fitting results for testudinidae, genera, Europe	20
10.	paleoTs object, Europe, continental	21
11.	Model-fitting results for testudinidae, genera, Europe, continental	22
12.	paleoTs object, Europe, insular	22
13.	Model-fitting results for testudinidae, genera, Europe, insular	22
14.	paleoTS object, Eurasia	23
15.	Model-fitting results for testudinidae, genera, Eurasia	24
16.	paleoTS object, Eurasia, continental	25
17.	Model-fitting results for testudinidae, genera, Eurasia, continental	25
18.	paleoTS object, Eurasia, insular	26
19.	Model-fitting results for testudinidae, genera, Eurasia, insular	27
20.	Genera per time bins	34
21.	Genera overview	37
22.	General statistics	39
23.	Data set, fossil	41
24.	Data set, extant	74

#### 1. Introduction

- Body size as a trait (read Smith, Smith & Lyons) and over time why is it interesting? is there an optimal body size for every organism? how can it be determined? (-> stasis??)
- evolutionary models (read Gene Hunt's paper and Posada, 2003) -> make sense of evolutionary modes: \* stasis \* unbiased random walk \* generalized random walk
  - body size in tortoises (why not use biomass? not necessary) distribution of tortoises (?)
- giant tortoises well suited for drifting on ocean currents (Meylan, 2000)
  - OR mammal megafauna extinctions -> giant tortoises
  - human and climatic influence
- purpose of this work: determine body size trends in tortoises and identify evolutionary mode (if possible). what lead to extinction?

#### 2. Material & Methods

#### 2.1. Data collection

I collected data on body size of fossil testudinids from the Miocene until recent times. The body size data set includes 30 genera, comprising over 100 fossil species. The majority of the data was obtained from the primary literature (Table 23). To find relevant publications, I relied mostly on the references listed in the FosFarBase (Böhme and Ilg, 2003), the Paleobiology Database (PDBD), and the review by Rhodin et al. (2015). Furthermore, the FosFarBase provided fossil occurrences of testudinids all over the world, including their exact localities and age (Table ??, Fig. 1), which were used to get an overview over the availability of body size data. The FosFarBase (last accessed 23.03.2017) contained 769 testudinid occurrences between the Eocene (33.9 - 56 mya) and the Holocene from 647 localities (Fig. 1). Of those, 641 occurrences from 534 localities were of relevant Age (Miocene to Holocene). The final body size data set, however, includes 376 data records from 193 localities, of which 106 localities are present in the FosFarBase.

put table only on CD?




Figure 1: Map displaying all fossil occurrences of testudinids from the Eocene to the Holocene as according to the FosFarBase, with color indicating whether body size data was available (blue) or not (black).

For extant taxa, I measured dry material (n = 67) from the collection of the Museum für Naturkunde zu Berlin (MFN). In addition, body size data (n = 173) from the literature was included (Table 24).

#### 2.2. Body size estimation

Body size is reported as straight carapace length (SCL) in mm. Where SCL was not available from the primary literature, it was estimated (n = 254) either from plastron length (PL) or appendicular elements (Table 23). For carapace length estimations based on plastron length, the measurements from the MFN collection material were used to calculate the ratio between SCL and PL. Since the SC/PL ratio was similar for all species (SCL/PL between 0.95 - 1.47), a single general ratio (SCL/PL = 1.1) was calculated for all testudinids and hence used for the SCL estimations unless stated otherwise (Table 23). For estimations based on femora and humeri, the ratios based on data provided by Hutterer et al. (1998) and Franz et al. (2001), respectively, were used. A number of publications did not state measurements but instead provided scaled figures of the fossil remains, from which either SCL directly or PL, humeri, or femora lengths for estimating SCL could be measured.

do I need
to provide
statistics
for that?
do I need
this sentence at
all?

#### 2.3. Analyses

All subsequent analyses were performed with R 3.4.1 (R Core Team, 2017), including the packages dplyr (Wickham et al., 2017) to prepare the data for the analysis and ggplot2 (Wickham, 2009) to create figures. The R package vegan (Oksanen et al., 2017) was used to create Sampling Accumulation Curves (most commonly referred to as Species Accumulation Curves), which show the increase in individuals, species or genera per sampling unit and are therefore used to determine if sampling is sufficient or not in terms of covering diversity and richness. Since the data set relies on literature, references were used as a sampling unit (x-axis). Sampling Accumulation Curves were created on species as well as genus level, since genera of fossil testudinids are relatively well resolved by now whereas determination on the species level is still somewhat obscure in some cases, because fossil species are frequently based on single individuals that are often fragmentary as well (Brattstrom, 1961). Since genera were better sampled than species (Fig. 4, 16 (a) - (b)), all subsequent analysis were performed on the generic level. Additional Sampling Accumulation Curves for the continents were created (Fig. ?? (c) - (i)), to check if subsequent analyses could be applied to these subgroups.

explain
what
species
accumulation
curves
are and
give refer-

ence

#### 2.3.1. Data strucure and statistics

Histograms and boxplots of the entire data set and several subgroups (fossil vs. modern, insular vs. continental) were created to explore the structure of the data set. General parameters like mean, median, variance, and with the package moments (Komsta and Novomestky, 2015) skewness and kurtosis were calculated (Table 22). The Wilcoxon Rank Sum Test (unpaired data) was used to test for differences between two subgroups. To be able to compare different subgroups, a subsample (1000 repeats) of the respective larger subgroup was taken to compare equal sample sizes.




Figure 2: Scatterplot of carapace length over time, indicating insular (triangle) and continental (circles) and colour indicating continents. Lines indicate stratigraphic stages which were used as time bins, the dashed line is the border between the two stages of the Lower Miocene, which were consideres as one time bin.

#### 2.3.2. Body size trends over time

To investigate trends in body size over time, the R package paleoTS (Hunt, 2015) was used. Data were split into time bins according to stratigraphic stages (Table 1, Fig. 2), although the two stages of the Lower Miocene were considered as one time bin, because the last bin otherwise would have contained only 2 data records. To prevent sampling bias and because Sampling Accumulation Curves showed that the genus level was well sampled in contrast to species level, the mean SCL per genus was calculated before the timescale analysis. The

paleoTS plots were created, wich display the mean trait over time and can be fitted to different evolutionary models: stasis, which ...., generalized random walk (GRW), which .... or unbiased random walk (URW), which..... The Akaike Weight Criterion (AICc) indicates which model is best supported -> see Catalina's Paper and Hunt's papers

## 3. Results




Figure 3: Map displaying all localities for which body size data for testudinids was available in the literature. Size of points denotes sample size, color denotes approximate age.

Table 1: Time ranges, mean age per bin, corresponding stratigraphic stages and epochs, and respective sample sizes (on individual, species and genus level).)

96114316461):)						
Age Range [mya] Mean Age [mya]	Mean Age [mya]	Stages	Epochs	n (Individuals)	n (Species) n (Genera)	n (Genera)
0 - 0.0117	0.00585	Modern	Modern	254	99	48
0.0117 - 0.126	0.06885	Upper Pleistocene	Upper Pleistocene	20	48	∞
0.126 - 0.781	0.45350	Middle Pleistocene	Middle Pleistocene	53	13	7
0.781 - 1.81	1.29350	Lower Pleistocene	Lower Pleistocene	22	27	12
1.81 - 2.59	2.19700	Gelasian	Lower Pleistocene	33	15	တ
2.59 - 3.6	3.09400	Piacencian	Upper Pliocene	24	15	10
3.6 - 5.33	4.46600	Zanclean	Lower Pliocene	31	17	∞
5.33 - 7.25	6.28900	Messinian	Upper Miocene	12	თ	9
7.25 - 11.6	9.42700	Tortonian	Upper Miocene	46	20	თ
11.6 - 13.8	12.71400	Serravallian	Middle Miocene	27	∞	9
13.8 - 16	14.89500	Langhian	Middle Miocene	18	<del>1</del>	თ
16 - 23	19.50000	Burdigalian/Aquitanian	Lower Miocene	31	15	O




Figure 4: Sampling Accumulation Curve of fossil genera per reference.




Figure 5: Body size distribution. Bimodally distributed.





Figure 6: Boxplots of mean CL per time bin, including mean and sd CL for each genus (as pointrange).



Wilcoxon Rank Sum Test (unpaired data):

modern < fossil (P =  $2.6739885 \times 10^{-7}$ )

Wilcoxon Rank Sum Test (unpaired data):

continental < insular (P =  $1.5123 \times 10^{-6}$ )

## 4. paleoTS analysis

#### 4.1. all (continental and insular)

Table 2: paleoTS object, all data

nn	vv	mm	tt
4	102306.64	401.9641	0.0000005
18	42607.58	314.1859	0.0058500
8	64620.11	506.3265	0.0688500
7	155241.85	516.4053	0.4535000
12	147507.20	593.8669	1.2935000
8	580540.76	971.8850	2.1970000
9	271043.73	658.0826	3.0940000
8	187937.61	785.0792	4.4660000
4	584378.85	1141.9375	6.2890000
9	195766.19	703.9570	9.4270000
6	285258.36	628.3020	12.7140000
7	169914.58	687.9619	14.8950000

tt	mm	VV	nn
19.5000000	441.5420	78467.65	9




Figure 7: paleoTS plot with genus mean, all

Table 3: Model-fitting results for testudinidae, genera, all

	logL	K	AICc	Akaike.wt
GRW	-81.31790	2	167.9691	0.161
URW	-82.05721	1	166.5144	0.332
Stasis	-80.16802	2	165.6694	0.507

## 4.2. continental (excluding insular species)

Table 4: paleoTS object, continental

tt	mm	vv	nn
0.0000005	233.1680	8331.753	3
0.0058500	241.7917	13004.928	15
0.0688500	397.4606	50619.392	6
0.4535000	416.9341	200982.124	5
1.2935000	346.8484	66240.066	7
2.1970000	1103.1067	595507.933	7
3.0940000	725.4156	414253.291	6
4.4660000	771.3833	259173.082	6
6.2890000	1054.4375	531455.932	4
9.4270000	703.9570	195766.185	9
12.7140000	628.3020	285258.362	6
14.8950000	687.9619	169914.577	7
19.5000000	441.5420	78467.646	9




Figure 8: paleoTS plot with genus mean, continental

Table 5: Model-fitting results for testudinidae, genera, continental

	logL	K	AICc	Akaike.wt
GRW	-82.26287	2	169.8591	0.300
URW	-83.12577	1	168.6515	0.548
Stasis	-82.93984	2	171.2130	0.152

#### 4.3. insular (excluding continental)

Table 6: paleoTS object, insular

		-	$\overline{}$
tt	mm	vv	nn
0.000005	860.9268	0.00	1
0.0058500	379.5354	68570.44	12
0.0688500	727.5938	14997.58	4

tt	mm	vv	nn
0.4535000	748.8333	142649.08	3
1.2935000	829.6744	112964.44	6
2.1970000	1178.3333	821158.33	3
3.0940000	449.4375	27058.77	4
4.4660000	826.1667	15196.06	2
6.2890000	1850.0000	0.00	1




Figure 9: paleoTS plot with genus mean, insular

Table 7: Model-fitting results for testudinidae, genera, insular

	logL	K	AICc	Akaike.wt
GRW	-68.57344	2	143.5469	0
URW	-75.76576	1	154.1982	0
Stasis	-60.41581	2	127.2316	1

## 4.4. per continent

#### 4.4.1. Europe, genera

Table 8: paleoTS object, Europe

Table 6. paice to object, Europe				
tt	vv	nn	mm	
0.00585	3338.406	2	148.8559	
0.06885	138802.333	3	616.6667	
0.45350	89203.953	3	377.8167	
1.29350	218431.974	5	697.3717	
2.19700	1110050.000	2	895.0000	
3.09400	39433.333	3	453.3333	
4.46600	159317.256	5	1215.8667	
6.28900	875495.281	2	838.3750	
9.42700	263434.389	6	800.0508	
12.71400	351634.528	5	653.9625	
14.89500	223154.375	5	772.0000	
19.50000	183706.682	5	533.8533	
	·			

Table 9: Model-fitting results for testudinidae, genera, Europe

	logL	K	AICc	Akaike.wt
GRW	-84.14010	2	173.7802	0.006
URW	-85.90727	1	174.2590	0.005
Stasis	-79.01365	2	163.5273	0.990




Figure 10: Genera, Europe

Table 10: paleoTs object, Europe, continental

mm	nn	vv	tt
149.5381	2	3450.8267	0.00585
187.0000	1	0.0000	0.06885
205.4750	2	198.0050	0.45350
204.9292	2	23.1767	1.29350
1420.0000	1	0.0000	2.19700
232.5000	1	0.0000	3.09400
1475.6667	3	57926.3333	4.46600
663.3750	2	473607.7812	6.28900
800.0508	6	263434.3893	9.42700
653.9625	5	351634.5281	12.71400
772.0000	5	223154.3750	14.89500
533.8533	5	183706.6821	19.50000

Table 11: Model-fitting results for testudinidae, genera, Europe, continental

	logL	K	AICc	Akaike.wt
GRW	-87.93137	2	181.3627	0.009
URW	-92.56882	1	187.5821	0.000
Stasis	-83.21073	2	171.9215	0.991

#### Europe, genera, continental

Table 12: paleoTs object, Europe, insular

mm	nn	vv	tt
187.5077	1	0.00	0.00585
831.5000	2	684.50	0.06885
722.5000	1	0.00	0.45350
835.0833	4	168423.36	1.29350
1005.0000	2	1462050.00	2.19700
451.6667	3	40558.33	3.09400
826.1667	2	15196.06	4.46600
1850.0000	1	0.00	6.28900

Table 13: Model-fitting results for testudinidae, genera, Europe, insular

	logL	K	AICc	Akaike.wt
GRW	-67.12192	2	141.2438	0.000
URW	-57.51634	1	117.8327	0.074
Stasis	-52.89638	2	112.7928	0.926

Europe, genera, insular




Figure 12: paleoTS, genera, Europe, insular

#### 4.4.2. Eurasia, genera

Table 14: paleoTS object, Eurasia

	•	•	
nn	vv	mm	tt
1	0.000	137.2637	0.0000005
5	9760.467	236.8217	0.0058500
4	122579.333	530.0000	0.0688500
3	89203.953	377.8167	0.4535000
7	162641.142	777.5579	1.2935000
5	562217.222	909.6667	2.1970000
5	381770.000	892.0000	3.0940000
6	296417.219	1048.0556	4.4660000
3	849651.021	1208.9167	6.2890000
6	263434.389	800.0508	9.4270000
5	351634.528	653.9625	12.7140000

tt	mm	vv	nn
14.8950000	772.0000	223154.375	5
19.5000000	513.8533	162399.349	5




Figure 13: paleoTS, genera, Eurasia

Table 15: Model-fitting results for testudinidae, genera, Eurasia

	logL	K	AICc	Akaike.wt
GRW	-85.25195	2	175.8372	0.149
URW	-85.39072	1	173.1814	0.562
Stasis	-84.58890	2	174.5111	0.289

#### 4.4.3. Eurasia, genera, continental

Table 16: paleoTS object, Eurasia, continental

tt	mm	vv	nn
0.0000005	137.2637	0.000	1
0.0058500	238.0120	9654.865	5
0.0688500	228.5000	3444.500	2
0.4535000	205.4750	198.005	2
1.2935000	595.5388	191487.404	4
2.1970000	1044.5833	442006.250	4
3.0940000	1110.8333	581102.083	3
4.4660000	1159.0000	439728.667	4
6.2890000	1092.2500	788605.188	3
9.4270000	800.0508	263434.389	6
12.7140000	653.9625	351634.528	5
14.8950000	772.0000	223154.375	5
19.5000000	513.8533	162399.349	5

Table 17: Model-fitting results for testudinidae, genera, Eurasia, continental

	logL	K	AICc	Akaike.wt
GRW	-82.20698	2	169.7473	0.222
URW	-82.42344	1	167.2469	0.776
Stasis	-87.19538	2	179.7241	0.002




Figure 14: paleoTS, genera, Eurasia, continental

#### 4.4.4. Eurasia, genera, insular

Table 18: paleoTS object, Eurasia, insular

tt	mm	vv	nn
0.0000005	137.2637	0.000	1
0.0058500	271.4596	5668.485	4
0.0688500	644.3333	105436.333	3
0.4535000	722.5000	0.000	1
1.2935000	882.0356	105684.077	6
2.1970000	953.6667	652233.889	5
3.0940000	891.0000	383430.000	5
4.4660000	620.4444	134562.926	3
6.2890000	1900.0000	5000.000	2
19.5000000	800.0000	0.000	1




Figure 15: paleoTS, genera, Eurasia, insular

Table 19: Model-fitting results for testudinidae, genera, Eurasia, insular

	logL	K	AICc	Akaike.wt
GRW	-69.56419	2	145.1284	0.193
URW	-71.67437	1	145.9202	0.130
Stasis	-68.31026	2	142.6205	0.677

#### 5. References

- Böhme, M. and Ilg, A. (2003). FosFARbase. *Available at www. wahre- staerke. com/. Accessed October*, 10:2011.
- Brattstrom, B. H. (1961). Some New Fossil Tortoises from Western North America with remarks on the Zoogeography and Paleoecology of Tortoises. *Journal of Paleontology*, 35(3):543–560.
- Franz, R., Carlson, L. A., Owen, R. D. R. D., and Steadman, D. (2001). Fossil tortoises from the Turks and Caicos Islands, BWI. In *Proceedings of the 8th Symposium on the Natural History of the Bahamas. Gerace Research Center, San Salvador, Bahamas*, pages 27–31.
- Hunt, G. (2015). paleoTS: Analyze Paleontological Time-Series.
- Hutterer, R., García-Talavera, F., López-Martínez, N., Michaux, J., Hutterer, F., García-Talavera, F., López-Martínez, N., and Michaux, J. (1998). New chelonian eggs fom the tertiary of Lanzarote and Fuerteventura and a review of fossil tortoises of the Canary Islands (Reptilia, Testudinidae).
- Komsta, L. and Novomestky, F. (2015). *moments: Moments, cumulants, skewness, kurtosis and related tests.*
- Oksanen, J., Blanchet, F. G., Friendly, M., Kindt, R., Legendre, P., McGlinn, D., Minchin, P. R., O'Hara, R. B., Simpson, G. L., Solymos, P., Stevens, M. H. H., Szoecs, E., and Wagner, H. (2017). *vegan: Community Ecology Package*.
- R Core Team (2017). *R: A Language and Environment for Statistical Computing*. R Foundation for Statistical Computing, Vienna, Austria.
- Rhodin, A. G. J., Thomson, S., Georgalis, G. L., Karl, H.-V. V., Danilov, I. G., Takahashi, A., de la Fuente, M. S., Bourque, J. R., Delfino, M., Bour, R., Iverson, J., Shaffer, B., van Dijk, P. P., and Others (2015). Turtles and tortoises of the world during the rise and global spread of humanity: first checklist and review of extinct Pleistocene and Holocene chelonians. *Chelonian Research Monographs*, 5(April):1–66.
- Wickham, H. (2009). ggplot2: elegant graphics for data analysis. Springer New York.

Wickham, H., Francois, R., Henry, L., and Müller, K. (2017). *dplyr: A Grammar of Data Manipulation*.

#### **Appendix A Sampling Accumulation Curves**




Figure 16: Sampling Accumulation Curves: (a) - (b) Species are not sufficiently sampled, regardless of sampling unit. (c) - (i) Sampling Accumulation Curves on generic level per continent. Only Europe (h) and Eurasia (i) are sufficiently sampled.

#### **Appendix B** Histograms




Figure 17: Visual test for normal distribution. In case of normally distributed data, the black circles should follow the red line, which is not the case for either raw data (a) nor logtransformed data (b). Therefore, data is not normally distributed.




Figure 18: Histograms for several subgroups of the dataset.

## **Appendix C** Boxplots




Figure 19: Boxplots fossil vs. modern, continental vs. insular species.

```
Wilcoxon Rank Sum Test (unpaired data):
```

modern continental < fossil continental (P =  $4.8532266 \times 10^{-8}$ )

modern insular < fossil insular (P = 0.0018564)

Kruskal-Wallis-Test:

Continent means differ (P =  $1.0833256 \times 10^{-6}$ ) (still have to look into the details. . . )




Figure 20: Boxplot: body size on different continents, genera summarised

## Appendix D Random Sampling




Figure 21: Random sampling for several subgroups. For (a), (c), and (d) the random sample reflects the real sample, for (b) this is not the case.




Figure 22: Random sampling for different continents. All random samples reflect the real sample.

## Appendix E Tables

Table 20: Overview over genera (modern and fossil) per time bin, with sample sizes and mean CL.

EpochBins	Genus	n	meanCL
Modern	Aldabrachelys	12	974.5833
Modern	Astrochelys	14	366.2143
Modern	Centrochelys	3	493.3333
Modern	Chelonoidis	45	531.5178
Modern	Chersina	15	176.2667
Modern	Cylindraspis	5	724.0000
Modern	Geochelone	8	252.1250
Modern	Gopherus	23	302.4839
Modern	Hesperotestudo	1	250.0000
Modern	Homopus	7	139.2857
Modern	Indotestudo	16	242.9875
Modern	Kinixys	15	213.0667
Modern	Malacochersus	2	166.5000
Modern	Manouria	9	380.7778
Modern	Psammobates	17	113.4118
Modern	Pyxis	16	124.1875
Modern	Stigmochelys	6	405.3333
Modern	Testudo	39	197.5436
Upper Pleistocene	Centrochelys	1	850.0000
Upper Pleistocene	Chelonoidis	11	693.1818
Upper Pleistocene	Eurotestudo	1	187.0000
Upper Pleistocene	gen.	1	813.0000
Upper Pleistocene	Geochelone	2	475.0000
Upper Pleistocene	Gopherus	22	294.1545
Upper Pleistocene	Hesperotestudo	10	468.2760
Upper Pleistocene	Indotestudo	1	270.0000
Middle Pleistocene	Centrochelys	4	722.5000
Middle Pleistocene	Chelonoidis	1	1139.0000

EpochBins	Genus	n	meanCL
Middle Pleistocene	Eurotestudo	4	195.5250
Middle Pleistocene	Geochelone	1	170.0000
Middle Pleistocene	Gopherus	33	307.0721
Middle Pleistocene	Hesperotestudo	5	882.0000
Middle Pleistocene	Testudo	5	198.7400
Lower Pleistocene	Centrochelys	4	762.5000
Lower Pleistocene	Cheirogaster	2	857.0000
Lower Pleistocene	Chelonoidis	3	716.6667
Lower Pleistocene	Eurotestudo	4	201.5250
Lower Pleistocene	gen.	1	900.0000
Lower Pleistocene	Geochelone	1	340.0000
Lower Pleistocene	Gopherus	13	316.8077
Lower Pleistocene	Hesperotestudo	16	323.0562
Lower Pleistocene	Megalochelys	5	1041.8800
Lower Pleistocene	Psammobates	1	107.8000
Lower Pleistocene	Testudo	6	259.1667
Lower Pleistocene	Titanochelon	1	1300.0000
Gelasian	Centrochelys	1	2050.0000
Gelasian	Eurotestudo	1	150.0000
Gelasian	Gopherus	15	185.7467
Gelasian	Hesperotestudo	2	1000.0000
Gelasian	Manouria	1	900.0000
Gelasian	Megalochelys	3	1683.3333
Gelasian	Testudo	6	166.0000
Gelasian	Titanochelon	2	1640.0000
Piacencian	Aldabrachelys	3	1333.3333
Piacencian	Centrochelys	1	610.0000
Piacencian	Chelonoidis	4	442.7500
Piacencian	Gopherus	1	885.5000
Piacencian	Hesperotestudo	5	211.1600

EpochBins	Genus	n	meanCL
Piacencian	Homopus	1	90.0000
Piacencian	Megalochelys	2	1600.0000
Piacencian	Testudo	3	230.0000
Piacencian	Titanochelon	1	520.0000
Zanclean	Caudochelys	2	805.5000
Zanclean	Centrochelys	3	913.3333
Zanclean	Cheirogaster	1	739.0000
Zanclean	Ergilemys	2	209.0000
Zanclean	Geochelone	6	741.0000
Zanclean	Hesperotestudo	1	195.8000
Zanclean	Testudo	5	1377.0000
Zanclean	Titanochelon	6	1300.0000
Messinian	Hesperotestudo	2	941.0000
Messinian	Megalochelys	2	1950.0000
Messinian	Testudo	4	176.7500
Messinian	Titanochelon	2	1500.0000
Tortonian	"Hadrianus"	1	1000.0000
Tortonian	Cheirogaster	3	1288.3333
Tortonian	gen.	3	660.0000
Tortonian	Geochelone	3	741.3333
Tortonian	Gopherus	6	354.0000
Tortonian	Hesperotestudo	4	439.9750
Tortonian	Paleotestudo	3	233.6667
Tortonian	Testudo	20	218.3050
Tortonian	Titanochelon	2	1400.0000
Serravallian	Cheirogaster	2	1250.0000
Serravallian	gen.	1	270.0000
Serravallian	Gopherus	1	500.0000
Serravallian	Paleotestudo	19	206.5789
Serravallian	Testudo	3	190.2333

EpochBins	Genus	n	meanCL
Serravallian	Titanochelon	1	1353.0000
Langhian	Caudochelys	1	339.9000
Langhian	Chelonoidis	3	553.3333
Langhian	Ergilemys	1	1000.0000
Langhian	gen.	1	1000.0000
Langhian	Paleotestudo	2	272.5000
Langhian	Testudo	2	337.5000
Langhian	Titanochelon	4	1312.5000
Burdigalian/Aquitanian	Caudochelys	1	334.0000
Burdigalian/Aquitanian	gen.	1	270.0000
Burdigalian/Aquitanian	Geochelone	4	652.5000
Burdigalian/Aquitanian	Impregnochelys	1	620.0000
Burdigalian/Aquitanian	Mesocherus	5	180.0000
Burdigalian/Aquitanian	Namibchersus	9	518.1111
Burdigalian/Aquitanian	Paleotestudo	2	146.1500
Burdigalian/Aquitanian	Testudo	6	252.1167
Burdigalian/Aquitanian	Titanochelon	1	1001.0000

Table 21: General overview over genera, with sample sizes and mean CL.

Genus	n	meanCL
"Hadrianus"	1	1000.0000
Aldabrachelys	15	1046.3333
Astrochelys	14	366.2143
Caudochelys	4	571.2250
Centrochelys	17	804.1176
Cheirogaster	8	1102.2500
Chelonoidis	67	571.0940
Chersina	15	176.2667
Cylindraspis	5	724.0000
Ergilemys	3	472.6667

Genus	n	meanCL
Eurotestudo	10	192.5200
gen.	8	654.1250
Geochelone	25	510.2800
Gopherus	114	298.0361
Hesperotestudo	46	465.3296
Homopus	8	133.1250
Impregnochelys	1	620.0000
Indotestudo	17	244.5765
Kinixys	15	213.0667
Malacochersus	2	166.5000
Manouria	10	432.7000
Megalochelys	12	1446.6167
Mesocherus	5	180.0000
Namibchersus	9	518.1111
Paleotestudo	26	210.1269
Psammobates	18	113.1000
Pyxis	16	124.1875
Stigmochelys	6	405.3333
Testudo	99	269.2465
Titanochelon	20	1315.2000

Table 22: General statistics of body size data: all, per time bin, insular and continental, per continent (all referring to CL: min, max, variance, mean, logmean, median, logmedian, skewness, logskewness, kurosis, logkurtosis

Variable	all	Modern	Upper Pleistocene	Middle Pleistocene	Lower Pleistocene	Gelasian	Piacencian	Zanclean	Messinian	Tortonian	Serravallian	Langhian	Burdigalian/Aquitanian	Modern	Fossil	continental	insular
logku	2.73	2.69	2.56	5.52	2.40	2.11	1.99	2.30	1.29	2.54	5.30	1.18	2.26	2.69	2.42	3.74	2.05
kurt	8.00	5.87	3.61	12.24	5.76	3.12	2.50	3.56	1.49	3.74	7.77	1.55	3.45	5.87	6.11	10.57	3.95
logsk	69.0	0.58	0.24	1.52	0.73	0.93	0.14	-0.40	-0.22	0.81	1.77	0.03	0.45	0.58	0.68	1.07	-0.27
skew	2.14	1.83	1.20	3.03	1.74	1.31	1.00	1.1	0.26	1.49	2.49	0.30	1.27	1.83	1.83	2.65	1.02
logmed	2.4	2.4	2.5	2.5	2.4	2.3	2.6	2.9	2.9	2.4	2.3	2.8	2.5	2.4	2.5	2.4	2.7
med	270.5	242.0	334.7	292.9	263.0	194.9	428.0	857.5	916.0	250.0	220.0	700.0	302.4	242.0	285.6	250.0	500.0
logm	2.5	2.4	2.6	2.5	2.5	2.5	2.6	2.9	2.8	2.5	2.4	2.8	2.5	2.4	2.6	2.5	2.6
mean	437.2	330.3	445.9	387.1	463.5	555.2	610.6	955.2	948.9	462.7	337.7	747.9	406.8	330.3	511.7	392.9	578.5
var	164537.80	67485.50	99.06969	97910.83	161948.82	411224.51	270535.82	476162.71	602611.21	175470.12	126060.40	230451.33	76288.76	67485.50	219004.66	157808.79	160834.35
max	2500	1300	1250	1800	2000	2050	1600	2500	2100	1540	1500	1600	1100	1300	2500	2500	2000
min	80.00	80.00	102.44	132.00	107.80	118.90	90.00	176.00	140.00	107.00	111.00	270.00	113.00	80.00	90.00	81.00	80.00
nCL	616	253	49	53	22	31	21	26	10	45	27	<del>1</del>	30	253	363	469	147

e	n-con	n-ins	no:	SU		Я		ď
kurt logku Variable	modern-con	modern-ins	fossil-con	fossil-ins	Africa	America	Asia	Europe
logku	2.98	1.77	2.96	3.18	2.48	2.91	2.24	2.34
	8.09	2.47	7.25	4.02	7.97	6.79	3.61	6.30
logsk	0.29	0.01	0.96	-0.40	0.68	0.75	0.85	2.4 1.86 0.81
skew	1.92	0.82	2.11	1.1	2.10	1.92	1.43	1.86
med logmed skew logsk	2.3	2.5	2.4	2.9	2.3	2.5	2.4	
	221.0	353.0	270.0	750.0	193.5	302.2	280.0	245.0
logm	2.3	2.6	2.5	2.8	2.4	2.5	2.6	2.5
mean	244.0	471.5	467.9	780.0	347.7	415.0	585.5	491.2
var	17009.02	118641.09	212116.79	180825.40	112417.26	82209.71	2100 323123.20	173 107.00 2500 254222.84
min max	830	1300	2500	2000	2050	1800		2500
min	81.00	80.00	90.00	150.00	80.00	102.44	150.00	107.00
nCL	157	96	312	21	142	242	29	173

≓
SS
S
.0
Ψ-
ئيد
ŝ
S
ന
ata
Œ
$\Box$
က
$\alpha$
4
<u>_</u>
Д
.ਜ
-

Genus	Taxon	CL	estimated	EpochBins	Age	Age Island	Con
Astrochelys	Astrochelys radiata	395.00	Ε	Modern	0.000001	>	Africa
Kinixys	Kinixys belliana	162.00	E	Modern	0.000001	드	Africa
Psammobates	Psammobates geometricus	107.00	E	Modern	0.000001	_	Africa
Kinixys	Kinixys belliana	157.00	E	Modern	0.000001	_	Africa
Aldabrachelys	Aldabrachelys gigantea	870.00	E	Modern	0.000001	>	Africa
Kinixys	Kinixys belliana	174.00	Ε	Modern	0.000001	_	Africa
Stigmochelys	Stigmochelys pardalis	345.00	Ε	Modern	0.000001	_	Africa
Psammobates	Psammobates geometricus	92.00	Ε	Modern	0.000001	_	Africa
Chersina	Chersina angulata	179.30	E	Modern	0.000001	_	Africa
Chersina	Chersina angulata	170.00	Ε	Modern	0.000001	_	Africa
Testudo	Testudo kleinmanni	144.00	Ε	Modern	0.000001	_	Africa
Malacochersus	Malacochersus tornieri	153.00	Ε	Modern	0.000001	_	Africa
Psammobates	Psammobates oculifer	119.00	٤	Modern	0.000001	_	Africa
Kinixys	Kinixys homeana	193.00	Ε	Modern	0.000001	_	Africa
Cylindraspis	Cylindraspis vosmaeri	500.00	Ε	Modern	0.000001	>	Africa
Homopus	Homopus aerolatus	88.00	٤	Modern	0.000001	<b>_</b>	Africa
Stigmochelys	Stigmochelys pardalis	405.00	٤	Modern	0.000001	<b>C</b>	Africa
Chersina	Chersina angulata	162.00	٤	Modern	0.000001	L	Africa

Genus	Taxon	CL	estimated	EpochBins	Age	Island	Con
Kinixys	Kinixys belliana	180.00	Ε	Modern	0.000001	C	Africa
Astrochelys	Astrochelys radiata	285.00	E	Modern	0.000001	>	Africa
Kinixys	Kinixys erosa	400.00	Ε	Modern	0.000001	⊑	Africa
Astrochelys	Astrochelys radiata	242.00	E	Modern	0.000001	>	Africa
Aldabrachelys	Aldabrachelys gigantea	810.00	Ε	Modern	0.000001	>	Africa
Pyxis	Pyxis planicauda	126.00	E	Modern	0.000001	>	Africa
Cylindraspis	Cylindraspis indica	00.009	٤	Modern	0.000001	>	Africa
Psammobates	Psammobates tentorius	111.00	E	Modern	0.000001	⊑	Africa
Kinixys	Kinixys erosa	164.00	E	Modern	0.000001	⊑	Africa
Kinixys	Kinixys erosa	271.00	Ε	Modern	0.000001	⊑	Africa
Indotestudo	Indotestudo travancorica	224.00	E	Modern	0.000001	⊑	Africa
Psammobates	Psammobates oculifer	101.00	Ε	Modern	0.000001	⊆	Africa
Homopus	Homopus signatus	94.00	Ε	Modern	0.000001	<b>_</b>	Africa
Kinixys	Kinixys belliana	194.00	E	Modern	0.000001	⊑	Africa
Kinixys	Kinixys belliana	230.00	Ε	Modern	0.000001	<b>_</b>	Africa
Stigmochelys	Stigmochelys pardalis	720.00	Ε	Modern	0.000001	⊆	Africa
Kinixys	Kinixys homeana	223.00	Ε	Modern	0.000001	<b>_</b>	Africa
Kinixys	Kinixys lobatsiana	200.00	E	Modern	0.000001	<b>C</b>	Africa
Kinixys	Kinixys natalensis	160.00	Е	Modern	0.000001	C	Africa

Genus	Taxon	CL	estimated	EpochBins	Age	Island	Con
Chersina	Chersina angulata	202.00	Ε	Modern	0.000001	_	Africa
Chersina	Chersina angulata	351.00	Ε	Modern	0.000001	>	Africa
Homopus	Homopus femoralis	168.00	Ε	Modern	0.000001	C	Africa
Centrochelys	Centrochelys sulcata	215.00	Ε	Modern	0.000001	_	Africa
Astrochelys	Astrochelys yniphora	307.00	Ε	Modern	0.000001	>	Africa
Chersina	Chersina angulata	181.00	Ε	Modern	0.000001	_	Africa
Psammobates	Psammobates tentorius	145.00	Ε	Modern	0.000001	_	Africa
Stigmochelys	Stigmochelys pardalis	315.00	٤	Modern	0.000001	_	Africa
Pyxis	Pyxis planicauda	160.00	Ε	Modern	0.000001	>	Africa
Psammobates	Psammobates antiquorum	107.80	Ε	Lower Pleistocene	1.800000	_	Africa
Stigmochelys	Stigmochelys pardalis	350.00	Ε	Modern	0.000001	_	Africa
Aldabrachelys	Aldabrachelys abrupta	1000.00	шо	Modern	0.002000	>	Africa
Chersina	Chersina angulata	181.90	Ε	Modern	0.000001	>	Africa
Psammobates	Psammobates tentorius	116.00	Ε	Modern	0.000001	>	Africa
Astrochelys	Astrochelys yniphora	415.00	Ε	Modern	0.000001	>	Africa
Aldabrachelys	Aldabrachelys gigantea	770.00	E	Modern	0.000001	>	Africa
Chersina	Chersina angulata	160.00	E	Modern	0.000001	_	Africa
Chersina	Chersina angulata	148.00	E	Modern	0.000001	<b>C</b>	Africa
Aldabrachelys	Aldabrachelys gigantea	720.00	E	Modern	0.000001	>	Africa

Genus	Taxon	CL	estimated	EpochBins	Age	Island	Con
Astrochelys	Astrochelys yniphora	426.00	Ε	Modern	0.000001	>	Africa
Astrochelys	Astrochelys radiata	334.00	E	Modern	0.000001	>	Africa
Centrochelys	Centrochelys sulcata	830.00	E	Modern	0.000001	_	Africa
Pyxis	Pyxis arachnoides	144.00	٤	Modern	0.000001	>	Africa
Pyxis	Pyxis arachnoides	86.00	E	Modern	0.000001	>	Africa
Pyxis	Pyxis arachnoides	154.00	E	Modern	0.000001	>	Africa
Pyxis	Pyxis arachnoides	110.00	E	Modern	0.000001	>	Africa
Namibchersus	Namibchersus namaquensis	254.00	Ε	Burdigalian/Aquitanian	18.000000	⊑	Africa
Pyxis	Pyxis planicauda	132.00	Ε	Modern	0.000001	>	Africa
Homopus	Homopus boulengeri	110.00	Ε	Modern	0.000001	_	Africa
Pyxis	Pyxis planicauda	134.00	Ε	Modern	0.000001	>	Africa
Pyxis	Pyxis planicauda	120.00	Ε	Modern	0.000001	>	Africa
Homopus	Homopus solus	109.00	Ε	Modern	0.000001	<b>_</b>	Africa
Centrochelys	Centrochelys sulcata	435.00	E	Modern	0.000001	⊑	Africa
Pyxis	Pyxis arachnoides	110.00	Ε	Modern	0.000001	>	Africa
Pyxis	Pyxis arachnoides	80.00	Ε	Modern	0.000001	>	Africa
Astrochelys	Astrochelys radiata	305.00	Ε	Modern	0.000001	>	Africa
Stigmochelys	Stigmochelys pardalis	297.00	E	Modern	0.000001	<b>_</b>	Africa
Namibchersus	Namibchersus aff. namaquensis	1100.00	mo	Burdigalian/Aquitanian	17.250000	<b>_</b>	Africa

Genus	Taxon	CL	estimated	EpochBins	Age	Island	Con
Aldabrachelys	Aldabrachelys gigantea	875.00	Ε	Modern	0.000001	>	Africa
Namibchersus	Namibchersus aff. namaquensis	550.00	шо	Burdigalian/Aquitanian	17.250000	⊏	Africa
Chersina	Chersina angulata	166.40	E	Modern	0.000001	⊑	Africa
Chersina	Chersina angulata	171.60	E	Modern	0.000001	>	Africa
Chersina	Chersina angulata	136.00	E	Modern	0.000001	⊏	Africa
Geochelone	Geochelone stromeri	425.00	E	Zanclean	4.466000	⊑	Africa
Testudo	Testudo sp.	184.00	mf	Gelasian	2.500000	⊑	Africa
Geochelone	Geochelone stromeri	350.00	Ε	Zanclean	4.466000	_	Africa
Namibchersus	Namibchersus namaquensis	264.00	٤	Burdigalian/Aquitanian	19.500000	⊆	Africa
Pyxis	Pyxis arachnoides	150.00	Ε	Modern	0.000001	>	Africa
Psammobates	Psammobates oculifer	103.00	٤	Modern	0.000001	⊑	Africa
Psammobates	Psammobates oculifer	105.00	٤	Modern	0.000001	<b>_</b>	Africa
Psammobates	Psammobates geometricus	118.00	Ε	Modern	0.000001	<b>_</b>	Africa
Psammobates	Psammobates geometricus	105.00	٤	Modern	0.000001	⊑	Africa
Testudo	Testudo oughlamensis	120.00	шо	Gelasian	2.500000	⊑	Africa
Astrochelys	Astrochelys radiata	355.00	٤	Modern	0.000001	>	Africa
Aldabrachelys	Aldabrachelys gigantea	800.00	٤	Modern	0.000001	>	Africa
Namibchersus	Namibchersus aff. namaquensis	440.00	шо	Burdigalian/Aquitanian	17.250000	<b>_</b>	Africa
Chersina	Chersina angulata	153.50	Ε	Modern	0.000001	⊑	Africa

Genus	Taxon	CL	estimated	EpochBins	Age	Island	Con
Cylindraspis	Cylindraspis triserrata	1100.00	Ε	Modern	0.000001	>	Africa
Astrochelys	Astrochelys yniphora	486.00	Ε	Modern	0.000001	>	Africa
Chersina	Chersina angulata	161.30	Ε	Modern	0.000001	>	Africa
Aldabrachelys	"Aldabrachelys" laetoliensis	1000.00	шо	Piacencian	2.703000	ㄷ	Africa
Geochelone	Geochelone sp.	1446.00	eh	Tortonian	8.476000	ᄓ	Africa
Kinixys	Kinixys sp.	268.00	ef	Modern	0.009500	C	Africa
Aldabrachelys	Aldabrachelys grandidieri	1240.00	Ε	Modern	0.001500	>	Africa
Testudo	Testudo aff. kenitrensis	142.00	m	Gelasian	2.500000	C	Africa
Testudo	Testudo sp.	200.00	m	Gelasian	2.500000	C	Africa
Aldabrachelys	Aldabrachelys gigantea	1190.00	Ε	Modern	0.000001	>	Africa
Psammobates	Psammobates tentorius	95.00	Ε	Modern	0.000001	C	Africa
Psammobates	Psammobates tentorius	81.00	Ε	Modern	0.000001	C	Africa
Pyxis	Pyxis planicauda	114.00	E	Modern	0.000001	>	Africa
Mesocherus	Mesocherus orangeus	160.00	ОШ	Burdigalian/Aquitanian	17.250000	<b>C</b>	Africa
Pyxis	Pyxis planicauda	148.00	E	Modern	0.000001	>	Africa
Psammobates	Psammobates oculifer	111.00	E	Modern	0.000001	<b>C</b>	Africa
Geochelone	Geochelone crassa	865.00	m	Zanclean	4.145000	_	Africa
Pyxis	Pyxis arachnoides	111.00	E	Modern	0.000001	>	Africa
Impregnochelys	Impregnochelys pachytectis	620.00	Ε	Burdigalian/Aquitanian	19.500000	C	Africa

Genus	Taxon	CL	estimated	EpochBins	Age	Island	Con
Mesocherus	Mesocherus orangeus	200.00	шо	Burdigalian/Aquitanian	17.250000	_	Africa
Namibchersus	Namibchersus namaquensis	815.00	E	Burdigalian/Aquitanian	18.000000	Ц	Africa
Chersina	Chersina angulata	120.00	E	Modern	0.000001	Ц	Africa
Namibchersus	Namibchersus namaquensis	300.00	E	Burdigalian/Aquitanian	19.500000	ᄆ	Africa
Aldabrachelys	Aldabrachelys gigantea	1140.00	E	Modern	0.000001	>	Africa
Astrochelys	Astrochelys radiata	400.00	E	Modern	0.000001	>	Africa
Aldabrachelys	Aldabrachelys grandidieri	1250.00	шо	Modern	0.001500	>	Africa
Astrochelys	Astrochelys yniphora	446.00	E	Modern	0.000001	>	Africa
Cylindraspis	Cylindraspis peltastes	420.00	E	Modern	0.000001	>	Africa
Psammobates	Psammobates geometricus	165.00	E	Modern	0.000001	C	Africa
Mesocherus	Mesocherus orangeus	180.00	шо	Burdigalian/Aquitanian	17.250000	۵	Africa
Psammobates	Psammobates oculifer	147.00	E	Modern	0.000001	C	Africa
Cylindraspis	Cylindraspis inepta	1000.00	Ε	Modern	0.000001	>	Africa
Centrochelys	Centrochelys atlantica	400.00	шо	Lower Pleistocene	1.300000	>	Africa
Aldabrachelys	Aldabrachelys gigantea	1030.00	E	Modern	0.000001	>	Africa
Homopus	Homopus aerolatus	300.00	E	Modern	0.000001	ᄆ	Africa
Psammobates	Psammobates oculifer	107.00	Ε	Modern	0.000001	۵	Africa
Namibchersus	Namibchersus namaquensis	470.00	E	Burdigalian/Aquitanian	18.000000	C	Africa
Astrochelys	Astrochelys yniphora	370.00	E	Modern	0.000001	>	Africa

Genus	Taxon	CL	estimated	EpochBins	Age	Island	Con
Centrochelys	Centrochelys marocana	2050.00	шо	Gelasian	2.500000	C	Africa
Kinixys	Kinixys spekii	220.00	E	Modern	0.000001	⊑	Africa
Homopus	Homopus fenestratus	90.00	шо	Piacencian	3.056500	⊑	Africa
Malacochersus	Malacochersus tornieri	180.00	E	Modern	0.000001	⊑	Africa
Homopus	Homopus signatus	106.00	E	Modern	0.000001	⊑	Africa
Mesocherus	Mesocherus orangeus	180.00	шо	Burdigalian/Aquitanian	17.250000	⊑	Africa
Testudo	Testudo kenitrensis	132.00	шо	Middle Pleistocene	0.453500	⊑	Africa
Mesocherus	Mesocherus orangeus	180.00	шо	Burdigalian/Aquitanian	17.250000	⊑	Africa
Astrochelys	Astrochelys yniphora	361.00	E	Modern	0.000001	>	Africa
Namibchersus	Namibchersus namaquensis	470.00	E	Burdigalian/Aquitanian	18.000000	⊑	Africa
Geochelone	Geochelone elegans	208.00	٤	Modern	0.000001	⊑	Asia
Geochelone	Geochelone elegans	245.00	E	Modern	0.000001	⊑	Asia
Geochelone	Geochelone elegans	221.00	E	Modern	0.000001	⊑	Asia
Geochelone	Geochelone elegans	220.00	٤	Modern	0.000001	>	Asia
Geochelone	Geochelone elegans	221.00	E	Modern	0.000001	⊑	Asia
Geochelone	Geochelone platynota	222.00	E	Modern	0.000001	⊑	Asia
Indotestudo	Indotestudo forstenii	202.00	Ε	Modern	0.000001	>	Asia
Megalochelys	Megalochelys sondaari	1000.00	Э	Lower Pleistocene	1.350000	>	Asia
Indotestudo	Indotestudo forstenii	309.00	E	Modern	0.000001	>	Asia

Genus	Taxon	CL	estimated	EpochBins	Age	Island	Con
Megalochelys	Megalochelys atlas	1650.00	ОШ	Gelasian	2.000000	>	Asia
Indotestudo	Indotestudo forstenii	199.00	٤	Modern	0.000001	>	Asia
Indotestudo	Indotestudo elongata	244.20	٤	Modern	0.000001	_	Asia
Indotestudo	Indotestudo travancorica	244.20	٤	Modern	0.000001	_	Asia
Testudo	Testudo graeca	300.00	Ε	Modern	0.000001	C	Asia
Manouria	Manouria impressa	165.00	٤	Modern	0.000001	С	Asia
Indotestudo	Indotestudo elongata	276.00	٤	Modern	0.000001	_	Asia
Indotestudo	Indotestudo elongata	235.00	Ε	Modern	0.000001	_	Asia
Indotestudo	Indotestudo elongata	208.00	٤	Modern	0.000001	_	Asia
Indotestudo	Indotestudo elongata	166.00	٤	Modern	0.000001	С	Asia
Manouria	Manouria impressa	350.00	Ε	Modern	0.000001	_	Asia
Testudo	Testudo graeca	250.00	Ε	Modern	0.000001	_	Asia
Testudo	Testudo graeca	280.00	Ε	Modern	0.000001	>	Asia
Manouria	Manouria emys	212.00	Ε	Modern	0.000001	_	Asia
Manouria	Manouria emys	445.00	٤	Modern	0.000001	С	Asia
Manouria	Manouria emys	330.00	٤	Modern	0.000001	С	Asia
Megalochelys	Megalochelys atlas	2000.00	шо	Gelasian	2.190500	_	Asia
Testudo	Testudo changshanesis	330.00	ОШ	Lower Pleistocene	1.684500	_	Asia
Indotestudo	Indotestudo forstenii	200.50	E	Modern	0.000001	>	Asia

Genus	Taxon	CL	estimated	EpochBins	Age	Island	Con
Testudo	Testudo horsfieldii	280.00	٤	Modern	0.000001	_	Asia
Megalochelys	Megalochelys sondaari	818.00	oe ec	Lower Pleistocene	1.350000	>	Asia
Indotestudo	Indotestudo travancorica	249.70	Ε	Modern	0.000001	ᆮ	Asia
Manouria	Manouria punjabiensis	900.006	шо	Gelasian	2.190500	C	Asia
Megalochelys	Megalochelys sp.	1200.00	ev	Lower Pleistocene	0.900000	>	Asia
Indotestudo	Indotestudo elongata	270.00	Ε	Upper Pleistocene	0.037000	_	Asia
Ergilemys	Ergilemys oskarkuhni	220.00	Ε	Zanclean	3.950000	_	Asia
Megalochelys	Megalochelys atlas	1600.00	шо	Piacencian	3.094000	_	Asia
Geochelone	Geochelone platynota	300.00	Ε	Modern	0.000001	_	Asia
Aldabrachelys	Aldabrachelys ? sp.	1500.00	шо	Piacencian	3.000000	_	Asia
Indotestudo	Indotestudo travancorica	219.60	Ε	Modern	0.000001	_	Asia
Megalochelys	Megalochelys sp.	191.40	Ε	Lower Pleistocene	1.684500	>	Asia
Manouria	Manouria oyamai	450.00	шо	Modern	0.011000	>	Asia
Indotestudo	Indotestudo elongata	219.60	Ε	Modern	0.000001	_	Asia
Megalochelys	Megalochelys atlas	1800.00	Ε	Messinian	5.423000	_	Asia
Testudo	Testudo transcaucasia	150.00	шо	Gelasian	2.190500	_	Asia
Megalochelys	Megalochelys atlas	1600.00	шо	Piacencian	3.094000	_	Asia
Manouria	Manouria emys	00.009	Ε	Modern	0.000001	<b>C</b>	Asia
Indotestudo	Indotestudo travancorica	331.00	Ε	Modern	0.000001	_	Asia

Sing	Tavon	5	petimitad	EnochBins	ΔΩ	Puelal	000
	layor	3	collinated	מווס	S S S S S S S S S S S S S S S S S S S	2 2	
Geochelone	Geochelone sp.	800.00	ev	Burdigalian/Aquitanian	16.500000	⊆	Asia
Manouria	Manouria impressa	275.00	Ε	Modern	0.000001	⊑	Asia
Indotestudo	Indotestudo elongata	360.00	Ε	Modern	0.000001	⊑	Asia
Manouria	Manouria emys	00.009	٤	Modern	0.000001	⊑	Asia
Ergilemys	Ergilemys oskarkuhni	198.00	Ε	Zanclean	3.950000	⊑	Asia
Megalochelys	Megalochelys sp.	2000.00	٤	Lower Pleistocene	1.684500	>	Asia
Megalochelys	Megalochelys atlas	1400.00	шо	Gelasian	2.000000	>	Asia
Geochelone	Geochelone elegans	380.00	Ε	Modern	0.000001	<b>C</b>	Asia
gen.	gen. indet.	900.006	шо	Lower Pleistocene	1.684500	⊏	Asia
Testudo	Testudo ranovi	200.00	шо	Gelasian	2.190500	⊆	Asia
Aldabrachelys	Aldabrachelys ? sp.	1500.00	шо	Piacencian	3.000000	⊑	Asia
Megalochelys	Megalochelys atlas	2100.00	шо	Messinian	5.423000	<b>C</b>	Asia
Chelonoidis	Chelonoidis sp.	550.00	Ε	Modern	0.001000	>	America
Gopherus	Gopherus morafkai	299.00	Ε	Modern	0.000001	<b>C</b>	America
Hesperotestudo	Hesperotestudo bermudae	500.00	Ε	Middle Pleistocene	0.310000	>	America
Chelonoidis	Chelonoidis monensis	500.00	٤	Upper Pleistocene	0.064500	>	America
Chelonoidis	Chelonoidis alburyorum	453.00	٤	Piacencian	3.201500	>	America
Chelonoidis	Chelonoidis marcanoi	614.00	eh	Upper Pleistocene	0.069000	>	America
Chelonoidis	Chelonoidis marcanoi	767.00	eh	Upper Pleistocene	0.0690.0	>	America

Genus	Taxon	CL	estimated	EpochBins	Age	Island	Con
Gopherus	Gopherus flavomarginatus	450.00	Е	Lower Pleistocene	1.050000	u	America
Chelonoidis	Chelonoidis alburyorum	428.00	Ε	Piacencian	3.201500	>	America
Chelonoidis	Chelonoidis marcanoi	778.00	eh	Upper Pleistocene	0.0690.0	>	America
Chelonoidis	Chelonoidis sombrerensis	990.00	Ε	Upper Pleistocene	0.0690.0	>	America
Geochelone	Geochelone sp.	340.00	шо	Lower Pleistocene	1.050000	L	America
Hesperotestudo	Hesperotestudo sp.	1500.00	шо	Lower Pleistocene	0.966000	L	America
Gopherus	Gopherus flavomarginatus	400.00	E	Modern	0.000001	L	America
Chelonoidis	Chelonoidis alburyorum	466.00	٤	Piacencian	3.201500	>	America
Chelonoidis	Chelonoidis sp.	00.009	шо	Upper Pleistocene	0.0690.0	>	America
Chelonoidis	Chelonoidis sp.	400.00	шо	Upper Pleistocene	0.0690.0	>	America
Gopherus	Gopherus berlandieri	195.00	E	Lower Pleistocene	1.050000	C	America
Chelonoidis	Chelonoidis sp.	440.00	шо	Modern	0.001000	>	America
Chelonoidis	Chelonoidis marcanoi	530.00	eh	Upper Pleistocene	0.0690.0	>	America
Chelonoidis	Chelonoidis cubensis	1139.00	eĮ	Middle Pleistocene	0.393500	>	America
Chelonoidis	Chelonoidis sp.	800.008	шо	Lower Pleistocene	1.357000	>	America
Gopherus	Gopherus berlandieri	240.00	E	Modern	0.000001	C	America
Chelonoidis	Chelonoidis sp.	00.099	шо	Modern	0.001000	>	America
Chelonoidis	Chelonoidis sp.	512.00	шо	Modern	0.001000	>	America
Chelonoidis	Chelonoidis sp.	854.00	шо	Modern	0.001000	>	America

Genus	Taxon	CL	estimated	EpochBins	Age	Age Island	Con
Chelonoidis	Chelonoidis sp.	750.00	шо	Lower Pleistocene	1.357000	<b>&gt;</b>	America
Chelonoidis	Chelonoidis alburyorum	424.00	Ε	Piacencian	3.201500	>	America
Chelonoidis	Chelonoidis sp.	550.00	шо	Modern	0.001000	>	America
Gopherus	Gopherus donlaloi	580.00	шо	Modern	0.000175	C	America
Hesperotestudo	Hesperotestudo bermudae	270.00	E	Middle Pleistocene	0.310000	>	America
Gopherus	Gopherus berlandieri	256.30	E	Lower Pleistocene	1.050000	C	America
Chelonoidis	Chelonoidis sp.	00.009	шо	Lower Pleistocene	1.357000	>	America
Ergilemys	Ergilemys sp.	1000.00	E	Langhian	14.000000	C	Europe
Testudo	Testudo graeca	195.00	mf	Lower Pleistocene	1.770000	C	Europe
Eurotestudo	Eurotestudo aff. hermanni	194.70	mf	Middle Pleistocene	0.740000	C	Europe
Centrochelys	Centrochelys burchardi	940.00	шо	Middle Pleistocene	0.435000	>	Europe
Titanochelon	Titanochelon bacharidisi	1164.00	E	Zanclean	3.950000	C	Europe
Paleotestudo	Paleotestudo antiqua	159.50	٤	Serravallian	13.000000	C	Europe
Testudo	Testudo horsfieldii	111.00	E	Modern	0.000001	C	Europe
Testudo	Testudo marginata	210.00	E	Lower Pleistocene	1.720000	C	Europe
Testudo	Testudo graeca	178.20	E	Modern	0.000001	C	Europe
Testudo	Testudo graeca	200.00	mf	Messinian	5.500000	C	Europe
Testudo	Testudo lunellensis	260.70	mf	Middle Pleistocene	0.450000	C	Europe
Testudo	Testudo sp.	500.00	ОШ	Zanclean	3.900000	C	Europe

Genus	Taxon	CL	estimated	EpochBins	Age	Island	Con
Testudo	Testudo sp.	200.00	mf	Messinian	6.165000	u	Europe
Testudo	Testudo hermanni	143.50	Ε	Modern	0.000001	>	Europe
Pyxis	Pyxis arachnoides	108.00	Ε	Modern	0.000001	⊏	Europe
Eurotestudo	Eurotestudo hermanni	237.60	mf	Middle Pleistocene	0.600000	⊑	Europe
Testudo	Testudo marginata	246.00	Ε	Modern	0.000001	⊑	Europe
Paleotestudo	Paleotestudo sp.	179.30	E	Burdigalian/Aquitanian	16.550000	⊑	Europe
Centrochelys	Centrochelys burchardi	500.00	ОШ	Middle Pleistocene	0.435000	>	Europe
Testudo	Testudo graeca	167.00	Ε	Messinian	5.500000	⊑	Europe
Testudo	Testudo marginata	290.00	Ε	Modern	0.000001	⊑	Europe
Paleotestudo	Paleotestudo antiqua	191.00	mf	Serravallian	13.600000	⊑	Europe
Testudo	Testudo hermanni	130.00	Ε	Modern	0.000001	⊑	Europe
Testudo	Testudo hermanni	138.50	Ε	Modern	0.000001	⊑	Europe
Testudo	Testudo kalksburgensis	230.00	Ε	Burdigalian/Aquitanian	19.965000	⊑	Europe
Testudo	Testudo marginata	250.00	Ε	Modern	0.000001	>	Europe
Testudo	Testudo marginata	242.50	Ε	Modern	0.000001	>	Europe
Cheirogaster	Cheirogaster sp.	925.00	eĮ	Lower Pleistocene	0.965000	>	Europe
Testudo	Testudo marginata	246.00	Ε	Modern	0.000001	<b>_</b>	Europe
Testudo	Testudo horsfieldii	123.00	Ε	Modern	0.000001	⊑	Europe
Testudo	Testudo marginata	246.70	Ε	Modern	0.000001	c	Europe

Genus	Taxon	CL	estimated	EpochBins	Age	Age Island	Con
Testudo	Testudo marginata	241.70	٤	Modern	0.000001	<b>C</b>	Europe
Testudo	Testudo hermanni	195.00	٤	Modern	0.000001	>	Europe
Testudo	Testudo hermanni	250.00	Ε	Modern	0.000001	ᄆ	Europe
Paleotestudo	Paleotestudo antiqua	203.00	Ε	Serravallian	12.150000	ᄆ	Europe
Testudo	Testudo horsfieldii	114.00	Ε	Modern	0.000001	Ц	Europe
Testudo	Testudo horsfieldii	132.00	٤	Modern	0.000001	П	Europe
Centrochelys	Centrochelys robusta	1200.00	ev	Lower Pleistocene	1.300000	>	Europe
Testudo	Testudo hermanni	183.30	٤	Modern	0.000001	>	Europe
Testudo	Testudo hermanni	196.00	٤	Modern	0.000001	ᆮ	Europe
Testudo	Testudo hermanni	176.90	٤	Modern	0.000001	П	Europe
Titanochelon	Titanochelon bacharidisi	900.006	шо	Zanclean	3.950000	ᄆ	Europe
gen.	gen. indet.	1000.00	шо	Langhian	14.700000	ᆮ	Europe
gen.	gen. indet.	270.00	шо	Serravallian	12.200000	ㄷ	Europe
Paleotestudo	Paleotestudo cf. antiqua	113.00	mf	Burdigalian/Aquitanian	17.300000	ᆮ	Europe
Testudo	Testudo graeca	194.60	٤	Modern	0.000001	ㄷ	Europe
Testudo	Testudo lunellensis	231.00	ev	Middle Pleistocene	0.453500	ᆮ	Europe
Testudo	Testudo lunellensis	176.00	шо	Middle Pleistocene	0.453500	<b>_</b>	Europe
Testudo	Testudo hermanni	168.30	Ε	Modern	0.000001	>	Europe
Testudo	Testudo sp.	2500.00	mf	Zanclean	3.900000	ᄆ	Europe

Genus	Taxon	CL	estimated	EpochBins	Age	Age Island	Con
Testudo	Testudo burgenlandica	275.00	Ε	Tortonian	8.750000	C	Europe
Testudo	Testudo kalksburgensis	275.00	Ε	Langhian	14.500000	⊑	Europe
Titanochelon	Titanochelon bolivari	1150.00	Ε	Messinian	6.289000	⊑	Europe
Paleotestudo	Paleotestudo cf. sp.	270.00	шо	Langhian	14.700000	⊑	Europe
gen.	gen. indet.	880.00	Ε	Tortonian	8.750000	⊑	Europe
Eurotestudo	Eurotestudo globosa	263.00	Ε	Lower Pleistocene	1.800000	⊑	Europe
Paleotestudo	Paleotestudo antiqua	195.00	mf	Serravallian	13.000000	⊑	Europe
Testudo	Testudo sp.	1200.00	mf	Zanclean	3.960000	⊑	Europe
Centrochelys	Centrochelys burchardi	650.00	шо	Middle Pleistocene	0.435000	>	Europe
Centrochelys	Centrochelys robusta	850.00	φ <b>\</b>	Lower Pleistocene	1.300000	>	Europe
Testudo	Testudo catalaunica	232.00	Ε	Serravallian	12.350000	⊑	Europe
Geochelone	Geochelone sp.	1000.00	Ε	Burdigalian/Aquitanian	16.650000	⊑	Europe
Geochelone	Geochelone s. I.	1750.00	шо	Zanclean	4.466000	_	Europe
Eurotestudo	Eurotestudo hermanni	170.50	mf	Middle Pleistocene	0.600000	⊑	Europe
Testudo	Testudo hermanni	160.00	Ε	Modern	0.000001	>	Europe
Testudo	Testudo hermanni	157.00	Ε	Modern	0.000001	>	Europe
gen.	gen. indet.	270.00	шо	Burdigalian/Aquitanian	16.400000	<b>_</b>	Europe
Testudo	Testudo hermanni	161.00	Ε	Modern	0.000001	<b>C</b>	Europe
Testudo	Testudo marginata	242.50	Ε	Modern	0.000001	>	Europe

Genus	Taxon	CL	estimated	EpochBins	Age	Island	Con
Centrochelys	Centrochelys robusta	1100.00	шо	Zanclean	4.917000	λ	Europe
Testudo	Testudo rectogularis	213.00	шо	Burdigalian/Aquitanian	16.370000	⊑	Europe
Testudo	Testudo kalksburgensis	225.00	шо	Burdigalian/Aquitanian	18.000000	⊑	Europe
Testudo	Testudo marginata	400.00	Ε	Modern	0.000001	⊑	Europe
Testudo	Testudo brevitesta	300.00	mf	Piacencian	2.600000	⊑	Europe
Testudo	Testudo sp.	232.10	٤	Tortonian	10.750000	⊑	Europe
Testudo	Testudo horsfieldii	136.00	Ε	Modern	0.000001	⊑	Europe
Titanochelon	Titanochelon cf. bolivari	1300.00	ev	Langhian	14.895000	⊑	Europe
Testudo	Testudo marginata	290.00	Ε	Lower Pleistocene	1.300000	>	Europe
Testudo	Testudo hermanni	147.00	٤	Modern	0.000001	⊑	Europe
Eurotestudo	Eurotestudo hermanni	187.00	mf	Upper Pleistocene	0.110500	⊑	Europe
Eurotestudo	Eurotestudo aff. hermanni	179.30	mf	Middle Pleistocene	0.740000	⊑	Europe
Titanochelon	Titanochelon cf. perpiniana	1001.00	шо	Burdigalian/Aquitanian	16.370000	_	Europe
Testudo	Testudo sp.	245.00	Ε	Tortonian	8.300000	⊆	Europe
Testudo	Testudo amiatae	140.00	шо	Messinian	5.815000	_	Europe
Cheirogaster	Cheirogaster cf. gymnesica	789.00	шо	Lower Pleistocene	1.800000	>	Europe
Eurotestudo	Eurotestudo hermanni	126.00	mf	Lower Pleistocene	1.150000	⊑	Europe
Paleotestudo	Paleotestudo antiqua	283.80	mf	Serravallian	12.500000	<b>C</b>	Europe
Cheirogaster	Cheirogaster sp.	1000.00	ОШ	Serravallian	12.200000	<b>C</b>	Europe

Genus	Taxon	CL	estimated	EpochBins	Age	Island	Con
Paleotestudo	Paleotestudo cf. sp.	270.00	шо	Serravallian	12.400000	Ľ	Europe
Paleotestudo	Paleotestudo antiqua	240.00	mf	Serravallian	13.600000	ㅁ	Europe
Paleotestudo	Paleotestudo antiqua	195.00	Ε	Serravallian	13.000000	ㅁ	Europe
Titanochelon	Titanochelon bolivari	1353.00	шо	Serravallian	12.500000	ㅁ	Europe
Testudo	Testudo hermanni	154.00	Ε	Modern	0.000001	ㅁ	Europe
Centrochelys	Centrochelys robusta	00.009	φ <b>Λ</b>	Lower Pleistocene	1.300000	>	Europe
Paleotestudo	Paleotestudo antiqua	185.00	mf	Serravallian	13.000000	_	Europe
Titanochelon	Titanochelon schafferi	2500.00	шо	Zanclean	4.466000	ᄃ	Europe
Testudo	Testudo promarginata	310.00	mf	Burdigalian/Aquitanian	18.000000	ㅁ	Europe
Paleotestudo	Paleotestudo antiqua	206.00	mf	Serravallian	13.000000	ㅁ	Europe
Testudo	Testudo steinheimensis	227.70	mf	Serravallian	13.000000	_	Europe
Paleotestudo	Paleotestudo antiqua	234.00	mf	Serravallian	13.600000	ᄃ	Europe
Centrochelys	Centrochelys robusta	850.00	шо	Upper Pleistocene	0.066000	>	Europe
Testudo	Testudo promarginata	230.00	mf	Burdigalian/Aquitanian	21.500000	ᄃ	Europe
Titanochelon	Titanochelon sp.	1420.00	шо	Gelasian	1.850000	_	Europe
Paleotestudo	Paleotestudo antiqua	240.00	Ε	Serravallian	13.000000	ᄃ	Europe
Titanochelon	Titanochelon aff. schafferi	1860.00	Ε	Gelasian	2.000000	>	Europe
Testudo	Testudo hermanni	200.00	Ε	Modern	0.000001	>	Europe
Testudo	Testudo steinheimensis	111.00	Ε	Serravallian	12.150000	ᄆ	Europe

Genus	Taxon	CL	estimated	EpochBins	Age	Age Island	Con
Titanochelon	Titanochelon perpiniana	1140.00	Е	Zanclean	3.900000	u	Europe
Testudo	Testudo cf. graeca	185.00	Ε	Zanclean	3.900000	L	Europe
Paleotestudo	Paleotestudo antiqua	145.00	mf	Serravallian	13.000000	ᄓ	Europe
Cheirogaster	Cheirogaster sp.	1170.00	Ε	Tortonian	10.250000	L	Europe
Testudo	Testudo cf. promarginata	250.00	Ε	Tortonian	8.300000	L	Europe
Titanochelon	Titanochelon bolivari	1100.00	шо	Langhian	15.000000	L	Europe
Centrochelys	Centrochelys robusta	790.00	eĮ	Zanclean	4.917000	>	Europe
Titanochelon	Titanochelon cf. bolivari	1600.00	ef	Langhian	14.895000	C	Europe
Eurotestudo	Testudo hermanni	133.10	mf	Lower Pleistocene	1.220000	C	Europe
Testudo	Testudo hermanni	176.60	E	Modern	0.000001	>	Europe
Testudo	Testudo s. s.	189.00	E	Tortonian	8.000000	C	Europe
Centrochelys	Centrochelys robusta	850.00	шо	Zanclean	4.917000	>	Europe
Testudo	Testudo lunellensis	194.00	mf	Middle Pleistocene	0.450000	L	Europe
Testudo	Testudo hermanni	173.00	E	Modern	0.000001	>	Europe
Paleotestudo	Paleotestudo antiqua	229.00	mf	Serravallian	13.000000	C	Europe
Cheirogaster	Cheirogaster sp.	1500.00	Φ	Serravallian	13.800000	C	Europe
Testudo	Testudo catalaunica	181.00	Ε	Tortonian	11.500000	C	Europe
gen.	gen. indet.	813.00	ef	Upper Pleistocene	0.012500	>	Europe
Titanochelon	Titanochelon cf. bolivari	1500.00	mf	Tortonian	9.433000	C	Europe

Genus	Taxon	CL	estimated	EpochBins	Age	Island	Con
Testudo	Testudo sp.	245.00	٤	Tortonian	8.300000	<b>C</b>	Europe
Paleotestudo	Paleotestudo antiqua	213.00	mf	Serravallian	13.600000	Г	Europe
Testudo	Testudo sp.	2500.00	mf	Zanclean	3.900000	С	Europe
Paleotestudo	Paleotestudo antiqua	180.00	E	Serravallian	13.000000	Ц	Europe
Paleotestudo	Paleotestudo sp.	270.00	mf	Tortonian	9.500000	ㅁ	Europe
Testudo	Testudo hermanni	220.00	mf	Lower Pleistocene	1.300000	ㅁ	Europe
Paleotestudo	Paleotestudo sp.	170.00	mf	Tortonian	9.500000	ᄆ	Europe
Paleotestudo	Paleotestudo antiqua	183.70	E	Serravallian	12.150000	ᄃ	Europe
Testudo	Testudo sp.	245.00	E	Tortonian	8.300000	ㅁ	Europe
Eurotestudo	Eurotestudo cf. hermanni	150.00	шо	Gelasian	2.000000	>	Europe
Cheirogaster	Cheirogaster gymnesica	739.00	eĮ	Zanclean	4.450000	>	Europe
Titanochelon	Titanochelon bolivari	1300.00	mf	Tortonian	9.500000	ᄃ	Europe
Testudo	Testudo graeca	210.00	mf	Tortonian	8.450000	_	Europe
Cheirogaster	Cheirogaster richardi	1155.00	шо	Tortonian	10.400000	ᄃ	Europe
Paleotestudo	Paleotestudo antiqua	275.00	mf	Langhian	15.000000	_	Europe
Testudo	Testudo cf. promarginata	250.00	E	Tortonian	8.300000	ᄃ	Europe
Titanochelon	Titanochelon bacharidisi	900.006	шо	Zanclean	3.950000	_	Europe
Titanochelon	Titanochelon bacharidisi	1196.00	E	Zanclean	3.950000	ᄃ	Europe
Paleotestudo	Paleotestudo antiqua	152.00	E	Serravallian	13.000000	ᄆ	Europe

Genus	Taxon	CL	estimated	EpochBins	Age	Age Island	Con
Cheirogaster	Cheirogaster sp.	1540.00	ef	Tortonian	8.300000	С	Europe
Testudo	Testudo sp.	245.00	E	Tortonian	8.300000	⊑	Europe
Paleotestudo	Paleotestudo antiqua	220.00	mf	Serravallian	13.000000	⊑	Europe
gen.	gen. indet.	00.099	E	Tortonian	8.750000	⊑	Europe
Testudo	Testudo pecorinii	225.00	٤	Piacencian	3.094000	>	Europe
Testudo	Testudo catalaunica	107.00	E	Tortonian	11.500000	⊑	Europe
Titanochelon	Titanochelon schafferi	1850.00	E	Messinian	6.250000	>	Europe
Testudo	Testudo catalaunica	175.00	٤	Tortonian	11.500000	⊑	Europe
Titanochelon	Titanochelon sp.	520.00	шо	Piacencian	2.600000	>	Europe
Testudo	Testudo promarginata	304.70	m	Burdigalian/Aquitanian	21.500000	⊑	Europe
Titanochelon	Titanochelon gymnesica	1300.00	ef	Lower Pleistocene	1.300000	>	Europe
Testudo	Testudo burgenlandica	112.00	E	Tortonian	8.750000	⊑	Europe
Centrochelys	Centrochelys vulcanica	610.00	шо	Piacencian	3.094000	>	Europe
Testudo	Testudo brevitesta	165.00	mf	Piacencian	2.600000	⊑	Europe
Testudo	Testudo sp.	245.00	Ε	Tortonian	8.300000	⊑	Europe
gen.	gen. indet.	440.00	Ε	Tortonian	8.750000	⊆	Europe
Testudo	Testudo sp.	245.00	E	Tortonian	8.300000	<b>_</b>	Europe
Eurotestudo	Eurotestudo hermanni	284.00	mf	Lower Pleistocene	1.350000	<b>C</b>	Europe
Testudo	Testudo hermanni	145.90	E	Modern	0.000001	>	Europe

Genus	Тахоп	CL	estimated	EpochBins	Age	Island	Con
Testudo	Testudo cf. promarginata	250.00	٤	Tortonian	8.300000	۵	Europe
Testudo	Testudo cf. promarginata	250.00	E	Tortonian	8.300000	_	Europe
Testudo	Testudo marginata	310.00	٤	Lower Pleistocene	1.300000	>	Europe
Testudo	Testudo cf. promarginata	250.00	Ε	Tortonian	8.300000	L	Europe
Paleotestudo	Paleotestudo sp.	261.00	mf	Tortonian	9.500000	_	Europe
Testudo	Testudo catalaunica	165.00	E	Tortonian	11.500000	L	Europe
"Hadrianus"	"Hadrianus sp."	1000.00	E	Tortonian	8.300000	_	Europe
Titanochelon	Titanochelon bolivari	1250.00	шо	Langhian	15.000000	_	Europe
Centrochelys	Centrochelys burchardi	800.00	E	Middle Pleistocene	0.435000	>	Europe
Gopherus	Gopherus polyphemus	217.90	шо	Lower Pleistocene	1.200000	_	America
Gopherus	Gopherus polyphemus	238.90	E	Modern	0.000001	_	America
Gopherus	Gopherus polyphemus	102.44	шо	Upper Pleistocene	0.0690.0	_	America
Gopherus	Gopherus polyphemus	327.60	шо	Upper Pleistocene	0.0690.0	_	America
Hesperotestudo	Hesperotestudo crassiscutata	284.90	E	Upper Pleistocene	0.0690.0	_	America
Gopherus	Gopherus polyphemus	276.60	шо	Lower Pleistocene	1.200000	C	America
Gopherus	Gopherus praecedens	360.00	шо	Upper Pleistocene	0.0690.0	_	America
Gopherus	Gopherus polyphemus	278.00	шо	Upper Pleistocene	0.0690.0	C	America
Gopherus	Gopherus sp.	236.70	шо	Gelasian	1.900000	_	America
Gopherus	Gopherus polyphemus	273.24	ОШ	Upper Pleistocene	0.069000	_	America

Genus	Taxon	CL	estimated	EpochBins	Age	Island	Con
Gopherus	Gopherus polyphemus	302.40	шо	Upper Pleistocene	0.0690.0	۵	America
Gopherus	Gopherus polyphemus	268.80	Ε	Modern	0.000001	>	America
Hesperotestudo	Hesperotestudo crassiscutata	425.00	шо	Upper Pleistocene	0.012000	L	America
Gopherus	Gopherus polyphemus	334.70	шо	Upper Pleistocene	0.0690.0	С	America
Gopherus	Gopherus polyphemus	300.00	Ε	Modern	0.000001	>	America
Gopherus	Gopherus polyphemus	350.00	шо	Upper Pleistocene	0.0690.0	_	America
Gopherus	Gopherus polyphemus	258.30	шо	Upper Pleistocene	0.0690.0	_	America
Hesperotestudo	Hesperotestudo crassiscutata	180.40	Ε	Upper Pleistocene	0.069000	С	America
Gopherus	Gopherus flavomarginatus	371.00	Ε	Modern	0.000001	_	America
Gopherus	Gopherus polyphemus	284.90	шо	Upper Pleistocene	0.0690.0	_	America
Hesperotestudo	Hesperotestudo crassiscutata	188.00	шо	Upper Pleistocene	0.012000	_	America
Gopherus	Gopherus ? sp.	500.00	Ε	Serravallian	11.850000	С	America
Hesperotestudo	Hesperotestudo crassiscutata	168.00	E	Lower Pleistocene	1.300000	С	America
Gopherus	Gopherus agassizii	400.00	Ε	Modern	0.000001	С	America
Hesperotestudo	Hesperotestudo orthopygia	1200.00	шо	Messinian	5.500000	_	America
Gopherus	Gopherus polyphemus	353.30	шо	Middle Pleistocene	0.400000	С	America
Gopherus	Gopherus sp.	202.80	ш	Lower Pleistocene	1.800000	С	America
Gopherus	Gopherus polyphemus	387.00	Ε	Modern	0.000001	С	America
Gopherus	Gopherus polyphemus	279.94	шо	Upper Pleistocene	0.069000	_	America

Genus	Taxon	CL	estimated	EpochBins	Age	Island	Con
Gopherus	Gopherus sp.	224.10	шо	Lower Pleistocene	1.800000	۵	America
Gopherus	Gopherus polyphemus	268.90	шо	Lower Pleistocene	1.200000	L	America
Hesperotestudo	Hesperotestudo sp.	639.00	Ε	Upper Pleistocene	0.060000	_	America
Gopherus	Gopherus flavomarginatus	281.00	E	Modern	0.000001	C	America
Gopherus	Gopherus polyphemus	252.56	шо	Upper Pleistocene	0.0690.0	L	America
Gopherus	Gopherus polyphemus	293.00	шо	Middle Pleistocene	0.400000	С	America
Gopherus	Gopherus polyphemus	155.50	шо	Upper Pleistocene	0.0690.0	_	America
Gopherus	Gopherus polyphemus	260.50	шо	Middle Pleistocene	0.400000	_	America
Gopherus	Gopherus polyphemus	256.44	шо	Middle Pleistocene	0.250000	_	America
Hesperotestudo	Hesperotestudo sp.	1000.00	шо	Gelasian	2.000000	_	America
Geochelone	Geochelone sp.	350.00	ef	Upper Pleistocene	0.0690.0	_	America
Gopherus	Gopherus sp.	181.00	шо	Gelasian	1.900000	C	America
Geochelone	Geochelone sp.	00.009	ОШ	Upper Pleistocene	0.012500	>	America
Gopherus	Gopherus polyphemus	303.00	Ε	Modern	0.000001	>	America
Gopherus	Gopherus polyphemus	342.00	Ε	Modern	0.000001	_	America
Gopherus	Gopherus sp.	256.08	Ε	Modern	0.000001	C	America
Gopherus	Gopherus sp.	180.90	ОШ	Gelasian	1.900000	C	America
Hesperotestudo	Hesperotestudo incisa	232.76	Ε	Upper Pleistocene	0.0690.0	C	America
Gopherus	Gopherus sp.	181.00	ОШ	Gelasian	1.900000	C	America

Genus	Taxon	CL	estimated	EpochBins	Age	Island	Con
Geochelone	Geochelone tedwhitei	440.00	Е	Burdigalian/Aquitanian	18.500000	u	America
Gopherus	Gopherus polyphemus	239.80	шо	Middle Pleistocene	0.250000	⊑	America
Hesperotestudo	Hesperotestudo sp.	974.00	də	Upper Pleistocene	0.060000	⊑	America
Gopherus	Gopherus polyphemus	260.11	шо	Upper Pleistocene	0.0690.0	⊑	America
Gopherus	Gopherus sp.	204.40	шо	Gelasian	1.900000	⊑	America
Hesperotestudo	Hesperotestudo crassiscutata	192.00	Ε	Lower Pleistocene	1.300000	⊑	America
Gopherus	Gopherus sp.	194.90	шо	Gelasian	1.900000	⊑	America
Gopherus	Gopherus polyphemus	391.90	ОШ	Upper Pleistocene	0.0690.0	⊆	America
Gopherus	Gopherus sp.	259.50	шо	Lower Pleistocene	1.800000	⊑	America
Geochelone	Geochelone sp.	170.00	m	Middle Pleistocene	0.700000	⊑	America
Gopherus	Gopherus sp.	230.10	ОШ	Lower Pleistocene	1.800000	_	America
Hesperotestudo	Hesperotestudo incisa	224.00	Ε	Lower Pleistocene	1.300000	⊑	America
Hesperotestudo	Hesperotestudo equicomes	340.00	ev	Middle Pleistocene	0.300000	⊑	America
Hesperotestudo	Hesperotestudo incisa	228.00	E	Lower Pleistocene	1.300000	<b>C</b>	America
Gopherus	Gopherus flavomarginatus	303.00	Ε	Modern	0.000001	_	America
Testudo	Testudo sp.	400.00	шо	Langhian	14.181000	⊑	America
Gopherus	Gopherus pertenuis	1050.00	ОШ	Lower Pleistocene	1.684500	_	America
Hesperotestudo	Hesperotestudo incisa	231.00	E	Lower Pleistocene	1.300000	<b>C</b>	America
Hesperotestudo	Hesperotestudo crassiscutata	327.00	E	Lower Pleistocene	1.300000	<b>C</b>	America

Genus	Taxon	CL	estimated	EpochBins	Age	Island	Con
Hesperotestudo	Hesperotestudo incisa	241.00	E	Lower Pleistocene	1.300000	Ľ	America
Hesperotestudo	Hesperotestudo incisa	250.00	Φ	Modern	0.007500	ᄕ	America
Gopherus	Gopherus polyphemus	352.00	шо	Upper Pleistocene	0.012000	_	America
Hesperotestudo	Hesperotestudo johnstoni	235.00	٤	Piacencian	3.350000	_	America
Gopherus	Gopherus polyphemus	274.30	шо	Middle Pleistocene	0.250000	С	America
Gopherus	Gopherus flavomarginatus	222.00	Ε	Modern	0.000001	С	America
Gopherus	Gopherus sp.	241.90	шо	Lower Pleistocene	1.800000	C	America
Gopherus	Gopherus sp.	216.37	Ε	Modern	0.000001	_	America
Hesperotestudo	Hesperotestudo sp.	1200.00	ev	Tortonian	9.500000	_	America
Gopherus	Gopherus polyphemus	257.80	шо	Middle Pleistocene	0.250000	_	America
Hesperotestudo	Hesperotestudo crassiscutata	282.70	Ε	Upper Pleistocene	0.0690.0	_	America
Hesperotestudo	Hesperotestudo campester	1000.00	шо	Gelasian	2.190500	_	America
Hesperotestudo	Hesperotestudo incisa	216.00	Ε	Lower Pleistocene	1.300000	_	America
Hesperotestudo	Hesperotestudo mlynarskii	203.50	Ε	Lower Pleistocene	1.250000	<b>C</b>	America
Geochelone	Geochelone sp.	880.00	Ε	Zanclean	4.500000	_	America
Gopherus	Gopherus polyphemus	431.48	шо	Upper Pleistocene	0.0690.0	_	America
Gopherus	Gopherus polyphemus	308.00	Ε	Modern	0.000001	_	America
Gopherus	Gopherus mohavetus	315.00	Ε	Tortonian	8.476000	<b>C</b>	America
Gopherus	Gopherus sp.	264.11	٤	Modern	0.000001	<b>C</b>	America

Genus	Taxon	CL	estimated	EpochBins	Age	Island	Con
Gopherus	Gopherus sp.	118.90	шо	Gelasian	1.900000	u	America
Gopherus	Gopherus polyphemus	337.30	шо	Middle Pleistocene	0.250000	⊑	America
Gopherus	Gopherus sp.	163.50	шо	Gelasian	1.900000	⊑	America
Caudochelys	Caudochelys rexroadensis	830.00	٤	Zanclean	4.550000	⊑	America
Hesperotestudo	Hesperotestudo riggsi	159.50	шо	Tortonian	7.600000	⊑	America
Gopherus	Gopherus polyphemus	306.00	шо	Middle Pleistocene	0.250000	⊑	America
Hesperotestudo	Hesperotestudo crassiscutata	561.00	E	Lower Pleistocene	1.250000	⊏	America
Geochelone	Geochelone sp.	176.00	Φ	Zanclean	5.000000	⊑	America
Gopherus	Gopherus sp.	218.80	шо	Gelasian	1.900000	⊑	America
Gopherus	Gopherus agassizi	252.00	Ε	Upper Pleistocene	0.025500	<b>_</b>	America
Hesperotestudo	Hesperotestudo crassiscutata	180.00	Ε	Lower Pleistocene	1.300000	⊏	America
Caudochelys	Caudochelys williamsi	334.00	Ε	Burdigalian/Aquitanian	17.750000	⊆	America
Hesperotestudo	Hesperotestudo incisa	290.40	E	Lower Pleistocene	1.300000	_	America
Gopherus	Gopherus sp.	245.40	ОШ	Lower Pleistocene	1.800000	<b>_</b>	America
Gopherus	Gopherus polyphemus	301.97	шо	Upper Pleistocene	0.0690.0	<b>_</b>	America
Hesperotestudo	Hesperotestudo incisa	212.00	E	Lower Pleistocene	1.300000	⊑	America
Gopherus	Gopherus sp.	188.30	ОШ	Gelasian	1.900000	_	America
Hesperotestudo	Hesperotestudo crassiscutata	1250.00	θΛ	Upper Pleistocene	0.012000	<b>_</b>	America
Gopherus	Gopherus polyphemus	350.83	шо	Middle Pleistocene	0.400000	<b>_</b>	America

Genus	Taxon	CL	estimated	EpochBins	Age	Island	Con
Hesperotestudo	Hesperotestudo riggsi	176.00	Е	Piacencian	3.000000	u	America
Gopherus	Gopherus polyphemus	304.70	шо	Middle Pleistocene	0.400000	П	America
Gopherus	Gopherus sp.	143.90	ОШ	Gelasian	1.900000	Ц	America
Hesperotestudo	Hesperotestudo sp.	176.00	mf	Piacencian	3.100000	ᄆ	America
Gopherus	Gopherus polyphemus	260.51	шо	Middle Pleistocene	0.400000	П	America
Gopherus	Gopherus sp.	241.56	Ε	Modern	0.000001	Ц	America
Hesperotestudo	Hesperotestudo orthopygia	682.00	ОШ	Messinian	5.500000	ᄆ	America
Hesperotestudo	Hesperotestudo wilsoni	226.00	Ε	Upper Pleistocene	0.018000	Ц	America
Gopherus	Gopherus sp.	211.31	Ε	Modern	0.000001	ᄆ	America
Gopherus	Gopherus polyphemus	304.20	шо	Upper Pleistocene	0.069000	C	America
Hesperotestudo	Hesperotestudo oelrichi	283.80	Ε	Piacencian	3.000000	ᆮ	America
Gopherus	Gopherus laticaudatus	375.00	шо	Middle Pleistocene	0.396350	C	America
Gopherus	Gopherus mohavetus	334.50	Ε	Tortonian	8.476000	C	America
Hesperotestudo	Hesperotestudo riggsi	159.50	шо	Tortonian	7.600000	C	America
Caudochelys	Caudochelys rexroadensis	781.00	Ε	Zanclean	4.550000	C	America
Gopherus	Gopherus polyphemus	267.00	шо	Middle Pleistocene	0.250000	C	America
Gopherus	Gopherus polyphemus	295.90	шо	Middle Pleistocene	0.400000	C	America
Hesperotestudo	Hesperotestudo riggsi	195.80	Ε	Zanclean	4.550000	C	America
Gopherus	Gopherus polyphemus	324.00	ОШ	Upper Pleistocene	0.069000	_	America

Genus	Taxon	CL	estimated	EpochBins	Age	Island	Con
Gopherus	Gopherus sp.	182.30	om	Gelasian	1.900000	_	America
Gopherus	Gopherus polyphemus	294.16	шо	Upper Pleistocene	0.0690.0	⊑	America
Hesperotestudo	Hesperotestudo alleni	240.90	Ε	Tortonian	10.950000	⊑	America
Gopherus	Gopherus polyphemus	283.41	ОШ	Middle Pleistocene	0.250000	⊑	America
Gopherus	Gopherus polyphemus	272.48	ОШ	Middle Pleistocene	0.250000	⊑	America
Hesperotestudo	Hesperotestudo riggsi	185.00	Ε	Piacencian	3.000000	⊑	America
Geochelone	Geochelone tedwhitei	370.00	Ε	Burdigalian/Aquitanian	18.500000	⊑	America
Gopherus	Gopherus ? sp.	500.00	Ε	Tortonian	10.100000	⊑	America
Gopherus	Gopherus sp.	209.60	шо	Gelasian	1.900000	⊏	America
Gopherus	Gopherus polyphemus	308.20	шо	Middle Pleistocene	0.400000	⊑	America
Gopherus	Gopherus polyphemus	314.60	шо	Middle Pleistocene	0.250000	_	America
Gopherus	Gopherus sp.	193.30	шо	Gelasian	1.900000	⊑	America
Gopherus	Gopherus sp.	188.70	шо	Gelasian	1.900000	<b>_</b>	America
Gopherus	Gopherus polyphemus	302.40	шо	Middle Pleistocene	0.250000	⊆	America
Gopherus	Gopherus polyphemus	292.00	шо	Middle Pleistocene	0.250000	_	America
Gopherus	Gopherus polyphemus	306.00	шо	Middle Pleistocene	0.250000	⊑	America
Hesperotestudo	Hesperotestudo turgida	230.00	шо	Lower Pleistocene	1.684500	⊑	America
Gopherus	Gopherus polyphemus	272.57	шо	Middle Pleistocene	0.400000	⊆	America
Gopherus	Gopherus polyphemus	322.63	ОШ	Middle Pleistocene	0.250000	<b>C</b>	America

Genus	Taxon	CL	estimated	EpochBins	Age	Island	Con
Gopherus	Gopherus flavomarginatus	278.00	Е	Modern	0.000001	u	America
Geochelone	Geochelone sp.	500.00	E	Tortonian	10.100000	_	America
Caudochelys	Caudochelys ducateli	339.90	Ε	Langhian	15.000000	드	America
Gopherus	Gopherus polyphemus	292.94	шо	Middle Pleistocene	0.250000	⊑	America
Gopherus	Gopherus polyphemus	348.70	шо	Middle Pleistocene	0.400000	드	America
Hesperotestudo	Hesperotestudo sp.	1500.00	шо	Middle Pleistocene	0.700000	_	America
Gopherus	Gopherus polyphemus	285.20	шо	Middle Pleistocene	0.250000	⊑	America
Gopherus	Gopherus mohavetus	412.50	E	Tortonian	8.476000	⊑	America
Hesperotestudo	Hesperotestudo sp.	1800.00	шо	Middle Pleistocene	0.700000	⊑	America
Gopherus	Gopherus polyphemus	285.60	шо	Middle Pleistocene	0.400000	_	America
Gopherus	Gopherus canyonensis	885.50	Ε	Piacencian	2.700000	_	America
Gopherus	Gopherus polyphemus	253.70	шо	Middle Pleistocene	0.250000	⊑	America
Gopherus	Gopherus polyphemus	293.57	ОШ	Middle Pleistocene	0.400000	<b>_</b>	America
Gopherus	Gopherus mohavetus	202.00	Ε	Tortonian	8.476000	_	America
Gopherus	Gopherus mohavetus	360.00	Ε	Tortonian	8.476000	_	America
Gopherus	Gopherus agassizi	445.00	шо	Middle Pleistocene	0.156000	_	America
Gopherus	Gopherus polyphemus	539.00	m	Middle Pleistocene	0.700000	_	America
Gopherus	Gopherus polyphemus	283.00	шо	Middle Pleistocene	0.250000	_	America
Hesperotestudo	Hesperotestudo mlynarskii	165.00	E	Lower Pleistocene	1.250000	<b>_</b>	America

Genus	Taxon	C	estimated	EpochBins	Age	Island	Con
Gopherus	Gopherus flavomarginatus	246.00	Ε	Modern	0.000001	Ľ	America
Chelonoidis	Chelonoidis chilensis	169.00	Ε	Modern	0.000001	_	America
Chelonoidis	Chelonoidis carbonaria	296.50	Ε	Modern	0.000001	드	America
Chelonoidis	Chelonoidis carbonaria	242.00	Ε	Modern	0.000001	_	America
Chelonoidis	Chelonoidis chilensis	200.00	Ε	Modern	0.000001	드	America
Chelonoidis	Chelonoidis carbonaria	253.00	Ε	Modern	0.000001	_	America
Chelonoidis	Chelonoidis denticulata	333.40	Ε	Modern	0.000001	드	America
Chelonoidis	Chelonoidis carbonaria	247.00	Ε	Modern	0.000001	_	America
Chelonoidis	Chelonoidis chilensis	186.00	Ε	Modern	0.000001	_	America
Chelonoidis	Chelonoidis chilensis	157.00	Ε	Modern	0.000001	_	America
Chelonoidis	Chelonoidis sp.	1000.00	шо	Upper Pleistocene	0.0690.0	_	America
Chelonoidis	Chelonoidis carbonaria	333.40	Ε	Modern	0.000001	_	America
Chelonoidis	Chelonoidis nigra	745.70	Ε	Modern	0.000001	>	America
Chelonoidis	Chelonoidis carbonaria	290.00	Ε	Modern	0.000001	>	America
Chelonoidis	Chelonoidis sp.	300.00	шо	Langhian	15.900000	_	America
Chelonoidis	Chelonoidis denticulata	365.00	Ε	Modern	0.000001	_	America
Chelonoidis	Chelonoidis chilensis	183.00	Ε	Modern	0.000001	_	America
Chelonoidis	Chelonoidis denticulata	317.00	Ε	Modern	0.000001	_	America
Chelonoidis	Chelonoidis chilensis	169.00	Ε	Modern	0.000001	_	America

Genus	Taxon	CL	estimated	EpochBins	Age	Island	Con
Chelonoidis	Chelonoidis hoodensis	813.00	Ε	Modern	0.000001	>	America
Chelonoidis	Chelonoidis phantastica	860.00	٤	Modern	0.000001	>	America
Chelonoidis	Chelonoidis lutzae	830.00	Ε	Upper Pleistocene	0.038500	C	America
Chelonoidis	Chelonoidis nigra	1300.00	Ε	Modern	0.000001	>	America
Chelonoidis	Chelonoidis becki	1050.00	٤	Modern	0.000001	>	America
Chelonoidis	Chelonoidis nigra	595.00	Ε	Modern	0.000001	>	America
Chelonoidis	Chelonoidis sp.	300.00	шо	Langhian	15.900000	C	America
Chelonoidis	Chelonoidis chilensis	450.00	٤	Modern	0.000001	_	America
Chelonoidis	Chelonoidis darwini	965.00	٤	Modern	0.000001	>	America
Chelonoidis	Chelonoidis nigra	731.30	٤	Modern	0.000001	>	America
Chelonoidis	Chelonoidis denticulata	616.00	٤	Upper Pleistocene	0.120000	_	America
Chelonoidis	Chelonoidis duncanensis	840.00	٤	Modern	0.000001	>	America
Chelonoidis	Chelonoidis denticulata	820.00	٤	Modern	0.000001	_	America
Chelonoidis	Chelonoidis abingdonii	980.00	٤	Modern	0.000001	>	America
Chelonoidis	Chelonoidis sp.	1060.00	ec	Langhian	15.900000	С	America
Chelonoidis	Chelonoidis nigra	588.00	Ε	Modern	0.000001	>	America
Chelonoidis	Chelonoidis carbonaria	189.00	Ε	Modern	0.000001	С	America
Chelonoidis	Chelonoidis chathamensis	890.00	Ε	Modern	0.000001	>	America
Chelonoidis	Chelonoidis chilensis	222.00	Ε	Modern	0.000001	⊑	America

Genus	Taxon	CL	CL estimated EpochBins	EpochBins	Age	Age Island Con	Con
Chelonoidis	Chelonoidis carbonaria	593.00	Ε	Modern	0.000001	<b>C</b>	America
Chelonoidis	Chelonoidis denticulata	333.00	Ε	Modern	0.000001	ᄆ	America
Chelonoidis	Chelonoidis nigra	610.00	Ε	Modern	0.000001	>	America
Chelonoidis	Chelonoidis vicina	1250.00	Ε	Modern	0.000001	>	America
Chelonoidis	Chelonoidis nigra	717.00	Ε	Modern	0.000001	>	America
Geochelone	Geochelone hesterna	278.00	Ε	Tortonian	8.500000	ᄃ	America
Chelonoidis	Chelonoidis denticulata	377.00	Ε	Modern	0.000001	<b>L</b>	America
Chelonoidis	Chelonoidis denticulata	466.00	Ε	Modern	0.000001	<b>_</b>	America
Chelonoidis	Chelonoidis carbonaria	226.00	Ш	Modern	0.000001	u	America

Genus	Taxon	CollNr	SCL	CCL	SCW	CCW	G	PL	ΡW	estimated	Island	Con	Reference
Kinixys	Kinixys belliana	ZMB 37388	162.0	16.20	22.5	15.5	21.5	164.0	12.6	Ε	_	Africa	freshly measured (MFN collection)
Aldabrachelys	Aldabrachelys gigantea	ZMB 51996	770.0	77.00	106.0	52.0	112.0	NA	ΑĀ	E	>	Africa	freshly measured (MFN collection)
Astrochelys	Astrochelys yniphora	1	426.0	42.60	NA	NA	Ϋ́	NA	NA	E	>	Africa	Pedrono, M., & Smith, L. L. (2013). Overview of the
Centrochelys	Centrochelys sulcata	ZMB 63203	215.0	21.50	29.5	16.5	27.0	214.0	14.8	E	С	Africa	freshly measured (MFN collection)
Malacochersus	Malacochersus tornieri	ZMB 63174	153.0	15.30	17.0	10.5	14.0	149.0	9.8	E	Ц	Africa	freshly measured (MFN collection)
Astrochelys	Astrochelys radiata	1	395.0	39.50	N	N	Ϋ́	NA	NA	Ε	>	Africa	Pedrono, M., & Smith, L. L. (2013). Overview of the
Pyxis	Pyxis arachnoides	ZMB 37616	110.0	11.00	15.0	8.0	14.0	75.0	9.7	E	>	Africa	freshly measured (MFN collection)
Kinixys	Kinixys homeana	ZMB 17747	193.0	19.30	25.0	14.0	21.0	175.0	11.8	E	_	Africa	freshly measured (MFN collection)
Aldabrachelys	Aldabrachelys gigantea	ZMB 47494	870.0	87.00	116.0	57.0	110.0	NA	ΑĀ	E	>	Africa	freshly measured (MFN collection)
Psammobates	Psammobates tentorius	ZMB 28782	111.0	11.10	15.0	8.5	14.0	95.0	7.9	E	С	Africa	freshly measured (MFN collection)
Psammobates	Psammobates oculifer	ZMB 25439	119.0	11.90	17.0	9.0	14.5	0.66	8.4	E	С	Africa	freshly measured (MFN collection)
Psammobates	Psammobates oculifer	ZMB 37472	107.0	10.70	15.0	8.4	13.5	106.0	∞	Ε	ч	Africa	freshly measured (MFN collection)
Astrochelys	Astrochelys yniphora	1	307.0	30.70	N	N	Α̈́	NA	Α̈́	Ε	>	Africa	Pedrono, M., & Smith, L. L. (2013). Overview of the
Homopus	Homopus aerolatus	ZMB 229	88.0	8.80	10.5	6.9	9.0	78.0	6.1	E	_	Africa	freshly measured (MFN collection)
Homopus	Homopus signatus	ZMB 63173	94.0	9.40	12.5	7.7	11.0	82.0	5.6	E	_	Africa	freshly measured (MFN collection)
Kinixys	Kinixys belliana	ZMB 63191	194.0	19.40	25.5	12.5	19.0	173.0	12	E	_	Africa	freshly measured (MFN collection)
Astrochelys	Astrochelys radiata	1	285.0	28.50	N	Ä	N A	N A	A A	E	>	Africa	Pedrono, M., & Smith, L. L. (2013). Overview of the
Kinixys	Kinixys belliana	ZMB 63192	174.0	17.40	24.5	11.5	20.5	143.0	1.1	E	L	Africa	freshly measured (MFN collection)
Kinixys	Kinixys belliana	ZMB 63193	157.0	15.70	21.0	6.6	16.5	141.0	9.4	E	_	Africa	freshly measured (MFN collection)
Aldabrachelys	Aldabrachelys gigantea	ZMB 37545	810.0	81.00	110.0	52.0	Α̈́	NA	Α̈́	Ε	>	Africa	freshly measured (MFN collection)
Chersina	Chersina angulata	ZMB 49400	162.0	16.20	21.5	10.9	17.5	170.0	9.2	E	_	Africa	freshly measured (MFN collection)
Chersina	Chersina angulata	ZMB 63181	170.0	17.00	23.0	11.4	19.0	169.0	10	E	_	Africa	freshly measured (MFN collection)
Chersina	Chersina angulata	ZMB 63183	120.0	12.00	17.0	8.6	15.5	118.0	7.3	E	_	Africa	freshly measured (MFN collection)
Chersina	Chersina angulata	ZMB 63182	136.0	13.60	18.0	6.6	16.0	138.0	80	E	_	Africa	freshly measured (MFN collection)
Kinixys	Kinixys erosa	ZMB 63190	164.0	16.40	21.0	11.2	16.5	163.0	10.6	E	_	Africa	freshly measured (MFN collection)
Centrochelys	Centrochelys sulcata	ZMB 37387	435.0	43.50	54.0	29.9	53.0	405.0	29.1	Ε	_	Africa	freshly measured (MFN collection)

Genus	Taxon	CollNr	SCL	CCL	SCW	CCW	CH	PL	PW	estimated	Island	Con	Reference
Indotestudo	Indotestudo travancorica	ZMB 37717	224.0	22.40	28.0	15.2	23.0	200.0	15.4	Е	u	Africa	freshly measured (MFN collection)
Stigmochelys	Stigmochelys pardalis	ZMB 37344	405.0	40.50	55.0	27.0	50.5	350.0	24.3	E	L	Africa	freshly measured (MFN collection)
Stigmochelys	Stigmochelys pardalis	ZMB 63235	315.0	31.50	43.5	23.4	39.0	298.0	22.1	Е	u	Africa	freshly measured (MFN collection)
Stigmochelys	Stigmochelys pardalis	ZMB 37495	297.0	29.70	41.5	21.4	36.0	271.0	19.2	Е	u	Africa	freshly measured (MFN collection)
Stigmochelys	Stigmochelys pardalis	ZMB 42400	345.0	34.50	46.5	24.0	40.0	285.0	21.3	E	п	Africa	freshly measured (MFN collection)
Stigmochelys	Stigmochelys pardalis	ZMB 63232	350.0	35.00	46.0	23.9	45.0	303.0	21.1	Е	u	Africa	freshly measured (MFN collection)
Psammobates	Psammobates geometricus	ZMB 192	92.0	9.20	13.5	7.1	13.0	0.89	6.3	Ε	۵	Africa	freshly measured (MFN collection)
Chersina	Chersina angulata	ı	181.9	18.19	A	N A	N A	Ν	Ą	Ε	>	Africa	Itescu, Y., Karraker, N. E., Raia, P., Pritchard, P. C.,
Aldabrachelys	Aldabrachelys gigantea	ZMB 47443	800.0	80.00	105.0	51.5	105.0	Ν	Ą	Ε	>	Africa	freshly measured (MFN collection)
Astrochelys	Astrochelys yniphora	ı	415.0	41.50	Ν Α	Ϋ́	Ϋ́	NA	A	Е	>	Africa	Pedrono, M., & Smith, L. L. (2013). Overview of the
Astrochelys	Astrochelys yniphora	ı	370.0	37.00	A	Ϋ́	Ϋ́	N	Ą	Ε	>	Africa	Pedrono, M., & Smith, L. L. (2013). Overview of the
Aldabrachelys	Aldabrachelys gigantea	ZMB 51995	1030.0	103.00	138.0	Ϋ́	Ϋ́	NA	A	Е	>	Africa	freshly measured (MFN collection)
Aldabrachelys	Aldabrachelys gigantea	ZMB ???	720.0	72.00	105.5	55.0	117.0	NA	A	Е	>	Africa	freshly measured (MFN collection)
Cylindraspis	Cylindraspis triserrata	ı	1100.0	110.00	Ν Α	Ϋ́	Ϋ́	NA	A	Е	>	Africa	Itescu, Y., Karraker, N. E., Raia, P., Pritchard, P. C.,
Cylindraspis	Cylindraspis vosmaeri	ı	500.0	50.00	ΑN	N A	N A	NA	Ą	E	>	Africa	Itescu, Y., Karraker, N. E., Raia, P., Pritchard, P. C.,
Astrochelys	Astrochelys radiata	ı	334.0	33.40	Ν Α	Ϋ́	Ϋ́	NA	A	E	>	Africa	Pedrono, M., & Smith, L. L. (2013). Overview of the
Astrochelys	Astrochelys radiata	ı	305.0	30.50	Ν Α	Ϋ́	Ϋ́	NA	A	E	>	Africa	Pedrono, M., & Smith, L. L. (2013). Overview of the
Centrochelys	Centrochelys sulcata	ı	830.0	83.00	Ν Α	Ϋ́	Ϋ́	NA	A	Е	u	Africa	Itescu, Y., Karraker, N. E., Raia, P., Pritchard, P. C.,
Psammobates	Psammobates geometricus	ZMB 186	105.0	10.50	13.5	7.4	13.0	0.06	6.9	Ε	L	Africa	freshly measured (MFN collection)
Astrochelys	Astrochelys radiata	ı	242.0	24.20	ΑN	Α̈́	Α̈́	NA	Ą	Ε	>	Africa	Pedrono, M., & Smith, L. L. (2013). Overview of the
Psammobates	Psammobates tentorius	ZMB 37627	116.0	11.60	15.0	9.4	14.5	117.0	8.9	E	>	Africa	freshly measured (MFN collection)
Psammobates	Psammobates tentorius	ZMB 50571	95.0	9.50	12.0	7.3	12.0	79.0	7	E	L	Africa	freshly measured (MFN collection)
Psammobates	Psammobates tentorius	ZMB 14766	81.0	8.10	10.5	8.9	10.0	0.79	5.9	E	۵	Africa	freshly measured (MFN collection)
Pyxis	Pyxis planicauda	ı	114.0	11.40	Ν Α	Ϋ́	Ϋ́	NA	A	E	>	Africa	Pedrono, M., & Smith, L. L. (2013). Overview of the
Pyxis	Pyxis planicauda	ı	134.0	13.40	ΑN	Α̈́	Α̈́	NA	Ą	Ε	>	Africa	Pedrono, M., & Smith, L. L. (2013). Overview of the
Pyxis	Pyxis planicauda	ı	120.0	12.00	ΑN	Α̈́	Α̈́	NA	Ą	Ε	>	Africa	Pedrono, M., & Smith, L. L. (2013). Overview of the
Psammobates	Psammobates oculifer	ZMB 16399	111.0	11.10	16.0	8.8	14.0	108.0	6.7	E	L	Africa	freshly measured (MFN collection)
Psammobates	Psammobates oculifer	ZMB 14772	101.0	10.10	15.0	8.0	14.0	98.0	7.3	E	L	Africa	freshly measured (MFN collection)

Genus	Taxon	CollNr	SCL	CCL	SCW	CCW	끙	PL	PW	estimated	Island	Con	Reference
Psammobates	Psammobates oculifer	ZMB 24261	103.0	10.30	14.0	8.2	13.5	100.0	7.8	٤	L	Africa	freshly measured (MFN collection)
Psammobates	Psammobates oculifer	ZMB 37623	105.0	10.50	14.5	7.9	13.5	93.0	7.4	Ε	_	Africa	freshly measured (MFN collection)
Kinixys	Kinixys belliana	ZMB 37489	180.0	18.00	24.0	12.0	20.5	176.0	11.8	Ε	_	Africa	freshly measured (MFN collection)
Pyxis	Pyxis planicauda	1	160.0	16.00	NA	N	N	Ϋ́	Ą	Ε	>	Africa	Itescu, Y., Karraker, N. E., Raia, P., Pritchard, P. C.,
Psammobates	Psammobates geometricus	ZMB 50568	107.0	10.70	15.0	7.9	14.5	79.0	7.3	Ε	Ц	Africa	freshly measured (MFN collection)
Aldabrachelys	Aldabrachelys gigantea	1	875.0	87.50	N	N A	Α	Ν	ΑN	Ε	>	Africa	Itescu, Y., Karraker, N. E., Raia, P., Pritchard, P. C.,
Aldabrachelys	Aldabrachelys gigantea	1	1190.0	119.00	N	N	Ą	Ϋ́	ΑĀ	Ε	>	Africa	Itescu, Y., Karraker, N. E., Raia, P., Pritchard, P. C.,
Chersina	Chersina angulata	1	202.0	20.20	N	Ν	Α	ΑN	ΑN	Ε	_	Africa	Itescu, Y., Karraker, N. E., Raia, P., Pritchard, P. C.,
Chersina	Chersina angulata	1	351.0	35.10	N	Ν	Α	Ϋ́	Ą	Ε	>	Africa	Itescu, Y., Karraker, N. E., Raia, P., Pritchard, P. C.,
Astrochelys	Astrochelys yniphora	1	446.0	44.60	NA	NA	NA	Ϋ́	Ϋ́	Ε	>	Africa	Itescu, Y., Karraker, N. E., Raia, P., Pritchard, P. C.,
Chersina	Chersina angulata	ZMB 37393	160.0	16.00	20.0	10.0	17.5	158.0	9.2	Ε	L	Africa	freshly measured (MFN collection)
Kinixys	Kinixys erosa	ZMB 50198	271.0	27.10	31.5	18.5	26.0	231.0	15.9	Ε	ч	Africa	freshly measured (MFN collection)
Chersina	Chersina angulata	ZMB 37392	181.0	18.10	22.5	11.6	19.0	177.0	9.7	Ε	L	Africa	freshly measured (MFN collection)
Psammobates	Psammobates oculifer	1	147.0	14.70	N	N A	N A	Ϋ́	Α̈́	Ε	_	Africa	Itescu, Y., Karraker, N. E., Raia, P., Pritchard, P. C.,
Psammobates	Psammobates tentorius	1	145.0	14.50	N	N A	N A	Ϋ́	Α̈́	Ε	_	Africa	Itescu, Y., Karraker, N. E., Raia, P., Pritchard, P. C.,
Pyxis	Pyxis arachnoides	1	150.0	15.00	N	N A	N A	Ν	¥ V	Ε	>	Africa	Itescu, Y., Karraker, N. E., Raia, P., Pritchard, P. C.,
Psammobates	Psammobates geometricus	ZMB 185	118.0	11.80	18.0	9.1	16.5	112.0	8.2	Ε	_	Africa	freshly measured (MFN collection)
Stigmochelys	Stigmochelys pardalis	1	720.0	72.00	NA	NA	A	Ϋ́	N A	Ε	_	Africa	Itescu, Y., Karraker, N. E., Raia, P., Pritchard, P. C.,
Chersina	Chersina angulata	1	179.3	17.93	NA	N A	A	Ϋ́	ΑĀ	Ε	L	Africa	Itescu, Y., Karraker, N. E., Raia, P., Pritchard, P. C.,
Astrochelys	Astrochelys radiata	1	355.0	35.50	N	Α	Α	Ϋ́	Ϋ́	Ε	>	Africa	Pedrono, M., & Smith, L. L. (2013). Overview of the
Pyxis	Pyxis planicauda	1	126.0	12.60	N	N A	N A	Ϋ́	Α̈́	Ε	>	Africa	Pedrono, M., & Smith, L. L. (2013). Overview of the
Testudo	Testudo kleinmanni	1	144.0	14.40	NA	NA	N	Ϋ́	Υ Y	Ε	_	Africa	Itescu, Y., Karraker, N. E., Raia, P., Pritchard, P. C.,
Cylindraspis	Cylindraspis indica	1	0.009	00.09	N	N A	N A	Ϋ́	Υ Y	Ε	>	Africa	Itescu, Y., Karraker, N. E., Raia, P., Pritchard, P. C.,
Astrochelys	Astrochelys yniphora	1	361.0	36.10	N	N A	N A	Ν	¥ V	Ε	>	Africa	Pedrono, M., & Smith, L. L. (2013). Overview of the
Astrochelys	Astrochelys yniphora	1	486.0	48.60	N	Α	Α	Ϋ́	Ϋ́	E	>	Africa	Pedrono, M., & Smith, L. L. (2013). Overview of the
Pyxis	Pyxis planicauda	1	148.0	14.80	NA	N A	A	Ϋ́	ΑĀ	Ε	>	Africa	Pedrono, M., & Smith, L. L. (2013). Overview of the
Pyxis	Pyxis arachnoides	1	111.0	11.10	N	N A	Α	Ν	Α̈́	E	>	Africa	Pedrono, M., & Smith, L. L. (2013). Overview of the
Pyxis	Pyxis arachnoides		110.0	11.00	N	N A	Α	Ν	Ą	Ε	>	Africa	Pedrono, M., & Smith, L. L. (2013). Overview of the

Genus	Taxon	CollNr	SCL	CCL	SCW	CCW	CH	PL	PW	estimated	Island	Con	Reference
Pyxis	Pyxis arachnoides		80.0	8.00	AN	A N	A N	A N	¥	E	>	Africa	Pedrono, M., & Smith, L. L. (2013). Overview of the
Kinixys	Kinixys lobatsiana	1	200.0	20.00	Ν	Ν	N	N	Ν	E	⊑	Africa	Itescu, Y., Karraker, N. E., Raia, P., Pritchard, P. C.,
Pyxis	Pyxis arachnoides	1	86.0	8.60	Ϋ́	Ϋ́	N A	N	Ą	Е	>	Africa	Pedrono, M., & Smith, L. L. (2013). Overview of the
Pyxis	Pyxis arachnoides	1	154.0	15.40	Ϋ́	Ν	N	N	ΑN	Е	>	Africa	Pedrono, M., & Smith, L. L. (2013). Overview of the
Kinixys	Kinixys homeana	1	223.0	22.30	Ν	Ν	N	N	Ą	E	⊑	Africa	Itescu, Y., Karraker, N. E., Raia, P., Pritchard, P. C.,
Homopus	Homopus femoralis	1	168.0	16.80	Ϋ́	Ϋ́	N A	N	ΑN	Е	_	Africa	Itescu, Y., Karraker, N. E., Raia, P., Pritchard, P. C.,
Pyxis	Pyxis planicauda	1	132.0	13.20	Ϋ́	Ϋ́	N A	N	Ą	Е	>	Africa	Pedrono, M., & Smith, L. L. (2013). Overview of the
Homopus	Homopus aerolatus	1	300.0	30.00	Ϋ́	Ϋ́	N A	N	Ą	Е	L	Africa	Itescu, Y., Karraker, N. E., Raia, P., Pritchard, P. C.,
Homopus	Homopus boulengeri	1	110.0	11.00	Ν	Ν	N	N	Ν	E	⊑	Africa	Itescu, Y., Karraker, N. E., Raia, P., Pritchard, P. C.,
Kinixys	Kinixys erosa	1	400.0	40.00	Ϋ́	Ν	N	N	ΑN	Е	<b>⊑</b>	Africa	Itescu, Y., Karraker, N. E., Raia, P., Pritchard, P. C.,
Chersina	Chersina angulata	ZMB 37479	148.0	14.80	20.0	10.1	17.0	142.0	9.5	Ε	⊑	Africa	freshly measured (MFN collection)
Psammobates	Psammobates geometricus	1	165.0	16.50	Ν	Ν	N	N	Ą	E	⊑	Africa	Itescu, Y., Karraker, N. E., Raia, P., Pritchard, P. C.,
Homopus	Homopus solus	1	109.0	10.90	Ϋ́	Ϋ́	N A	N	ΑN	Е	_	Africa	Itescu, Y., Karraker, N. E., Raia, P., Pritchard, P. C.,
Malacochersus	Malacochersus tornieri	1	180.0	18.00	Ϋ́	Ϋ́	N A	NA	Ą	E	_	Africa	Itescu, Y., Karraker, N. E., Raia, P., Pritchard, P. C.,
Chersina	Chersina angulata	1	153.5	15.35	Ϋ́	Ϋ́	N A	Ν	Α̈́	E	_	Africa	Itescu, Y., Karraker, N. E., Raia, P., Pritchard, P. C.,
Pyxis	Pyxis arachnoides	1	144.0	14.40	Ϋ́	Ϋ́	N A	NA	Ϋ́	E	>	Africa	Pedrono, M., & Smith, L. L. (2013). Overview of the
Kinixys	Kinixys belliana	1	230.0	23.00	Ϋ́	Ϋ́	N A	NA	Ϋ́	E	ㅁ	Africa	Itescu, Y., Karraker, N. E., Raia, P., Pritchard, P. C.,
Aldabrachelys	Aldabrachelys gigantea	1	1140.0	114.00	Ϋ́	Ϋ́	N A	NA	Ϋ́	E	>	Africa	Itescu, Y., Karraker, N. E., Raia, P., Pritchard, P. C.,
Astrochelys	Astrochelys radiata	1	400.0	40.00	Ϋ́	Ϋ́	N A	N	Ϋ́	E	>	Africa	Itescu, Y., Karraker, N. E., Raia, P., Pritchard, P. C.,
Chersina	Chersina angulata	1	166.4	16.64	Ϋ́	Ϋ́	N A	N	Ϋ́	E	_	Africa	Itescu, Y., Karraker, N. E., Raia, P., Pritchard, P. C.,
Chersina	Chersina angulata	1	171.6	17.16	Ϋ́	Ϋ́	N A	Ν	Α̈́	E	>	Africa	Itescu, Y., Karraker, N. E., Raia, P., Pritchard, P. C.,
Cylindraspis	Cylindraspis peltastes	1	420.0	42.00	Ϋ́	Ϋ́	N A	NA	Ϋ́	E	>	Africa	Itescu, Y., Karraker, N. E., Raia, P., Pritchard, P. C.,
Chersina	Chersina angulata	1	161.3	16.13	Ν	Ϋ́	N A	ΑN	Α	E	>	Africa	Itescu, Y., Karraker, N. E., Raia, P., Pritchard, P. C.,
Homopus	Homopus signatus	1	106.0	10.60	Ϋ́	Ϋ́	N A	NA	Ϋ́	E	ㅁ	Africa	Itescu, Y., Karraker, N. E., Raia, P., Pritchard, P. C.,
Kinixys	Kinixys spekii	1	220.0	22.00	Ν	Ϋ́	NA	ΑN	Ϋ́	E	_	Africa	Itescu, Y., Karraker, N. E., Raia, P., Pritchard, P. C.,
Cylindraspis	Cylindraspis inepta	1	1000.0	100.00	Ϋ́	Ϋ́	N A	N	Ϋ́	E	>	Africa	Itescu, Y., Karraker, N. E., Raia, P., Pritchard, P. C.,
Kinixys	Kinixys natalensis	1	160.0	16.00	Ν	Ϋ́	N A	ΑN	Α	E	_	Africa	Itescu, Y., Karraker, N. E., Raia, P., Pritchard, P. C.,
Geochelone	Geochelone elegans	ZMB 63222	208.0	20.80	29.5	14.6	28.5	199.0	13.3	Ε	ㅁ	Asia	freshly measured (MFN collection)

Genus	Taxon	CollNr	SCL	CCL	SCW	CCW	공	PL	PW	estimated	Island	Con	Reference
Geochelone	Geochelone elegans	ZMB 37523	245.0	24.50	32.0	16.6	32.0	228.0	14.6	E	ے	Asia	freshly measured (MFN collection)
Geochelone	Geochelone elegans	ZMB 63220	221.0	22.10	32.0	16.0	31.0	179.0	13.5	Ε	_	Asia	freshly measured (MFN collection)
Geochelone	Geochelone elegans	ZMB 63221	220.0	22.00	31.0	15.4	27.0	209.0	4	Ε	>	Asia	freshly measured (MFN collection)
Geochelone	Geochelone elegans	ZMB 63218	221.0	22.10	31.5	15.1	30.0	203.0	13.7	E	u	Asia	freshly measured (MFN collection)
Geochelone	Geochelone platynota	ZMB 6096	222.0	22.20	29.5	15.1	27.0	Ν	MA	Ε	L	Asia	freshly measured (MFN collection)
Manouria	Manouria emys	ı	0.009	00.09	N	N A	N A	Ν	Ą	Ε	_	Asia	Itescu, Y., Karraker, N. E., Raia, P., Pritchard, P. C.,
Indotestudo	Indotestudo forstenii	ı	202.0	20.20	N	N A	N A	Ν	N A	Ε	>	Asia	Itescu, Y., Karraker, N. E., Raia, P., Pritchard, P. C.,
Indotestudo	Indotestudo travancorica	ı	249.7	24.97	N	N A	N A	Ν	N A	Ε	_	Asia	Itescu, Y., Karraker, N. E., Raia, P., Pritchard, P. C.,
Indotestudo	Indotestudo forstenii	ı	309.0	30.90	N	Ϋ́	Ϋ́	Ν	ΑN	Ε	>	Asia	Itescu, Y., Karraker, N. E., Raia, P., Pritchard, P. C.,
Indotestudo	Indotestudo elongata	ı	360.0	36.00	NA	Ϋ́	Ϋ́	Ν	ΑŽ	Ε	u	Asia	Itescu, Y., Karraker, N. E., Raia, P., Pritchard, P. C.,
Indotestudo	Indotestudo forstenii	ı	199.0	19.90	NA	Ϋ́	Ϋ́	Ν	A	Ε	>	Asia	Itescu, Y., Karraker, N. E., Raia, P., Pritchard, P. C.,
Indotestudo	Indotestudo elongata	ı	244.2	24.42	NA	Ϋ́	Ϋ́	Ν	A	Ε	L	Asia	Itescu, Y., Karraker, N. E., Raia, P., Pritchard, P. C.,
Indotestudo	Indotestudo travancorica	ı	244.2	24.42	NA	Ϋ́	Ϋ́	Ν	Ϋ́	Ε	L	Asia	Itescu, Y., Karraker, N. E., Raia, P., Pritchard, P. C.,
Manouria	Manouria impressa	ZMB 63172	165.0	16.50	20.0	12.9	18.0	157.0	10.5	Ε	c	Asia	freshly measured (MFN collection)
Indotestudo	Indotestudo elongata	ZMB 50492	276.0	27.60	33.0	19.4	28.5	246.0	17.1	Ε	c	Asia	freshly measured (MFN collection)
Indotestudo	Indotestudo elongata	ZMB 63175	235.0	23.50	30.5	16.0	29.5	202.0	14.4	Ε	۵	Asia	freshly measured (MFN collection)
Indotestudo	Indotestudo elongata	ZMB 4174	208.0	20.80	26.0	13.4	20.0	180.0	11.6	Ε	_	Asia	freshly measured (MFN collection)
Indotestudo	Indotestudo elongata	ZMB 6106	166.0	16.60	21.0	11.3	18.0	151.0	11.3	Ε	_	Asia	freshly measured (MFN collection)
Manouria	Manouria emys	ı	0.009	00.09	N	N A	N A	Ν	Ā	E	_	Asia	Karl, H., & Staesche, U. (2007). Fossile Riesen-Lar
Testudo	Testudo graeca	ı	250.0	25.00	N	N A	N A	Ν	Ā	E	_	Asia	Itescu, Y., Karraker, N. E., Raia, P., Pritchard, P. C.,
Testudo	Testudo graeca	1	280.0	28.00	N	Ϋ́	Ϋ́	Ν	Ā	Ε	>	Asia	Itescu, Y., Karraker, N. E., Raia, P., Pritchard, P. C.,
Manouria	Manouria emys	ZMB 49049	212.0	21.20	26.5	16.5	25.0	Ν	A	Ε	c	Asia	freshly measured (MFN collection)
Manouria	Manouria emys	ZMB 37350	445.0	44.50	52.0	32.0	20.0	455.0	29.8	Ε	۵	Asia	freshly measured (MFN collection)
Manouria	Manouria emys	ZMB 37342	330.0	33.00	40.5	26.7	37.0	330.0	23.4	Ε	L	Asia	freshly measured (MFN collection)
Indotestudo	Indotestudo travancorica	ı	331.0	33.10	NA	Ϋ́	Ϋ́	Ν	A	Ε	L	Asia	Itescu, Y., Karraker, N. E., Raia, P., Pritchard, P. C.,
Indotestudo	Indotestudo travancorica	ı	219.6	21.96	NA	Ϋ́	Ϋ́	Ν	A	Ε	L	Asia	Itescu, Y., Karraker, N. E., Raia, P., Pritchard, P. C.,
Indotestudo	Indotestudo forstenii	ı	200.5	20.05	NA	Ϋ́	Ϋ́	Ν	A	Ε	>	Asia	Itescu, Y., Karraker, N. E., Raia, P., Pritchard, P. C.,
Testudo	Testudo horsfieldii	ı	280.0	28.00	N	N A	N A	Ν	N A	Ε	c	Asia	Itescu, Y., Karraker, N. E., Raia, P., Pritchard, P. C.,

Genus	Taxon	CollNr	SCL	CCL	SCW	CCW	ᆼ	Ч	PW	estimated	Island	Con	Reference
Manouria	Manouria impressa	1	350.0	35.00	NA	ΝΑ	NA	NA	AN	ш	u	Asia	Itescu, Y., Karraker, N. E., Raia, P., Pritchard, P. C.,
Geochelone	Geochelone elegans	ı	380.0	38.00	Ν	Ą	Ϋ́	Ν A	ΑN	Ε	⊑	Asia	Itescu, Y., Karraker, N. E., Raia, P., Pritchard, P. C.,
Manouria	Manouria impressa	1	275.0	27.50	Ν	Ϋ́	Ą	NA	Α̈́	E	п	Asia	Karl, H., & Staesche, U. (2007). Fossile Riesen-Lar
Indotestudo	Indotestudo elongata	1	219.6	21.96	Ν	Ϋ́	Ą	NA	Α̈́	E	п	Asia	Itescu, Y., Karraker, N. E., Raia, P., Pritchard, P. C.,
Geochelone	Geochelone platynota	ı	300.0	30.00	Ν	¥	Ϋ́	N	ΑN	E	L	Asia	Itescu, Y., Karraker, N. E., Raia, P., Pritchard, P. C.,
Testudo	Testudo graeca	1	300.0	30.00	Ν	¥	Ą	Ν	Ą	Ε	_	Asia	Itescu, Y., Karraker, N. E., Raia, P., Pritchard, P. C.,
Gopherus	Gopherus flavomarginatus	ı	400.0	40.00	Ν	Ą	Ϋ́	Ν A	ΑN	Ε	⊑	America	Itescu, Y., Karraker, N. E., Raia, P., Pritchard, P. C.,
Gopherus	Gopherus morafkai	1	299.0	29.90	Ν	¥	Ą	N	Ą	E	п	America	Itescu, Y., Karraker, N. E., Raia, P., Pritchard, P. C.,
Gopherus	Gopherus berlandieri	1	240.0	24.00	Ν	¥	Ą	N	Ą	E	п	America	Itescu, Y., Karraker, N. E., Raia, P., Pritchard, P. C.,
Testudo	Testudo horsfieldii	ZMB 63259	111.0	11.10	14.0	10.0	15.0	108.0	9.5	E	L	Europe	freshly measured (MFN collection)
Pyxis	Pyxis arachnoides	ZMB 37615	108.0	10.80	15.0	7.9	13.0	0.96	7.1	E	L	Europe	freshly measured (MFN collection)
Testudo	Testudo marginata	ı	241.7	24.17	Ν	¥	A A	N	Ϋ́	Ε	ᄆ	Europe	Willemsen, R. E., & Hailey, A. (2003). Sexual dimor
Testudo	Testudo horsfieldii	ZMB 63258	123.0	12.30	14.5	10.9	15.0	121.0	8.6	Ε	ᄆ	Europe	freshly measured (MFN collection)
Testudo	Testudo hermanni	ı	183.3	18.33	Ν	¥	N A	N	Α̈́	Ε	>	Europe	Itescu, Y., Karraker, N. E., Raia, P., Pritchard, P. C.,
Testudo	Testudo hermanni	ı	176.9	17.69	Ν	¥	N A	N	Α̈́	Ε	_	Europe	Willemsen, R. E., & Hailey, A. (2003). Sexual dimor
Testudo	Testudo horsfieldii	ZMB 63257	114.0	11.40	14.5	10.2	14.0	110.0	6.6	٤	۵	Europe	freshly measured (MFN collection)
Testudo	Testudo marginata	ı	246.7	24.67	Ν	Ϋ́	¥ N	N	¥.	Ε	_	Europe	Willemsen, R. E., & Hailey, A. (2003). Sexual dimor
Testudo	Testudo hermanni	ı	196.0	19.60	Ν	Ϋ́	A A	NA	Ϋ́	E	L	Europe	Willemsen, R. E., & Hailey, A. (2003). Sexual dimor
Testudo	Testudo hermanni	ı	143.5	14.35	Ν	¥	A A	N	Ϋ́	Ε	>	Europe	Itescu, Y., Karraker, N. E., Raia, P., Pritchard, P. C.,
Testudo	Testudo graeca	ı	194.6	19.46	Ν	¥	A A	N	Ϋ́	Ε	ᄆ	Europe	Willemsen, R. E., & Hailey, A. (2003). Sexual dimor
Testudo	Testudo hermanni	ı	200.0	20.00	Ν	¥	N A	N	Α̈́	Ε	>	Europe	Itescu, Y., Karraker, N. E., Raia, P., Pritchard, P. C.,
Testudo	Testudo hermanni	ı	250.0	25.00	Ν	A A	A A	NA	Ϋ́	Ε	ч	Europe	Itescu, Y., Karraker, N. E., Raia, P., Pritchard, P. C.,
Testudo	Testudo marginata	ī	246.0	24.60	N	Α̈́	ΝΑ	Ν	ΑĀ	٤	۵	Europe	Itescu, Y., Karraker, N. E., Raia, P., Pritchard, P. C.,
Testudo	Testudo marginata	ī	242.5	24.25	N	Α̈́	ΝΑ	Ν	ΑĀ	٤	>	Europe	Itescu, Y., Karraker, N. E., Raia, P., Pritchard, P. C.,
Testudo	Testudo marginata	I	246.0	24.60	Ν	A A	N A	Ν	Ϋ́	Ε	п	Europe	Itescu, Y., Karraker, N. E., Raia, P., Pritchard, P. C.,
Testudo	Testudo hermanni	ı	147.0	14.70	Ν	¥	A A	N	Ϋ́	Ε	ᄆ	Europe	Itescu, Y., Karraker, N. E., Raia, P., Pritchard, P. C.,
Testudo	Testudo marginata	ı	290.0	29.00	N	Α̈́	Ν	Ν	ΑĀ	٤	_	Europe	Itescu, Y., Karraker, N. E., Raia, P., Pritchard, P. C.,
Testudo	Testudo marginata	1	250.0	25.00	N	ΑΝ	A A	Ν	Ā	Ε	>	Europe	Itescu, Y., Karraker, N. E., Raia, P., Pritchard, P. C.,

Genus	Taxon	CollNr	SCL	CCL	SCW	CCW	끙	PL	PW	estimated	Island	Con	Reference
Testudo	Testudo hermanni		145.9	14.59	NA	A A	AA	NA	A A	E	>	Europe	Itescu, Y., Karraker, N. E., Raia, P., Pritchard, P. C.,
Testudo	Testudo graeca	ı	178.2	17.82	Ν Α	Ϋ́	A	ΑN	Ν	Ε	L	Europe	Willemsen, R. E., & Hailey, A. (2003). Sexual dimor
Testudo	Testudo marginata	1	400.0	40.00	Ν Α	Ϋ́	NA	ΑN	Ą	E	u	Europe	Itescu, Y., Karraker, N. E., Raia, P., Pritchard, P. C.,
Testudo	Testudo horsfieldii	ZMB 63255	136.0	13.60	18.0	13.0	16.5	129.0	12.2	E	u	Europe	freshly measured (MFN collection)
Testudo	Testudo horsfieldii	ZMB 63256	132.0	13.20	17.0	12.4	17.0	133.0	11.3	E	u	Europe	freshly measured (MFN collection)
Testudo	Testudo hermanni	1	168.3	16.83	Ν Α	Ϋ́	N	ΑN	Ą	Ε	>	Europe	Itescu, Y., Karraker, N. E., Raia, P., Pritchard, P. C.,
Testudo	Testudo hermanni	ı	160.0	16.00	Ν Α	Ϋ́	A	ΑN	Ν	Ε	>	Europe	Itescu, Y., Karraker, N. E., Raia, P., Pritchard, P. C.,
Testudo	Testudo hermanni	1	154.0	15.40	ΑN	Ϋ́	Ν	ΑN	Ą	Ε	L	Europe	Itescu, Y., Karraker, N. E., Raia, P., Pritchard, P. C.,
Testudo	Testudo hermanni	1	138.5	13.85	Ν A	Ϋ́	N	ΑN	Ą	E	u	Europe	Itescu, Y., Karraker, N. E., Raia, P., Pritchard, P. C.,
Testudo	Testudo hermanni	1	173.0	17.30	Ν Α	Ϋ́	NA	ΑN	Ą	E	>	Europe	Itescu, Y., Karraker, N. E., Raia, P., Pritchard, P. C.,
Testudo	Testudo marginata	ı	242.5	24.25	Ν Α	Ϋ́	Ν	ΑN	Ϋ́	E	>	Europe	Itescu, Y., Karraker, N. E., Raia, P., Pritchard, P. C.,
Testudo	Testudo hermanni	ı	195.0	19.50	Ν	¥	Α	ΑN	Ϋ́	E	>	Europe	Itescu, Y., Karraker, N. E., Raia, P., Pritchard, P. C.,
Testudo	Testudo hermanni	ı	157.0	15.70	Ν	¥	Α	ΑN	Ϋ́	E	>	Europe	Itescu, Y., Karraker, N. E., Raia, P., Pritchard, P. C.,
Testudo	Testudo hermanni	ı	176.6	17.66	Ν	¥	Ϋ́	Ν	Α̈́	E	>	Europe	Itescu, Y., Karraker, N. E., Raia, P., Pritchard, P. C.,
Testudo	Testudo hermanni	ı	130.0	13.00	Ν	¥	Ϋ́	Ν	Α̈́	E	c	Europe	Itescu, Y., Karraker, N. E., Raia, P., Pritchard, P. C.,
Testudo	Testudo hermanni	ı	161.0	16.10	N	N A	Ϋ́	Ν	Α	E	c	Europe	Itescu, Y., Karraker, N. E., Raia, P., Pritchard, P. C.,
Gopherus	Gopherus polyphemus	ı	300.0	30.00	Ν	¥.	Ϋ́	Ν	¥	E	>	America	Itescu, Y., Karraker, N. E., Raia, P., Pritchard, P. C.,
Gopherus	Gopherus sp.	MVZ 210020	Ϋ́	N	Ν Α	Ϋ́	Ν	219.6	Ϋ́	E	L	America	Biewer J., Sankey J., Hutchison H., Garber D., 2016
Gopherus	Gopherus sp.	MVZ 210003	Ν	N	Ν	¥	Α	192.1	Ϋ́	E	L	America	Biewer J., Sankey J., Hutchison H., Garber D., 2016
Gopherus	Gopherus polyphemus	ı	268.8	26.88	Ν	¥	Α	ΑN	Ϋ́	E	>	America	Itescu, Y., Karraker, N. E., Raia, P., Pritchard, P. C.,
Gopherus	Gopherus sp.	MVZ 120004	Ν	N	Ν	¥	Ϋ́	196.7	Α̈́	E	c	America	Biewer J., Sankey J., Hutchison H., Garber D., 2016
Gopherus	Gopherus sp.	MVZ 210009	Ϋ́	ΑΝ	Ν Α	ΑΝ	N A	232.8	Ϋ́	E	L	America	Biewer J., Sankey J., Hutchison H., Garber D., 2016
Gopherus	Gopherus sp.	MVZ 210010	N	N	N	N A	Ϋ́	240.1	Α	E	c	America	Biewer J., Sankey J., Hutchison H., Garber D., 2016
Gopherus	Gopherus agassizii	ı	400.0	40.00	N	N A	Ϋ́	Ν	Α	E	c	America	Itescu, Y., Karraker, N. E., Raia, P., Pritchard, P. C.,
Gopherus	Gopherus flavomarginatus	KU 39415	303.0	30.30	Ν	23.2	Α	ΑN	Ϋ́	E	L	America	Legler, 1959
Gopherus	Gopherus polyphemus	ı	308.0	30.80	Ν	¥	Α	ΑN	Ϋ́	E	L	America	Itescu, Y., Karraker, N. E., Raia, P., Pritchard, P. C.,
Gopherus	Gopherus polyphemus	ı	303.0	30.30	Ν	¥	Α	ΑN	Ϋ́	E	>	America	Itescu, Y., Karraker, N. E., Raia, P., Pritchard, P. C.,
Gopherus	Gopherus polyphemus		387.0	38.70	N A	N A	Ϋ́	NA	Ą	E	_	America	Itescu, Y., Karraker, N. E., Raia, P., Pritchard, P. C.,

Genus	Taxon	CollNr	SCL	CCL	SCW	ccw	CH	PL	ΡW	estimated	Island	Con	Reference
Gopherus	Gopherus polyphemus		342.0	34.20	A	¥ Y	A A	NA	¥	E		America	Itescu, Y., Karraker, N. E., Raia, P., Pritchard, P. C.,
Gopherus	Gopherus flavomarginatus	<b>USNM</b> 61253	222.0	22.20	N	16.6	Ϋ́	212.0	N A	Ε	_	America	Legler, 1959
Gopherus	Gopherus flavomarginatus	USNM 61254	371.0	37.10	NA	29.2	Ϋ́	358.0	ΑĀ	Ε	_	America	Legler, 1959
Gopherus	Gopherus polyphemus	1	238.9	23.89	NA	ΑN	Ϋ́	NA	ΑĀ	Ε	_	America	Itescu, Y., Karraker, N. E., Raia, P., Pritchard, P. C.,
Gopherus	Gopherus flavomarginatus	92609 WNSN	246.0	24.60	N	21.2	Ν	252.0	N A	Ε	_	America	Legler, 1959
Gopherus	Gopherus flavomarginatus	IU 42953	281.0	28.10	N	22.0	Ϋ́	Ν	ΑĀ	Ε	_	America	Legler, 1959
Gopherus	Gopherus flavomarginatus	IU 42954	278.0	27.80	N	21.4	Ϋ́	Ν A	N A	Ε	_	America	Legler, 1959
Chelonoidis	Chelonoidis nigra	USNM 51069	588.0	58.80	68.3	44.5	Ν	506.0	N A	Ε	>	America	Franz, R., & Franz, S. E. (2009). A new fossil land t
Chelonoidis	Chelonoidis nigra	USNM1 102904	610.0	61.00	67.5	44.4	Ϋ́	515.0	N A	Ε	>	America	Franz, R., & Franz, S. E. (2009). A new fossil land t
Chelonoidis	Chelonoidis carbonaria	1	593.0	59.30	NA	Ϋ́	Ϋ́	Ν	ΑĀ	Ε	_	America	Itescu, Y., Karraker, N. E., Raia, P., Pritchard, P. C.,
Chelonoidis	Chelonoidis abingdonii	1	0.086	98.00	NA	N A	Ϋ́	Ν	ΑĀ	Ε	>	America	Itescu, Y., Karraker, N. E., Raia, P., Pritchard, P. C.,
Chelonoidis	Chelonoidis denticulata	1	333.4	33.34	NA	¥	Š	Ν	ΑŽ	Ε	_	America	Itescu, Y., Karraker, N. E., Raia, P., Pritchard, P. C.,
Chelonoidis	Chelonoidis chilensis	UF33604	169.0	16.90	21.5	13.2	Š	161.0	ΑŽ	Ε	_	America	Franz, R., & Franz, S. E. (2009). A new fossil land t
Chelonoidis	Chelonoidis chilensis	UF33618	186.0	18.60	25.0	14.7	Ν Α	169.0	A A	Ε	c	America	Franz, R., & Franz, S. E. (2009). A new fossil land t
Chelonoidis	Chelonoidis nigra	ı	717.0	71.70	N	N A	Ν Α	Ν	A A	Ε	>	America	Itescu, Y., Karraker, N. E., Raia, P., Pritchard, P. C.,
Chelonoidis	Chelonoidis chilensis	UF33617	169.0	16.90	22.8	14.6	N A	162.0	N A	Ε	_	America	Franz, R., & Franz, S. E. (2009). A new fossil land t
Chelonoidis	Chelonoidis carbonaria	UF27384	242.0	24.20	31.7	15.5	Ϋ́	219.0	A A	Ε	_	America	Franz, R., & Franz, S. E. (2009). A new fossil land t
Chelonoidis	Chelonoidis carbonaria	UF33597	253.0	25.30	31.7	15.3	Ϋ́	215.0	ΑĀ	Ε	_	America	Franz, R., & Franz, S. E. (2009). A new fossil land t
Chelonoidis	Chelonoidis nigra	USNM1 222494	595.0	59.50	0.89	43.6	Ν Α	533.0	N A	Ε	>	America	Franz, R., & Franz, S. E. (2009). A new fossil land t
Chelonoidis	Chelonoidis carbonaria	1	333.4	33.34	NA	N A	Ϋ́	Ν	ΑĀ	Ε	_	America	Itescu, Y., Karraker, N. E., Raia, P., Pritchard, P. C.,
Chelonoidis	Chelonoidis carbonaria	UF5259	226.0	22.60	28.7	12.9	Ν Α	198.0	A A	Ε	c	America	Franz, R., & Franz, S. E. (2009). A new fossil land t
Chelonoidis	Chelonoidis becki	ı	1050.0	105.00	N	N A	Ν Α	Ν	A A	Ε	>	America	Itescu, Y., Karraker, N. E., Raia, P., Pritchard, P. C.,
Chelonoidis	Chelonoidis denticulata	UF33661	333.0	33.30	38.0	21.4	N A	305.0	N A	Ε	_	America	Franz, R., & Franz, S. E. (2009). A new fossil land t
Chelonoidis	Chelonoidis denticulata	UF61931	317.0	31.70	41.2	18.5	Ϋ́	291.0	ΑĀ	Ε	_	America	Franz, R., & Franz, S. E. (2009). A new fossil land t
Chelonoidis	Chelonoidis denticulata	UF33670	365.0	36.50	47.0	22.0	Ϋ́	326.0	ΑĀ	Ε	_	America	Franz, R., & Franz, S. E. (2009). A new fossil land t
Chelonoidis	Chelonoidis chilensis	UF33603	183.0	18.30	23.4	14.5	Ϋ́	166.0	ΑĀ	Ε	_	America	Franz, R., & Franz, S. E. (2009). A new fossil land t
Chelonoidis	Chelonoidis nigra	ı	731.3	73.13	N	N A	N A	Ν	N A	E	>	America	Itescu, Y., Karraker, N. E., Raia, P., Pritchard, P. C.,
Chelonoidis	Chelonoidis chilensis	1	200.0	20.00	N	Ą	Ν Α	Ν	A A	Ε	۵	America	Itescu, Y., Karraker, N. E., Raia, P., Pritchard, P. C.,

Genus	Taxon	CollNr	SCL	CCL	SCW	CCW	끙	٦ ٦	ΡW	estimated	Island	Con	Reference
Chelonoidis	Chelonoidis carbonaria	UF48278	247.0	24.70	33.9	15.5	N A	214.0	Ą	E	u	America	Franz, R., & Franz, S. E. (2009). A new fossil land t
Chelonoidis	Chelonoidis carbonaria	1	296.5	29.62	Ν	N	N A	Ν	Ą	E	Ц	America	Itescu, Y., Karraker, N. E., Raia, P., Pritchard, P. C.,
Chelonoidis	Chelonoidis carbonaria	1	290.0	29.00	Ν	N	Ϋ́	Ν	Ą	E	>	America	Itescu, Y., Karraker, N. E., Raia, P., Pritchard, P. C.,
Chelonoidis	Chelonoidis carbonaria	UF33596	189.0	18.90	24.7	12.1	Ϋ́	174.0	Ą	E	п	America	Franz, R., & Franz, S. E. (2009). A new fossil land t
Chelonoidis	Chelonoidis nigra	1	745.7	74.57	Ϋ́	N	N A	Ν	Ą	E	>	America	Itescu, Y., Karraker, N. E., Raia, P., Pritchard, P. C.,
Chelonoidis	Chelonoidis chathamensis	1	0.068	89.00	Ϋ́	N	N A	Ν	Ą	E	>	America	Itescu, Y., Karraker, N. E., Raia, P., Pritchard, P. C.,
Chelonoidis	Chelonoidis denticulata	UF19242	466.0	46.60	29.7	26.5	N A	410.0	Ą	E	Ц	America	Franz, R., & Franz, S. E. (2009). A new fossil land t
Chelonoidis	Chelonoidis denticulata	UF23231	377.0	37.70	47.1	23.8	N A	334.0	Ą	E	Ц	America	Franz, R., & Franz, S. E. (2009). A new fossil land t
Chelonoidis	Chelonoidis denticulata	1	820.0	82.00	Ν	N	N A	Ν	Ą	E	Ц	America	Itescu, Y., Karraker, N. E., Raia, P., Pritchard, P. C.,
Chelonoidis	Chelonoidis duncanensis	ı	840.0	84.00	N	Υ Y	Ϋ́	Ν	Α̈́	E	>	America	Itescu, Y., Karraker, N. E., Raia, P., Pritchard, P. C.,
Chelonoidis	Chelonoidis chilensis	1	222.0	22.20	Ν	N	N A	Ν	Ą	E	ч	America	Itescu, Y., Karraker, N. E., Raia, P., Pritchard, P. C.,
Chelonoidis	Chelonoidis chilensis	UF33600	157.0	15.70	20.8	11.9	Ä	145.0	¥	E	ч	America	Franz, R., & Franz, S. E. (2009). A new fossil land t
Chelonoidis	Chelonoidis phantastica	ı	860.0	86.00	Ν	N A	Ä	Ν	¥	E	>	America	Itescu, Y., Karraker, N. E., Raia, P., Pritchard, P. C.,
Chelonoidis	Chelonoidis vicina	ı	1250.0	125.00	Ν	N A	Ä	Ν	Α̈́	E	>	America	Itescu, Y., Karraker, N. E., Raia, P., Pritchard, P. C.,
Chelonoidis	Chelonoidis hoodensis	ı	813.0	81.30	Ν	N A	Ä	Ν	Α̈́	E	>	America	Itescu, Y., Karraker, N. E., Raia, P., Pritchard, P. C.,
Chelonoidis	Chelonoidis nigra	ı	1300.0	130.00	N	Υ Y	Ϋ́	Ν	Α̈́	E	>	America	Itescu, Y., Karraker, N. E., Raia, P., Pritchard, P. C.,
Chelonoidis	Chelonoidis darwini	ı	965.0	96.50	N	Υ Y	Ϋ́	Ν	Α̈́	E	>	America	Itescu, Y., Karraker, N. E., Raia, P., Pritchard, P. C.,
Chelonoidis	Chelonoidis chilensis	-	450.0	45.00	NA	NA	NA	NA	NA	ш	u	America	Itescu, Y., Karraker, N. E., Raia, P., Pritchard, P. C.,

## Acknowledgements

**Declaration of Authorship**