AI1110 - ASSIGNMENT 5

J. TUSHITA SHARVA - CS21BTECH11022

May 30, 2022



Outline

Question

Solution

Question

Papoulis Chapter 2, Example 2.10

In a fair die experiment, S represents the set consisting of the faces $f_1, f_2, f_3, f_4, f_5, f_6$. Determine the conditional probability of the event $\{f_2\}$, assuming the event *even* occurred.

Concepts

Formulas

$$\Pr(E) = \frac{n(E)}{n(S)} \tag{2.0.1}$$

$$Pr(E) = \frac{n(E)}{n(S)}$$

$$Pr(E|F) = \frac{Pr(E, F)}{Pr(F)}$$
(2.0.1)



Solution

Let X be a random variable which is used to denote the outcome of the dice. Then we have, $X \in \{f_1, f_2, f_3, f_4, f_6\}$ where in f_k , k denotes the outcome of the dice.

We must find the value of

$$Pr(X = f_2 \mid X = f_2 \cdot f_4 \cdot f_6)$$
 (2.0.3)

$$\Pr(X = k) = \begin{cases} \frac{1}{6}, & 1 \le k \le 6\\ 0, & otherwise \end{cases}$$
 (2.0.4)



$$\Pr(X = f_2 \mid X = f_2, f_4, f_6) = \frac{\Pr(X = f_2 \cdot f_2, f_4, f_6)}{\Pr(X = f_2, f_4, f_6)}$$
(2.0.5)

Since the events f_2 , f_4 , f_6 are mutually exclusive, we have

$$\Pr(X = f_2, f_4, f_6) = \Pr(X = f_2) + \Pr(X = f_4) + \Pr(X = f_6) \qquad (2.0.6)$$

$$\implies \Pr(X = f_2, f_4, f_6) = \frac{1}{6} + \frac{1}{6} + \frac{1}{6} \qquad (2.0.7)$$

$$\implies \Pr(X = f_2, f_4, f_6) = \frac{3}{6} = \frac{1}{2} \qquad (2.0.8)$$

We see that $\{f_2\}$ is a subset of $\{f_2, f_4, f_6\}$.

We know that if

$$A \subset B \ then, A \cdot B = A.$$
 (2.0.9)

Therefore

$$\Pr(X = f_2 \cdot f_2, f_4, f_6) = \Pr(f_2) = \frac{1}{6}$$
 (2.0.10)

Therefore, by substituting the results (2.0.9) and (2.0.11), in the equation (2.0.5), we have

$$\Pr(X = f_2 \mid X = f_2, f_4, f_6) = \frac{1/6}{1/2}$$
 (2.0.11)

$$\implies \Pr(X = f_2 \mid X = f_2, f_4, f_6) = \frac{1}{6} \times \frac{2}{1}$$
 (2.0.12)

$$\implies \Pr(X = f_2 \mid X = f_2, f_4, f_6) = \frac{1}{3}$$
 (2.0.13)

Source codes

Download Latex source code of this pdf from:

Click Here

Download document of this presentation at:

Click Here

To download this presentation, visit:

Click here