

AI1110 ASSIGNMENT 8

JANGA TUSHITA SHARVA (CS21BTECH11022)

Abstract—This document gives solution to Chapter 7, Question 7.20 from the Papoulis and Pillai Probability, Random Variables and Stochastic Processes Text Book.

Download Latex source code of this pdf from:

[link](#)

Download Presentation of this document at:

[link](#)

To download this document, visit:

[link](#)

Case 2: $x > y$ then,

$$\bar{A}_X B_Y = \{\phi\} \quad (2.0.7)$$

Therefore, we have $F_{XY}(x, y) = \Pr(A_X B_Y) =$

$$\begin{cases} y^n - (y - x)^n & x \leq y \\ y^n & x > y \end{cases}$$

1. QUESTION

We place at random n points in the interval $(0, 1)$ and we denote by x and y the distance from the origin to the first and last point respectively. Find $F(x)$, $F(y)$, $F(x, y)$.

2. ANSWER

The event $\tilde{x} \leq x$ occurs if there is at least one point in the interval $(0, x)$; The event $\tilde{y} \leq y$ occurs if all the points are in the interval $(0, \tilde{y})$.

Let $A_X =$ at least one point in $(0, x) = \tilde{x} \leq x$

Let $B_Y =$ no points in $(y, 1)$

$=$ all points in $(0, y) = \tilde{y} \leq y$

Hence for $0 \leq x \leq 1$, $0 \leq y \leq 1$, we have:

$$F_X(x) = \Pr(A_X) = 1 - \Pr(\bar{A}_X) = 1 - (1 - x)^n \quad (2.0.1)$$

$$F_Y(y) = \Pr(B_Y) = y^n \quad (2.0.2)$$

Further more, we have

$$\left\{ \tilde{x} \leq x, \tilde{y} \leq y \right\} = A_x \cdot B_y \quad (2.0.3)$$

$$A_X B_Y + \bar{A}_X B_Y = B_Y \quad (2.0.4)$$

Case 1: $x \leq y$ then,

$$\bar{A}_x B_Y = \{\text{all points in } (x, y)\} \quad (2.0.5)$$

$$\implies \Pr(\bar{A}_X B_Y) = (y - x)^n \quad (2.0.6)$$