

Probability 201-1-2391 ASSIGNMENT 5-answers
Asymptotic laws: The theorems of DeMoivre, Laplace, Bernoulli and
Poisson
instructor: Ronen Peretz,
math dept, BGU

Problem 1

Give an estimate, using normal approximation, to the probability that in a 1000 tosses of a coin with a probability 0.2 for an H, we will get at least 180 H's.

$$p(N(0, 1) \geq \frac{179.5 - 200}{\sqrt{160}}) = 1 - \Phi\left(\frac{-20.5}{\sqrt{160}}\right) = 0.9747.$$

Problem 2

A balanced coin is tossed 300 times. Compute the probability that it will show 140 times 'H':

(a) Using the Stirling asymptotic formula, and

$$p = \binom{300}{140} \left(\frac{1}{2}\right)^{300} \approx \frac{300^{300}}{140^{140} \cdot 160^{160}} \sqrt{\frac{300}{2\pi \cdot 140 \cdot 160}} \left(\frac{1}{2}\right)^{300}$$

(b) Assuming the normal distribution.

What is the relative deviation between the two methods?

$$\text{By DeMoivre } p \approx \frac{1}{\sqrt{75}} \phi\left(\frac{140 - 150}{\sqrt{75}}\right) = \frac{1}{\sqrt{75}\sqrt{2\pi}} e^{-(-10/\sqrt{75})^2/2}.$$

Problem 3

We do 15000 independent trials with the probability 1/3 for a success. Compute the probability that the number of successes is between 4950 and 5050.

$$n = 15000, p = \frac{1}{3}, q = \frac{2}{3}, \mu = np = 5000, \sigma = \sqrt{npq} = \frac{100}{\sqrt{3}}$$

$$a = 4950, b = 5050, X_a = \frac{4950 - 5000}{100/\sqrt{3}}, X_b = \frac{5050 - 5000}{100/\sqrt{3}}$$

So, answer (using Laplace Theorem): $p(4950 \leq m \leq 5050) \approx \Phi(X_b) - \Phi(X_a)$.

Problem 4

A balanced die is thrown 24000 times one after the other. The face 5 appeared in m of these throws. Estimate the probability of the event $3900 \leq m \leq 4050$. Use Laplace as in Problem 3.

Problem 5

Find the smallest natural number x that satisfies the condition: If in k out of 10000 deliveries male babies were born then

$$P(5000 - x \leq k \leq 5000 + x) \geq 0.95.$$

$$n = 10000, p = q = \frac{1}{2}, \mu = np = 5000, \sigma = \sqrt{npq} = 50.$$

$$\text{By Laplace : } p(5000 - x \leq k \leq 5000 + x) \approx \Phi(X_{5000+x}) - \Phi(X_{5000-x}).$$

$$\text{We have : } X_{5000-x} = \frac{(5000 - x) - 5000}{50} = -\frac{x}{50}, X_{5000+x} = \frac{x}{50}.$$

$$\text{SO : } 2\Phi\left(\frac{x}{50}\right) - 1 = 0.95, \text{ hence : } x = 50 \times \Phi^{-1}\left(\frac{1.95}{2}\right)$$

Problem 6

Find the probability that in a group of 600 people there will be m birthdays on the New Years day.

Problem 7

In a certain typing process the probability for a typo is 0.001. Each line contains 50 characters. Compute the probability for two typos to occur

- (a) Using the exact distribution.
- (b) Using Poisson's approximation.

What is the relative deviation between the two results?

Problem 8

There is 1 percent of sick people in a population. Estimate the probability that among 200 of the people we will find at least 4 sick.

Problem 9

A certain population has 0.5 percents color blinds. We choose out of the population n people in a blind sampling (a person might be chosen more than once). Find the minimal value of n if the probability to find a color blind man in this sample is at least 0.95.