

# **Week 11**

## **Artificial Intelligence (AI) for Investments**

---



# Lesson 1: Regression Modelling

# Introduction


- Regression modelling: background and motivation
- Types of data
- Ordinary least square (OLS) estimation
- Simple linear regression
- Multiple linear regression

# Introduction

- Key CLRM assumptions
- Violation of CLRM assumptions
- BLUE properties of OLS estimators
- Hypothesis testing with regression modelling
- Other non-linear functional forms

# Background and Motivation

# Amazon, Netflix Movie Recommendations



# Filtering SPAM from Mail and Messages





# Big Data Text Analysis

## Headlines

[More Headlines](#)

[COVID-19 news](#): See the latest coverage of the coronavirus >

### India should play a constructive role in Afghanistan: Pak ambassador to China

Hindustan Times • 3 hours ago



- **EXCLUSIVE | Haqqani Network Scion Anas Haqqani Says Taliban Won't 'Interfere' in Kashmir, Clarifies Pakista**

News18 • 6 hours ago

- **Taliban says Afghanistan 'free nation' as it hails US exit**

Al Jazeera English • 20 hours ago

- **India must rethink 'wait and watch' Afghan policy**

The Indian Express • 7 hours ago • Opinion

- **Does China really have the upper hand in Afghanistan?**

Moneycontrol.com • 8 hours ago • Opinion

[View Full coverage](#)



### Pakistani terrorists coming to Afghanistan: Ghani informed Biden on July 23

Hindustan Times • 3 hours ago



- **US President Joe Biden says US committed to safe passage for last 100-200 Americans left in Afghanistan**

Times of India • 15 hours ago

[View Full coverage](#)



### Afghanistan Crisis LIVE: India Announces First Formal Meet With Taliban, Biden Defends US Exit

NDTV • 1 hour ago



- **First formal meeting between India-Taliban: Afghanistan should not be used to spread terror in India**

TIMES NOW • 22 hours ago

[View Full coverage](#)



### Supertech's twin towers case: Take strict action against officials involved in irregularities, says UP CM Yogi

The Indian Express • 3 hours ago



- **Supreme Court orders demolition of Supertech Twin Towers | Oneindia News**

Oneindia News • 21 hours ago

[View Full coverage](#)





# Summary


Making the Computers Learn Without Being Explicitly Programmed

- Amazon, Netflix movie recommendations
- Filtering out spams
- Medical prognosis with health records
- Algorithmic trading, credit scoring models
- Making computers think like humans
- Handwriting recognition, natural language processing, web-click data

# Machine Learning Algorithms


# Supervised Learning

House price prediction problem (Regression problem)



# Supervised Learning

Credit default scoring (classification problem)



# Unsupervised Learning

Clustering problem (clustering problem: market segmentation)



# Unsupervised Learning

Clustering problem (clustering problem : news aggregation)

## Headlines

[More Headlines](#)

[COVID-19 news](#): See the latest coverage of the coronavirus

### India should play a constructive role in Afghanistan: Pak ambassador to China

Hindustan Times • 3 hours ago

- EXCLUSIVE | Haqqani Network Scion Anas Haqqani Says Taliban Won't 'Interfere' in Kashmir, Clarifies Pakista

News18 • 6 hours ago

- Taliban says Afghanistan 'free nation' as it hails US exit

Al Jazeera English • 20 hours ago

- India must rethink 'wait and watch' Afghan policy

The Indian Express • 7 hours ago • Opinion

- Does China really have the upper hand in Afghanistan?

Moneycontrol.com • 8 hours ago • Opinion

[View Full coverage](#)



### Pakistani terrorists coming to Afghanistan: Ghani informed Biden on July 23

Hindustan Times • 3 hours ago

- US President Joe Biden says US committed to safe passage for last 100-200 Americans left in Afghanistan

Times of India • 15 hours ago

[View Full coverage](#)



### Afghanistan Crisis LIVE: India Announces First Formal Meet With Taliban, Biden Defends US Exit

NDTV • 1 hour ago

- First formal meeting between India-Taliban: Afghanistan should not be used to spread terror in India

TIMES NOW • 22 hours ago

[View Full coverage](#)




### Supertech's twin towers case: Take strict action against officials involved in irregularities, says UP CM Yogi

The Indian Express • 3 hours ago

- Supreme Court orders demolition of Supertech Twin Towers | Oneindia News

Oneindia News • 21 hours ago

[View Full coverage](#)



# Summary

- Supervised learning algorithm comprise data with features and labels
- The algorithm is trained map the relationship between the features and labels
- Then it makes predictions/create-label on the new unlabeled data based on its features
- The unsupervised learnings algorithms comprise unlabeled data that only carries features
- The data is clustered in groups based on these features



# Types of Data

# Types of Data


**Cross-sectional data:** Observations about multiple individuals (units) collected over a single period

**Time-series data:** Observations of a single individual (unit) collected over multiple periods

**Panel or longitudinal data:** Observations about multiple individuals (units) collected over various time periods


# Types of Data

If information about A is collected over times  $t_1$ ,  $t_2$ ,  $t_3$  then it is time-series data




# Types of Data

If information about A, B, C, D, E, and F is collected at  $t_1$ , then it is cross-sectional data




# Types of Data

If information about A, B, C, D, E, and F is collected at  $t_1$ ,  $t_2$ ,  $t_3$ , etc. then it is panel/longitudinal data



# Summary



# Introduction to Simple Linear Regression



# Introduction to Simple Linear Regression

Consider a simple linear regression model provided :

$$Y = \beta_0 + \beta_1 X + u$$

- This is also a two-variable linear regression model or bivariate linear regression model
- Here 'Y' is the dependent /explained /response /predicted/regressand variable
- Here 'X' is the independent/explanatory/predictor/regressor variable

# Introduction to Simple Linear Regression

Consider a simple linear regression model provided :

$$Y = \beta_0 + \beta_1 X + u$$

- 'u' is the error term, residual term or disturbance term that represents unobserved factors other than 'X' that affect 'Y'; since 'u' is also the random or stochastic variable it has a probabilistic distribution
- Here,  $\beta_0$  is the constant term and  $\beta_1$  is called the slope term (Why?)
- This simple model aims to study the dependence of Y on X

# Regression vs. Causation vs. Correlation

While regression deals with the dependence of one variable over another, it does not imply causation

- Regression only establishes the statistical strength of the relation, the causation is established by theory

## Example of crop and rain

- A priori theoretical considerations are needed to imply causation
- In regression analysis, dependent variable is considered random or stochastic (i.e., with probability distribution), while explanatory variable is assumed to have fixed values

# Regression vs. Causation vs. Correlation

A closely associated concept is correlation, which establishes the degree of linear relationship between the two variables

- In correlation analysis, both the variables are treated in a similar manner and considered to be random

# Summary



# Expectations Operator

# Expectations Operator 'E'

Any random probabilistic variable is often represented through expectations operator

- Any random variable attains multiple values. For example, a coin-toss can obtain two values with 50% odds for any outcome
- Similarly, in regression any random value is assumed to be probabilistic in nature and its expected value is represented by  $E(Y)$



# Expectations Operator 'E'

For example, if there are 'n' possibilities of an event,  $y_1, y_2, y_3, \dots, y_n$  each with possibilities  $p_1, p_2, p_3, p_4 \dots p_n$ , then expectations operator is defined as

- $E(y) = p_1 * y_1 + p_2 * y_2 + p_3 * y_3 + \dots + p_n * y_n$
- This is also called probability weighted mean
- If all the probabilities are assumed to be equal then  $p_1 = p_2 = \dots = p_n = \frac{1}{n}$
- Then  $E(y) = \frac{1}{n} (y_1 + y_2 + y_3 + \dots + y_n)$ , i.e., simple average of Y's

# Summary

- We discussed the role of expectations operator ( $E$ ) in the context of stochastic random variable with a probability distribution
- In simple terms, expectations are probability weighted averages of stochastic random variable
- In case there is no a priori probabilities assigned to these variables, then the expectation is simple average of the stochastic random variable

# A Simple Example

# A Simple Example

Consider a simple example of family income and consumption expenditure shown below

$Y \downarrow \quad X \rightarrow$	80	100	120	140	160	180	200	220	240	260
Weekly family consumption expenditure $Y$ ,	55	65	79	80	102	110	120	135	137	150
	60	70	84	93	107	115	136	137	145	152
	65	74	90	95	110	120	140	140	155	175
	70	80	94	103	116	130	144	152	165	178
	75	85	98	108	118	135	145	157	175	180
	–	88	–	113	125	140	–	160	189	185
	–	–	–	115	–	–	–	162	–	191
Total	325	462	445	707	678	750	685	1043	966	1211
Conditional means of $Y$ , $E(Y X)$	65	77	89	101	113	125	137	149	161	173

# A Simple Example

Here population of 60 families is divided into 10 income (X) groups from 80-260 (independent or fixed variable)

- The corresponding consumption expenditure values (Y) are also shown
- For each given level of income (X), the conditional means  $E(Y/X)$  that is mean of Y for a given level of X is also provided

# A Simple Example

For example, at  $X=80$ , the mean of  $Y$  is 65, i.e.,  $E(Y/X=80)=65$ ; these are called conditional expectations or conditional means of  $Y$  given the value of  $X$

- As they depend on the conditioning variable  $X$
- The average of all  $Y$ 's, that is unconditional mean or unconditional expected value  $E(Y)=121.2$

# A Simple Example

This unconditional mean does not account for the level of income ( $X$ ) and is the prediction of  $Y$  (expected value) when there is no knowledge of  $X$

- However, if one has the knowledge of  $X$ , then one can improve the prediction by computing conditional mean of  $Y$ , i.e.,  $E(Y/X)$ , which is a more accurate prediction of  $Y$




# A Simple Example

Que: What is the best (mean) prediction of weekly expenditure of families with a weekly income of  $X = 140$ :  $Y = 101$

- Thus the knowledge of the income level may enable us to better predict the mean value of consumption expenditure than if we do not have that knowledge
- This is the essence of regression modelling


# A Simple Example

Que: What is the best (mean) prediction of weekly expenditure of families with a weekly income of  $X = 140$ :  $Y = 101$



# A Simple Example

Que: What is the best (mean) prediction of weekly expenditure of families with a weekly income of  $X = 140$ :  $Y = 101$



# Population and Sample Regression Function

# Concept of Population Regression Function

If we join these conditional mean values, we obtain what is known as the population regression line (PRL)

- More simply, it is the regression of  $Y$  on  $X$
- Geometrically, then, a population regression curve is simply the locus of the conditional means of the dependent variable for the fixed values of the explanatory variable

# Concept of Population Regression Function

Population regression function (PRF)

- $E(Y/X_i) = f(X_i)$
- In this case,  $f(X_i)$  is a linear function of  $X$
- The expression is also called population regression function

# Concept of Population Regression Function

More generally, for a two variable case:  $E(Y/X_i) = \beta_0 + \beta_1 X_i$

- Here it is important to note that linearity means linearity in parameters
- $E(Y/X_i) = \beta_0 + \beta_1^2 X_i$ ; this model is non-linear in parameters and will not be handled in linear regression modelling
- $E(Y/X_i) = \beta_0 + \beta_1 X_i^2$ , in contrast this model is non-linear in variables and can be handled under linear regression models

# Sample Regression Function (SRF)


Sample regression function is shown by adding “^” hat symbol, indicating the estimated values:  $\hat{Y}_i = \hat{\beta}_0 + \hat{\beta}_1 X_i$

- $\hat{\beta}_0$  is the estimator of  $\beta_0$ ;  $\hat{\beta}_1$  is the estimator of  $\beta_1$ , and  $\hat{Y}_i$  is the estimator of  $y_i$
- SRF is only an estimate of PRF
- Thus, SRF can over or underestimate PRF values



# Sample Regression Function (SRF)

Sample regression function (SRF):  $\hat{Y}_i = \hat{\beta}_0 + \hat{\beta}_1 X_i$



# Ordinary Least Square (OLS) Estimation

# Method of Ordinary Least Square Estimation (OLS)

Recall the SRF function:  $Y_i = \widehat{\beta}_0 + \widehat{\beta}_1 X_i + \widehat{\mu}_i$ ; where  $\widehat{Y}_i = \widehat{\beta}_0 + \widehat{\beta}_1 X_i$

- Here,  $\widehat{\mu}_i = Y_i - \widehat{Y}_i = Y_i - \widehat{\beta}_0 - \widehat{\beta}_1 X_i$
- The line fit should aim to minimize the square this error  $\widehat{\mu}_i$
- Concept of OLS suggests that the best cost function to minimize is as follows


# Method of Ordinary Least Square Estimation (OLS)

Recall the SRF function:  $Y_i = \widehat{\beta}_0 + \widehat{\beta}_1 X_i + \widehat{\mu}_i$ ; where  $\widehat{Y}_i = \widehat{\beta}_0 + \widehat{\beta}_1 X_i$

- $\sum \widehat{\mu}_i^2 = \sum (Y_i - \widehat{Y}_i)^2 = \sum (Y_i - \widehat{\beta}_0 - \widehat{\beta}_1 X_i)^2$
- That is we minimize squared residuals (why not just residuals or absolute residuals)

# Method of Ordinary Least Square Estimation (OLS)

- $\sum \hat{\mu}_i^2 = \sum (Y_i - \hat{Y}_i)^2 = \sum (Y_i - \hat{\beta}_0 - \hat{\beta}_1 X_i)^2$ : Minimize these squared residuals



# Method of Ordinary Least Square Estimation (OLS)

$$\sum \widehat{\mu}_i^2 = \sum (Y_i - \widehat{Y}_i)^2 = \sum (Y_i - \widehat{\beta}_0 - \widehat{\beta}_1 X_i)^2$$

- Obvious to note here that  $\sum \widehat{\mu}_i^2 = f(\widehat{\beta}_0, \widehat{\beta}_1)$
- Setting differential of  $\sum \widehat{\mu}_i^2 = 0$  that satisfies and double differential to positive for minima condition, one obtains the estimates that is,  $\widehat{\beta}_0$  and  $\widehat{\beta}_1$

# Method of Ordinary Least Square Estimation (OLS)

$$\sum \widehat{\mu}_i^2 = \sum (Y_i - \widehat{Y}_i)^2 = \sum (Y_i - \widehat{\beta}_0 - \widehat{\beta}_1 X_i)^2$$

- Thus, these estimators are called least square estimators
- The regression model such estimated is also called the Gaussian, standard, or classical linear regression model (CLRM),

# Method of Ordinary Least Square Estimation (OLS)

$$\sum \widehat{\mu}_i^2 = \sum (Y_i - \widehat{Y}_i)^2 = \sum (Y_i - \widehat{\beta}_0 - \widehat{\beta}_1 X_i)^2$$

- $\frac{\partial(\sum \widehat{\mu}_i^2)}{\partial \widehat{\beta}_0} = -2 \sum (Y_i - \widehat{\beta}_0 + \widehat{\beta}_1 X_i) = -2 \sum \widehat{\mu}_i$  (partial differential w.r.t. to  $\widehat{\beta}_0$ )
- $\frac{\partial(\sum \widehat{\mu}_i^2)}{\partial \widehat{\beta}_1} = -2 \sum (Y_i - \widehat{\beta}_0 + \widehat{\beta}_1 X_i) X_i = -2 \sum \widehat{\mu}_i X_i$  (partial differential w.r.t. to  $\widehat{\beta}_1$ )




# Method of Ordinary Least Square Estimation (OLS)

$$\sum \widehat{\mu}_i^2 = \sum (Y_i - \widehat{Y}_i)^2 = \sum (Y_i - \widehat{\beta}_0 - \widehat{\beta}_1 X_i)^2$$

- $\widehat{\beta}_1 = \frac{\sum (X_i - \bar{X})(Y_i - \bar{Y})}{\sum (X_i - \bar{X})^2}$  and  $\widehat{\beta}_0 = \bar{Y} - \widehat{\beta}_1 \bar{X}$
- However, to achieve this closed form solution CLRM-OLS makes certain assumptions

# Summary

- $\sum \hat{\mu}_i^2 = \sum (Y_i - \hat{Y}_i)^2 = \sum (Y_i - \hat{\beta}_0 - \hat{\beta}_1 X_i)^2$ : Minimize these squared residuals



# Introduction to Multiple Linear Regression

# Introduction to Multiple Linear Regression

We can generalize the two variable problem into multiple linear regression as  $Y = \beta_0 + \beta_1 X_1 + \beta_2 X_2 \dots + \beta_n X_n + u$

- $X_i$ 's represent the explanatory variables
- Here the coefficients  $\beta_1, \beta_2, \dots, \beta_n$  are called the partial regression coefficients

# Introduction to Multiple Linear Regression

Multiple linear regression  $Y = \beta_0 + \beta_1 X_1 + \beta_2 X_2 \dots + \beta_n X_n + u$

- Other aspects of the regression remain the same, including the properties of the error term, that is,  $u$
- Zero conditional mean of error:  $E(u_i | X_{1i}, X_{2i}, \dots, X_{ni}) = 0$  for each 'i'
- No serial correlation:  $cov(u_i, u_j) = 0$ ; Homoscedasticity:  $var(u_i) = \sigma^2$

# Introduction to Multiple Linear Regression

Multiple linear regression  $Y = \beta_0 + \beta_1 X_1 + \beta_2 X_2 \dots + \beta_n X_n + u$

- Zero correlation (or covariance) between  $u_i$  and  $X$ :  
 $cov(u_i, X_1) = cov(u_i, X_2) = \dots = cov(u_i, X_n) = 0$
- The model is correctly specified

# Introduction to Multiple Linear Regression


Multiple linear regression  $Y = \beta_0 + \beta_1 X_1 + \beta_2 X_2 \dots + \beta_n X_n + u$

- Lastly, one more condition is added; that is no exact linear relationship between  $X_i$  and  $X_j$ s ( $X_1, X_2 \dots X_n$ ):  $\alpha_1 X_1 + \alpha_2 X_2 + \alpha_3 X_3 + \dots \alpha_n X_n \neq 0$
- If such a relationship exists, the model will be affected by the problem of perfect multicollinearity, and will not run (i.e., indeterminate)

# Introduction to Multiple Linear regression

However, there may be instances of less than perfect collinearity across variables and can affect the estimation

- If the multicollinearity is not perfect but high, then the estimators have large variances (standard errors of estimate)






# Introduction to Multiple Linear regression

However, there may be instances of less than perfect collinearity across variables and can affect the estimation

- This makes the 't-values' low and high chances of failure to reject the null-hypothesis (wider confidence intervals), even though the  $R^2$  may be high



# Summary

- We discussed multiple linear regression model
- All the properties and discussions on simple linear regression model apply to multiple linear regression model
- Some important properties of simple and multiple linear regression included: (a) zero conditional mean of the error term; (b) error term should not be serially correlated; (c) variance of the error term should be constant: Homoscedasticity; (d) no correlation between the error term and the independent variable; (e) model should be correctly specified; (f) multicollinearity should be low

# Interpreting the Multiple Linear Regression

# Interpreting the Multiple Linear Regression

Similar to the two variable regression, the following expression

$$E(Y|X_1 \dots X_n) = \beta_0 + \beta_1 X_1 + \beta_2 X_2 \dots + \beta_n X_n$$

- Represents the conditional mean or expected value of  $Y$  given the fixed values of all the  $X_i$ 's
- The partial coefficient  $\beta_1$  is the effect of  $X_1$  on  $Y$ , net of any effect from other explanatory variables ( $X_i$ 's), or in other words, keeping all the  $X_i$ 's constant

# Interpreting the Multiple Linear Regression

Similar to the two variable regression, the following expression

$$E(Y|X_1 \dots X_n) = \beta_0 + \beta_1 X_1 + \beta_2 X_2 \dots + \beta_n X_n$$

- The definition of  $R^2 = \frac{ESS}{TSS} = 1 - \frac{RSS}{TSS}$ , which is same as earlier

- One also calculates adjusted- $R^2 = 1 - \frac{\frac{RSS}{n-k}}{\frac{TSS}{n-1}} = 1 - \frac{(MSS \text{ of } RSS)}{(MSS \text{ of } TSS)}$  ;

remember the dfs?

# Interpreting the Multiple Linear Regression

$$\text{Adjusted-}R^2 = 1 - \frac{\frac{RSS}{n-k}}{\frac{TSS}{n-1}} = 1 - \frac{(MSS \text{ of } RSS)}{(MSS \text{ of } TSS)}$$

- Or  $\text{Adjusted-}R^2 = 1 - (1 - R^2) * (n-1)/(n-k)$
- Adjusted- $R^2$  penalizes addition of more variables. So if the  $R^2$  is inflated just by adding the number of variables, rather than their quality, then Adjusted- $R^2$  can identify the same

# Interpreting the Multiple Linear Regression

In the OLS estimation each parameter  $(\widehat{\beta}_0, \widehat{\beta}_1)$  is estimated with some error

- The square-root of the variance of the estimated parameter indicates that error in estimation or the precession of the estimate

# Summary

- The interpretation of multiple linear regression model broadly remains similar to the bivariate regression model
- The coefficients are partial coefficients that measure the impact of independent variable on dependent variable, keeping other variables constant
- The explanatory power of the model is measured using the  $R^2$  measure
- An improvement of over the  $R^2$  measure is adjusted-  $R^2$  measure which penalizes the addition of variables in the model
- Lower standard errors of the coefficients increases the power and efficiency of the mode
- OLS estimators are the best estimators in the class of linear estimators




# Key CLRM Assumptions

# Key CLRM Assumptions

The Gaussian, standard, or classical linear regression model (CLRM), makes 10 key assumptions

- Assumption 1: The regression model is linear in the parameters  $(\hat{\beta}_0, \hat{\beta}_1, \dots)$

Linear in Parameters




# Key CLRM Assumptions

**Assumption 2:** Values taken by the regressor  $X$  are considered fixed in repeated samples. More technically,  $X$  is assumed to be non-stochastic

# Key CLRM Assumptions



Assumption 3: Zero conditional mean of disturbance ( $u_i$ ): given the value of  $X$ , the mean, or expected, value of the random disturbance term  $u_i$  is zero.  $E(u_i/X_i)=0$



# Key CLRM Assumptions

**Assumption 4:** Homoscedasticity or equal variance of  $u_i$ . Given the value of  $X$ , the variance of  $u_i$  is the same for all observations. That is, the conditional variances of  $u_i$  are identical.  $var(u_i/x_i) = E[u_i - E(u_i|X_i)]^2 = constant = \sigma^2$

- Heteroscedastic variance =  $var(u_i/x_i) = \sigma_i^2$



# Key CLRM Assumptions


**Assumption 5:** No autocorrelation between the disturbances. Given any two  $X$  values,  $X_i$  and  $X_j$  ( $i \neq j$ ), the correlation between any two  $u_i$  and  $u_j$  ( $i \neq j$ ) is zero.

$$\begin{aligned}\text{Symbolically, } Cov(u_i, u_j | X_i, X_j) &= E \left[ [u_i - E(u_i) | X_i] [u_j - E(u_j) | X_j] \right]^2 \\ &= E[(u_i | X_i)(u_j | X_j)] = 0\end{aligned}$$

# Key CLRM Assumptions

Assumption 5: No autocorrelation between the disturbances

- (a) Positive autocorrelation
- (b) negative autocorrelation
- (c) No autocorrelation



# Key CLRM Assumptions

Assumption 6: Zero covariance between  $u_i$  and  $X_i$ , or  $E(u_i X_i) = 0$ .

- $Cov(u_i, X_i) = E[(u_i - E(u_i))(X_i - E(X_i))]$
- By definition:  $E(u_i)=0$ ;  $E(u_i X_i) - E(u_i)E(X_i)$
- $Cov(u_i, X_i) = E(u_i X_i) = 0$
- That is,  $u_i$  and  $X_i$  are not correlated



# Key CLRM Assumptions

- **Assumption 7:** The number of observations must be greater than the number of parameters to be estimated
- **Assumption 8:** The  $X$  values (independent variable) must have some finite variance
- **Assumption 9:** The regression model is correctly specified
- **Assumption 10:** There is no perfect multicollinearity, i.e., no perfect linear relationships among the explanatory variables




# Summary

- In this video we reviewed and summarized the ten (10) key CLRM assumptions

# BLUE Properties of OLS Estimators


# BLUE Properties of OLS Estimators

Biased and Unbiased estimators



# BLUE Properties of OLS Estimators

Efficient and inefficient estimators




# Summary

- OLS estimators are BEST in linear class of estimators
- They are best, linear, unbiased, and efficient estimators
- Thus, they are also consistent estimators: for large samples, OLS estimators converge to true population parameters

# Classical Normal Linear Regression Model (CNLRM) and Hypothesis testing I


# A Few Words on Normal Distribution





# A Few Words on Normal Distribution

## Standard Normal Distribution



# Classical Normal Linear Regression Model (CNLRM)

The estimation of sample parameters is not complete without hypothesis testing  $(\widehat{\beta}_0, \widehat{\beta}_1)$

- It is important to draw inferences about population parameters using sample estimates, more clearly, we would like the estimated parameters to be as close as possible to population parameters
- It must be noted that the randomness in the beta (coefficient) estimates is introduced by  $\mu_i$  (error term): **How?**
- Thus, these sample coefficient estimates also have a probability distribution [as one take different samples from population, one gets different estimates]

# The Normality Assumption of the Error Term $\mu_i$

To make any inference about the probability distribution of the estimate, we need to make some assumption about the distribution of the error term  $\mu_i$

- The CNLRM assumes that  $\mu_i$  is distributed normally with the following:
- *Mean* =  $E(u_i) = 0$
- *Variance* =  $E[u_i - E(u_i)]^2 = E(u_i)^2 = \sigma^2$
- *Covariance* =  $E[u_i - E(u_i)][u_j - E(u_j)] = E(u_i, u_j) = 0; i \neq j$
- These assumptions are summarised as  $u_i \sim N(0, \sigma^2)$

# Properties of OLS Estimators under Normality

Normal distributions are very easily defined with just two parameters, i.e., mean and variance of the population

- Under the normality assumption, OLS estimates are unbiased, efficient, consistent (estimates converge to their to population values as sample size increases)  $Y_i = \widehat{\beta}_0 + \widehat{\beta}_1 X_i + \widehat{\mu}_i$ ; where  $\widehat{Y}_i = \widehat{\beta}_0 + \widehat{\beta}_1 X_i$
- Mean:  $E(\widehat{\beta}_1) = \beta_1$ ; Variance =  $var(\widehat{\beta}_1) = \sigma_{\widehat{\beta}_1}^2$ ; then  $\widehat{\beta}_1 \sim N(\beta_1, \sigma_{\widehat{\beta}_1}^2)$

# Properties of OLS Estimators under Normality


Normal distributions are very easily defined with just two parameters, i.e., mean and variance of the population

- By the properties of standard normal distribution  $Z = \frac{\widehat{\beta}_1 - \beta_1}{\sigma_{\widehat{\beta}_1}}$
- Where  $\mathbf{Z} \sim \mathbf{N}(\mathbf{0}, \mathbf{1})$ :  $\mathbf{Z}$  is normally distributed with mean of 0, and SD=1

# Properties of OLS Estimators under Normality: Summary

Normal distributions are very easily defined with just two parameters, i.e., mean and variance of the population

Mean:  $E(\hat{\beta}_1) = \beta_1$ ; Variance =  $var(\hat{\beta}_1) = \sigma_{\hat{\beta}_1}^2$ ; then  $\hat{\beta}_1 \sim N(\beta_1, \sigma_{\hat{\beta}_1}^2)$



# Classical Normal Linear Regression Model (CNLRM) and Hypothesis testing II

# Interval Estimation and Hypothesis Testing

While in repeated sampling the point estimate  $\widehat{\beta}_1$  converges to true population parameter, i.e.,  $E(\widehat{\beta}_1) = \beta_1$ , but the accuracy of this point estimate is important: How reliable is this estimate

- This is so because the single estimate differs from true value; this reliability of the estimate is measured by its standard error




# Interval Estimation and Hypothesis Testing

In the OLS estimation each parameter  $(\widehat{\beta}_0, \widehat{\beta}_1)$  is estimated with some error

- The square-root of the variance of the estimated parameter indicates that error in estimation or the precession of the estimate

# Interval Estimation and Hypothesis Testing

In statistics we configure the confidence interval around the estimate



# Interval Estimation and Hypothesis Testing

For example, if you hypothesize that the population parameter =  $\beta_1$ ; then you set-up a confidence interval  $[1 - \alpha]$  around the estimate  $\hat{\beta}_1$

- If the estimate does not fall in this interval, then you can reject your hypothesis at 5% significance level
- Practically, you hypothesize that coefficient is zero. That is, the X variable does not have any impact on the Y variable. Then you set-up a confidence interval around that zero value

# Interval Estimation and Hypothesis Testing

Then that range can be written as  $(\pm t_{\alpha/2})$ ; if the estimated value falls outside this value, then with a given level of confidence ( $1 - \alpha$  generally 90%, 95%, or 99%) or significance level 10%, 5%, or 1% you reject the hypothesis and state that the variable has a significant relationship

# Interval Estimation and Hypothesis Testing

Null hypothesis:  $H_0$ : The true population parameter is  $\beta_1$  (= '0' in most cases)

- Alternate hypothesis  $H_1$ : The true population parameter is not  $\beta_1$
- Decision rule: Construct  $[1 - \alpha]$  confidence interval for the population parameter  $\beta_1$ ; if the estimate falls outside this value, you reject the null  $H_0$  (don't say you accept the null hypothesis). If the estimated parameter falls inside this range, you can not reject the null


# Interval Estimation and Hypothesis Testing

So if you hypothesized that  $\beta_1=0$  (i.e., no impact of X on Y) and the estimate falls in the confidence interval [this is checked by looking at the t-value of the estimate] then you say that you fail to reject the null and there is no relationship between X and Y

- What if it falls outside

# Interval Estimation and Hypothesis Testing

Interpretation:  $(1-\alpha)\%$  of the times the true parameter will fall within the confidence interval



# Other Functional Forms and Non-linear Transformations



# Other Functional Forms and Non-linear Transformations

- Log-linear or log-log model:  $Y_i = \beta_1 X_i^{\beta_2} * e^{u_i}$ ; take natural log and transform the model as below
- $\ln(Y_i) = \ln(\beta_1) + \beta_2 \ln(X_i) + u_i$  or alternatively
- $Y_i' = \alpha + \beta_2 X_i' + u_i$ : The model is now linear in parameters  $\alpha$  and  $\beta_2$
- The interpretation goes as follows:  $\beta_2$  measures percentage change in  $Y_i$  for a given percentage change in  $X_i$

# Other Functional Forms and Non-linear Transformations

- Log-lin model:  $Y_t = Y_0(1 + r)^t$  ; take natural log and transform the model as below
- $\ln(Y_t) = \beta_1 + t\beta_2$
- This is a semi-log model, and  $\beta_2$  measures proportional change in  $Y_t$  for a given absolute change in  $t$
- Vice-versa interpretation goes for Lin-log model below (absolute change in  $Y_t$  for a % or relative change in  $X_t$ ).
- $Y_t = \beta_1 + \ln(X_t) \beta_2$

# Summary and Concluding Remarks

# Summary and Concluding Remarks

- Among supervised learning algorithms, regression algorithm is a very important tool employed in the finance domain for applications such as forecasting security prices or credit scoring
- Regression algorithms can be run with only two variables (one independent and one dependent) : simple regression or with more than two variables: Multiple regression
- They key variables in a regression include a dependent variable, one or more independent variables, coefficients of these variables, and an error term

# Summary and Concluding Remarks

- The error term accounts for the variation in the dependent variable that can not be explained by the model (independent variables)
- While regression analysis can provide the statistical significance of the relationship, the direction of causality should come a priori from the theoretical underpinnings (rain vs. crop example)
- OLS is the most often employed method to estimate a regression model, which involves minimizing residual sum of squares

# Summary and Concluding Remarks

- OLS estimation of regression involves 10 key assumptions
- The most important assumptions include linearity in parameters, exogeneity of independent variables, zero conditional mean of the error (residual) term, homoscedasticity of error variances, absence of multicollinearity, no autocorrelation across error terms, no correlation between error and dependent variables
- If these assumptions are held then OLS estimators are referred to as BLUE, that is best linear unbiased and efficient estimates

# Summary and Concluding Remarks

- The statistical significance of OLS estimators is determined through hypothesis testing of coefficients individually
- This requires normality assumption of the error (residuals)
- Very often the model is not linear and may require some kind of transformation to make it linear, which can be subsequently estimated through OLS
- However, the interpretation of coefficients also change with such transformations



# Thanks!