

# Using the 2-MASS Photometric Redshift Survey to optimize LIGO Follow-Up Observations

Elisa Antolini<sup>1</sup>, Jeremy S. Heyl<sup>1\*2</sup>

<sup>1</sup>*Dipartimento di Fisica e Geologia, Università degli Studi di Perugia, I-06123 Perugia, Italia*

<sup>2</sup>*Department of Physics and Astronomy, University of British Columbia, 6224 Agricultural Road, Vancouver, BC V6T 1Z1, Canada*

Accepted —. Received —; in original form —

## ABSTRACT

The initial source localizations from LIGO-Virgo will cover large areas of sky. Because astrophysical sources of gravitational radiation are likely to reside in galaxies, it would make sense to search first in regions where the LIGO-Virgo probability is large and where the density of galaxies is large as well. Under the Bayesian prior assumption that the probability of a gravitational-wave event from a given region of space is proportional to the density of galaxies within the probed volume, one can calculate an improved localization of the position of the source simply by multiplying the LIGO-Virgo skymap by the density of galaxies in the range of redshifts. We propose using the 2-MASS Photometric Redshift Galaxy Catalogue for this purpose and demonstrate that using it can dramatically reduce the search region for electromagnetic counterparts.

## 1 INTRODUCTION

In the near future LIGO and Virgo should begin to detect gravitational wave events from the local Universe. During these initial years of gravitational astronomy, the localization of the candidate events on the sky will be poor with the ninety-percent confidence regions covering hundreds or even thousands of square degrees. Finding an electromagnetic counterpart to these candidate gravitational-wave events will be crucial to understanding what produces them, interpretation of the signal and to provide tests of general relativity. The ideas of how the electromagnetic counterparts would appear are varied and uncertain. Over the large search regions and over the span of days or weeks, many electromagnetic transients typically occur, and with the wide variety of models it will be difficult to associate unambiguously a particular electromagnetic event with a candidate gravitational-wave event.

The purpose of this letter is to present a strategy to alleviate both of these issues; that is, to reduce both the search region and the time required to plan and begin observations. We follow the spirit of Gehrels et al. (2015) to develop a galaxy catalogue to guide the observational plan. However, our goal here is to develop a nearly complete catalogue at the expense of having less accurate estimates of the redshifts of the galaxies within the catalogue. The accuracy of the galaxy distances needs to be only as good as the distance estimates of the gravitational-wave events. Additionally we will outline a straightforward and rapid technique to generate a nearly optimal observing plan to follow up the events rapidly (i.e. within a few seconds of the trigger).

## 2 GALAXY CATALOGUES

## 3 BAYESIAN APPROACH TO FOLLOW-UP

Because we will be interested in the rapid follow-up of candidate gravitational-wave events, we will focussed on the rapid Bayesian reconstruction outlined by Singer & Price (2015), BAYESTAR. At the most basic level, BAYESTAR yields a probability map on the sky in the form of a HEALPix map where each pixel contains the probability  $P(d|m)$  that a particular model (i.e. position on the sky) will yield the data (i.e. the observed strains on the LIGO and Virgo interferometers). To plan an observing strategy one would like the probability of a particular model (i.e. position on the sky) given the data. We have from Bayes's theorem

$$P(\text{position}|\text{data}) = \frac{P(\text{position})P(\text{data}|\text{position})}{P(\text{data})}. \quad (1)$$

If we make the additional mild assumption that gravitational-waves originate from nearby galaxies, the probability of a given position on the sky naturally is proportional to the surface density of galaxies in that direction within the distance range determined from the modelling of the gravitational waveform. Of course, these distance estimates will usually have large uncertainties so the distance range over which to integrate the galaxy density distribution will also be large, so highly accurate redshift information is not needed to construct  $P(\text{position})$ .


Furthermore, because we will ultimately be interested in which fields to observe (not which particular galaxies), accurate positions are not required in the construction of  $P(\text{position})$ . It is natural to sample  $P(\text{position})$  also as a HEALPix grid with each pixel covering about the same solid angle as the field of view of the telescope of interest or the BAYESTAR map (a HEALPix level of 512 or about 50

square arcminutes per pixel), so positions no more accurate than arcminutes are required. The key to generate the observing plan rapidly is to calculate the required galaxy density maps beforehand in principle at the desired resolution (this optimization only speeds the process up slightly) for the distance ranges of interest. With the arrival of an alert, all that is required is to calculate Eq. (1) using the HEALPix maps, resample to the scale of the telescope, renormalize the probability, sort the pixels from most likely to least and output the positions to cover a given amount of cumulative probability.

This work was supported by the Natural Sciences and Engineering Research Council of Canada, the Canadian Foundation for Innovation, the British Columbia Knowledge Development Fund and the Bertha and Louis Weinstein Research Fund at the University of British Columbia.

## REFERENCES

- Gehrels, N., Cannizzo, J. K., Kanner, J., Kasliwal, M. M., Nissanke, S., & Singer, L. P. 2015, ArXiv e-prints  
 Singer, L. P. & Price, L. R. 2015, ArXiv e-prints



**Figure 1.** The number of fields required to cover the given fraction of the probability region for a simulated LIGO detection (red curves without the galaxy map, green curves with a smoothed galaxy, blue curves with a raw galaxy map). The upper solid curves use a healpix map with about 200,000 cells, the dashed curves have about 50,000 cells, the dotted curves have about 12,000 cells and lower solid curves have about 3,000 cells, corresponding 0.2, 0.8, 3.2 and 13 square-degree fields of view. The redshift range of the galaxy map in the upper panel is  $0.03 < z < 0.04$  and  $0.01 < z < 0.05$  in the lower panel.