Array Efficiency Analysis for 2024 / 2025

Author: Marwan Negm

Date: 2026-1-06

Relevant Links:

- Monday Post
- DR0
- The following code is mostly taken from 2024 efficiency analysis, but streamlined and generalised to have fewer plots

Imports

In [1]:
```
import scripts.string_power_functions as sf
import matplotlib.dates as mdates
from datetime import datetime, timedelta
import pytz
import matplotlib.pyplot as plt
import numpy as np
import pvlib
from scipy.signal import butter, filtfilt
```

Data Acquisition

In [2]:
```
## Parameters: ##
# Location Information
latitude = 37.002923
longitude = -86.3686424

# Surface Area
total_surface_area = 4

# Dates: Somewhat important that you format it "year-month-day"
# 2024:
start_date_str = "2024-07-16"
end_date_str = "2024-07-18"

# 2025:
# start_date_str = "2025-07-01"
# end_date_str = "2025-07-03"

start_date = list(map(int, start_date_str.split("-"))) # List of integers representir
end_date = list(map(int, end_date_str.split("-")))

# Times
```

```
year = start_date[0]

month = start_date[1]

day1 = start_date[2]
day2 = start_date[2] + 1
day3 = start_date[2] + 2

# Finding location for pvLib
location = pvlib.location.Location(latitude, longitude, tz='America/Chicago', altituc

# Find the start and end times and localise it to UTC
start_time: datetime = pytz.timezone('UTC').localize(datetime(year, month, start_date
end_time: datetime   = pytz.timezone('UTC').localize(datetime(year, month, end_date[2

print(f"Start time is {start_time}; End time is {end_time}")

# For 2025 : ["A", "B", "C"], "OutputCurrent", "OutputVoltage"
# For 2024 : ["1", "2"], "CurrentSensor", "VoltSensor"

total_power_series = sf.get_total_power(start_time, end_time, ["1", "2"], "CurrentSer
```

```
Start time is 2024-07-16 12:00:00+00:00; End time is 2024-07-19 06:00:00+00:00
CurrentSensor1
VoltSensor1
CurrentSensor2
VoltSensor2
```

Analysis

This block shifts the data by 7 hours because InlfuxDB is in the wrong timezone:

In [3]:
```
shifted_power_series = total_power_series.copy()

shifted_power_series._start = total_power_series._start + timedelta(hours=(7)) #Time
shifted_power_series._stop = total_power_series._stop + timedelta(hours=(7))
```

Generate the new irradiance series the existing time series:

In [4]:
```
date_time_x_axis = shifted_power_series.datetime_x_axis

new_irradiance_series = location.get_clearsky(date_time_x_axis)
```

Preform a low pass filter to reduce noise on the data:

This is where my understanding fails slightly, parameters were selected arbitrarily until enough noise was cut off.

In [5]:
```
N = len(shifted_power_series)  # Number of data points
sampling_rate = 1/shifted_power_series._period  # Sampling rate (in Hz)
```

```
# Perform FFT
fft_result = np.fft.fft(shifted_power_series)
# Frequency axis (for single-sided spectrum)
frequencies = np.fft.fftfreq(N, d=shifted_power_series._period)  # Positive frequenci
fft_magnitude = np.abs(fft_result)  # Magnitude of the FFT (single-sided)

# Filter parameters #
cutoff_freq = 1  # Cutoff frequency in Hz
fs = 20000  # Sampling rate in Hz
order = 4  # Filter order

# Design the low-pass filter
nyq = 0.5 * fs
normal_cutoff = cutoff_freq / nyq
b, a = butter(order, normal_cutoff, btype='lowpass')

filtered_power_series = filtfilt(b, a, shifted_power_series)

# (OPTIONAL) Plot the FFT
# plt.plot(frequencies, fft_magnitude)
# plt.xlabel('Frequency (Hz)')
# plt.ylabel('Magnitude')
# plt.title('FFT of Signal')
# plt.yscale('log')
# plt.grid(True)
# plt.show()
```

Graph the power series

This is done to see how the data looks before and after the main processing step (low pass filter)

In [6]:

```
# Graph of the power series
plt.plot(date_time_x_axis, shifted_power_series/total_surface_area, label = 'Array Po
plt.plot(date_time_x_axis, filtered_power_series/total_surface_area, label = 'Array P
plt.plot(date_time_x_axis, new_irradiance_series['ghi'], label = 'Solar Irradiance')

plt.xticks(rotation = 45, ha='right')
plt.title(f"Array Power and Solar Irradiance in FSGP {year}")

plt.grid()

plt.legend(loc = 'upper right')

plt.ylabel("Solar Irradiance (W/m²)")
plt.xlabel("Time")

plt.show()

# Graph of residuals
plt.plot(date_time_x_axis, (filtered_power_series - shifted_power_series)/total_surfa
plt.xticks(rotation = 45, ha='right')
plt.ylabel("Residuals between lowpass and actual")
```

```
plt.xlabel("Time")
plt.show()
```





Mask the Data and Extract Peak Efficiency

Using numpy, the data was masked based on day and only when the irradiance was greater than 750 watts. This was done to make sure to only include times when there solar cells are getting enough power to be accurately measured. All code which excludes day 2 is commented out.

In [7]:
```
mask = new_irradiance_series['ghi'] > 750 # Only show data where the irradiance is gr

# Day masks, maintains the previous irradiance mask
mask_day1 = np.logical_and(np.logical_and((mask), date_time_x_axis > datetime(year, m
mask_day2 = np.logical_and(np.logical_and((mask), date_time_x_axis > datetime(year, m
mask_day3 = np.logical_and(np.logical_and((mask), date_time_x_axis > datetime(year, m

mask_without_day_2 = np.logical_or(mask_day1, mask_day3)

#Filter both the x and y data using the mask
x_filtered = date_time_x_axis[mask]
y_filtered = (filtered_power_series[mask] / total_surface_area) / new_irradiance_seri

y_filtered_without_day_2 = (filtered_power_series[mask_without_day_2] / total_surface
```

Find Peak and Average efficiencies

Finds the average and peak efficiencies, again all code which excludes day 2 is commented out.

In [11]:
```
# Find average and peak efficiencies
average_efficiency = np.mean(y_filtered)
peak_efficiency = np.max(y_filtered)

average_efficiency_without_day_2 =  np.mean(y_filtered_without_day_2)
print(f"Average efficiency without day 2 is {average_efficiency_without_day_2*100:.2f

print(f"Average efficiency is {average_efficiency*100:.2f}%")
print(f"Peak efficiency is {peak_efficiency*100:.2f}%")

## Plot the filtered data ##
plt.scatter(x_filtered, y_filtered, s=2)
plt.grid()

plt.ylabel("Efficiency (W/m²)")
plt.xlabel("Time")

plt.xticks(rotation = 45, ha='right')

plt.title(f"Efficiency vs. Solar Irradiance in FSGP {year}")

plt.show()

#Plotting efficiency by day. Used in 2024 to show how weird day 2 is
plt.scatter(new_irradiance_series['ghi'][mask_day1], (filtered_power_series[mask_day1
plt.scatter(new_irradiance_series['ghi'][mask_day2], (filtered_power_series[mask_day2
plt.scatter(new_irradiance_series['ghi'][mask_day3], (filtered_power_series[mask_day3

plt.axhline(average_efficiency, label='Average Efficiency', ls = ':', color = 'navajo
plt.axhline(average_efficiency_without_day_2, label='Average Efficiency (Day 2 excluc

plt.legend(loc = 'upper left')
plt.grid()
plt.ylim(0, 0.27)

plt.ylabel("Efficiency")
plt.xlabel("Solar Irradiance (W/m²)")

plt.title(f"Array Efficiency by Day in FSGP {year}")

plt.show()
```

```
Average efficiency without day 2 is 16.16%
Average efficiency is 13.77%
Peak efficiency is 23.01%
```


Efficiency vs. Solar Irradiance in FSGP 2024
Efficiency (W/m²)
0.20
0.15
0.10
0.05
07-16 18
07-17 00
07-17 06
07-17 12
07-17 18
07-18 00
07-18 06
07-18 12
07-18 18
Time


Array Efficiency by Day in FSGP 2024
Day 1 Efficiencies
Day 2 Efficiencies
Day 3 Efficiencies
Average Efficiency
Average Efficiency (Day 2 excluded)
Efficiency
0.25
0.20
0.15
0.10
0.05
0.00
750
775
800
825
850
875
900
925
Solar Irradiance (W/m²)

Plot to show the change in data from the Lowpass filter

In [12]:
```
plt.hist((shifted_power_series[mask]/total_surface_area)/new_irradiance_series['ghi']
plt.hist((filtered_power_series[mask] / total_surface_area) / new_irradiance_series['

plt.legend()

plt.ylabel("Frequency")
plt.xlabel("Efficiency")

plt.title(f"Effective Filtering on Frequency of Efficiency for FSGP {year}")

plt.show()
```



Plots by day

In [ ]:
```
def showDayPlot(mask, label: str):
    # Capture the mappable object returned by plt.scatter
    scatter = plt.scatter(new_irradiance_series['ghi'][mask], (filtered_power_series[
    plt.title(f'Power vs. Irradiance for {label}')
    #plt.scatter(new_irradiance_series['ghi'][mask], new_irradiance_series['ghi'][mas

    # Create the color bar and store the object
    cbar = plt.colorbar()

    # Apply date formatting to the color bar's y-axis #
```

```
    # Get the AutoDateLocator to find appropriate tick locations
    locator = mdates.AutoDateLocator()
    cbar.ax.yaxis.set_major_locator(locator)

    # Apply a DateFormatter
    cbar.ax.yaxis.set_major_formatter(mdates.ConciseDateFormatter(locator))

    plt.xlim(0,1000)
    plt.ylim(0,270)

    plt.grid()

    plt.ylabel("Solar Power (W/m2)")
    plt.xlabel("Solar Irradiance (W/m2)")

    plt.show()

showDayPlot(mask, 'All Days')
showDayPlot(mask_day1, 'Day 1')
showDayPlot(mask_day2, 'Day 2')
showDayPlot(mask_day3, 'Day 3')
```

```
C:\Users\XPS\AppData\Local\Temp\ipykernel_27028\3007184895.py:22: UserWarning: No arti
sts with labels found to put in legend.  Note that artists whose label start with an u
nderscore are ignored when legend() is called with no argument.
  plt.legend(loc = 'upper left')
```




Power vs. Irradiance for Day 1
2024-Jul-16
Solar Power (W/m2)
Solar Irradiance (W/m2)
0
50
100
150
200
250
0
200
400
600
800
1000
20:15
20:00
19:45
19:30
19:15
19:00
18:45
18:30
18:15


Power vs. Irradiance for Day 2
2024-Jul-17
Solar Power (W/m2)
Solar Irradiance (W/m2)
0
50
100
150
200
250
0
200
400
600
800
1000
20:15
20:00
19:45
19:30
19:15
19:00
18:45
18:30
18:15


Power vs. Irradiance for Day 3
2024-Jul-18
Solar Power (W/m2)
Solar Irradiance (W/m2)
0
50
100
150
200
250
0
200
400
600
800
1000
20:15
20:00
19:45
19:30
19:15
19:00
18:45
18:30
18:15