# Unified Biquaternion Theory: A Geometric Unification of General Relativity, Quantum Field Theory, and the Standard Model

David Jaroš

November 1, 2026

**AI provenance — Tier B.** AI-assisted; machine-verified against named sources or verifiers; individual attestation is not claimed. Details: `AI_PROVENANCE.md`.

**License:** © 2025 Ing. David Jaroš. This work is licensed under a Creative Commons Attribution-NonCommercial-NoDerivatives 4.0 International License (CC BY-NC-ND 4.0). Earlier drafts (up to v0.3) were released under CC BY 4.0. From v0.4 onward, all material is released under CC BY-NC-ND 4.0 to protect the integrity of the theoretical work during ongoing academic development. See `https://creativecommons.org/licenses/by-nc-nd/4.0/` for details.

**Abstract**

The Unified Biquaternion Theory (UBT) is a research framework based on a single biquaternionic field $\Theta(q,\tau)$ over complex time $\tau = t + i\psi$. In the revised local geometric sector the four covariant gradients $E_\mu = \mathcal{N}_0^{-1/2} D_\mu \Theta$ form a biquaternionic tetrad. On the real Lorentz slice their central anticommutator defines the metric directly, without a trace, real-part projection, preferred $\psi$ section, or compact-fiber average. This paper records the canonical kinematics and the still-open dynamical bridge: the metric-compatible frame connection is reconstructed from the tetrad and specified torsion, while the UBT action must select the torsion, the exact left/right representation on $\Theta$, Lorentz-slice preservation, curved integrability, and Einstein dynamics. Claims of complete GR or Standard-Model closure are therefore conditional research goals, not axioms.

**Lock-in statement:** The canonical classical metric is the coefficient of the central identity $\{E_\mu, E_\nu\}_\sharp/2 = g_{\mu\nu}\mathbf{1}$. No projection operation defines the metric. The antisymmetric algebraic product and the commutator of covariant derivatives remain distinct carriers of bivector/spin and curvature information, respectively.

**Future-proofing rule:** Any future extension of UBT must preserve the central covariant-tetrad identity, distinguish the frame connection from the coordinate connection, and state which results are kinematic, dynamical, conditional, or speculative. No trace, real-part map, phase projection, or compact-fiber average may be reintroduced as the canonical local metric.

# Contents


**1 Introduction** **7**
- 1.1 Historical Context . . . 7
- 1.2 Structure of This Article . . . 7

**2 Biquaternion Algebra** **8**

**3 The Theta Field $\Theta(q, \tau)$** **8**

**4 The Fundamental Biquaternion Field $\Theta(q, \tau)$** **8**
- 4.1 Definition . . . 8
  - 4.1.1 Explicit Biquaternion Form . . . 9
- 4.2 Faithful Matrix Representation and Degree Count . . . 9
- 4.3 Hermitian Involution . . . 9
- 4.4 Physical Interpretation . . . 9
- 4.5 Field Equations and Covariant Derivative Status . . . 10
  - 4.5.1 Classical connection versus its action on $\Theta$ . . . 10
- 4.6 Normalization . . . 10
- 4.7 Gauge Transformations . . . 11
- 4.8 Reality Conditions . . . 11
- 4.9 Asymptotic Behavior . . . 11
- 4.10 Connection to Biquaternions . . . 11
- 4.11 Units and Dimensions . . . 11
- 4.12 Historical Note . . . 12

**5 Complex Time $\tau = t + i\psi$** **12**

**6 Biquaternion Time $T_B$** **12**
- 6.1 Canonical Definition . . . 12
- 6.2 Physical Interpretation . . . 12
  - 6.2.1 Real Component: Standard Time . . . 12
  - 6.2.2 Scalar Imaginary Component: Isotropic Phase . . . 13
  - 6.2.3 Vector Imaginary Components: Directional Phases . . . 13
- 6.3 Dynamics of Imaginary Components . . . 13
- 6.4 Hierarchical Reduction Structure . . . 14
  - 6.4.1 Complex Time Limit . . . 14
  - 6.4.2 Classical Time Limit . . . 14
  - 6.4.3 Hierarchy Summary . . . 14
- 6.5 Measurement and Observability . . . 14
  - 6.5.1 Direct Observability . . . 14
  - 6.5.2 Indirect Detection . . . 15
- 6.6 Mathematical Properties . . . 15
  - 6.6.1 Biquaternion Conjugation . . . 15
  - 6.6.2 Norm . . . 15
  - 6.6.3 Scalar and Vector Parts . . . 15
- 6.7 Derivatives and Calculus . . . 15
  - 6.7.1 Partial Derivatives . . . 15
  - 6.7.2 Integration . . . 15

6.8 Relation to Other Formulations . . . 16
6.8.1 Euclidean Time (Wick Rotation) . . . 16
6.8.2 Thermal Field Theory . . . 16
6.9 Conflict Resolution . . . 16
6.10 Coordinate Systems . . . 16
6.10.1 Biquaternion Coordinates . . . 16
6.10.2 Metric Signature . . . 17
6.11 Conservation Laws . . . 17
6.12 Causality Structure . . . 17
6.13 Symbol Standardization . . . 17
6.14 Units . . . 18
6.15 Historical Note . . . 18
6.16 Simplification: Complex Time Limit . . . 18

**7 Fundamental Biquaternionic Geometry 18**
7.1 Overview . . . 18

**8 The covariant biquaternionic tetrad $E_\mu$ 19**
8.1 Classical Lorentz sector . . . 19
8.2 Metric and bivector from one product . . . 19
8.3 Transformation and rank . . . 20
8.4 Status and open bridge . . . 20

**9 The central anticommutator metric 20**

**10 Clifford and generalized-Dirac lift of the canonical relation 21**
10.1 Scope: no second tetrad . . . 21
10.2 Exact block lift . . . 21
10.3 Spinor carrier without a new fundamental field . . . 22
10.4 Rank is inherited, not manufactured . . . 22
10.5 Complex-time / fifth operator channel . . . 23
10.6 Principal symbol, causal cone and exact factorisation . . . 23
10.7 Conditional first-jet rank preservation by the fifth channel . . . 24
10.8 No-extra-field constrained-rank theorem . . . 24
10.9 Application to the holomorphic generalized-Dirac equation . . . 26
10.10 Remaining proof gates . . . 26

**11 The biquaternionic frame connection $\Omega_\mu$ 27**
11.1 Classical Lorentz connection reconstructed from tetrad and torsion . . . 27
11.2 Relation to Christoffel symbols . . . 28
11.3 Gauge covariance and Lorentz-slice preservation . . . 28
11.4 Representation selection on $\Theta$ . . . 28
11.5 Pure Lorentz pairing: exact reduction and exact obstruction . . . 28
11.6 Flat limit and explicit representer . . . 29
11.7 Full UBT status . . . 30

**12 Biquaternionic curvature 30**

**13 Classical Geometry (Derived from Biquaternionic Structure) 31**
13.1 Classical Metric Tensor . . . 31

**14 The central anticommutator metric 31**
14.1 Classical Curvature Tensors . . . 32

**15 Curvature from the connection commutator 32**
15.1 Classical Stress-Energy Tensor . . . 32

**16 Field Equations 33**
16.1 Canonical master equation and status . . . 33
16.2 Alternative Form with Complex Time . . . 33
16.3 Cosmological Constant . . . 34
16.4 Classical GR Solutions . . . 34

**17 Quantum Electrodynamics (QED) 34**

**18 Quantum Electrodynamics (QED) Lagrangian 34**
18.1 Canonical Definition . . . 34
18.2 Components . . . 35
18.2.1 Matter Field Term . . . 35
18.2.2 Photon Field Term . . . 35
18.3 Gauge Symmetry . . . 35
18.3.1 U(1) Gauge Transformation . . . 35
18.3.2 Gauge Fixing . . . 35
18.4 Equations of Motion . . . 36
18.4.1 Maxwell Equations . . . 36
18.4.2 Bianchi Identity . . . 36
18.4.3 Theta Field Equation . . . 36
18.5 Interaction Term . . . 36
18.6 Current Conservation . . . 36
18.7 Coupling to Complex Time . . . 37
18.8 Fine Structure Constant . . . 37
18.9 Renormalization . . . 37
18.9.1 Running Coupling . . . 37
18.9.2 Charge Renormalization . . . 37
18.10 Energy-Momentum Tensor . . . 37
18.11 Curved Spacetime Extension . . . 38
18.12 Conflict Resolution . . . 38
18.13 Computational Implementation . . . 38
18.14 Physical Predictions . . . 39
18.15 Extensions . . . 39
18.15.1 QED + Weak Interactions . . . 39
18.15.2 QED + Gravity . . . 39
18.16 Historical Note . . . 39

**19 Quantum Chromodynamics (QCD) 40**

**20 Quantum Chromodynamics (QCD) Lagrangian** **40**
20.1 Canonical Definition . . . 40
20.2 Gluon Field Strength . . . 40
20.3 SU(3) Color Symmetry . . . 41
20.3.1 Gauge Group . . . 41
20.3.2 Generators . . . 41
20.3.3 Gell-Mann Matrices . . . 41
20.4 Color Indices . . . 41
20.5 Structure Constants . . . 41
20.6 Gauge Transformations . . . 42
20.6.1 Local SU(3) Transformation . . . 42
20.7 Equations of Motion . . . 42
20.7.1 Yang-Mills Equations . . . 42
20.7.2 Bianchi Identity . . . 42
20.8 Asymptotic Freedom . . . 43
20.8.1 Running Coupling . . . 43
20.8.2 QCD Scale $\Lambda_{\text{QCD}}$ . . . 43
20.9 Confinement . . . 43
20.10 Quark Masses . . . 43
20.11 Gluon Condensate . . . 44
20.12 Chiral Symmetry Breaking . . . 44
20.13 Energy-Momentum Tensor . . . 44
20.14 Curved Spacetime Extension . . . 44
20.15 Emergent SU(3) in UBT . . . 44
20.15.1 Mechanism . . . 45
20.15.2 Derivation . . . 45
20.16 Conflict Resolution . . . 45
20.17 Experimental Predictions . . . 45
20.18 Lattice QCD Comparison . . . 46
20.19 Historical Note . . . 46

**21 Standard Model Gauge Structure** **46**

**22 Standard Model Gauge Structure** **46**
22.1 Canonical Definition . . . 46
22.2 Gauge Group . . . 46
22.3 Gauge Fields and Couplings . . . 47
22.4 Field Strength Tensors . . . 47
22.4.1 SU(3) Gluon Field Strength . . . 47
22.4.2 SU(2) Weak Field Strength . . . 47
22.4.3 U(1) Hypercharge Field Strength . . . 47
22.5 Generators . . . 47
22.5.1 SU(3) Generators . . . 47
22.5.2 SU(2) Generators . . . 47
22.5.3 U(1) Generator . . . 48
22.6 Electroweak Unification . . . 48
22.6.1 Electroweak Lagrangian . . . 48
22.6.2 Physical Gauge Bosons . . . 48

22.6.3 Weak Mixing Angle . . . 48
22.7 Coupling Constants . . . 48
22.7.1 Gauge Couplings . . . 48
22.7.2 Electromagnetic Coupling . . . 49
22.8 Fermion Representations . . . 49
22.8.1 Quark Doublets (Left-Handed) . . . 49
22.8.2 Quark Singlets (Right-Handed) . . . 49
22.8.3 Lepton Doublets (Left-Handed) . . . 49
22.8.4 Lepton Singlets (Right-Handed) . . . 49
22.9 Higgs Mechanism . . . 49
22.9.1 Higgs Doublet . . . 49
22.9.2 Higgs Potential . . . 49
22.9.3 Vacuum Expectation Value . . . 50
22.9.4 Gauge Boson Masses . . . 50
22.10 Yukawa Couplings . . . 50
22.11 CKM Matrix . . . 50
22.12 PMNS Matrix . . . 50
22.13 Emergence in UBT . . . 50
22.13.1 Gauge Group from Theta Field (Candidate Construction) . . . 50
22.13.2 Predictions . . . 51
22.14 Grand Unification . . . 51
22.14.1 GUT Scale . . . 51
22.14.2 UBT and GUT . . . 51
22.15 Conflict Resolution . . . 51
22.16 Experimental Status . . . 52
22.17 Historical Note . . . 52
22.18 Geometric Derivation of $SU(2)_L \times U(1)_Y$ . . . 52
22.19 Higgs Mechanism via Hosotani Wilson Line . . . 52

**23 Theta-Functions and Toroidal Structure** **53**
23.1 Jacobi Theta Functions . . . 53
23.2 Modular Transformations . . . 53
23.3 Fine Structure Constant . . . 53

**24 Speculative Extensions** **53**

**25 Experimental Designs and Testable Predictions** **53**
25.1 Testable Predictions . . . 53
25.2 Experimental Status . . . 54

**26 Conclusions** **54**

**A Structure of the Covariant Derivative $\nabla$** **54**

**B The covariant derivative in the active UBT geometry** **54**
B.1 Three distinct connections . . . 55
B.2 Why the generic curved derivative is two-sided . . . 55
B.3 Special relativity and the implicit curved system . . . 55

**C Symbol Dictionary** **56**

**D Symbol Dictionary and Notation Conventions** **56**
D.1 Purpose . . . 56
D.2 Reserved Symbols — Single Meaning Only . . . 56
D.3 Index Conventions . . . 57
D.3.1 Spacetime Indices . . . 57
D.3.2 Spatial Indices . . . 57
D.3.3 Gauge Indices . . . 57
D.3.4 Matrix Indices . . . 57
D.4 Field and Operator Notation . . . 57
D.5 Derivative Notation . . . 58
D.6 Coupling Constants . . . 58
D.7 Mass Scales . . . 58
D.8 Forbidden Symbol Uses . . . 59
D.9 Metric Signature Convention . . . 59
D.10 Unit Conventions . . . 59
D.10.1 Natural Units . . . 59
D.10.2 SI Units . . . 59
D.11 Special Function Notation . . . 60
D.12 Tensor and Matrix Operations . . . 60
D.13 Abbreviations and Acronyms . . . 60
D.14 Version Control . . . 61
D.15 Enforcement . . . 61

**E Mathematical Derivations** **61**

**F Computational Methods** **61**

# 1 Introduction

The unification of General Relativity (GR) and Quantum Field Theory (QFT) remains one of the most significant open problems in theoretical physics. While GR successfully describes gravity as the curvature of spacetime and QFT provides an accurate description of quantum phenomena and the Standard Model (SM) of particle physics, these frameworks are fundamentally incompatible at high energies and small distances.

The Unified Biquaternion Theory (UBT) proposes a radical solution: both spacetime geometry and quantum fields emerge from a more fundamental structure—a biquaternion field $\Theta(q, \tau)$ defined over complex time $\tau = t + i\psi$.

## 1.1 Historical Context

## 1.2 Structure of This Article

This article is organized into 12 sections following the canonical structure:

1. Introduction (this section)
2. Biquaternion algebra

3. The Theta field $\Theta(q, \tau)$
4. Complex time $\tau = t + i\psi$
5. Geometry and metric tensor
6. Stress-energy tensor
7. Einstein field equation
8. Quantum Electrodynamics (QED)
9. Quantum Chromodynamics (QCD)
10. Theta-functions and toroidal structure
11. Psychons and consciousness (speculative; see `speculative_extensions/`)
12. Experimental designs and testable predictions

# 2 Biquaternion Algebra

# 3 The Theta Field $\Theta(q, \tau)$

# 4 The Fundamental Biquaternion Field $\Theta(q, \tau)$

## 4.1 Definition

The fundamental field of Unified Biquaternion Theory is a **biquaternion**:

$$\boxed{\Theta(q, \tau) \in \mathcal{B} = \mathbb{H} \otimes \mathbb{C}} \tag{1}$$

where:

- $q \in \mathcal{B}$ is a biquaternion coordinate (full coordinate model: 8 real parameters)
- $\tau = t + i\psi$ is the canonical complex time
- Biquaternion time $T_B = t + i\psi + j\chi + k\xi$ is a deprecated/historical extension (see `speculative_extensions/`)
- $\mathcal{B} = \mathbb{H} \otimes \mathbb{C}$ denotes the biquaternion algebra
- $\mathbb{H}$ = quaternions with units $\{1, i, j, k\}$

**Coordinate bookkeeping note.** In this repository, $q$ can denote either (1) the full biquaternionic coordinate $q = z^\mu e_\mu \in \mathbb{C} \otimes \mathbb{H}$, or (2) the real-projected GR coordinate $q_{\text{GR}} = x^\mu e_\mu$ used in the classical GR recovery chain. The canonical complex-time coordinate/fiber is $\tau = z^0 = t + i\psi$. The GR-sector recovery assumes classical complex holomorphy only in tau.

### 4.1.1 Explicit Biquaternion Form

The field $\Theta$ is written explicitly as:

$$\Theta(q,\tau) = \Theta_0(q,\tau) + \Theta_1(q,\tau)i + \Theta_2(q,\tau)j + \Theta_3(q,\tau)k \tag{2}$$

where $\Theta_a(q,\tau) \in \mathbb{C}$ $(a = 0,1,2,3)$ are **complex-valued** component fields.

**Important**: $\Theta$ is a biquaternion (NOT a matrix). Matrix representations are computational tools only.

## 4.2 Faithful Matrix Representation and Degree Count

A biquaternion has four complex coefficients and therefore eight real components:

$$\Theta = \Theta_0 + \Theta_1\mathbf{i} + \Theta_2\mathbf{j} + \Theta_3\mathbf{k}, \qquad \Theta_a \in \mathbb{C}. \tag{3}$$

A faithful computational representation is a constrained $2 \times 2$ complex matrix,

$$\rho(\Theta) = \begin{pmatrix} \Theta_0 + i\Theta_1 & \Theta_2 + i\Theta_3 \\ -\Theta_2 + i\Theta_3 & \Theta_0 - i\Theta_1 \end{pmatrix}. \tag{4}$$

Thus

$$\dim_{\mathbb{C}} \mathbb{B} = 4, \qquad \dim_{\mathbb{R}} \mathbb{B} = 8. \tag{5}$$

A generic $4 \times 4$ or $8 \times 8$ complex matrix is not a biquaternion. Matrix-valued flavour or Standard-Model extensions, when used, must be denoted separately (for example $\Theta_{\text{ext}} \in \text{Mat}_n(\mathbb{B})$) and must not be silently substituted for the canonical core field.

## 4.3 Hermitian Involution

For variational and metric formulas we use the full Hermitian involution $\ddagger$, which combines complex conjugation with quaternion conjugation and corresponds to matrix Hermitian conjugation in the representation (4):

$$\rho(\Theta^{\ddagger}) = \rho(\Theta)^{\dagger}. \tag{6}$$

The notation must distinguish this operation from quaternion conjugation without complex conjugation.

## 4.4 Physical Interpretation

The field $\Theta(q,\tau)$ encodes:

1. **Geometric structure**: The covariant tetrad $E_\mu = \mathcal{N}_0^{-1/2} D_\mu \Theta$ emerges from $\Theta$. On the classical Lorentz slice its central anticommutator satisfies $\frac{1}{2}(E_\mu^\sharp E_\nu + E_\nu^\sharp E_\mu) = g_{\mu\nu}\mathbf{1}$, so no trace, real-part, or fiber projection defines the metric
2. **Matter content**: Fermion fields and masses arise from internal structure
3. **Gauge fields**: Electromagnetic and weak/strong interactions emerge from phase structure
4. **Internal auxiliary sector**: Imaginary time component $\psi$ provides dynamics for an internal auxiliary field (speculative interpretations in `speculative_extensions/`)

## 4.5 Field Equations and Covariant Derivative Status

A compact candidate master equation used in UBT is

$$D^\dagger D\Theta(q,\tau) = \kappa\mathcal{T}(q,\tau). \tag{7}$$

Its precise source functional, independent variables, boundary data, and common-action origin remain part of the dynamical closure programme. The geometric first jet entering the metric is

$$E_\mu = \mathcal{N}_0^{-1/2} D_\mu\Theta. \tag{8}$$

### 4.5.1 Classical connection versus its action on $\Theta$

Write $E_\mu = ie_\mu{}^0\mathbf{1} + e_\mu{}^k\mathbf{e}_k$. For every nondegenerate tetrad and specified torsion, the metric-compatible Lorentz-frame connection is kinematically unique,

$$\omega = \mathring{\omega}(e) + K(T). \tag{9}$$

For $T = 0$ this is the Levi–Civita spin connection. The affine connection $\Gamma^\rho{}_{\mu\nu}$ and frame connection $\omega_\mu{}^a{}_b$ are related by tetrad compatibility; neither is a real-part projection of the other.

This reconstruction does not by itself fix how the connection and internal gauge structure act on the biquaternionic field. A naive one-sided regular action is excluded for generic invertible torsion-free curved configurations. The active algebra-native candidate is the two-sided derivative

$$\boxed{D_\mu\Theta = \partial_\mu\Theta + A_\mu\Theta - \Theta B_\mu,} \tag{10}$$

with

$$[D_\mu, D_\nu]\Theta = F^A_{\mu\nu}\Theta - \Theta F^B_{\mu\nu}. \tag{11}$$

The exact decomposition of $A_\mu, B_\mu$ into Lorentz and Standard-Model connections must be derived from the same UBT action. In the invertible torsion-free branch, integrability requires the curvature-intertwiner relation $F^A_{\mu\nu} = \Theta F^B_{\mu\nu}\Theta^{-1}$; this permits nonzero curvature but does not prove existence of a curved solution.

In Cartesian inertial Minkowski gauge one may choose $\Gamma = \omega = \Omega = 0$, hence $D_\mu = \partial_\mu$. The affine field

$$\Theta_{\mathrm{SR}} = \Theta_0 + \sqrt{\mathcal{N}_0}(ix^0\mathbf{1} + x^k\mathbf{e}_k) \tag{12}$$

generates the constant Minkowski tetrad and has vanishing second spacetime derivatives.

The remaining dynamical gaps are torsion selection, the exact paired left/right lift, curved-space existence and uniqueness of $E_\mu = D_\mu\Theta/\sqrt{\mathcal{N}_0}$, Lorentz-slice preservation by the full dynamics, and derivation of Einstein equations from the canonical action.

## 4.6 Normalization

The field satisfies the normalization condition:

$$\mathrm{Tr}(\Theta^\dagger\Theta) = \mathrm{const.} \tag{13}$$

for bound states. The constant depends on the physical system under consideration.

## 4.7 Gauge Transformations

Under local gauge transformations:

$$\Theta(q,\tau) \to U(q,\tau)\Theta(q,\tau)V^\dagger(q,\tau) \tag{14}$$

where $U, V \in U(n)$ are unitary matrices encoding:

- $U(1)$ electromagnetic gauge symmetry
- $SU(2)$ weak isospin symmetry
- $SU(3)$ color symmetry

## 4.8 Reality Conditions

For physical observables, we impose:

$$\mathrm{Re}\,\mathrm{Tr}(\Theta^\dagger \mathcal{O}\Theta) \in \mathbb{R} \tag{15}$$

for any observable operator $\mathcal{O}$.

## 4.9 Asymptotic Behavior

At spatial infinity ($|q| \to \infty$):

$$\Theta(q,\tau) \to \Theta_0 + O(1/|q|) \tag{16}$$

where $\Theta_0$ is the vacuum configuration.

For temporal asymptotics (large $|t|$), boundary conditions depend on the specific physical scenario (scattering, bound states, etc.).

## 4.10 Connection to Biquaternions

The field can be expressed in biquaternion basis:

$$\Theta = \sum_{A=0}^{3}\sum_{B=0}^{3} \theta_{AB}(q,\tau)\,\sigma_A \otimes \sigma_B \tag{17}$$

where $\sigma_A$ are Pauli matrices (with $\sigma_0 = I$) and $\theta_{AB}$ are complex scalar fields.

This representation makes the biquaternionic structure explicit.

## 4.11 Units and Dimensions

In natural units ($\hbar = c = 1$):

$$[\Theta] = [\text{mass}]^1 = [\text{length}]^{-1} \tag{18}$$

This ensures dimensional consistency with standard field theory.

## 4.12 Historical Note

This definition supersedes all previous versions found in the repository, including:

- 4D biquaternion representation (old preprint)
- Alternative spinor formulations
- Conflicting matrix dimensions

**This is the canonical version.** All other formulations should be considered historical or deprecated.

# 5 Complex Time $\tau = t + i\psi$

# 6 Biquaternion Time $T_B$

## 6.1 Canonical Definition

The fundamental time coordinate in Unified Biquaternion Theory is a **biquaternion**:

$$\boxed{T_B = t + i\psi + j\chi + k\xi} \tag{19}$$

where:

- $t \in \mathbb{R}$ is the **real time coordinate** (standard physical time)
- $\psi, \chi, \xi \in \mathbb{R}$ are **imaginary time components**
- $i, j, k$ are quaternion units satisfying $i^2 = j^2 = k^2 = ijk = -1$

Equivalently, using vector notation:

$$T_B = t + i(\psi + \mathbf{v} \cdot \boldsymbol{\sigma}) = t + i\psi + j\chi + k\xi \tag{20}$$

where $\mathbf{v} = (\chi, \xi, 0)$ and $\boldsymbol{\sigma} = (j, k, ij)$ are quaternion vector components.

**Note**: The canonical UBT time coordinate is $\tau = t + i\psi$ (complex time). Biquaternion time $T_B = t + i\psi + j\chi + k\xi$ is a deprecated/historical extension preserved here for reference. It is not part of the canonical minimal theory.

## 6.2 Physical Interpretation

### 6.2.1 Real Component: Standard Time

The real part $t$ represents:

- Ordinary physical time
- Observable temporal evolution
- Causality structure of spacetime
- Time measured by physical clocks

### 6.2.2 Scalar Imaginary Component: Isotropic Phase

The scalar imaginary part $\psi$ represents:

- **Isotropic phase structure** of the $\Theta$ field
- **Internal auxiliary sector** (speculative; see `speculative_extensions/`)
- **Scalar dark energy** contributions
- **Universal quantum coherence** (direction-independent)

### 6.2.3 Vector Imaginary Components: Directional Phases

The vector imaginary parts $(\chi, \xi)$ represent:

- **Directional phase structures** in spacetime
- **Torsion effects** and spin-orbit coupling
- **Anisotropic dark matter** distributions
- **Directional phase modes** (collective internal sector)

**Critical**: All components $(\psi, \chi, \xi)$ are **dynamical variables** with physical consequences, not mathematical artifacts.

## 6.3 Dynamics of Imaginary Components

The imaginary time components have their own dynamics. The full equations emerge from variation of the action, but schematically:

$$\Box\psi + m_\psi^2\psi = J_\psi^{\text{scalar}} \tag{21}$$

$$\nabla^2\mathbf{v} + m_v^2\mathbf{v} = \mathbf{J}_v^{\text{vector}} \tag{22}$$

where:

- $\Box = \partial_\mu\partial^\mu$ is the d'Alembertian operator
- $m_\psi, m_v$ are effective mass scales
- $J_\psi, \mathbf{J}_v$ are sources coupling to matter/internal sector
- $\mathbf{v} = (\chi, \xi, 0)$ is the vector imaginary component

These equations couple through the full $\Theta$ field dynamics.

## 6.4 Hierarchical Reduction Structure

### 6.4.1 Complex Time Limit

When the vector components are negligible ($\|\mathbf{v}\|^2 \ll \psi^2$), or when there is directional isotropy:

$$\boxed{\chi, \xi \to 0 \quad \Rightarrow \quad T_B \to \tau = t + i\psi} \tag{23}$$

This **complex time limit** is valid for:

- Spherically symmetric systems
- Weakly coupled systems
- Weakly-coupled isotropic sector
- Isotropic cosmological backgrounds

### 6.4.2 Classical Time Limit

When all imaginary components vanish:

$$\psi, \chi, \xi \to 0 \quad \Rightarrow \quad T_B \to t \quad \Rightarrow \quad \text{UBT reduces to standard GR/QFT} \tag{24}$$

This ensures:

- Einstein equations recovered exactly
- Standard Model preserved
- All experimental tests of GR/QFT satisfied

### 6.4.3 Hierarchy Summary

$$\boxed{T_B = t + i\psi + j\chi + k\xi \quad \xrightarrow{\|\mathbf{v}\| \to 0} \quad \tau = t + i\psi \quad \xrightarrow{\psi \to 0} \quad t} \tag{25}$$

**Biquaternion time** (full) → **Complex time** (isotropic limit) → **Classical time** (GR limit)

## 6.5 Measurement and Observability

### 6.5.1 Direct Observability

The imaginary components $(\psi, \chi, \xi)$ are **not directly observable** in classical measurements because:

- Ordinary matter couples only to real metric $g_{\mu\nu} := \mathrm{Re}(\mathcal{G}_{\mu\nu})$ where $\mathcal{G}_{\mu\nu}$ is the biquaternionic metric
- Classical observables involve $\mathrm{Re}\,\mathrm{Tr}(\ldots)$
- Phase information is hidden in quantum coherence

### 6.5.2 Indirect Detection

The imaginary time components may in principle leave observable traces through:

- Quantum entanglement signatures
- Dark-sector interactions (*speculative/open*)
- Anisotropic dark-sector contributions via vector components (*speculative/open*)
- Torsion effects in strong gravity (vector components)

*Consciousness/psychon interpretations are speculative and not part of core UBT. See* `speculative_extensions/consciousness/` *for that material.*

## 6.6 Mathematical Properties

### 6.6.1 Biquaternion Conjugation

The full conjugate combines complex and quaternion conjugation:

$$T_B^\dagger = t - i\psi - j\chi - k\xi \tag{26}$$

### 6.6.2 Norm

$$|T_B|^2 = T_B \cdot T_B^\dagger = t^2 + \psi^2 + \chi^2 + \xi^2 \tag{27}$$

### 6.6.3 Scalar and Vector Parts

$$T_B = t + i\psi_{\text{scalar}} + \mathbf{v}_{\text{vector}} \tag{28}$$

where $\mathbf{v}_{\text{vector}} = j\chi + k\xi$.

## 6.7 Derivatives and Calculus

### 6.7.1 Partial Derivatives

For a function $f(T_B) = f(t, \psi, \chi, \xi)$:

$$\frac{\partial f}{\partial T_B} = \frac{1}{4}\left(\frac{\partial f}{\partial t} - i\frac{\partial f}{\partial \psi} - j\frac{\partial f}{\partial \chi} - k\frac{\partial f}{\partial \xi}\right) \tag{29}$$

In the complex time limit ($\chi, \xi \to 0$), this reduces to:

$$\frac{\partial f}{\partial \tau} = \frac{1}{2}\left(\frac{\partial f}{\partial t} - i\frac{\partial f}{\partial \psi}\right) \tag{30}$$

### 6.7.2 Integration

Biquaternion time integration:

$$\int dT_B = \int dt + i\int d\psi + j\int d\chi + k\int d\xi \tag{31}$$

For physical observables, we typically integrate over real time only:

$$\mathcal{O}_{\text{phys}} = \int dt\, \text{Re}\, \mathcal{O}(T_B)|_{\psi,\chi,\xi=f(t)} \tag{32}$$

## 6.8 Relation to Other Formulations

### 6.8.1 Euclidean Time (Wick Rotation)

Biquaternion time $T_B$ is **NOT** the same as Euclidean time from Wick rotation:

- **Wick rotation**: $t \to -it_E$ (analytical continuation)
- **UBT biquaternion time**: $T_B = t + i\psi + j\chi + k\xi$ (physical extension)

In UBT, *all* components $t, \psi, \chi, \xi$ are real, physical parameters.

### 6.8.2 Thermal Field Theory

At finite temperature $T$, the scalar component $\psi$ may be related to thermal time:

$$\psi \sim \beta = \frac{1}{k_B T} \tag{33}$$

but this is a special case, not a general requirement. The vector components $(\chi, \xi)$ are independent of thermal effects.

## 6.9 Conflict Resolution

This definition supersedes the following conflicting versions:

1. **Drift-diffusion Fokker-Planck variant**: imaginary components as stochastic variables only
2. **Toroidal variant with $\theta$-functions**: time as modular parameter only
3. **Hermitized variant (Appendix F)**: imaginary components as purely mathematical
4. **Complex time only variants**: missing directional structure

**Canonical resolution**: $T_B = t + i\psi + j\chi + k\xi$ where *all imaginary components* are **dynamical physical fields** with equations of motion. Complex time $\tau = t + i\psi$ is the limiting case valid for isotropic systems.

## 6.10 Coordinate Systems

### 6.10.1 Biquaternion Coordinates

The full coordinate set is:

$$(q, T_B) = (q^0, q^1, q^2, q^3, t, \psi, \chi, \xi) \tag{34}$$

where $q^\mu$ $(\mu = 0, 1, 2, 3)$ are biquaternion spatial coordinates.
In the complex time limit:

$$(q, \tau) = (q^0, q^1, q^2, q^3, t, \psi) \tag{35}$$

### 6.10.2 Metric Signature

The extended metric in full biquaternion time has signature:

$$\text{signature}(g_{AB}) = (+,-,-,-,-,+,+,+) \quad \text{or similar} \tag{36}$$

depending on convention. The imaginary time coordinates typically have mixed signature from spatial coordinates.

## 6.11 Conservation Laws

Energy-momentum conservation in biquaternion time:

$$\frac{\partial T^{\mu\nu}}{\partial T_B} = 0 \tag{37}$$

This expands to separate conservation laws for each component:

$$\frac{\partial T^{\mu\nu}}{\partial t} = \text{sources from } \psi, \chi, \xi \tag{38}$$

$$\frac{\partial T^{\mu\nu}}{\partial \psi} = \text{phase sector coupling} \tag{39}$$

$$\frac{\partial T^{\mu\nu}}{\partial \chi}, \frac{\partial T^{\mu\nu}}{\partial \xi} = \text{torsion/anisotropic sources} \tag{40}$$

## 6.12 Causality Structure

The causal structure is determined by:

$$ds^2 = g_{\mu\nu}(T_B) dx^\mu dx^\nu \tag{41}$$

where $g_{\mu\nu}$ is the central covariant-tetrad metric of the canonical complex-time theory; the $T_B$ construction in this historical extension does not redefine it. Light cones may rotate in the extended time space, but physical causality (in real time $t$) is preserved through appropriate boundary conditions.

## 6.13 Symbol Standardization

| Symbol | Meaning |
|---|---|
| $T_B$ | Biquaternion time coordinate (deprecated/historical extension) |
| $\tau$ | Complex time coordinate (isotropic limit) |
| $t$ | Real time component |
| $\psi$ | Scalar imaginary time component |
| $\chi, \xi$ | Vector imaginary time components |
| $T_B^\dagger$ | Biquaternion conjugate of $T_B$ |

**Forbidden uses**:

- $\psi$ for wavefunction (use $\Psi$ instead)
- $\psi$ for spinor (use $\psi_{\text{spinor}}$ if needed)
- $\tau$ for proper time (use $s$ or $\lambda$)
- $T$ for temperature (use $\mathcal{T}$ or specify $T_{\text{temp}}$)

## 6.14 Units

In natural units ($\hbar = c = 1$):

$$[t] = [\psi] = [\chi] = [\xi] = [\text{length}] = [\text{time}] \tag{42}$$

All components have dimension of length (or inverse energy).

## 6.15 Historical Note

The canonical UBT time coordinate is complex time $\tau = t + i\psi$. Biquaternion time $T_B = t + i\psi + j\chi + k\xi$ is a deprecated/historical extension. The imaginary component $\psi$ is a dynamical field; extensions $\chi, \xi$ are not part of the canonical minimal theory.

Complex time $\tau = t + i\psi$ is the canonical UBT time coordinate. The earlier biquaternion extensions $\chi, \xi$ may appear in exploratory research tracks but are not canonical. This document is preserved for reference; any new canonical content should use $\tau$.

## 6.16 Simplification: Complex Time Limit

**When to use complex time**: For systems with directional isotropy or weak coupling, the complex time limit $\tau = t + i\psi$ is sufficient. This includes:

- Spherically symmetric gravitational systems
- Cosmological models with isotropy
- Individual weakly-coupled sector (isotropic modes)
- Weak-field approximations

**Biquaternion time $T_B$ (deprecated extension)**: Extensions $\chi, \xi$ have been proposed for strong torsion effects, anisotropic dark matter, and non-Abelian dynamics, but these are speculative and belong in `research_tracks/` or `speculative_extensions/`.

The canonical formulation uses $\tau = t + i\psi$. Biquaternion time $T_B$ is a deprecated extension for research purposes only.

# 7 Fundamental Biquaternionic Geometry

## 7.1 Overview

The local classical geometry is carried by

$$E_\mu = \mathcal{N}_0^{-1/2} D_\mu \Theta$$

and the exact central identity

$$\frac{1}{2}(E_\mu^\sharp E_\nu + E_\nu^\sharp E_\mu) = g_{\mu\nu}\mathbf{1}.$$

The symmetric product defines the metric, the antisymmetric algebraic product is a bivector candidate, and the commutator $[D_\mu, D_\nu]$ is the curvature of the frame connection. These three structures must not be conflated.

**Canonical geometric hierarchy**

1. Fundamental field: $\Theta(q,\tau) \in \mathbb{B}$.
2. Covariant first jet: $E_\mu = \mathcal{N}_0^{-1/2} D_\mu \Theta$.
3. Metric: $\{E_\mu, E_\nu\}_\sharp/2 = g_{\mu\nu}\mathbf{1}$ on the Lorentz slice.
4. Classical frame connection: for specified torsion, $\omega = \mathring{\omega}(e) + K(T)$ is uniquely determined by the tetrad; the UBT action must select the torsion and its lift to $(A_\mu, B_\mu)$.
5. Coordinate connection: $\Gamma^\rho{}_{\mu\nu}$, related to $\Omega_\mu$ by tetrad compatibility.
6. Curvature: $[D_\mu, D_\nu]$ in the relevant representation.

**Status:** local metric rank, connection reconstruction from specified $(e,T)$, the torsion-free Levi–Civita branch, and flat affine representers are proved. Torsion dynamics, curved integrability, and Einstein dynamics remain open.

**Matter-sector caution**

The tetrad, metric, connection, and curvature files below implement the current geometric core. The stress-energy and exotic-regime files retain older candidate projection language and are not part of the proved metric theorem.

# 8 The covariant biquaternionic tetrad $E_\mu$

The fundamental UBT field is $\Theta(q,\tau) \in \mathbb{B}$. The local geometric object is its normalized covariant first derivative

$$\boxed{E_\mu := \mathcal{N}_0^{-1/2} D_\mu \Theta,} \tag{43}$$

where $\mathcal{N}_0 > 0$ is a fixed unit-setting constant. No local denominator is allowed.

## 8.1 Classical Lorentz sector

Let $i$ be the commuting complex unit and $\mathbf{e}_k$ the quaternion units. The real Lorentz slice is

$$W_L = \{ix^0\mathbf{1} + x^k\mathbf{e}_k \mid x^a \in \mathbb{R}\}. \tag{44}$$

Quaternion conjugation is denoted by $\sharp$ and acts as $(z^0\mathbf{1} + z^k\mathbf{e}_k)^\sharp = z^0\mathbf{1} - z^k\mathbf{e}_k$. The classical tetrad sector is defined by $E_\mu \in W_L$ and $\det(e_\mu{}^a) \neq 0$, where

$$E_\mu = ie_\mu{}^0\mathbf{1} + e_\mu{}^k\mathbf{e}_k. \tag{45}$$

## 8.2 Metric and bivector from one product

On $W_L$ the symmetrised product is central:

$$\boxed{\frac{1}{2}(E_\mu^\sharp E_\nu + E_\nu^\sharp E_\mu) = g_{\mu\nu}\mathbf{1},} \tag{46}$$

with

$$g_{\mu\nu} = e_\mu{}^a e_\nu{}^b \eta_{ab}, \qquad \eta_{ab} = \mathrm{diag}(-1,1,1,1). \tag{47}$$

The metric is therefore not obtained by a trace, real-part projection, phase projection, or compact-fiber average. It is the unique coefficient of the unit element because the full symmetrised product is already central.

The antisymmetric companion

$$\Sigma_{\mu\nu} := \frac{1}{2}(E_\mu^\sharp E_\nu - E_\nu^\sharp E_\mu) \tag{48}$$

is a biquaternionic bivector carrying local oriented-plane/spin information. It is distinct from the connection curvature $[D_\mu, D_\nu]$.

### 8.3 Transformation and rank

Under coordinate changes $E_\mu$ transforms as a covector. Under local Lorentz-frame changes, its coefficients transform by a local Lorentz matrix; Eq. (46) is invariant.

The map $e_\mu{}^a \mapsto g_{\mu\nu}$ has rank ten at every nondegenerate tetrad. Its six-dimensional kernel consists of local Lorentz rotations and boosts. A proof and exact symbolic verification are given in `canonical/gr_closure/covariant_tetrad_rank_theorem.tex` and `tools/verify_covariant_tetrad_rank.py`. The same $E_\mu$ also has an injective block lift to curved Clifford operators $\Gamma_\mu$ satisfying $\{\Gamma_\mu, \Gamma_\nu\}/2 = g_{\mu\nu} I_4$; see `canonical/geometry/biquaternion_`
No second spinor-current tetrad is introduced.

### 8.4 Status and open bridge

The local metric rank problem is solved at the tetrad level. The metric-compatible frame connection is also kinematically reconstructed from the tetrad and specified torsion, $\omega = \mathring{\omega}(e) + K(T)$; in the torsion-free GR branch this is the unique Levi–Civita spin connection. Every constant Lorentz tetrad has the explicit affine representer $\Theta = \Theta_0 + \sqrt{\mathcal{N}_0} E_\mu x^\mu$ in the flat inertial branch.

For curved configurations, however, it remains to derive torsion and the exact paired left/right connection from the UBT action and to solve the implicit system $E_\mu = D_\mu \Theta / \sqrt{\mathcal{N}_0}$. A naive one-sided regular action is generically flat for invertible $\Theta$ under torsion-free compatibility; the two-sided bimodule derivative avoids this obstruction but does not yet prove curved-space existence.

## 9 The central anticommutator metric

Let $E_\mu = \mathcal{N}_0^{-1/2} D_\mu \Theta$ be the covariant UBT tetrad. Define first the full ordered biquaternionic geometric tensor

$$\mathfrak{G}_{\mu\nu} := E_\mu^\sharp E_\nu \in \mathbb{C} \otimes \mathbb{H}. \tag{49}$$

Historical UBT notes sometimes called $\mathfrak{G}_{\mu\nu}$ the "biquaternionic metric". Canonical GR terminology reserves *metric* for its symmetric central channel, because the full ordered tensor is not symmetric and has no canonical noncommutative determinant or Levi–Civita inverse. No algebraic information is discarded by this terminology.

On the classical Lorentz slice $W_L$, the fundamental quadratic identity is

$$\boxed{\mathfrak{G}_{\mu\nu} = E_\mu^\sharp E_\nu = g_{\mu\nu}\mathbf{1} + \Sigma_{\mu\nu},} \tag{50}$$

where

$$g_{\mu\nu}\mathbf{1} = \frac{1}{2}(E_\mu^\sharp E_\nu + E_\nu^\sharp E_\mu), \tag{51}$$

$$\Sigma_{\mu\nu} = \frac{1}{2}(E_\mu^\sharp E_\nu - E_\nu^\sharp E_\mu). \tag{52}$$

The first term is symmetric and central; the second is antisymmetric and biquaternionic. The classical metric is therefore defined intrinsically, not by applying a trace, real-part operator, phase projector, or fiber average to a noncentral object.

In components,

$$g_{\mu\nu} = e_\mu{}^a e_\nu{}^b \eta_{ab}, \qquad \eta_{ab} = \mathrm{diag}(-1,1,1,1). \tag{53}$$

Nondegeneracy is equivalent to invertibility of $e_\mu{}^a$. The local Lorentzian signature follows immediately.

The full product in Eq. (126) remains relevant: its antisymmetric part may carry spin/bivector information. No part is discarded, but no identification with a gauge field is claimed without a transformation and dynamical derivation.

# 10 Clifford and generalized-Dirac lift of the canonical relation

## 10.1 Scope: no second tetrad

The starting point remains the canonical UBT tetrad

$$E_\mu = \mathcal{N}_0^{-1/2} D_\mu \Theta \in W_L \tag{54}$$

and the central metric identity

$$\frac{1}{2}(E_\mu^\sharp E_\nu + E_\nu^\sharp E_\mu) = g_{\mu\nu}\mathbf{1}. \tag{55}$$

The construction below introduces no independent vielbein and no spinor-current replacement for $E_\mu$. It is an injective operator lift of the same four biquaternionic covectors.

## 10.2 Exact block lift

Use the Pauli representation

$$\mathbf{e}_k = -i_{\mathrm{c}}\sigma_k, \qquad E = i_{\mathrm{c}} e^0 \mathbf{1} + e^k \mathbf{e}_k, \qquad e^a \in \mathbb{R}, \tag{56}$$

where $i_{\mathrm{c}}$ is the commuting complex unit. Regard $E$ as a $2\times 2$ complex matrix and define

$$\boxed{\mathcal{C}(E) := \begin{pmatrix} 0 & E \\ E^\sharp & 0 \end{pmatrix} \in \mathrm{Mat}(4,\mathbb{C}).} \tag{57}$$

The map $\mathcal{C} : W_L \to \mathrm{Mat}(4,\mathbb{C})$ is real-linear and injective.

**Theorem (canonical relation implies the curved Clifford relation).** For all $E, F \in W_L$,

$$\frac{1}{2}\{\mathcal{C}(E), \mathcal{C}(F)\} = H_L(E, F) I_4, \tag{58}$$

where

$$H_L(E, F)\mathbf{1} = \frac{1}{2}(E^\sharp F + F^\sharp E). \tag{59}$$

Consequently, with

$$\Gamma_\mu := \mathcal{C}(E_\mu), \tag{60}$$

Eq. (55) is equivalent to

$$\boxed{\frac{1}{2}\{\Gamma_\mu, \Gamma_\nu\} = g_{\mu\nu} I_4.} \tag{61}$$

*Proof.* Direct block multiplication gives

$$\frac{1}{2}\{\mathcal{C}(E), \mathcal{C}(F)\} = \frac{1}{2}\begin{pmatrix} EF^\sharp + FE^\sharp & 0 \\ 0 & E^\sharp F + F^\sharp E \end{pmatrix}. \tag{62}$$

For Lorentz paravectors both diagonal expressions equal $2H_L(E, F)\mathbf{1}$. Hence the block matrix is $H_L(E, F) I_4$. Injectivity follows because either off-diagonal block recovers $E$. □

## 10.3 Spinor carrier without a new fundamental field

As a complex vector space, $\mathbb{B}$ has dimension four. Fix any complex-linear isomorphism

$$\mathrm{vec} : \mathbb{B} \longrightarrow \mathbb{C}^4, \qquad \Psi := \mathrm{vec}(\Theta). \tag{63}$$

The symbol $\Psi$ therefore denotes the same four-complex-component UBT field in a column representation; it is not an additional matter field. The lifted matrices $\Gamma_\mu = \mathcal{C}(E_\mu)$ act on this carrier. A different choice $\mathrm{vec}' = U\,\mathrm{vec}$, $U \in GL(4, \mathbb{C})$, gives

$$\Psi' = U\Psi, \qquad \Gamma'_\mu = U\Gamma_\mu U^{-1}, \tag{64}$$

and leaves the Clifford relation, the metric and the characteristic determinant invariant. Thus the block representation is a faithful operator realisation of the canonical biquaternionic relation, not a change of physical field content.

## 10.4 Rank is inherited, not manufactured

The operator lift adds no geometric degrees of freedom. The metric is still that of the tetrad coefficients $e_\mu{}^a$,

$$g = e\eta e^T, \qquad \eta = \mathrm{diag}(-1, 1, 1, 1). \tag{65}$$

At every invertible tetrad, the differential $\delta e \mapsto \delta g$ is surjective onto symmetric $4 \times 4$ matrices. Indeed, for an arbitrary symmetric $h$, the choice

$$\delta e = \frac{1}{2} h\, g^{-1} e \tag{66}$$

obeys $\delta g = h$. The kernel consists precisely of

$$\delta e = eX, \qquad X\eta + \eta X^T = 0, \tag{67}$$

which is the six-dimensional Lorentz Lie algebra. Therefore

$$\boxed{\operatorname{rank} D_e g = 10, \qquad \dim\ker D_e g = 6.} \tag{68}$$

Because $\mathcal{C}$ is injective, the same rank statement holds when the four $E_\mu$ are represented by the four $\Gamma_\mu$.

This theorem concerns the map $E \mapsto g$. The stronger on-shell statement that the complete UBT equation admits enough solutions of $E_\mu = D_\mu\Theta/\sqrt{\mathcal{N}_0}$ remains a separate dynamical theorem.

## 10.5 Complex-time / fifth operator channel

Define the grading matrix

$$\Gamma_* := \begin{pmatrix} I_2 & 0 \\ 0 & -I_2 \end{pmatrix}. \tag{69}$$

Every matrix $\mathcal{C}(E)$ is off diagonal, so

$$\{\Gamma_*, \mathcal{C}(E)\} = 0, \qquad \Gamma_*^2 = I_4. \tag{70}$$

Thus one may form either

$$\Gamma_\psi = \Gamma_*, \qquad \Gamma_\psi^2 = +I_4, \tag{71}$$

or

$$\Gamma_\psi = i_{\mathrm{c}}\Gamma_*, \qquad \Gamma_\psi^2 = -I_4, \tag{72}$$

according to the signature assigned to the $\psi$ channel. This establishes an exact algebraic fifth channel compatible with the four operators derived from $E_\mu$; it does not by itself decide whether $\psi$ is a physical fifth coordinate, a compact fibre, or an internal complex-time direction.

The corresponding generalized first-order operator has the candidate form

$$\boxed{\mathcal{D}_{\mathrm{UBT}}[\Theta]\Psi = i_{\mathrm{c}}\Gamma^\mu[D\Theta]\nabla_\mu\Psi + i_{\mathrm{c}}\Gamma_\psi D_\psi\Psi - \mathcal{M}[\Theta, D\Theta]\Psi.} \tag{73}$$

Equation (73) is an architectural candidate until it is derived from the UBT action. The proved statement is the Clifford lift and its compatibility with the central metric relation.

## 10.6 Principal symbol, causal cone and exact factorisation

Let

$$\Gamma^\mu := g^{\mu\nu}\Gamma_\nu \tag{74}$$

and, for a real cotangent vector $\xi = \xi_\mu dx^\mu$, define the first-order principal symbol

$$\sigma_4(\xi) := \Gamma^\mu\xi_\mu. \tag{75}$$

The curved Clifford relation immediately gives

$$\boxed{\sigma_4(\xi)^2 = g^{\mu\nu}\xi_\mu\xi_\nu I_4.} \tag{76}$$

Since $\sigma_4(\xi)$ is block off diagonal and traceless, its characteristic polynomial is

$$\chi_{\sigma_4}(\lambda) = \left(\lambda^2 - g^{\mu\nu}\xi_\mu\xi_\nu\right)^2, \tag{77}$$

and therefore

$$\boxed{\det \sigma_4(\xi) = \left(g^{\mu\nu}\xi_\mu\xi_\nu\right)^2 .} \tag{78}$$

Thus the characteristic cone of the canonical generalized-Dirac lift is exactly the null cone of the metric already defined by the UBT relation. No second causal structure is introduced.

Including a fifth anticommuting channel with $\Gamma_\psi^2 = \varepsilon I_4$, $\varepsilon = \pm 1$, gives

$$\sigma_5(\xi, \xi_\psi) := \sigma_4(\xi) + \Gamma_\psi \xi_\psi, \tag{79}$$

and the exact factorisation

$$\boxed{\sigma_5(\xi, \xi_\psi)^2 = \left(g^{\mu\nu}\xi_\mu\xi_\nu + \varepsilon\xi_\psi^2\right) I_4.} \tag{80}$$

Equation (80) is algebraic. Its physical interpretation depends on whether $\psi$ is treated as a geometric coordinate, a compact fibre coordinate, or an internal complex-time spectral direction.

## 10.7 Conditional first-jet rank preservation by the fifth channel

The fifth matrix is invertible. If $\Gamma_\psi^2 = \varepsilon I_4$, then

$$\Gamma_\psi^{-1} = \varepsilon \Gamma_\psi. \tag{81}$$

Consider a first-order equation written in $\psi$-normal form,

$$\Gamma_\psi D_\psi \Psi + \mathcal{F}(\Psi, E_0, E_1, E_2, E_3) = 0, \tag{82}$$

where $\mathcal{F}$ contains no $D_\psi\Psi$. For every prescribed value of $\Psi$ and every prescribed Lorentz-admissible four-tetrad $E_\mu$, the unique pointwise solution is

$$\boxed{D_\psi \Psi = -\varepsilon \Gamma_\psi \mathcal{F}(\Psi, E_0, E_1, E_2, E_3).} \tag{83}$$

Consequently, at the purely algebraic first-jet level, the equation manifold of (82) projects surjectively onto the four spacetime tetrad slots $E_\mu$. Under these stated hypotheses the equation therefore does not lower the already proved rank ten of $E \mapsto g$.

This is a conditional pointwise theorem, not yet the full UBT on-shell theorem. Canonical UBT also uses holomorphy in $\tau = t + i_c\psi$. A strict Cauchy–Riemann condition relates $D_\psi\Psi$ to the real-time derivative and removes its independent first-jet status. The decisive remaining task is to show either that the covariant complex-time equation possesses a compatible normal form after holomorphy is imposed, or directly that its constrained jet projection onto the ten metric directions remains surjective.

## 10.8 No-extra-field constrained-rank theorem

Strict holomorphy removes the independent $D_\psi\Psi$ slot, but it does not remove the value of the original field itself. The correct pointwise rank problem therefore uses only the original UBT variables

$$\Psi = \mathrm{vec}(\Theta) \in \mathbb{C}^4, \qquad e = (e_\mu{}^a) \in \mathbb{R}^{16}, \tag{84}$$

and no additional field.

Let the real form of the four-complex-component generalized-Dirac equation be

$$F(\Psi, e) = 0 \in \mathbb{R}^8. \tag{85}$$

At a solution, write its Jacobian blocks as

$$F_\Psi : \mathbb{R}^8 \to \mathbb{R}^8, \qquad F_e : \mathbb{R}^{16} \to \mathbb{R}^8. \tag{86}$$

The admissible tetrad variations are exactly

$$\mathcal{A} := \left\{ \delta e \in \mathbb{R}^{16} : \ F_e \delta e \in \operatorname{im} F_\Psi \right\}. \tag{87}$$

Let

$$K := \ker D_e g, \qquad \dim K = 6, \tag{88}$$

be the Lorentz-frame kernel proved above.

**Theorem (exact constrained metric rank).** For every nondegenerate tetrad and every smooth constraint (85),

$$\boxed{\operatorname{rank}\left(D_e g\big|_{\mathcal{A}}\right) = \dim(\mathcal{A} + K) - 6.} \tag{89}$$

Consequently the full metric rank ten is retained if and only if

$$\boxed{\mathcal{A} + K = \mathbb{R}^{16}.} \tag{90}$$

*Proof.* The map $D_e g$ has kernel $K$ and induces an isomorphism from $\mathbb{R}^{16}/K$ onto the ten-dimensional space of symmetric metric variations. The image of $\mathcal{A}$ is therefore isomorphic to $(\mathcal{A} + K)/K$. Its dimension is $\dim(\mathcal{A} + K) - \dim K$, which gives (89). □

**Corollary (original-field absorption).** If

$$\operatorname{rank}_{\mathbb{R}} F_\Psi = 8, \tag{91}$$

then $\operatorname{im} F_\Psi = \mathbb{R}^8$, so $\mathcal{A} = \mathbb{R}^{16}$. For every tetrad variation there is a unique field-value variation

$$\delta\Psi = -F_\Psi^{-1} F_e \delta e \tag{92}$$

that remains tangent to the equation manifold. Hence

$$\boxed{\operatorname{rank}\left(D_e g\big|_{F=0}\right) = 10.} \tag{93}$$

No compensator, auxiliary field, second biquaternion, fibre average, or split jet has been introduced: the eight real directions used here are the value of the original $\Theta$ field.

**No-go without field-value absorption.** If $F_\Psi = 0$ and $F_e$ has eight independent real rows, then $\dim \mathcal{A} = 8$, and therefore

$$\operatorname{rank}\left(D_e g\big|_{\mathcal{A}}\right) \le 8. \tag{94}$$

Thus four independent complex first-order equations acting only on the sixteen tetrad components cannot generically retain metric rank ten. They can avoid this conclusion only if their real rank is at most six or if the removed directions are transverse only to Lorentz gauge, as expressed exactly by (90).

## 10.9 Application to the holomorphic generalized-Dirac equation

Use the covariant complex-time condition

$$D_\psi \Psi = i_{\mathrm{c}} D_0 \Psi \tag{95}$$

and the canonical identification

$$D_\mu \Psi = \sqrt{\mathcal{N}_0}\ \mathrm{vec}(E_\mu). \tag{96}$$

Then the candidate first-order equation

$$i_{\mathrm{c}} \Gamma^\mu(e) D_\mu \Psi + i_{\mathrm{c}} \Gamma_\psi D_\psi \Psi - \mathcal{M}\Psi = 0 \tag{97}$$

becomes the pointwise constraint

$$\mathcal{M}\Psi = \sqrt{\mathcal{N}_0}\,[i_{\mathrm{c}} \Gamma^\mu(e)\,\mathrm{vec}(E_\mu) - \Gamma_\psi\,\mathrm{vec}(E_0)]\,. \tag{98}$$

The right-hand side depends on the tetrad first jet, while the left-hand side uses the value of the same master field.

If $\mathcal{M} = mI_4$ with $m \neq 0$, then the complex determinant is $m^4$ and the real Jacobian has rank eight. Equation (98) uniquely determines $\Psi$ for every Lorentz-admissible tetrad, so the equation manifold is locally a graph over all sixteen tetrad slots and the metric rank is ten.

More generally, for the scalar–pseudoscalar block

$$\mathcal{M} = m_s I_4 + i_{\mathrm{c}} m_p \Gamma_*, \tag{99}$$

one has

$$\mathcal{M}^{-1} = \frac{m_s I_4 - i_{\mathrm{c}} m_p \Gamma_*}{m_s^2 + m_p^2}, \qquad \det_{\mathbb{C}} \mathcal{M} = (m_s^2 + m_p^2)^2, \tag{100}$$

so the realified determinant is $(m_s^2 + m_p^2)^4$. It therefore preserves rank ten whenever $(m_s, m_p) \neq (0, 0)$.

**Status and limitation.** Equations (89)– (93) are unconditional finite-dimensional rank theorems. Their application to (98) is conditional only on the finalized UBT equation having an invertible original-field Jacobian $F_\Psi$; a nonzero scalar or scalar–pseudoscalar zero-order block is an explicit sufficient realization. Deriving that block and its normalization from the canonical UBT action, and proving local solutions of the resulting implicit first-order PDE, remain dynamical tasks. The rank obstruction itself is thereby reduced to the exact transversality condition (90), without adding variables.

## 10.10 Remaining proof gates

A complete closure requires:

1. derivation of $\mathcal{D}_{\mathrm{UBT}}$ and $\mathcal{M}$ from one covariant UBT action over $\tau = t + i_{\mathrm{c}}\psi$;
2. proof that the implicit cycle $\Theta \to E \to \Gamma \to \omega(E) \to D\Theta$ has local solutions;
3. preservation of the Lorentz slice and nondegeneracy under evolution;

4. derivation from the canonical action of a generalized-Dirac zero-order block whose original-field Jacobian $F_\Psi$ is invertible, or an alternative proof of the exact transversality condition $\mathcal{A} + K = \mathbb{R}^{16}$;

5. local existence and integrability of the resulting implicit holomorphic first-order PDE (the pointwise constrained rank is proved under the preceding transversality condition);

6. derivation of the Einstein low-energy dynamics and the quantum limits.

The historical spinor-current tetrad is not used in these gates.

# 11 The biquaternionic frame connection $\Omega_\mu$

A connection specifies how local Lorentz/biquaternionic frames are compared at neighbouring spacetime points. The UBT tetrad is

$$E_\mu = \mathcal{N}_0^{-1/2} D_\mu \Theta. \tag{101}$$

The action of $D_\mu$ must always be stated explicitly; left, right, adjoint, and two-sided actions are not interchangeable.

## 11.1 Classical Lorentz connection reconstructed from tetrad and torsion

Write $E_\mu = i e_\mu{}^0 \mathbf{1} + e_\mu{}^k \mathbf{e}_k$. Let

$$T^a = de^a + \omega^a{}_b \wedge e^b = \frac{1}{2} T^a{}_{bc} e^b \wedge e^c. \tag{102}$$

For every nondegenerate tetrad and specified torsion, there is exactly one metric-compatible Lorentz connection,

$$\boxed{\omega^a{}_b = \mathring{\omega}^a{}_b(e) + K^a{}_b(T),} \tag{103}$$

where

$$\mathring{\omega}_\mu{}^a{}_b = e_b{}^\nu \left( \mathring{\Gamma}^\rho{}_{\mu\nu}(g) e_\rho{}^a - \partial_\mu e_\nu{}^a \right) \tag{104}$$

and, with the displayed torsion convention,

$$\boxed{K_{abc} = \frac{1}{2}(T_{cab} - T_{abc} - T_{bca}).} \tag{105}$$

Thus the connection has no residual kinematic ambiguity once $(e, T)$ are fixed. In the torsion-free classical GR branch, $T = 0$, $K = 0$, and $\omega = \mathring{\omega}(e)$.

The biquaternionic/spin connection is the lift

$$\Omega_\mu = \rho_{\mathrm{spin}*}(\omega_\mu) = \frac{1}{4} \omega_{\mu ab} \Sigma^{ab} \tag{106}$$

for the chosen representation. The complete proof is in `canonical/gr_closure/gap_10omega_connecti`

## 11.2 Relation to Christoffel symbols

The coordinate and frame descriptions are joined by the tetrad postulate,

$$\partial_\mu e_\nu{}^a - \Gamma^\rho{}_{\mu\nu} e_\rho{}^a + \omega_\mu{}^a{}_b e_\nu{}^b = 0. \tag{107}$$

$\Gamma^\rho{}_{\mu\nu}$ transports coordinate indices, while $\omega$ and $\Omega$ transport local Lorentz/biquaternionic indices. They are related but not identical: they encode the same geometry in different representations. A componentwise identification such as $\Gamma = \mathrm{Re}\,\Omega$ is generally incorrect.

## 11.3 Gauge covariance and Lorentz-slice preservation

Under $e \mapsto \Lambda e$,

$$\omega \mapsto \Lambda\omega\Lambda^{-1} - d\Lambda\,\Lambda^{-1}. \tag{108}$$

This is a local Lorentz gauge change, not a new physical degree of freedom. Since every metric-compatible $\omega_\mu$ lies in $\mathfrak{so}(1,3)$, its parallel transport preserves $\eta_{ab}$ and the real Lorentz slice, including branches with contorsion. Hence GAP-10L-CONN is closed. The Lorentz slice is also the fixed set of $\mathcal{J}X = -\overline{X}^\sharp$, so every unique $\mathcal{J}$-equivariant evolution with fixed data and sources preserves it (GAP-10L-SYM, closed conditionally). GAP-10L-DYN is narrowed to verifying these equivariance and well-posedness hypotheses for the complete $\Theta$ equations.

## 11.4 Representation selection on $\Theta$

A naive one-sided action

$$D^L_\mu \Theta = \partial_\mu \Theta + A_\mu \Theta \tag{109}$$

obeys $[D^L_\mu, D^L_\nu]\Theta = F^L_{\mu\nu}\Theta$. Under torsion-free tetrad compatibility, invertible $\Theta$ would then force $F^L = 0$. Therefore this one-sided regular representation is not the generic curved-GR route.

The algebra-native viable form is the two-sided bimodule derivative

$$\boxed{D_\mu \Theta = \partial_\mu \Theta + A_\mu \Theta - \Theta B_\mu,} \tag{110}$$

for which

$$\boxed{[D_\mu, D_\nu]\Theta = F^A_{\mu\nu}\Theta - \Theta F^B_{\mu\nu}.} \tag{111}$$

Torsion-free antisymmetric compatibility requires $F^A_{\mu\nu}\Theta = \Theta F^B_{\mu\nu}$; if $\Theta$ is invertible, the two curvatures are conjugate rather than zero. This removes the one-sided flatness obstruction, but the general pair is not thereby derived from the action.

## 11.5 Pure Lorentz pairing: exact reduction and exact obstruction

Let $W_L = \{X \mid X^\ddagger = -X\}$ be the Lorentz slice. Requiring the bimodule derivative to preserve $W_L$ for every $X \in W_L$ gives

$$B_\mu = -A^\ddagger_\mu + c_\mu \mathbf{1}, \qquad c_\mu \in \mathbb{R}. \tag{112}$$

Requiring in addition preservation of the Lorentz quadratic form removes the real dilation and reduces the connection action, modulo a common central one-form that cancels identically, to

$$\boxed{D_\mu X = \partial_\mu X + \Omega_\mu X + X\Omega_\mu^\ddagger,} \qquad \boxed{A_\mu = \Omega_\mu, \quad B_\mu = -\Omega_\mu^\ddagger.} \tag{113}$$

Thus $A_\mu$ and $B_\mu$ are not two new gravitational fields in this branch; they are the left and right representations of the one already reconstructed spin connection.

This economical reduction has two sharply different branches. Because Eq. (113) commutes with the Lorentz involution, a nondegenerate **torsion-free** solution $D_\mu\Theta = \sqrt{\mathcal{N}_0}E_\mu$ implies a spacetime vector $V$ obeying

$$\boxed{\mathring{\nabla}_\mu V^\nu = \delta_\mu{}^\nu,} \qquad \mathcal{L}_V g = 2g, \qquad R^\rho{}_{\sigma\mu\nu}V^\sigma = 0. \tag{114}$$

This concurrent-vector condition excludes the non-flat Schwarzschild vacuum exterior with nonzero mass and generic torsion-free Einstein geometries. Therefore GAP-10I-PAIR-GR is closed as a torsion-free no-go.

It is not a no-go for the same one-connection pair with composite torsion. On a sufficiently small non-null Gaussian patch, define

$$W_{\mu\nu} = g_{\mu\nu} - \mathring{\nabla}_\mu V_\nu, \qquad K_{\nu\mu\rho} = \frac{W_{\mu\nu}V_\rho - V_\nu W_{\mu\rho}}{V^2}, \tag{115}$$

with $V_\mu = \frac{1}{2}\partial_\mu V^2$. Then $K$ is metric-compatible and $\nabla_\mu^{(\mathring{\Gamma}+K)}V^\nu = \delta_\mu{}^\nu$. Taking $\Theta = \sqrt{\mathcal{N}_0}V^a\mathbf{u}_a$ gives exactly $D_\mu\Theta = \sqrt{\mathcal{N}_0}E_\mu$. This closes GAP-10I-TORSION-LOCAL locally without independent $A_\mu, B_\mu$ fields.

A relative remainder

$$A_\mu = \Omega_\mu + P_\mu, \qquad B_\mu = -\Omega_\mu^\ddagger + Q_\mu \tag{116}$$

is therefore optional for local kinematics, though it remains a possible torsion-free completion. If used, $P_\mu X - XQ_\mu$ must be derived from the canonical action and shown composite or auxiliary rather than propagating. A common central constant or one-form cannot remove the torsion-free obstruction because it cancels. The proofs are in `canonical/gr_closure/gap_10i_paired_connection_audit.tex` and `canonical/gr_closure/gap_10i_torsionful_local_representer.tex`.

## 11.6 Flat limit and explicit representer

In a flat inertial torsion-free sector,

$$\Gamma^\rho{}_{\mu\nu} = 0, \qquad \Omega_\mu = 0, \qquad D_\mu = \partial_\mu. \tag{117}$$

The explicit field

$$\Theta_{\mathrm{SR}} = \Theta_0 + \sqrt{\mathcal{N}_0}(ix^0\mathbf{1} + x^k\mathbf{e}_k) \tag{118}$$

generates the Minkowski tetrad and has vanishing second spacetime derivatives.

### 11.7 Full UBT status

The kinematic and sharply scoped conditional subproblems are:

- GAP-10Ω-KIN/GR: specified $(e, T)$ uniquely determine $\omega$, with the Levi–Civita torsion-free branch;
- GAP-10T-PALATINI (conditional): in the minimal first-order branch the 24-component Cartan torsion equation is algebraic and invertible;
- GAP-10L-CONN and GAP-10L-SYM (conditional): compatible transport and unique equivariant evolution preserve the Lorentz slice;
- GAP-10I-PRESCRIBED: prescribed $(E, A, B)$ obey the exact augmented-holonomy existence criterion.

The full-theory questions are narrowed rather than closed:

- GAP-10T-SPIN is closed conditionally for the direct fixed-background Palatini variation, while GAP-10T-FLAT-NOGO and GAP-10T-PAIRING-NOGO are closed as no-go results;
- GAP-10T-DYN: compute the full composite $\Theta$ variation and derive the selected first-order branch, a canonical torsion cancellation or translational/relative completion, and normalization from UBT;
- GAP-10I-PAIR-KIN is closed; GAP-10I-PAIR-GR is closed as a torsion-free no-go; GAP-10I-TORSION-LOCAL is closed locally;
- GAP-10I-2S is optional for local kinematics and remains a possible torsion-free composite/auxiliary completion;
- GAP-10I-CURVED has local kinematics closed, while action selection, physical torsion constraints, regularity and global continuation remain;
- GAP-10L-DYN: verify equivariance and well-posed uniqueness for the full dynamics.

## 12 Biquaternionic curvature

For the general two-sided biquaternionic derivative

$$D_\mu \Theta = \partial_\mu \Theta + A_\mu \Theta - \Theta B_\mu, \tag{119}$$

define

$$F^A_{\mu\nu} = \partial_\mu A_\nu - \partial_\nu A_\mu + [A_\mu, A_\nu], \tag{120}$$

$$F^B_{\mu\nu} = \partial_\mu B_\nu - \partial_\nu B_\mu + [B_\mu, B_\nu]. \tag{121}$$

Then

$$\boxed{[D_\mu, D_\nu]\Theta = F^A_{\mu\nu}\Theta - \Theta F^B_{\mu\nu}.} \tag{122}$$

The one-sided formula is recovered only as the special case $B_\mu = 0$.

In the torsion-free compatible branch, antisymmetric integrability requires

$$F^A_{\mu\nu}\Theta = \Theta F^B_{\mu\nu}. \tag{123}$$

For invertible $\Theta$, the curvatures may be nonzero but must be conjugate:

$$F^A_{\mu\nu} = \Theta F^B_{\mu\nu}\Theta^{-1}. \tag{124}$$

This is why the two-sided action avoids the generic flatness obstruction of a naive one-sided regular representation.

The Lorentz frame curvature built from the uniquely reconstructed $\omega = \mathring{\omega}(e) + K(T)$ represents the corresponding Riemann–Cartan geometry. For $T = 0$ it is the usual Riemann curvature of the Levi–Civita connection. The pure Lorentz identification is exactly $A_\mu = \Omega_\mu$, $B_\mu = -\Omega^\ddagger_\mu$ modulo a cancelling central term. With $K = 0$ this generated-tetrad branch is a concurrent-vector no-go for the non-flat Schwarzschild vacuum exterior. It is not a no-go for the same pairing with arbitrary metric-compatible torsion: an explicit composite-contortion construction supplies a local single-$\Theta$ representer for every smooth Lorentzian tetrad. What remains open is canonical action selection and physical admissibility of that torsion (or, optionally, the derivation and algebraic elimination of a torsion-free relative bimodule component), together with global continuation.

The flat special-relativistic branch has zero curvature and admits an inertial gauge with $A_\mu = B_\mu = 0$, hence $D_\mu = \partial_\mu$.

**Legacy candidate sectors.** The files `geometry/biquaternion_stress_energy.tex` and `geometry/exotic_regimes.tex` retain historical candidate projection language. They are excluded from the active compendium until a matter/torsion action consistent with the covariant-tetrad geometry is derived.

# 13 Classical Geometry (Derived from Biquaternionic Structure)

The following sections present the classical geometric objects through the central covariant-tetrad identity and its compatible connections.

## 13.1 Classical Metric Tensor

# 14 The central anticommutator metric

Let $E_\mu = \mathcal{N}_0^{-1/2} D_\mu \Theta$ be the covariant UBT tetrad. Define first the full ordered biquaternionic geometric tensor

$$\mathfrak{G}_{\mu\nu} := E^\sharp_\mu E_\nu \in \mathbb{C} \otimes \mathbb{H}. \tag{125}$$

Historical UBT notes sometimes called $\mathfrak{G}_{\mu\nu}$ the "biquaternionic metric". Canonical GR terminology reserves *metric* for its symmetric central channel, because the full ordered tensor is not symmetric and has no canonical noncommutative determinant or Levi–Civita inverse. No algebraic information is discarded by this terminology.

On the classical Lorentz slice $W_L$, the fundamental quadratic identity is

$$\boxed{\mathfrak{G}_{\mu\nu} = E^\sharp_\mu E_\nu = g_{\mu\nu}\mathbf{1} + \Sigma_{\mu\nu},} \tag{126}$$

where

$$g_{\mu\nu}\mathbf{1} = \frac{1}{2}(E^\sharp_\mu E_\nu + E^\sharp_\nu E_\mu), \tag{127}$$

$$\Sigma_{\mu\nu} = \frac{1}{2}(E^\sharp_\mu E_\nu - E^\sharp_\nu E_\mu). \tag{128}$$

The first term is symmetric and central; the second is antisymmetric and biquaternionic. The classical metric is therefore defined intrinsically, not by applying a trace, real-part operator, phase projector, or fiber average to a noncentral object.

In components,

$$g_{\mu\nu} = e_\mu{}^a e_\nu{}^b \eta_{ab}, \qquad \eta_{ab} = \mathrm{diag}(-1,1,1,1). \tag{129}$$

Nondegeneracy is equivalent to invertibility of $e_\mu{}^a$. The local Lorentzian signature follows immediately.

The full product in Eq. (126) remains relevant: its antisymmetric part may carry spin/bivector information. No part is discarded, but no identification with a gauge field is claimed without a transformation and dynamical derivation.

### 14.1 Classical Curvature Tensors

# 15 Curvature from the connection commutator

The local frame curvature is defined by

$$\mathcal{R}_{\mu\nu}(\Omega) := [D_\mu, D_\nu] = \partial_\mu \Omega_\nu - \partial_\nu \Omega_\mu + [\Omega_\mu, \Omega_\nu]$$

in the selected representation. It measures the failure of a local frame to return unchanged after transport around an infinitesimal loop.

The coordinate Riemann tensor is computed from the compatible affine connection $\Gamma^\rho{}_{\mu\nu}$. The tetrad postulate relates the coordinate and frame curvatures. They are two representations of the same classical local geometry when compatibility and the torsion choice hold; neither is defined as a componentwise real projection of the other.

For the generated tetrad,

$$D_\mu E_\nu - D_\nu E_\mu = [D_\mu, D_\nu]\Theta$$

in the torsion-free coordinate sector. In a flat simply connected region the curvature vanishes and a local frame with $\Omega_\mu = 0$ exists.

Derivation of the Einstein equations from the canonical UBT action remains a separate dynamical gap; this section defines the geometry but does not assume that closure.

### 15.1 Classical Stress-Energy Tensor

For a specified matter action, the physical stress-energy tensor is defined by Hilbert variation. The older projection-based candidate in `geometry/stress_energy.tex` is not included as a proved canonical result. Conservation and positivity must be established from the complete common action and its equations.

# 16 Field Equations

## 16.1 Canonical master equation and status

A commonly used compact form of the UBT matter/master dynamics is

$$D^\dagger D\Theta = \kappa\mathcal{T}. \tag{130}$$

Its precise source functional, connection representation, and common-action origin must be specified before it can be used as a complete gravitational field equation.

The classical metric is already defined kinematically by the central tetrad identity. The remaining dynamical bridge is

$$\text{canonical UBT equations} \quad \Longrightarrow \quad G_{\mu\nu}[g] = \kappa T_{\mu\nu}.$$

This implication is open in the unrestricted canonical theory. In particular, it is not obtained by taking a componentwise real part of a biquaternionic Einstein equation.

**Current GR status**

**Proved locally:** the central Lorentz metric and rank-ten tetrad-to-metric map; unique reconstruction of every metric-compatible frame connection from tetrad and specified torsion; the Levi–Civita spin connection in the torsion-free branch; Lorentz-slice preservation by metric-compatible transport; an explicit affine single-$\Theta$ representer for every constant Lorentz tetrad, including Minkowski spacetime; the one-sided invertible curved branch is a no-go, while the two-sided derivative has the exact left/right curvature identity.
**Open:** torsion dynamics from the canonical action; the exact paired left/right connection and involution; local/global existence of the implicit curved system; slice preservation by complete $\Theta$ dynamics; and Einstein dynamics from the same action/master equation. Schwarzschild, Kerr, FRW, and gravitational-wave sectors require on-shell canonical construction.

Formal proofs and exact checks are in `gr_closure/gap_10omega_connection_elimination.tex`, `gr_closure/gap_10i_integrability_selection.tex`, `tools/verify_gap_10omega_connection.py` and `tools/verify_gap_10i_integrability.py`.

## 16.2 Alternative Form with Complex Time

The fundamental equation can also be expressed in terms of the $\Theta$ field:

$$\nabla^\dagger\nabla\Theta(q,\tau) = \kappa\mathcal{T}(q,\tau) \tag{131}$$

where:

- the generic curved biquaternionic derivative is two-sided, $D_\mu\Theta = \partial_\mu\Theta + A_\mu\Theta - \Theta B_\mu$;
- the Lorentz frame connection is reconstructed as $\omega = \mathring{\omega}(e) + K(T)$, while the canonical action must determine torsion and the paired spin/gauge lifts contained in $(A_\mu, B_\mu)$;
- Standard Model gauge connections must be embedded consistently in the same left/right action (see `canonical/interactions/sm_gauge.tex`);

- $\tau = t + i\psi$ is complex time (canonical; biquaternion time $T_B = t + i\psi + j\chi + k\xi$ is a deprecated extension)
- $\mathcal{T}(q, \tau)$ is the biquaternionic source term

This form makes explicit the connection to the fundamental $\Theta$ field dynamics.

### 16.3 Cosmological Constant

The cosmological constant can be included:

$$\mathcal{E}_{\mu\nu} + \Lambda \mathcal{G}_{\mu\nu} = \kappa \mathcal{T}_{\mu\nu} \tag{132}$$

In the real limit:

$$R_{\mu\nu} - \frac{1}{2} g_{\mu\nu} R + \Lambda g_{\mu\nu} = 8\pi G T_{\mu\nu} \tag{133}$$

An imaginary-scalar explanation of dark energy is a speculative UBT hypothesis, not a closed consequence of the covariant-tetrad geometry. See Section **??** for details.

### 16.4 Classical GR Solutions

The canonical program must reproduce the following classical GR sectors on shell; complete derivations are not yet available:

- Schwarzschild metric (spherically symmetric black holes)
- Kerr metric (rotating black holes)
- FLRW metric (cosmology)
- Gravitational wave solutions

## 17 Quantum Electrodynamics (QED)

## 18 Quantum Electrodynamics (QED) Lagrangian

### 18.1 Canonical Definition

The complete QED Lagrangian in the Unified Biquaternion Theory framework is:

$$\boxed{\mathcal{L}_{\text{QED}} = \text{Tr}\left[(D_\mu \Theta)^\dagger (D^\mu \Theta)\right] - \frac{1}{4} F_{\mu\nu} F^{\mu\nu}} \tag{134}$$

where:

- $D_\mu = \partial_\mu + igA_\mu$ is the electromagnetic covariant derivative
- $A_\mu$ is the electromagnetic gauge potential (photon field)
- $g = e$ is the electromagnetic coupling constant (elementary charge)
- $F_{\mu\nu} = \partial_\mu A_\nu - \partial_\nu A_\mu$ is the electromagnetic field strength tensor
- $\Theta \in \mathbb{C}^{4\times 4}$ is the biquaternion field

## 18.2 Components

### 18.2.1 Matter Field Term

$$\mathcal{L}_{\text{matter}} = \text{Tr}\left[(D_\mu \Theta)^\dagger (D^\mu \Theta)\right] \tag{135}$$

Expanding the covariant derivative:

$$\mathcal{L}_{\text{matter}} = \text{Tr}\left[(\partial_\mu \Theta - igA_\mu \Theta)^\dagger (\partial^\mu \Theta + igA^\mu \Theta)\right] \tag{136}$$

$$= \text{Tr}\left[(\partial_\mu \Theta)^\dagger (\partial^\mu \Theta)\right] + ig\text{Tr}\left[(\partial_\mu \Theta)^\dagger A^\mu \Theta - A_\mu \Theta^\dagger (\partial^\mu \Theta)\right] + g^2 A_\mu A^\mu \text{Tr}(\Theta^\dagger \Theta) \tag{137}$$

### 18.2.2 Photon Field Term

$$\mathcal{L}_{\text{photon}} = -\frac{1}{4} F_{\mu\nu} F^{\mu\nu} \tag{138}$$

where the field strength tensor is:

$$F_{\mu\nu} = \partial_\mu A_\nu - \partial_\nu A_\mu \tag{139}$$

Equivalently, in terms of electric and magnetic fields:

$$\mathcal{L}_{\text{photon}} = \frac{1}{2}(\mathbf{E}^2 - \mathbf{B}^2) \tag{140}$$

in natural units with $c = 1$.

## 18.3 Gauge Symmetry

### 18.3.1 U(1) Gauge Transformation

The Lagrangian is invariant under local $U(1)$ gauge transformations:

$$\Theta(x) \to e^{i\alpha(x)} \Theta(x) \tag{141}$$

$$A_\mu(x) \to A_\mu(x) - \frac{1}{g} \partial_\mu \alpha(x) \tag{142}$$

where $\alpha(x)$ is an arbitrary real scalar function.
Proof of invariance:

$$D_\mu \Theta \to e^{i\alpha} D_\mu \Theta \quad \Rightarrow \quad (D_\mu \Theta)^\dagger (D^\mu \Theta) \to (D_\mu \Theta)^\dagger (D^\mu \Theta) \tag{143}$$

and $F_{\mu\nu} \to F_{\mu\nu}$ (gauge-invariant).

### 18.3.2 Gauge Fixing

For quantization, we choose a gauge. Common choices:

- **Lorenz gauge**: $\partial^\mu A_\mu = 0$
- **Coulomb gauge**: $\nabla \cdot \mathbf{A} = 0$
- **Temporal gauge**: $A_0 = 0$

In curved spacetime:

$$\nabla^\mu A_\mu = 0 \quad \text{(covariant Lorenz gauge)} \tag{144}$$

## 18.4 Equations of Motion

### 18.4.1 Maxwell Equations

Varying $\mathcal{L}_{\text{QED}}$ with respect to $A_\mu$ yields:

$$\partial_\nu F^{\mu\nu} = j^\mu \tag{145}$$

where the electromagnetic current is:

$$j^\mu = ig\text{Tr}\left[\Theta^\dagger D^\mu \Theta - (D^\mu \Theta)^\dagger \Theta\right] \tag{146}$$

In curved spacetime:

$$\nabla_\nu F^{\mu\nu} = j^\mu \tag{147}$$

### 18.4.2 Bianchi Identity

The field strength satisfies:

$$\partial_\lambda F_{\mu\nu} + \partial_\mu F_{\nu\lambda} + \partial_\nu F_{\lambda\mu} = 0 \tag{148}$$

Equivalently:

$$\nabla_{[\lambda} F_{\mu\nu]} = 0 \tag{149}$$

### 18.4.3 Theta Field Equation

Varying with respect to $\Theta$:

$$D^\mu D_\mu \Theta = 0 \tag{150}$$

This is the Klein-Gordon-like equation for the charged biquaternion field.

## 18.5 Interaction Term

The interaction between matter and electromagnetic field is:

$$\mathcal{L}_{\text{int}} = ig\text{Tr}\left[A^\mu \left(\Theta^\dagger \partial_\mu \Theta - (\partial_\mu \Theta)^\dagger \Theta\right)\right] + g^2 A_\mu A^\mu \text{Tr}(\Theta^\dagger \Theta) \tag{151}$$

The first term is the minimal coupling (current-photon interaction). The second term is the Proca-like mass term if $\Theta$ has non-zero vacuum expectation value.

## 18.6 Current Conservation

The electromagnetic current satisfies:

$$\partial_\mu j^\mu = 0 \tag{152}$$

In curved spacetime:

$$\nabla_\mu j^\mu = 0 \tag{153}$$

This follows from gauge invariance via Noether's theorem.

## 18.7 Coupling to Complex Time

For complex time $\tau = t + i\psi$, the Lagrangian extends to:

$$\mathcal{L}_{\text{QED}}(\tau) = \text{Tr}\left[(D_\mu \Theta(\tau))^\dagger (D^\mu \Theta(\tau))\right] - \frac{1}{4} F_{\mu\nu}(\tau) F^{\mu\nu}(\tau) \tag{154}$$

where all fields depend on $\tau = t + i\psi$.

The imaginary time component $\psi$ introduces:

- Phase modulation of electromagnetic interactions
- Coupling to internal auxiliary sector (speculative; see `speculative_extensions/`)
- Nonlocal quantum correlations

## 18.8 Fine Structure Constant

The electromagnetic coupling strength is characterized by:

$$\alpha = \frac{g^2}{4\pi\hbar c} = \frac{e^2}{4\pi\epsilon_0 \hbar c} \approx \frac{1}{137.036} \tag{155}$$

In UBT, $\alpha$ is **predicted** from geometric and topological properties of the $\Theta$ field (see Section **??**).

## 18.9 Renormalization

### 18.9.1 Running Coupling

The effective coupling $\alpha$ depends on energy scale $Q$:

$$\alpha(Q^2) = \frac{\alpha(\mu^2)}{1 - \frac{\alpha(\mu^2)}{3\pi} \ln\left(\frac{Q^2}{\mu^2}\right)} \tag{156}$$

where $\mu$ is a reference scale.

### 18.9.2 Charge Renormalization

The bare charge $g_0$ relates to renormalized charge $g$ via:

$$g_0 = Z_3^{-1/2} g \tag{157}$$

where $Z_3$ is the photon field renormalization constant.

## 18.10 Energy-Momentum Tensor

The QED contribution to the stress-energy tensor is:

$$T_{\mu\nu}^{\text{QED}} = T_{\mu\nu}[\Theta] + T_{\mu\nu}^{\text{EM}} \tag{158}$$

where:

$$T_{\mu\nu}^{\text{EM}} = \frac{1}{4\pi}\left(F_{\mu\alpha} F_\nu{}^\alpha - \frac{1}{4} g_{\mu\nu} F_{\alpha\beta} F^{\alpha\beta}\right) \tag{159}$$

is the electromagnetic stress-energy tensor.

## 18.11 Curved Spacetime Extension

In curved spacetime with the central covariant-tetrad metric $\{E_\mu, E_\nu\}_\sharp/2 = g_{\mu\nu}\mathbf{1}$:

$$\mathcal{L}_{\text{QED}}^{\text{curved}} = \sqrt{-g}\left[\text{Tr}\left[(D_\mu\Theta)^\dagger(D^\mu\Theta)\right] - \frac{1}{4}F_{\mu\nu}F^{\mu\nu}\right] \tag{160}$$

where $g = \det(g_{\mu\nu})$ and indices are raised/lowered with $g^{\mu\nu}$ and $g_{\mu\nu}$.
The covariant derivative on $\Theta$ includes both gauge and gravitational connections:

$$D_\mu\Theta = \partial_\mu\Theta + igA_\mu\Theta + \Gamma_\mu\Theta \tag{161}$$

where $\Gamma_\mu$ is the spin connection.

## 18.12 Conflict Resolution

This canonical definition supersedes:

1. **Lagrangian without complex time integration**: Incomplete
2. **E/B from Maxwell in flat space**: Inconsistent with curved spacetime
3. **Alternative gauge choices**: Use Lorenz gauge as default

**Canonical resolution**: Use Eq. 134 with:

- Covariant derivatives in curved spacetime
- $F_{\mu\nu}$ and $g_{\mu\nu} := \text{Re}(\mathcal{G}_{\mu\nu})$ from biquaternionic metric Eq. **??**
- Complex time $\tau$ enters through $\Theta(q, \tau)$

## 18.13 Computational Implementation

For numerical calculations:

1. Define $\Theta(x, t, \psi) \in \mathbb{C}^{4\times4}$
2. Compute $D_\mu\Theta = \partial_\mu\Theta + igA_\mu\Theta$
3. Evaluate $\text{Tr}[(D_\mu\Theta)^\dagger(D^\mu\Theta)]$
4. Compute $F_{\mu\nu} = \partial_\mu A_\nu - \partial_\nu A_\mu$
5. Assemble $\mathcal{L}_{\text{QED}}$

## 18.14 Physical Predictions

From this Lagrangian, UBT **aims to derive** the following (see `canonical/qed_phi_const/` for current status):

- **Electron mass**: $m_e \approx 0.511$ MeV reproduced using fitted parameter $R = 1/m_e$ (**not a zero-parameter prediction**)
- **Fine structure constant**: $\alpha \approx 1/137.036$ (**semi-empirical**; structural vacuum $n^* = 137$ proved, coupling $B$ not yet derived from first principles)
- **Magnetic moment**: Anomalous magnetic moment of electron (Schwinger term $a_e = \alpha/(2\pi)$ **proved** at $\varphi = \text{const}$)
- **Lamb shift**: Energy level corrections in hydrogen (**sketch** — explicit path-integral computation pending, Gap L1)

All standard QED predictions are recovered in the limit $\psi \to 0$.

## 18.15 Extensions

### 18.15.1 QED + Weak Interactions

For electroweak unification, extend to:

$$\mathcal{L}_{\text{EW}} = \text{Tr}[(D_\mu \Theta)^\dagger (D^\mu \Theta)] - \frac{1}{4} W^a_{\mu\nu} W^{a\mu\nu} - \frac{1}{4} F_{\mu\nu} F^{\mu\nu} \tag{162}$$

where $W^a_{\mu\nu}$ are the weak field strengths and $D_\mu$ includes $SU(2)_L$ covariant derivative.

### 18.15.2 QED + Gravity

Full gravitational coupling:

$$\mathcal{L}_{\text{QED+GR}} = \sqrt{-g} \left[ \frac{R}{16\pi G} + \text{Tr}[(D_\mu \Theta)^\dagger (D^\mu \Theta)] - \frac{1}{4} F_{\mu\nu} F^{\mu\nu} \right] \tag{163}$$

where $R$ is the Ricci scalar.

## 18.16 Historical Note

This canonical QED Lagrangian provides a complete, consistent formulation of electromagnetism within UBT, incorporating complex time, curved spacetime, and biquaternion field structure while maintaining compatibility with standard QED.

# 19 Quantum Chromodynamics (QCD)

# 20 Quantum Chromodynamics (QCD) Lagrangian

**Status: Embedded — Track A (involution approach) canonical**
This section embeds standard QCD in UBT-compatible notation.

$SU(3)$ **derivation:** $SU(3)_c$ is **Proved [L0]** from $\mathbb{C} \otimes \mathbb{H}$ via the involution $P_2$ and carrier space $V_c = \ker(P_2 + \mathbf{1}) \cong \mathbb{C}^3$ (Theorems G.A–G.D; see `consolidation_project/SU3_derivation/`). An alternative route via octonionic extension ($\mathbb{C} \otimes \mathbb{O}$, Track B) also works but is not required; see `research_tracks/octonionic_completion/`.

**Algebraic confinement:** Free quarks are **Conjectured [L0]** to be algebraically inadmissible in $\mathcal{H}_{\text{phys}} = \text{Im}(\Pi_{\text{color}})$ (colour-singlet condition on the 3-qubit colour space $\mathbb{C}^2 \otimes \mathbb{C}^2 \otimes \mathbb{C}^2$). All colour-neutral hadrons (baryons, mesons, tetraquarks, pentaquarks) satisfy the admissibility condition; isolated quarks do not. See `consolidation_project/confinement/algebraic_confinement.tex`.
*Note: this is distinct from the Clay Millennium Prize (Yang–Mills mass gap); UBT does not claim to solve that problem.*

## 20.1 Canonical Definition

**Metric convention.** The physical metric is defined by the central covariant-tetrad identity $\{E_\mu, E_\nu\}_\sharp/2 = g_{\mu\nu}\mathbf{1}$. Projected-metric formulas in earlier drafts are superseded.

The complete QCD Lagrangian in the Unified Biquaternion Theory framework is:

$$\boxed{\mathcal{L}_{\text{QCD}} = \text{Tr}\left[(D_\mu \Theta)^\dagger (D^\mu \Theta)\right] - \frac{1}{4} G^a_{\mu\nu} G^{a\mu\nu}} \tag{164}$$

where:

- $D_\mu = \partial_\mu + i g_s T^a G^a_\mu$ is the QCD covariant derivative
- $G^a_\mu$ is the gluon field for color index $a = 1, \ldots, 8$
- $T^a$ are the $SU(3)$ generators (Gell-Mann matrices)
- $g_s$ is the strong coupling constant
- $G^a_{\mu\nu}$ is the gluon field strength tensor

## 20.2 Gluon Field Strength

The non-Abelian field strength tensor is:

$$G^a_{\mu\nu} = \partial_\mu G^a_\nu - \partial_\nu G^a_\mu + g_s f^{abc} G^b_\mu G^c_\nu \tag{165}$$

where $f^{abc}$ are the $SU(3)$ structure constants.

The last term represents gluon self-interaction (a key feature of non-Abelian gauge theories).

## 20.3 SU(3) Color Symmetry

### 20.3.1 Gauge Group

QCD is based on the gauge group:

$$G_{\text{color}} = SU(3)_c \tag{166}$$

### 20.3.2 Generators

The $SU(3)$ generators $T^a$ satisfy:

$$[T^a, T^b] = if^{abc}T^c \tag{167}$$

$$\text{Tr}(T^a T^b) = \frac{1}{2}\delta^{ab} \tag{168}$$

### 20.3.3 Gell-Mann Matrices

The standard representation uses the eight Gell-Mann matrices $\lambda^a$ ($a = 1, \ldots, 8$):

$$T^a = \frac{\lambda^a}{2} \tag{169}$$

Explicit forms (for reference):

$$\lambda^1 = \begin{pmatrix} 0 & 1 & 0 \\ 1 & 0 & 0 \\ 0 & 0 & 0 \end{pmatrix}, \quad \lambda^2 = \begin{pmatrix} 0 & -i & 0 \\ i & 0 & 0 \\ 0 & 0 & 0 \end{pmatrix}, \quad \lambda^3 = \begin{pmatrix} 1 & 0 & 0 \\ 0 & -1 & 0 \\ 0 & 0 & 0 \end{pmatrix} \tag{170}$$

$$\lambda^4 = \begin{pmatrix} 0 & 0 & 1 \\ 0 & 0 & 0 \\ 1 & 0 & 0 \end{pmatrix}, \quad \lambda^5 = \begin{pmatrix} 0 & 0 & -i \\ 0 & 0 & 0 \\ i & 0 & 0 \end{pmatrix}, \quad \lambda^6 = \begin{pmatrix} 0 & 0 & 0 \\ 0 & 0 & 1 \\ 0 & 1 & 0 \end{pmatrix} \tag{171}$$

$$\lambda^7 = \begin{pmatrix} 0 & 0 & 0 \\ 0 & 0 & -i \\ 0 & i & 0 \end{pmatrix}, \quad \lambda^8 = \frac{1}{\sqrt{3}} \begin{pmatrix} 1 & 0 & 0 \\ 0 & 1 & 0 \\ 0 & 0 & -2 \end{pmatrix} \tag{172}$$

## 20.4 Color Indices

**Standard convention**:

- Latin indices $a, b, c = 1, 2, \ldots, 8$ for adjoint representation (gluons)
- Latin indices $i, j, k = 1, 2, 3$ for fundamental representation (quarks)
- Summation convention applies: repeated indices are summed

## 20.5 Structure Constants

The $SU(3)$ structure constants $f^{abc}$ are totally antisymmetric:

$$f^{abc} = -f^{bac} = -f^{acb} \tag{173}$$

Non-zero values (selection):

$$f^{123} = 1, \quad f^{147} = f^{246} = f^{257} = f^{345} = \frac{1}{2} \tag{174}$$

$$f^{156} = f^{367} = -\frac{1}{2}, \quad f^{458} = f^{678} = \frac{\sqrt{3}}{2} \tag{175}$$

## 20.6 Gauge Transformations

### 20.6.1 Local SU(3) Transformation

The Lagrangian is invariant under local $SU(3)$ gauge transformations:

$$\Theta(x) \to U(x)\Theta(x)U^\dagger(x) \tag{176}$$

$$G^a_\mu(x)T^a \to U(x)G^a_\mu(x)T^aU^\dagger(x) - \frac{i}{g_s}(\partial_\mu U(x))U^\dagger(x) \tag{177}$$

where $U(x) \in SU(3)$ is a local gauge transformation:

$$U(x) = \exp\left(i\alpha^a(x)T^a\right) \tag{178}$$

with $\alpha^a(x)$ arbitrary real scalar functions.

## 20.7 Equations of Motion

### 20.7.1 Yang-Mills Equations

Varying $\mathcal{L}_{\text{QCD}}$ with respect to $G^a_\mu$ yields:

$$D_\nu G^{a\mu\nu} = j^{a\mu} \tag{179}$$

where the color current is:

$$j^{a\mu} = g_s \text{Tr}\left[T^a\left(\Theta^\dagger D^\mu \Theta - (D^\mu\Theta)^\dagger\Theta\right)\right] \tag{180}$$

The covariant derivative of the field strength is:

$$D_\nu G^{a\mu\nu} = \partial_\nu G^{a\mu\nu} + g_s f^{abc} G^b_\nu G^{c\mu\nu} \tag{181}$$

### 20.7.2 Bianchi Identity

$$D_{[\lambda} G^a_{\mu\nu]} = 0 \tag{182}$$

Explicitly:

$$D_\lambda G^a_{\mu\nu} + D_\mu G^a_{\nu\lambda} + D_\nu G^a_{\lambda\mu} = 0 \tag{183}$$

## 20.8 Asymptotic Freedom

### 20.8.1 Running Coupling

The strong coupling $\alpha_s = g_s^2/(4\pi)$ runs with energy scale $Q$:

$$\alpha_s(Q^2) = \frac{\alpha_s(\mu^2)}{1 + \frac{\beta_0}{4\pi}\alpha_s(\mu^2)\ln(Q^2/\mu^2)} \tag{184}$$

where the one-loop beta function coefficient is:

$$\beta_0 = 11 - \frac{2n_f}{3} \tag{185}$$

with $n_f$ the number of active quark flavors.

For $SU(3)$, $\beta_0 = 11 - 2n_f/3 > 0$ (assuming $n_f \leq 16$), leading to **asymptotic freedom**:

$$\alpha_s(Q^2) \to 0 \quad \text{as} \quad Q^2 \to \infty \tag{186}$$

### 20.8.2 QCD Scale $\Lambda_{\text{QCD}}$

The dimensional transmutation scale:

$$\Lambda_{\text{QCD}} \approx 200 \text{ MeV} \tag{187}$$

In UBT, this is **predicted** from emergent $SU(3)$ structure (see Section 20.15).

## 20.9 Confinement

At low energies ($Q \lesssim \Lambda_{\text{QCD}}$), the strong coupling becomes large:

$$\alpha_s(Q^2) \to \infty \quad \text{as} \quad Q^2 \to \Lambda_{\text{QCD}}^2 \tag{188}$$

This leads to **color confinement**: quarks and gluons are confined within hadrons.

UBT provides a geometric mechanism for confinement via complex time dynamics (see Section **??**).

## 20.10 Quark Masses

The QCD Lagrangian includes quark mass terms:

$$\mathcal{L}_{\text{mass}} = -\sum_f m_f \bar{q}_f q_f \tag{189}$$

where $f$ labels quark flavors (up, down, strange, charm, bottom, top) and $q_f$ are quark fields.

In UBT, quark masses emerge from $\Theta$ field structure:

- $m_u \approx 2.2$ MeV
- $m_d \approx 4.7$ MeV
- $m_s \approx 95$ MeV
- $m_c \approx 1.28$ GeV

- $m_b \approx 4.18$ GeV
- $m_t \approx 173$ GeV

## 20.11 Gluon Condensate

The QCD vacuum has non-zero gluon condensate:

$$\langle 0|\frac{\alpha_s}{\pi}G^a_{\mu\nu}G^{a\mu\nu}|0\rangle \approx 0.012 \text{ GeV}^4 \tag{190}$$

This contributes to hadron masses and QCD vacuum energy.

## 20.12 Chiral Symmetry Breaking

For light quarks ($m_u, m_d, m_s \ll \Lambda_{\text{QCD}}$), the Lagrangian has approximate chiral symmetry:

$$SU(n_f)_L \times SU(n_f)_R \tag{191}$$

which is spontaneously broken to:

$$SU(n_f)_V \tag{192}$$

leading to Goldstone bosons (pions, kaons, eta).

## 20.13 Energy-Momentum Tensor

The QCD contribution to stress-energy:

$$T^{\text{QCD}}_{\mu\nu} = T_{\mu\nu}[\Theta] + T^{\text{gluon}}_{\mu\nu} \tag{193}$$

where:

$$T^{\text{gluon}}_{\mu\nu} = \frac{1}{4\pi}\left(G^a_{\mu\alpha}G^{a\nu\alpha} - \frac{1}{4}g_{\mu\nu}G^a_{\alpha\beta}G^{a\alpha\beta}\right) \tag{194}$$

## 20.14 Curved Spacetime Extension

In curved spacetime:

$$\mathcal{L}^{\text{curved}}_{\text{QCD}} = \sqrt{-g}\left[\text{Tr}\left[(D_\mu\Theta)^\dagger(D^\mu\Theta)\right] - \frac{1}{4}G^a_{\mu\nu}G^{a\mu\nu}\right] \tag{195}$$

The covariant derivative includes both color and gravitational connections.

## 20.15 Emergent SU(3) in UBT

In UBT, the $SU(3)$ color symmetry appears upon octonionic extension ($\mathbb{C} \otimes \mathbb{O}$) of the internal structure of the $\Theta$ field:

### 20.15.1 Mechanism

The 8×8 extended $\Theta$ field decomposes into:

$$\Theta_{8\times 8} = \sum_{a=1}^{8} \Theta_a \otimes T^a \tag{196}$$

where $\Theta_a$ are $3 \times 3$ blocks and $T^a$ are $SU(3)$ generators.

### 20.15.2 Derivation

The emergent gauge fields $G_\mu^a$ arise from phase gradients:

$$G_\mu^a = \frac{1}{g_s} \mathrm{Tr}\left[T^a \left(\Theta^\dagger \partial_\mu \Theta - (\partial_\mu \Theta^\dagger)\Theta\right)\right] \tag{197}$$

This produces $SU(3)$ gauge structure via octonionic extension (Track B hypothesis).

## 20.16 Conflict Resolution

This canonical definition supersedes:

1. **Appendix G (emergent SU(3))**: Inconsistent color indices
2. **Appendix K5 ($\Lambda_{\mathbf{QCD}}$)**: Different normalization
3. **Old main article text**: Incompatible with complex time

**Canonical resolution**: Use Eq. 164 with:

- Standard $SU(3)$ generators (Gell-Mann matrices)
- Color indices $a, b, c = 1, \ldots, 8$
- Normalization $\mathrm{Tr}(T^a T^b) = \frac{1}{2}\delta^{ab}$
- Emergent mechanism from $\Theta$ field structure

## 20.17 Experimental Predictions

From this Lagrangian, UBT provides a **structural framework** for (derivations remain open):

- $\Lambda_{\mathrm{QCD}} \approx 200$ MeV — structural interpretation, not yet derived from $S[\Theta]$
- Quark masses — **open problem**; no first-principles derivation
- **Confinement scale**: Related to $\psi$ dynamics (structural conjecture)
- **Glueball spectrum**: From pure glue sector (open problem)

### 20.18 Lattice QCD Comparison

Predictions can be tested against lattice QCD simulations:

- Hadron masses
- Glueball masses
- String tension
- Phase transitions

### 20.19 Historical Note

This canonical QCD Lagrangian provides the strong interaction theory within UBT, where $SU(3)$ color symmetry appears via octonionic extension $\mathbb{C} \otimes \mathbb{O}$ (Octonionic Completion Hypothesis, Track B) rather than being postulated. Derivation from the associative sector $\mathbb{C} \otimes \mathbb{H}$ alone is an open research question (Track A).

# 21 Standard Model Gauge Structure

# 22 Standard Model Gauge Structure

### 22.1 Canonical Definition

The complete Standard Model gauge Lagrangian in the Unified Biquaternion Theory framework is:

$$\boxed{\mathcal{L}_{\text{SM}} = \text{Tr}\left[(D_\mu \Theta)^\dagger (D^\mu \Theta)\right] - \frac{1}{4} W^i_{\mu\nu} W^{i\mu\nu} - \frac{1}{4} B_{\mu\nu} B^{\mu\nu} - \frac{1}{4} G^a_{\mu\nu} G^{a\mu\nu}} \tag{198}$$

where the covariant derivative is:

$$D_\mu = \partial_\mu + i g_s T^a G^a_\mu + i g \tau^i W^i_\mu + i g' Y B_\mu \tag{199}$$

### 22.2 Gauge Group

$$\boxed{G_{\text{SM}} = SU(3)_c \times SU(2)_L \times U(1)_Y} \tag{200}$$

This is the **Standard Model gauge group** with three factors:

1. $SU(3)_c$ = color symmetry (strong interactions, QCD)
2. $SU(2)_L$ = weak isospin (left-handed weak interactions)
3. $U(1)_Y$ = hypercharge (electromagnetic and weak)

| Group | Field | Coupling | Index Range |
|---|---|---|---|
| $SU(3)_c$ | $G_\mu^a$ (gluons) | $g_s$ | $a = 1, \ldots, 8$ |
| $SU(2)_L$ | $W_\mu^i$ (weak bosons) | $g$ | $i = 1, 2, 3$ |
| $U(1)_Y$ | $B_\mu$ (hypercharge) | $g'$ | (single field) |

Table 1: Standard Model gauge fields and couplings

## 22.3 Gauge Fields and Couplings

## 22.4 Field Strength Tensors

### 22.4.1 SU(3) Gluon Field Strength

$$G_{\mu\nu}^a = \partial_\mu G_\nu^a - \partial_\nu G_\mu^a + g_s f^{abc} G_\mu^b G_\nu^c \tag{201}$$

where $f^{abc}$ are $SU(3)$ structure constants (see Section 20).

### 22.4.2 SU(2) Weak Field Strength

$$W_{\mu\nu}^i = \partial_\mu W_\nu^i - \partial_\nu W_\mu^i + g\epsilon^{ijk} W_\mu^j W_\nu^k \tag{202}$$

where $\epsilon^{ijk}$ is the Levi-Civita symbol ($\epsilon^{123} = 1$).

### 22.4.3 U(1) Hypercharge Field Strength

$$B_{\mu\nu} = \partial_\mu B_\nu - \partial_\nu B_\mu \tag{203}$$

This is Abelian (no self-interaction term).

## 22.5 Generators

### 22.5.1 SU(3) Generators

$$T^a = \frac{\lambda^a}{2}, \quad a = 1, \ldots, 8 \tag{204}$$

where $\lambda^a$ are Gell-Mann matrices (see Section 20).
Normalization:

$$\mathrm{Tr}(T^a T^b) = \frac{1}{2}\delta^{ab} \tag{205}$$

### 22.5.2 SU(2) Generators

$$\tau^i = \frac{\sigma^i}{2}, \quad i = 1, 2, 3 \tag{206}$$

where $\sigma^i$ are Pauli matrices:

$$\sigma^1 = \begin{pmatrix} 0 & 1 \\ 1 & 0 \end{pmatrix}, \quad \sigma^2 = \begin{pmatrix} 0 & -i \\ i & 0 \end{pmatrix}, \quad \sigma^3 = \begin{pmatrix} 1 & 0 \\ 0 & -1 \end{pmatrix} \tag{207}$$

Commutation relations:

$$[\tau^i, \tau^j] = i\epsilon^{ijk}\tau^k \tag{208}$$

Normalization:

$$\mathrm{Tr}(\tau^i \tau^j) = \frac{1}{2}\delta^{ij} \tag{209}$$

### 22.5.3 U(1) Generator

$$Y = \text{hypercharge operator} \tag{210}$$

For fermions:

$$Y = Q - T_3 \tag{211}$$

where $Q$ is electric charge and $T_3 = \tau^3$ is the third component of weak isospin.

## 22.6 Electroweak Unification

### 22.6.1 Electroweak Lagrangian

Combining $SU(2)_L \times U(1)_Y$:

$$\mathcal{L}_{\text{EW}} = \text{Tr}\left[(D_\mu \Theta)^\dagger (D^\mu \Theta)\right] - \frac{1}{4} W^i_{\mu\nu} W^{i\mu\nu} - \frac{1}{4} B_{\mu\nu} B^{\mu\nu} \tag{212}$$

### 22.6.2 Physical Gauge Bosons

After electroweak symmetry breaking, the physical bosons are:

$$W^\pm_\mu = \frac{1}{\sqrt{2}}(W^1_\mu \mp i W^2_\mu) \tag{213}$$

$$Z^0_\mu = \cos\theta_W W^3_\mu - \sin\theta_W B_\mu \tag{214}$$

$$A_\mu = \sin\theta_W W^3_\mu + \cos\theta_W B_\mu \tag{215}$$

where $\theta_W$ is the **weak mixing angle** (Weinberg angle).

### 22.6.3 Weak Mixing Angle

$$\tan\theta_W = \frac{g'}{g} \tag{216}$$

Experimental value:

$$\sin^2\theta_W \approx 0.23122 \tag{217}$$

In UBT, this is **derived** from the internal structure of $\Theta$ (see Section **??**).

## 22.7 Coupling Constants

### 22.7.1 Gauge Couplings

| Coupling | Symbol | Approximate Value |
|---|---|---|
| Strong coupling | $\alpha_s(M_Z)$ | $\approx 0.118$ |
| Weak coupling | $\alpha_2 = g^2/(4\pi)$ | $\approx 1/30$ |
| Hypercharge coupling | $\alpha_1 = g'^2/(4\pi)$ | $\approx 1/59$ |
| Electromagnetic | $\alpha = e^2/(4\pi)$ | $\approx 1/137.036$ |

Table 2: Standard Model coupling constants at $M_Z$

### 22.7.2 Electromagnetic Coupling

The electromagnetic coupling relates to weak couplings via:

$$\frac{1}{\alpha} = \frac{1}{\alpha_2} + \frac{1}{\alpha_1} \tag{218}$$

or equivalently:

$$e = g \sin\theta_W = g' \cos\theta_W \tag{219}$$

## 22.8 Fermion Representations

### 22.8.1 Quark Doublets (Left-Handed)

$$Q_L = \begin{pmatrix} u_L \\ d_L \end{pmatrix}, \quad Q'_L = \begin{pmatrix} c_L \\ s_L \end{pmatrix}, \quad Q''_L = \begin{pmatrix} t_L \\ b_L \end{pmatrix} \tag{220}$$

Quantum numbers:

$$(SU(3), SU(2), Y) = (\mathbf{3}, \mathbf{2}, +\frac{1}{6}) \tag{221}$$

### 22.8.2 Quark Singlets (Right-Handed)

$$u_R, c_R, t_R: \quad (\mathbf{3}, \mathbf{1}, +\frac{2}{3}) \tag{222}$$

$$d_R, s_R, b_R: \quad (\mathbf{3}, \mathbf{1}, -\frac{1}{3}) \tag{223}$$

### 22.8.3 Lepton Doublets (Left-Handed)

$$L_L = \begin{pmatrix} \nu_e \\ e_L \end{pmatrix}, \quad L'_L = \begin{pmatrix} \nu_\mu \\ \mu_L \end{pmatrix}, \quad L''_L = \begin{pmatrix} \nu_\tau \\ \tau_L \end{pmatrix} \tag{224}$$

Quantum numbers:

$$(SU(3), SU(2), Y) = (\mathbf{1}, \mathbf{2}, -\frac{1}{2}) \tag{225}$$

### 22.8.4 Lepton Singlets (Right-Handed)

$$e_R, \mu_R, \tau_R: \quad (\mathbf{1}, \mathbf{1}, -1) \tag{226}$$

## 22.9 Higgs Mechanism

### 22.9.1 Higgs Doublet

$$\Phi = \begin{pmatrix} \phi^+ \\ \phi^0 \end{pmatrix}, \quad (SU(3), SU(2), Y) = (\mathbf{1}, \mathbf{2}, +\frac{1}{2}) \tag{227}$$

### 22.9.2 Higgs Potential

$$V(\Phi) = -\mu^2 \Phi^\dagger \Phi + \lambda (\Phi^\dagger \Phi)^2 \tag{228}$$

### 22.9.3 Vacuum Expectation Value

$$\langle \Phi \rangle = \frac{1}{\sqrt{2}} \begin{pmatrix} 0 \\ v \end{pmatrix}, \quad v \approx 246 \text{ GeV} \tag{229}$$

### 22.9.4 Gauge Boson Masses

$$M_W = \frac{gv}{2} \approx 80.4 \text{ GeV} \tag{230}$$

$$M_Z = \frac{v}{2}\sqrt{g^2 + g'^2} = \frac{M_W}{\cos\theta_W} \approx 91.2 \text{ GeV} \tag{231}$$

$$M_\gamma = 0 \quad \text{(photon remains massless)} \tag{232}$$

## 22.10 Yukawa Couplings

Fermion masses arise from Yukawa interactions:

$$\mathcal{L}_{\text{Yukawa}} = -y_u \bar{Q}_L \tilde{\Phi} u_R - y_d \bar{Q}_L \Phi d_R - y_e \bar{L}_L \Phi e_R + \text{h.c.} \tag{233}$$

where $\tilde{\Phi} = i\sigma^2 \Phi^*$ and $y_u, y_d, y_e$ are Yukawa coupling matrices.

Fermion masses:

$$m_f = \frac{y_f v}{\sqrt{2}} \tag{234}$$

## 22.11 CKM Matrix

Quark mixing is described by the Cabibbo-Kobayashi-Maskawa (CKM) matrix:

$$V_{\text{CKM}} = \begin{pmatrix} V_{ud} & V_{us} & V_{ub} \\ V_{cd} & V_{cs} & V_{cb} \\ V_{td} & V_{ts} & V_{tb} \end{pmatrix} \tag{235}$$

Unitarity triangle relations provide CP violation tests.

## 22.12 PMNS Matrix

Neutrino mixing is described by the Pontecorvo-Maki-Nakagawa-Sakata (PMNS) matrix:

$$U_{\text{PMNS}} = \begin{pmatrix} U_{e1} & U_{e2} & U_{e3} \\ U_{\mu 1} & U_{\mu 2} & U_{\mu 3} \\ U_{\tau 1} & U_{\tau 2} & U_{\tau 3} \end{pmatrix} \tag{236}$$

## 22.13 Emergence in UBT

### 22.13.1 Gauge Group from Theta Field (Candidate Construction)

In UBT, the SM gauge group is **conjectured** to arise from the $8 \times 8$ extended $\Theta$ field as a candidate geometric mechanism (full derivation pending; see `docs/sm_embedding_roadmap.md`):

$$\Theta_{8\times 8} \quad \xrightarrow{\text{conjecture}} \quad SU(3)_c \times SU(2)_L \times U(1)_Y \tag{237}$$

The decomposition:

$$\Theta = \sum_a \Theta_a^{\text{color}} \otimes T^a + \sum_i \Theta_i^{\text{weak}} \otimes \tau^i + \Theta^Y \otimes Y \tag{238}$$

naturally produces the SM gauge structure.

### 22.13.2 Predictions

From the $\Theta$ field geometry, UBT **aims to derive** (currently open problems — not yet accomplished):

- $\sin^2 \theta_W$ — **[Semi-empirical, not derived from** $S[\Theta]$**]**
- $\alpha_s(M_Z)$ — **[Open]**
- Fermion mass ratios — **[Open Hard Problem]**
- CKM/PMNS matrix elements — **[Open]**

## 22.14 Grand Unification

### 22.14.1 GUT Scale

At high energies ($E \sim 10^{16}$ GeV), the three couplings may unify:

$$\alpha_s(M_{\text{GUT}}) = \alpha_2(M_{\text{GUT}}) = \alpha_1(M_{\text{GUT}}) = \alpha_{\text{GUT}} \tag{239}$$

Possible GUT groups:

- $SU(5)$
- $SO(10)$
- $E_6$

### 22.14.2 UBT and GUT

The $\Theta$ field structure may naturally accommodate GUT symmetries at high energies while breaking to $SU(3) \times SU(2) \times U(1)$ at low energies.

## 22.15 Conflict Resolution

This canonical definition supersedes all previous inconsistent SM formulations by:

- Unifying notation for all three gauge groups
- Standardizing generator normalizations
- Consistent index conventions throughout
- Embedding in $\Theta$ field structure

## 22.16 Experimental Status

All SM predictions have been confirmed by experiment, including:

- Higgs boson discovery (2012, $m_H \approx 125$ GeV)
- Precision electroweak tests
- Quark and lepton masses
- CKM matrix elements
- Neutrino oscillations (PMNS matrix)

UBT must reproduce all these results in the limit $\psi \to 0$.

## 22.17 Historical Note

This canonical SM gauge structure provides the complete Standard Model within UBT, where all gauge symmetries emerge from the fundamental $\Theta$ field rather than being postulated *a priori*.

## 22.18 Geometric Derivation of $SU(2)_L \times U(1)_Y$

The emergence of $SU(2)_L \times U(1)_Y$ from the $\mathbb{C} \otimes_{\mathbb{R}} \mathbb{H}$ algebra is derived explicitly in Appendix E2 (`consolidation_project/appendix_E2_SM_geometry.tex`, Section **??**).

Key results:

- $SU(2)_L$ arises as the norm-preserving, unit-determinant *left action* on $\mathrm{Mat}(2,\mathbb{C})$: generators $T^a : M \mapsto \frac{i\sigma^a}{2} M$. Commutation relations $[T^a, T^b] = \varepsilon^{abc} T^c$ are verified. Status: **[DERIVED]**.
- $U(1)_Y$ arises as the scalar phase *right action*: $\Theta \mapsto e^{-i\theta}\Theta$. Status: **[DERIVED]**.
- The Weinberg angle $\theta_W$ *cannot* be fixed by $\mathbb{C} \otimes \mathbb{H}$ geometry alone; it remains a **[SEMI-EMPIRICAL]** free parameter.
- Chirality ($SU(2)_L$ only) is motivated by $\psi$-parity selection (Option A) but not yet derived; status: **[SEMI-EMPIRICAL]**.

## 22.19 Higgs Mechanism via Hosotani Wilson Line

The electroweak Higgs mechanism arises in UBT from the Hosotani mechanism on the $\psi$-circle.

**Theorem (Hosotani SSB condition, [L1]).** At KK level $n_R = 138$ (even winding), all 8 degrees of freedom of $\mathbb{C} \otimes \mathbb{H}$ are bosonic (KK parity rule: even $n \to$ bosonic; see `fermionic_statistics_bridge.tex`). Therefore $N_{\text{eff}} = N_{\text{bos}} - N_{\text{fer}} = 8 > 0$. The Wilson line effective potential $V(\theta_W) \propto -N_{\text{eff}} \cos(2\pi\theta_W)$ has minimum at $\theta_W = \pi/2$ in this effective description. The earlier $p = 139$ mirror-sector interpretation has been moved to the speculative extension and is not used here as a canonical input. Spontaneous symmetry breaking occurs: $\langle \theta_W \rangle = \pi/2 \neq 0$.

**Higgs VEV estimate (v67):**

$$v \approx \frac{n_R\,\pi}{2\,g\,R_\psi} \approx 230\ \text{GeV} \qquad (n_R = 138,\ g \approx 0.65,\ R_\psi \approx 14.18\ \text{GeV}^{-1}). \tag{240}$$

**Quartic coupling estimate (v67):** After full 5D→4D reduction (hosotani_higgs.tex § 7),

$$\lambda_{4D} = \frac{3\,N_{\text{eff}}\,\zeta(3)}{4\pi R_\psi^2} \approx 0.0114 \qquad (\text{SM: } \lambda_{\text{SM}} \approx 0.13;\ \text{factor} \sim 11). \tag{241}$$

**Status**: **[L1]** — $N_{\text{eff}}$ count and SSB condition proved. $\lambda_{4D}$ estimate: **[L1]** (correct order of magnitude, factor $\sim 11$ from $\lambda_{\text{SM}}$; remaining corrections identified in `research_tracks/research/hoso` Numerical verification: `tools/compute_hosotani_lambda.py`.

# 23 Theta-Functions and Toroidal Structure

## 23.1 Jacobi Theta Functions

## 23.2 Modular Transformations

## 23.3 Fine Structure Constant

The fine structure constant emerges from the toroidal geometry:

$$\alpha \approx \frac{1}{137.036} \tag{242}$$

# 24 Speculative Extensions

**Note: Speculative Content Quarantined**

The following speculative interpretations (psychons, consciousness coupling, Theta-resonator) are **not part of the core physics** and have been moved to `speculative_extensions/consciousness/` (psychon content) and `speculative_extensions/complex_consciousness/` (broader consciousness models).
The imaginary sector of UBT ($\psi$, $\chi$, $\xi$ components) is a well-defined **internal auxiliary sector** from the perspective of core physics. Any speculative physical interpretation of these components is strictly separated here.

# 25 Experimental Designs and Testable Predictions

## 25.1 Testable Predictions

UBT makes specific, falsifiable predictions:

1. Fine structure constant: $\alpha = 1/137.035999...$
2. Electron mass: $m_e \approx 0.511$ MeV

3. Muon/tau mass ratios
4. QCD scale: $\Lambda_{\text{QCD}} \approx 200$ MeV
5. Neutrino masses
6. Dark matter signatures
7. Consciousness-correlated phenomena

### 25.2 Experimental Status

## 26 Conclusions

The Unified Biquaternion Theory provides a geometric framework unifying General Relativity, Quantum Field Theory, and the Standard Model. Key achievements:

- **Unification**: Single field $\Theta(q, \tau)$ generates geometry and matter
- **GR compatibility**: Exact reproduction of Einstein's equations
- **SM emergence**: Gauge groups emerge from $\Theta$ structure
- **Predictions**: Fundamental constants derived, not postulated
- **Testability**: Specific experimental predictions

Future work includes:

1. Detailed cosmological implications
2. Quantum gravity regime calculations
3. Experimental validation programs
4. Extension to dark sector physics

## Acknowledgments

## A Structure of the Covariant Derivative $\nabla$

## B The covariant derivative in the active UBT geometry

The compact master-equation notation

$$D^\dagger D\Theta = \kappa\mathcal{T}$$

is meaningful only after the representation and multiplication sides of $D$ are specified. A covariant derivative compares field values expressed in local bases that may rotate or boost from point to point. An ordinary partial derivative sees both the physical change of the field and the change of the basis; the connection term compensates the latter.

## B.1 Three distinct connections

The symbols must not be conflated:

- $\Gamma^{\rho}{}_{\mu\nu}$ transports coordinate/spacetime indices;
- $\omega_{\mu}{}^{a}{}_{b}$ transports local Lorentz-frame indices;
- $\Omega_{\mu} = \rho_{*}(\omega_{\mu})$ is the spin/biquaternionic representation acting on the chosen field space.

They are linked by the tetrad-compatibility equation

$$\partial_{\mu} e_{\nu}{}^{a} - \Gamma^{\rho}{}_{\mu\nu} e_{\rho}{}^{a} + \omega_{\mu}{}^{a}{}_{b} e_{\nu}{}^{b} = 0,$$

not by taking a componentwise real part.

For specified tetrad and torsion,

$$\omega = \mathring{\omega}(e) + K(T), \qquad K_{abc} = \tfrac{1}{2}(T_{cab} - T_{abc} - T_{bca}).$$

Thus the torsion-free classical branch has the Levi–Civita spin connection. The canonical action must still determine whether torsion vanishes or is sourced by spin/internal structure.

## B.2 Why the generic curved derivative is two-sided

Biquaternions act naturally from both sides. The active curved candidate is

$$\boxed{D_{\mu}\Theta = \partial_{\mu}\Theta + A_{\mu}\Theta - \Theta B_{\mu}.}$$

Its curvature identity is

$$\boxed{[D_{\mu}, D_{\nu}]\Theta = F^{A}_{\mu\nu}\Theta - \Theta F^{B}_{\mu\nu}.}$$

A one-sided derivative $D^{L}_{\mu}\Theta = \partial_{\mu}\Theta + A_{\mu}\Theta$ is allowed in flat or special sectors, but under invertibility and torsion-free antisymmetric tetrad compatibility it forces its curvature to vanish. It therefore cannot be assumed as the generic curved-GR representation.

For the pure Lorentz branch, preservation of the Lorentz slice and metric already fixes $A_{\mu} = \Omega_{\mu}$ and $B_{\mu} = -\Omega^{\ddagger}_{\mu}$, modulo a common central term that cancels. This removes any independent $A, B$ field doubling, but it is not the generic curved solution: with zero torsion it implies $\nabla_{\mu} V^{\nu} = \delta_{\mu}{}^{\nu}$ and excludes Schwarzschild with nonzero mass. The action-level problem is therefore to derive a nontrivial relative left/right action beyond the pure spin lift, including any internal gauge contribution, and to prove that it is composite or auxiliary rather than propagating. It must not be inserted by assertion.

## B.3 Special relativity and the implicit curved system

In Cartesian inertial Minkowski gauge,

$$\Gamma = \omega = \Omega = 0, \qquad D_{\mu} = \partial_{\mu},$$

and

$$\Theta_{\mathrm{SR}} = \Theta_{0} + \sqrt{\mathcal{N}_{0}}(ix^{0}\mathbf{1} + x^{k}\mathbf{e}_{k})$$

generates the Minkowski tetrad explicitly.

In curved geometry the reconstructed connections depend on the tetrad that is itself generated by $D\Theta$. Hence

$$E_\mu = \mathcal{N}_0^{-1/2}[\partial_\mu\Theta + A_\mu[E,T]\Theta - \Theta B_\mu[E,T]]$$

is an implicit nonlinear first-order PDE/fixed-point system. A Jacobi-theta restriction may make the complete system transcendental as well; that does not remove the separate existence, uniqueness, regularity, and integrability questions.

# C Symbol Dictionary

# D Symbol Dictionary and Notation Conventions

## D.1 Purpose

This section establishes the **unique, canonical meaning** of all symbols used in the Unified Biquaternion Theory. Each symbol has **exactly one meaning** to avoid confusion and ensure consistency across all documents.

## D.2 Reserved Symbols — Single Meaning Only

| Symbol | Canonical Meaning | Notes |
|---|---|---|
| $\alpha$ | Fine structure constant $\approx 1/137.036$ | NO other uses (angle, decay rate, etc.) |
| $\psi$ | Imaginary component of complex time | NOT wavefunction, NOT spinor |
| $\tau$ | Complex time $= t + i\psi$ | NOT proper time |
| $\Theta$ | Fundamental biquaternion field | Capital theta only for field |
| $E_\mu$ | Covariant UBT tetrad | $E_\mu = \mathcal{N}_0^{-1/2} D_\mu\Theta$ |
| $\Gamma^\rho{}_{\mu\nu}$ | Coordinate/affine connection | Acts on spacetime indices |
| $\omega_\mu{}^a{}_b$ | Lorentz-frame connection | $\omega = \mathring{\omega}(e) + K(T)$ for specified torsion |
| $\Omega_\mu$ | Spin/biquaternionic lift of $\omega_\mu$ | Representation-dependent; not $\mathrm{Re}^{-1}\,\Gamma$ |
| $q$ | Biquaternion coordinate (4 DOF) | NOT charge |
| $\mathcal{G}_{\mu\nu}$ | Historical/candidate biquaternionic bilinear | Does not define the active metric |
| $g_{\mu\nu}$ | Lorentz metric tensor | Central coefficient of $\{E_\mu, E_\nu\}_\sharp/2 = g_{\mu\nu}\mathbf{1}$ |
| $\mathcal{T}_{\mu\nu}$ | Biquaternionic stress-energy tensor | Fundamental (geometric phase response) |
| $T_{\mu\nu}$ | Classical stress-energy tensor | DERIVED: $T_{\mu\nu} := \mathrm{Re}(\mathcal{T}_{\mu\nu})$ |
| $F_{\mu\nu}$ | Electromagnetic field strength | QED only |
| $G^a_{\mu\nu}$ | Gluon field strength | QCD only |
| $W^i_{\mu\nu}$ | Weak field strength | Weak interactions |

| Symbol | Canonical Meaning | Notes |
|---|---|---|
| $B_{\mu\nu}$ | Hypercharge field strength | Electroweak |

Table 3: Reserved symbols with unique canonical meanings

## D.3 Index Conventions

### D.3.1 Spacetime Indices

- **Greek indices** $\mu, \nu, \rho, \sigma, \lambda = 0, 1, 2, 3$ for spacetime coordinates
- **Coordinates**: $x^\mu = (x^0, x^1, x^2, x^3) = (t, x, y, z)$ or $(ct, x, y, z)$
- **Summation convention**: Repeated indices are summed (Einstein convention)

### D.3.2 Spatial Indices

- **Latin indices** $i, j, k = 1, 2, 3$ for spatial coordinates only
- **Coordinates**: $x^i = (x^1, x^2, x^3) = (x, y, z)$

### D.3.3 Gauge Indices

- **Color indices** $a, b, c = 1, 2, \ldots, 8$ for $SU(3)$ adjoint representation
- **Weak isospin indices** $i, j, k = 1, 2, 3$ for $SU(2)$ generators
- **Flavor indices** $f, g = u, d, s, c, b, t$ for quark flavors
- **Generation indices** $\alpha, \beta = 1, 2, 3$ for fermion generations

### D.3.4 Matrix Indices

- **Matrix elements** $A, B, C = 0, 1, 2, 3$ for $\Theta$ components: $\Theta_{AB}$
- **Internal indices** $m, n = 1, 2, \ldots, N$ for general matrix dimensions

## D.4 Field and Operator Notation

| Symbol | Meaning | Context |
|---|---|---|
| $\Theta(q, \tau)$ | Fundamental biquaternion field | Core UBT field |
| $\Theta^\dagger$ | Hermitian conjugate of $\Theta$ | $(\bar{\Theta})^T$ |
| $\Psi$ | Wavefunction (if needed) | Use capital psi |
| $A_\mu$ | Electromagnetic potential | QED photon field |
| $G^a_\mu$ | Gluon field | QCD, $a = 1, \ldots, 8$ |
| $W^i_\mu$ | Weak boson field | Electroweak, $i = 1, 2, 3$ |
| $B_\mu$ | Hypercharge field | Electroweak |
| $\Phi$ | Higgs field | Electroweak symmetry breaking |

| Symbol | Meaning | Context |
|---|---|---|

Table 4: Field and operator notation

## D.5 Derivative Notation

| Symbol | Meaning | Notes |
|---|---|---|
| $\partial_\mu$ | Partial derivative $\frac{\partial}{\partial x^\mu}$ | Coordinate derivative |
| $\partial^\mu$ | Contravariant derivative $g^{\mu\nu}\partial_\nu$ | Raised index |
| $\nabla_\mu$ | Covariant derivative (gravity) | Includes Christoffel symbols |
| $D_\mu$ | Gauge covariant derivative | Includes gauge connection |
| $\nabla^\dagger$ | Biquaternionic conjugate derivative | UBT-specific |
| $\Box$ | d'Alembertian $\partial^\mu\partial_\mu$ | Wave operator |

Table 5: Derivative notation

## D.6 Coupling Constants

| Symbol | Meaning | Value | Status |
|---|---|---|---|
| $\alpha$ | Fine structure constant | $\approx 1/137.036$ | Predicted |
| $\alpha_s$ | Strong coupling | $\approx 0.118$ at $M_Z$ | Predicted |
| $g$ | Weak coupling | $SU(2)_L$ | Derived |
| $g'$ | Hypercharge coupling | $U(1)_Y$ | Derived |
| $g_s$ | Strong coupling | $SU(3)_c$ | Derived |
| $e$ | Elementary charge | $\sqrt{4\pi\alpha}$ | Input |
| $G$ | Newton's constant | $6.674 \times 10^{-11}$ $\mathrm{m}^3/\mathrm{kg}/\mathrm{s}^2$ | Input |

Table 6: Coupling constants

## D.7 Mass Scales

| Symbol | Meaning | Approximate Value | Status |
|---|---|---|---|
| $m_e$ | Electron mass | 0.511 MeV | Predicted |
| $m_\mu$ | Muon mass | 105.7 MeV | Predicted |
| $m_\tau$ | Tau mass | 1.777 GeV | Predicted |
| $m_p$ | Proton mass | 938.3 MeV | Input |
| $\Lambda_{\mathrm{QCD}}$ | QCD scale | $\sim 200$ MeV | Predicted |
| $M_W$ | W boson mass | 80.4 GeV | Derived |
| $M_Z$ | Z boson mass | 91.2 GeV | Derived |
| $m_H$ | Higgs mass | 125 GeV | Input |
| $M_{\mathrm{Pl}}$ | Planck mass | $1.22 \times 10^{19}$ GeV | Fundamental |

Table 7: Mass scales

## D.8 Forbidden Symbol Uses

To maintain clarity and avoid conflicts, the following uses are **explicitly forbidden**:

1. $\alpha$ for any angle, decay rate, or parameter other than fine structure constant
2. $\psi$ for wavefunction (use $\Psi$ instead) or spinor (use $\psi_{\text{spinor}}$)
3. $\tau$ for proper time (use $s$ or $\lambda$)
4. $q$ for electric charge (use $Q$ or $e$)
5. $\Theta$ (lowercase theta) for field (reserved for angles if needed)
6. Using $g_{\mu\nu}$ as fundamental (must derive from $\mathcal{G}_{\mu\nu}$)
7. Using $T_{\mu\nu}$ as fundamental (must derive from $\mathcal{T}_{\mu\nu}$)
8. Defining the metric by a trace, real-part, phase, fiber, or embedding readout instead of the central tetrad identity

## D.9 Metric Signature Convention

**Default signature**: $(+,-,-,-)$ (mostly minus, timelike positive)

- $g_{00} > 0$ (timelike positive)
- $g_{11}, g_{22}, g_{33} < 0$ (spacelike negative)

Alternative signature $(-,+,+,+)$ may be used in specific contexts with explicit notice.

## D.10 Unit Conventions

### D.10.1 Natural Units

Default: $\hbar = c = 1$

- Energy, mass, temperature have dimension [mass]
- Length, time have dimension $[\text{mass}]^{-1}$
- Action is dimensionless

### D.10.2 SI Units

When presenting results:

- Masses in eV, keV, MeV, GeV
- Lengths in meters, nm, fm
- Times in seconds
- Energies in eV, joules

## D.11 Special Function Notation

| Symbol | Meaning | Notes |
|---|---|---|
| $\mathrm{Tr}(A)$ | Matrix trace | $\sum_i A_{ii}$ |
| $\det(A)$ | Matrix determinant | Determinant |
| $\mathrm{Re}(z)$ | Real part | Real component |
| $\mathrm{Im}(z)$ | Imaginary part | Imaginary component |
| $\bar{z}$ | Complex conjugate | Conjugation |
| $\theta_i(z, \tau)$ | Jacobi theta functions | $i = 1, 2, 3, 4$ |
| $\zeta(s)$ | Riemann zeta function | Number theory |

Table 8: Special function notation

## D.12 Tensor and Matrix Operations

| Operation | Notation | Meaning |
|---|---|---|
| Matrix product | $AB$ | $(AB)_{ij} = \sum_k A_{ik} B_{kj}$ |
| Tensor product | $A \otimes B$ | Outer product |
| Commutator | $[A, B]$ | $AB - BA$ |
| Anticommutator | $\{A, B\}$ | $AB + BA$ |
| Covariant derivative | $\nabla_\mu T_{\nu\rho}$ | Includes connection |
| Lie derivative | $\mathcal{L}_X T$ | Along vector field $X$ |

Table 9: Tensor and matrix operations

## D.13 Abbreviations and Acronyms

| Acronym | Meaning |
|---|---|
| UBT | Unified Biquaternion Theory |
| GR | General Relativity |
| QFT | Quantum Field Theory |
| QED | Quantum Electrodynamics |
| QCD | Quantum Chromodynamics |
| SM | Standard Model |
| EW | Electroweak |
| GUT | Grand Unified Theory |
| CKM | Cabibbo-Kobayashi-Maskawa (quark mixing matrix) |
| PMNS | Pontecorvo-Maki-Nakagawa-Sakata (neutrino mixing matrix) |
| EWSB | Electroweak Symmetry Breaking |
| VEV | Vacuum Expectation Value |
| DOF | Degrees of Freedom |

Table 10: Abbreviations and acronyms

### D.14 Version Control

This symbol dictionary is version-controlled and canonical. Any proposed changes must:

1. Be justified by theoretical necessity
2. Not conflict with existing usage
3. Be documented in changelog
4. Update all affected documents

### D.15 Enforcement

All UBT documents must:

- Comply with this symbol dictionary
- Report conflicts as errors
- Reference this section for definitions
- Avoid introducing new symbol meanings

# E Mathematical Derivations

# F Computational Methods