

Comparative Risk Assessment of the Burden of Disease from Climate Change

Diarmid Campbell-Lendrum^{1,2} and Rosalie Woodruff³

¹Department of Public Health and Environment, World Health Organization, Geneva, Switzerland; ²London School of Hygiene and Tropical Medicine, London, United Kingdom; ³National Centre for Epidemiology and Population Health, Australian National University, Canberra, Australia

The World Health Organization has developed standardized comparative risk assessment methods for estimating aggregate disease burdens attributable to different risk factors. These have been applied to existing and new models for a range of climate-sensitive diseases in order to estimate the effect of global climate change on current disease burdens and likely proportional changes in the future. The comparative risk assessment approach has been used to assess the health consequences of climate change worldwide, to inform decisions on mitigating greenhouse gas emissions, and in a regional assessment of the Oceania region in the Pacific Ocean to provide more location-specific information relevant to local mitigation and adaptation decisions. The approach places climate change within the same criteria for epidemiologic assessment as other health risks and accounts for the size of the burden of climate-sensitive diseases rather than just proportional change, which highlights the importance of small proportional changes in disease such as diarrhea and malnutrition that cause a large burden. These exercises help clarify important knowledge gaps such as a relatively poor understanding of the role of nonclimatic factors (socioeconomic and other) that may modify future climatic influences and a lack of empirical evidence and methods for quantifying more complex climate-health relationships, which consequently are often excluded from consideration. These exercises highlight the need for risk assessment frameworks that make the best use of traditional epidemiologic methods and that also fully consider the specific characteristics of climate change. These include the long-term and uncertain nature of the exposure and the effects on multiple physical and biotic systems that have the potential for diverse and widespread effects, including high-impact events. *Key words:* burden of disease, climate change, national, quantitative comparative risk assessment, regional. *Environ Health Perspect* 114:1935–1941 (2006). doi:10.1289/ehp.8432 available via <http://dx.doi.org/> [Online 11 July 2006]

The process of climate change, including both increases in global average temperatures ("global warming") and changes in other climate characteristics such as the spatial and temporal distribution of precipitation, has important implications for human health. It is important to describe, measure, and predict the health effects of climate change for two reasons. First, this provides a fuller picture of the consequences of mitigating, or failing to mitigate, emissions of greenhouse gases that are the main anthropogenic contribution to climate change. The long persistence of these gases in the atmosphere means that current mitigation activities (or lack of them) will have consequences for all natural and human systems over coming decades and centuries. They should ideally be informed by measures of the overall size and global distribution of likely health effects of climate change throughout suitably long periods to be considered alongside other impacts such as on biodiversity (Parmesan and Yohe 2003; Thomas et al. 2004a). Even imperfect estimates of the full range of global impacts can provide useful information, provided they are accompanied by clear descriptions of the associated assumptions and uncertainties. Second, quantitative studies can help inform policies to adapt to climate changes that are now inevitable because of both natural variability and past greenhouse gas emissions. Such

actions typically affect the national or subnational level and require information on the likelihood and expected magnitude of specific health impacts in the local context, allowing for the more appropriate allocation of resources to prevent harm from effects such as extreme weather-related events and changes in disease distributions.

Recent comparisons of natural and anthropogenic influences on regional climate (Stott et al. 2004) have demonstrated that human activity increased the probability of a specific past climate event, with severe health consequences (> 44,000 deaths in the European heat wave of summer 2003) (Kosatsky 2005). However, estimating the full range of effects of climate change on health over long time scales presents additional challenges to epidemiologic methods. These include the absence of an appropriate comparison group, the long period over which human actions affect the climate, the large number of health outcomes potentially affected by climatic change, and the numerous nonclimatic influences on each of these outcomes. For these reasons it is misleading simply to observe long-term trends in climate-related diseases and to attribute these changes directly to anthropogenic climate change (e.g., Kovats et al. 2001; McMichael and Githeko 2001; Reiter 2001). The most plausible estimates of future climate change

impacts are instead based on empirically observed relationships between weather or climate conditions and health effects, either in space and/or in time, or, for infectious diseases, on models that capture a detailed understanding of the effects of climate on the biologic processes that determine disease transmission (Rogers and Randolph 2000; Small et al. 2003). Projections of global climate models can be linked to these relationships to indicate how future climate change may influence the level of health outcomes—such as changes in the population living in areas with climates suitable for the transmission of malaria parasites or dengue virus (e.g., Hales et al. 2002; Martens 1998; Rogers and Randolph 2000) or the numbers of people exposed to coastal flooding (Nicholls et al. 1999). These models already provide useful quantitative measures of future risk. However, the results of these models are difficult to relate directly to inform decisions on mitigation (e.g., greenhouse gas emission reduction strategies) because *a*) many do not attempt to account for changes in nonclimatic influences such as economic development (and hence the ability to protect against disease risk), and *b*) the model outcomes are often indirectly related to health, and then only to specific diseases. It is therefore difficult to judge the overall magnitude of the likely health impacts of climate change, either globally or in a specific country (e.g., the combined health effect of a projected 10% increase of the population exposed to coastal flooding, a 20% increase in population living in areas suitable for dengue transmission, and

This article is part of the mini-monograph "Climate Change and Human Health: National Assessments of Impacts and Adaptation."

Address correspondence to D. Campbell-Lendrum, Department of Public Health and Environment, WHO, Geneva CH-1211, Switzerland. Telephone: 41-22-781-4261. Fax: 41-22-791-1383. E-mail: Campbell.lendrum@who.int

We are grateful to T. McMichael, S. Kovats, P. Wilkinson, S. Edwards, F. Tanser, D. Le Sueur, M. Schlesinger, N. Andronova, R. Nicholls, T. Wilson, S. Hales, M. Livermore, A. Prüss-Ustün, C. Corvalán, and B. Menne for contributions to the global assessment exercise, and to T. McMichael, N. Nicholls, K. Hennessy, S. Hales, T. Kjellstrom, and A. Woodward for Oceania.

The views expressed in this article are those of the authors and do not necessarily reflect the position of the World Health Organization.

The authors declare they have no competing financial interests.

Received 17 June 2005; accepted 10 May 2006.

a 5% drop in deaths in cold extremes) and compare these with other threats.

These concerns can be addressed partly by using a standard framework for comparison across risk factors and diseases. The World Health Organization (WHO) has recently developed an approach for comparative risk assessment that has been applied to estimate the current and future disease burden from 25 diverse risk factors, including climate change, in a comparable and transparent manner (Ezzati et al. 2002, 2004; Murray and Lopez 1997; WHO 2002). The assessment generated estimates of the numbers of deaths and disability-adjusted life years (DALYs) attributable to each risk factor in the year 2000, along with expected changes in exposures and associated relative risks of disease outcomes, for several time points between 2000 and 2030. A similar approach has been applied to estimate the effects of climate change on health within the Oceania region in the Pacific Ocean for 2020 and 2050. This assessment principally focused on the impacts on Australian populations, although quantitative estimates were generated for a subset of the health outcomes for New Zealand and Pacific Island countries.

This article outlines the comparative risk assessment approach and discusses its main advantages and limitations for its use in assessments of the health impacts of climate change at the national level. The comparative risk assessment method involves four stages: *a*) identifying health outcomes sensitive to climatic influences, *b*) quantifying the dose-response relationship for a baseline climate period, *c*) defining future exposure scenarios, and *d*) estimating the burden of disease that is attributable to a risk factor (i.e., relative to the risk if climate were unaffected by human actions) and the burden that is avoidable by plausible reductions in the risk factor. Each of these steps requires more detailed decisions, for example, selection between various possible scenarios for future greenhouse gas emissions and associated climate change, or between alternative models describing the relationships between climate and individual health outcomes. These are described only briefly here; more detail on the specific methods used in both the global and the regional assessment are reported by McMichael et al. (2003b, 2004).

Identifying Climate-Sensitive Health Outcomes

Time-series studies and geographic comparisons provide good evidence that a range of health impacts are sensitive to variations in meteorologic conditions of a scale comparable to the climate changes that are expected over the coming century or so (i.e., a 1.4–5.8°C increase in global mean temperatures, changes

in regional patterns of rainfall, and potential increases in the frequency of severe storms [Intergovernmental Panel on Climate Change (IPCC) 2001a; Knutson and Tuleya 2004]. These impacts include deaths and morbidity associated with weather extremes such as heat waves, cold waves, and floods, the incidence of all infectious diseases transmitted by insects and other invertebrate vectors or caused by pathogens that replicate in food or water, and the effects of malnutrition, particularly in poor populations that rely on subsistence farming (McMichael et al. 2003a). These therefore provide an initial list of climate-sensitive health outcomes that should be considered in comparative risk assessment exercise. It should be noted that taking this disease-by-disease approach already tends toward conservative estimates of the full range of health effects. Climate can influence an even wider range of diseases through multiple pathways (e.g., Gommers et al. 2004), and climate is such a major influence on all ecosystem functions that climate change, and particularly sudden "threshold" shifts, may lead to the emergence of new disease threats that are not currently foreseeable.

Quantitative Estimation of Climate–Health Relationships

Comparative risk assessment requires quantitative models of the climatic effects for health outcomes (relevant to the study population) or sufficient reliable disease and environmental data to allow their construction. These models are usually generated based on measurements of the health effects of observed variations in climate in time [e.g., the effect of unusually hot or cold days on disease rates (Checkley et al. 2000; Hajat et al. 2005)] or space (Hales et al. 2002; Rogers and Randolph 2000), or both (e.g., Kuhn et al. 2003; Singh et al. 2001). The extrapolation of short-term or geographic relationships between climate and disease to the process of long-term climate change is probably one of the most important sources of uncertainty in the process because impacts from more gradual processes may be either less severe (e.g., because of gradual adaptation) or more severe (e.g., because of long-term stress leading to irreversible changes

in food-producing ecosystems) than expected. Also, some projected climatic conditions such as heat waves are of a duration or intensity not previously experienced by many (researched) human populations, which presents challenges to the estimation of future risk.

If the assessment seeks to make direct comparisons with disease burdens attributable to other risk factors (as was the case in the global assessment), then it is usually necessary to use a summary measure of population health such as the DALY (Murray 1994) to combine effects of both mortality and morbidity from the various health impacts. This further restricts assessment to diseases with well-characterized and quantified disease burdens (e.g., cases of diarrhea), excluding many likely outcomes of climate change that are relevant to health (e.g., populations suffering increased water stress; Arnell 1999) but that do not yet have well-defined relationships to disease risk. Where different models exist for the same health outcome, final selection should be made on the basis of *a*) validation against historical data, *b*) plausibility of biological assumptions, and *c*) plausibility of extrapolation to other regions. The outcomes considered in the global and regional assessment are shown in Table 1.

Defining Exposure Scenarios

The global and regional analyses used climate models to define alternative plausible distributions of the risk factor in geographic regions over several decades. Many risk factors for health can potentially be altered over relatively short periods (i.e., up to several years). In these cases the focus is usually on calculating current attributable and avoidable burdens to indicate the benefits of addressing the risk factor in the near future. In contrast, decisions on greenhouse gas emissions influence global climate over many decades, and the impacts on human societies are therefore likely to be increasingly evident over similar periods. In this case the full implications of policy change are made clear only when exposure and associated effects are considered over the medium to long term.

A logical baseline "exposure" for comparison would consist of a climate not yet affected by any human activities. This is commonly

Table 1. Health outcomes quantified in the global and Oceania comparative risk assessments.

Type of outcome	Outcome measure	Health measure	Assessment
		Global	Regional
Direct impact of heat and cold	Cardiovascular disease deaths	Incidence	✓ ✓
Foodborne disease	Diarrhea episodes	Incidence	✓ ✓
Waterborne disease	Diarrhea episodes	Incidence	✓
Vectorborne disease	Malaria cases; dengue cases	Incidence	✓ ✓
Natural disasters ^a	Fatal unintentional injuries	Incidence	✓ ✓
	Population displacement	Prevalence	✓ ✓
Risk of malnutrition	Nonavailability of recommended daily calorie intake	Prevalence	✓

✓, risk assessment conducted.

^aAll natural disaster outcomes are separately attributed to coastal floods, or inland floods and landslides.

approximated by using the previous World Meteorological Organization-approved standard period from 1961 to 1990 as the baseline (World Meteorological Organization 1989). The IPCC has concluded that climate changes since around the middle of this period (i.e., 1975) are at least partly attributable to human action (IPCC 2001b). Therefore, this baseline tends to produce conservative estimates of attributable future risk.

Exposure scenarios are based on global climate scenarios: internally consistent representations of future climatic conditions. These are generated by applying a range of levels of anthropogenic "forcings" (most important, greenhouse gas emissions) to computer models representing human and natural influences on the global climate. The output data consist of grid maps of variables, such as temperature, precipitation, and humidity, at a greater or lesser spatial resolution. The global assessment, for example, applied three "scenarios" of future greenhouse gas emission levels (Arnell et al. 2002): *a*) continuing on an unmitigated trajectory approximately following the IPCC IS92a scenario in which effective atmospheric carbon dioxide (CO_2) concentration rises at 1% per year after 1990 (Arnell et al. 2002); *b*) stabilization of CO_2 concentrations at 750 ppm (approximately double preindustrial concentrations; Carbon Dioxide Information Analysis Center 2003) by 2110 (scenario s750); or *c*) stabilization of CO_2 concentrations at 550 ppm by 2170 (scenario s550).

temperature under each climate scenario were given by "shifting" these distributions according to the projected changes in future mean monthly temperature per spatial unit. The resulting relative risks therefore represent net annual deaths, the balance of increasing risks from high temperatures and decreasing risks from low temperatures. The global study included an adjustment for adaptation to increasing summer temperatures. In both cases the relative risk estimates are used to calculate only attributable deaths but not DALYs because the contribution of a relatively short-term advance of deaths in highly vulnerable individuals to the total duration of life lost is highly uncertain.

Impacts on diarrheal disease. Dose-response relationships were derived from time-series studies of temperature variations and diarrhea incidence in developing countries (Checkley et al. 2000; Singh et al. 2001). Relative risks were calculated by multiplying the projected increase in temperature by the derived exposure-response relationship. In the global assessment the resulting relative risks were applied to WHO estimates of the overall current burden from diarrhea in developing regions, to estimate attributable diarrhea deaths and DALYs from climate change in 2000 and relative risk estimates for years to 2030 (adjusting for effects of economic development). In the Oceania assessment the relative risks from studies conducted in developing countries were applied to remote Aboriginal populations (which suffer rates of diarrheal disease similar to those of developing countries) to estimate increased diarrhea cases out to 2050.

Impacts on malnutrition (global assessment only). Existing crop models were used to estimate the effect of projected changes in temperature, rainfall, and CO₂ on future yields of grain, cereals, and soybeans (Parry et al. 1999). These crop yield estimates are part of a world food trade model that accounts for the effects of market forces and government policies on prices, trade, and trends in agricultural and technologic conditions. The model estimates the proportion of the population in each region that has access to sufficient

food to avoid undernourishment (Food and Agriculture Organisation 1987) within each climate scenario.

Natural disasters caused by extreme weather and sea-level rise: coastal floods, inland floods, and mudslides. Relative risks of health impacts from rising sea levels were derived from published models that assess the contribution of projected sea-level rise, topography, and population distribution to estimate numbers of people likely to be exposed to flooding in the future (Hoozemans and Hulsburgen 1995; Nicholls et al. 1999). A new model was developed to estimate the effect of increasingly extreme rainfall events on the impact of inland floods and mudslides on human health. This model was based on an *a priori* assumption that inland flood/landslide frequency is proportional to the frequency with which monthly rainfall exceeds the highest value that might occur once every 10 years, under baseline (i.e., 1961–1990) climate conditions. The assessment in Australia was able to take advantage of a high-quality historical rainfall record, which provided a longer time series to estimate the baseline mean and variability distribution than could be collected at the global level. The future change in frequency of such extreme events was mapped against future population estimates to give the per capita change in risk of experiencing such an extreme weather event. The relative risk for each geographic region was applied to the baseline rate of flood death and injury [derived from reports catalogued in the Emergency Events database (EM-DAT)] [Office of U.S. Foreign Disaster Assistance/Center for Research on the Epidemiology of Disasters (OFDA/CRED) 2001]. The models of flood risk from sea-level rise incorporate the adaptive effect of improved flood protection, assumed to correlate with increasing gross domestic product (GDP) over time. In the global analysis an equivalent adjustment was made for the effect that an increasing GDP was assumed to have on protecting against inland floods.

Vectorborne diseases. Several approaches exist for examining the effect of climate

change on vectorborne diseases, based either on observed relationships between climate conditions and vector development in laboratory or field studies (biological or empirical models) or on correlations between the geographic distribution of disease or vectors and climate variables (statistical models). In the global assessment a validated biological model of the influence of climate on the distribution of *falciparum* malaria in Africa (Tanser et al. 2003) was used to estimate the relative change in population exposed to transmission throughout the world under the alternative climate scenarios. In the Oceania assessment, a locally derived biological model was used to estimate future changes in climatic regions suitable for maintenance of the malaria parasite and vectors within Australia (Bryan et al. 1996), and a global statistical model (Hales et al. 2002) was used to predict regions where dengue transmission could occur in Australia and the increase in regions suitable for dengue transmission in Pacific Island nations. The level of spatial detail in the final models enabled a preliminary estimate of the future costs that might be needed to retain a similar level of protection against these diseases in northern Australia.

Estimates obtained using the methods described above are summarized in Table 2 for the Oceania assessment and Table 3 for one region within the global assessment.

Discussion

There is a general consensus in the scientific literature that human actions are contributing to climate change (IPCC 2001b; Oreskes 2004). Many diseases of public health significance are highly sensitive to climate variability and are likely to be affected by the observed and predicted trend toward warmer and more variable climate conditions (e.g., McMichael and Githeko 2001; McMichael et al. 2003a; National Research Council 2001; Patz et al. 2005). Therefore, to inform mitigation decisions, policy makers are increasingly interested in the likely direction and size of these health effects and their interaction with other nonclimatic influences. To plan adaptation

Table 2. Summary of main findings of the Oceania (for 2050) risk assessment.

Exposure	Health impact estimated	Baseline health impact	Future health impact
Temperature extremes (cold and heat)	Attributable mortality in > 65-year-old age group	1,100 deaths per year (across 10 cities); temperate cities have higher rates of heat deaths than tropical cities	Annual mortality range from 1,400 to 2,000, depending on scenario; increase in heat deaths will significantly outweigh decrease in cold deaths
Rainfall (inland)	Annual incidence of deaths and injuries	Average annual death rate in Australia (1970–2001) was 0.41/million (state rates varied from 0.05 to 3.1); the injury rate was 1.9/million (range, 0.1–8.7) Imported cases only	Predicted annual death rate of 0.53–0.61/million (state rates vary from 0.06 to 4.8); the injury rate was 1.99/million (range, 0.22–13.77)
Temperature and rainfall	Population living in a potential malaria transmission zone	Dengue not established, but local outbreaks from infected travelers occur in far northeast Australia in most years	Substantial southeastern expansion of the malaria zone
Vapor pressure	Population living in a potential dengue transmission zone	Aboriginal people living in remote arid communities have high levels of diarrheal disease	Substantial southeastern and westward expansion of the dengue zone
Temperature	Annual incidence of diarrheal disease		A 10% (5–18%) increase in the annual number of diarrheal hospital admissions among Aboriginal children

measures, these policy makers also require information on the most important disease threats that climate change may bring to specific populations.

Ideally, policymakers considering a particular decision should have quantitative estimates of the full range of effects on human health, over the duration of effect of the decision (e.g., actions to reduce greenhouse gas emissions should be considered over the entire period for which greenhouse gases persist in the atmosphere and influence climate). The comparative risk assessment provides a framework in which to work toward this aim. It is based on standard epidemiologic methods, that is, the definition of a theoretical minimum exposure to the risk factor, measurement of current and projected future exposure levels, consideration of the strength of evidence of an association between the risk factor and various health states, measurement of the relative risk of suffering the disease under alternative exposures, and adjustment for the effects of confounders or effect modifiers. The methods are therefore transparent and open to challenge and further refinement. For example, recent studies of the effect of climate change on measures of malaria risk at national (Hartman et al. 2002) or regional (Small et al. 2003) levels could potentially be used to reestimate or to provide sensitivity measures of this specific impact. The global assessment shows how summary measures of population health, such as DALYs, further allow the different potential health effects of climate change to be aggregated into a single metric, so that the total (measurable) effect can be compared with that of other health risk factors. The Oceania regional assessment demonstrates how different health impacts can be represented separately, illustrating variation in risk between geographic regions and subpopulations, which may be more transparent and intelligible for policy makers.

Such assessments therefore have important advantages. First, by aiming at a comprehensive assessment, they give a better representation of the health consequences of climate change than studies of single disease outcomes in restricted populations. Second,

they help to identify the relative public health burden of different climate-sensitive diseases. The global assessment, for example, showed that relatively small proportional increases in risk for climate-sensitive diseases such as diarrhea and malnutrition may cause very large increases in the total future disease burden.

The attempt to carry out a full accounting of the health impacts of climate change rapidly clarifies significant knowledge gaps. Most of the climate-health models estimate the effects of changing mean values of a climate condition, usually temperature, whereas there is increasing evidence that less predictable changes in extreme values (e.g., Zhou et al. 2004), particularly of precipitation (e.g., Small et al. 2003), may be more important for many diseases. The outputs of many models relevant to such assessments [e.g., predictions of changes in the land area suitable for malaria transmission (Thomas et al. 2004b), population exposed to malaria (Rogers and Randolph 2000), or per capita duration of exposure (Tanser et al. 2003)] are linked only indirectly to disease rates and therefore represent only very approximate measures of the impacts on the burden of clinical disease. Finally, there is still only a limited understanding of the interactions between climate and many important diseases, such as the effects of both temperature and precipitation on diarrhea incidence across different populations. Many plausible or even probable mechanisms by which climate change may affect health have not been modeled quantitatively and have therefore not been included in these assessments. These include, but are not restricted to, changes in outdoor air pollution and aeroallergen (pollen) levels (Beggs 2004), the effect of melting snows and glaciers on floods and landslides, changes in the distribution and transmission of other infectious diseases (particularly vector-borne diseases), the rate of recovery of the ozone hole [affecting exposure to ultraviolet radiation (Shindell et al. 1998)], indirect effects on food production acting through plant pests and diseases, population displacement and destruction of health infrastructure in natural disasters, and the risk of conflict over natural resources.

More generally, quantitative methods generally disregard low-probability but high-impact outcomes. There is increasing evidence from the paleo record and among climatologists (e.g., Hoerling and Kumar 2003) to support a hypothesis that the projected levels of greenhouse gas emissions could lead to a "regime shift" in future climate (driven by, e.g., large releases of methane from the ocean floor, burning and deforestation in the Amazonian rainforest, or a shutdown of the Gulf Stream). Future research in the health area may reasonably assess the health risks—and adaptive requirements—that an abrupt climate change might provoke.

Challenges to Be Addressed

The particular characteristics of climate change cause an additional range of methodologic issues that may be more difficult to resolve. Compared with more traditional risk factors, actions to mitigate or adapt to climate change affect human health through a much wider variety of mechanisms and over much longer periods. Models vary in the extent to which they account for changes in nonclimatic confounders such as the protective effect of adaptation, socioeconomic development, and technologic advances or, conversely, increased vulnerability through population aging and inequity in income or health care provision. Improved health surveillance data, more detailed epidemiologic analysis, and collaboration with nonhealth disciplines should help to narrow these uncertainties. The comparative risk assessment framework described here attempts to estimate only the consequences of changing levels of the risk factor rather than the total effect of any intervention to reduce the risk factor. Examples include ignoring the health co-benefits of reduced air pollution (Cifuentes et al. 2001a, 2001b) or, conversely, the possibility that interventions that reduce fossil fuel consumption may suppress economic development and therefore health status.

There are several levels of uncertainty inherent in the process of estimating climate change health risks. Because anthropogenic climate change is a long-term phenomenon that is superimposed onto natural climate

Table 3. Example of findings of the global (for 2030) risk assessment for one WHO subregion (AfricaE: those sub-Saharan African countries with high child and very high adult mortality).

Exposure	Health impact estimated	Baseline regional situation in 2000	Estimated relative risks attributable to climate change under unmitigated emissions scenario
Rainfall (inland)	Annual incidence of mortality from inland flooding	Average 230 deaths/year reported from 1980 through 1999 ^a	1.86 (1.2–2.44)
Sea-level rise and coastal flooding	Annual incidence of mortality from coastal flooding	No deaths reported in 1980–1999 ^a	1.18 (1.09–1.35)
Temperature and rainfall	Annual incidence of <i>falciparum</i> malaria	More than 420,000 deaths/year ^b	1.14 (1.1–1.28)
Temperature and rainfall	Annual incidence of malnutrition	More than 900,000 deaths/year from malnutrition-related conditions ^b	1.02 (1.1–1.05)
Temperature	Annual incidence of diarrheal disease	More than 430,000 deaths/year ^b	1.08 (0.99–1.06)

The effects of temperature extremes on cardiovascular disease deaths are not presented here because of considerations of short-term mortality displacement (see text).

^aBaseline data derived from OFDA/CRED (2001). ^bBaseline data derived from WHO (2002).

variability, there will always be some uncertainty around the attribution of health impacts, particularly at the national or regional level. Two major types of uncertainty surround the estimation of the exposure measurement. First, we still have incomplete knowledge about how the climate system will respond to continuing change in the composition of gases in the atmosphere. Second, we cannot know in advance what social, technologic,

Appendix.

The IPCC SRES (Nakicenovic et al. 2000) has approved four different storylines that describe plausible relationships between greenhouse-gas-emission driving forces (economic growth, technology, etc.) and the future concentration of gases in atmosphere. These SRES scenarios provide the basis for quantification of future emissions.

The A1 storyline describes a world of very rapid economic growth, a global population that peaks around 2050 and declines thereafter, and the rapid introduction of new and more efficient technologies. Major underlying themes are convergence among regions, capacity building, and increased cultural and social interactions, with a substantial reduction in regional differences in per capita income. Within A1, there are three main subgroups, distinguished by their technologic emphases: fossil fuel intensive (A1FI), predominantly nonfossil energy sources (A1T), and balanced across all energy sources (A1B).

The A2 storyline describes a world of regional self-reliance and preservation of local identities. Fertility patterns across regions converge slowly, which results in increasing global population. Per capita economic growth and technologic change are more fragmented and slower than other storylines.

The B1 storyline describes a convergent world with the same population as in A1 but with rapid changes in economic structures toward a service and information economy, an emphasis on global solutions to economic, social, and environmental sustainability, and the introduction of clean and resource-efficient technologies.

The B2 storyline has an emphasis on local solutions to economic, social, and environmental sustainability. Global population increases at a rate lower than A2, with intermediate economic development and less rapid and more diverse technologic change than in A1 and B1.

In all the scenarios, the effect of specific climate initiatives to reduce emissions [e.g., the Kyoto Protocol to the United Nations Framework Convention on Climate Change (United Nations 1998)] is not included.

demographic, and behavioral changes will occur in human societies over coming decades. For these reasons, results should not be reported in a way that suggests a higher probability to the central estimate of a series of scenarios.

Conclusions

The comparative risk assessment framework is a potentially useful approach to presenting policy-relevant quantitative estimates of the risks that climate change poses to health, at both the global and the national level. In presenting these findings to decision makers, it is important to make clear the limitations of these assessments: quantitative estimates are unavoidably uncertain, changes in nonclimatic factors will influence both the baseline rates of disease and their sensitivity to climate effects, and many of the mechanisms by which climate change may affect health are not currently modeled, likely leading to an underestimation rather than an overestimation of health threats.

Given limited public health resources in many countries for risk assessments, it is important to reduce duplication in this work. Global assessments capture the scale of the future climate-change-related disease burden and highlight regions most at risk for particular health outcomes. This can provide a prompt for specialized regional assessments, which in many regions may give enough accurate information to use at the national level—particularly if countries within the region share similar climatic and economic conditions (e.g., estimated mosquito-borne disease burden in countries with equivalent risks and control programs). National assessments can provide the highest level of spatial resolution and hence the opportunity to quantify variation in risk between administrative divisions. They can also be useful to focus political and community awareness on what is now increasingly recognized as a serious public health issue.

REFERENCES

- Arnell NW. 1998. Climate change and global water resources. *Glob Environ Change* 9:S31–S51.
- Arnell NW, Cannell MGR, Hulme M, Mitchell JFB, Kovats RS, Nicholls RJ, et al. 2002. The consequences of CO₂ stabilization for the impacts of climate change. *Clim Change* 53:413–446.
- Beggs PJ. 2004. Impacts of climate change on aeroallergens: past and future. *Clin Exp Allergy* 34:1507–1513.
- Bryan JH, Foley JD, Sutherland RW. 1998. Malaria transmission and climate change in Australia. *Med J Aust* 164:345–347.
- Carbon Dioxide Information Analysis Center. 2003. Trends Online: A Compendium of Data on Global Change. Available: <http://cdiac.ornl.gov/> [accessed 14 February 2006].
- Checkley W, Epstein LD, Gilman RH, Figueroa D, Cama RI, Patz JA, et al. 2000. Effects of El Niño and ambient temperature on hospital admissions for diarrhoeal diseases in Peruvian children. *Lancet* 355:442–450.
- Cifuentes L, Borja-Aburto VH, Gouveia N, Thurston G, Davis DL. 2001a. Assessing the health benefits of urban air pollution reductions associated with climate change mitigation (2000–2020): Santiago, São Paulo, Mexico City, and New York City. *Environ Health Perspect* 109(suppl 3):419–425.
- Cifuentes L, Borja-Aburto VH, Gouveia N, Thurston G, Davis DL. 2001b. Climate change: Hidden health benefits of greenhouse gas mitigation. *Science* 293:1257–1259.
- Ezzati M, Lopez A, Rodgers A, Murray C, eds. 2004. Comparative Quantification of Health Risks: Global and Regional Burden of Disease due to Selected Major Risk Factors. Geneva:World Health Organization.
- Ezzati M, Lopez AD, Rodgers A, Vander Hoorn S, Murray CJ. 2002. Selected major risk factors and global and regional burden of disease. *Lancet* 360:1347–1360.
- Food and Agriculture Organization. 1987. Fifth World Food Survey. Rome:United Nations Food and Agriculture Organization.
- Gommes R, de Guerry J, Glantz MH, Hsu L-N. 2004. Climate and HIV/AIDS: A Hotspots Analysis for Early Warning Rapid Response Systems. Bangkok:United Nations Development Programme South East Asia.
- Hajat S, Armstrong BG, Gouveia N, Wilkinson P. 2005. Mortality displacement of heat-related deaths: a comparison of Delhi, São Paulo, and London. *Epidemiology* 16:613–620.
- Hales S, de Wet N, Maindonald J, Woodward A. 2002. Potential effect of population and climate changes on global distribution of dengue fever: an empirical model. *Lancet* 360:830–834.
- Hartman J, Ebi KL, McConnell JK, Chan N, Weyant J. 2002. Climate suitability for stable malaria transmission in Zimbabwe under different climate change scenarios. *Glob Change Hum Health* 3:2–14.
- Hoerling M, Kumar A. 2003. The perfect ocean for drought. *Science* 299:691–694.
- Hooymans FMJ, Hulsgiesen CH. 1995. Sea-level rise: a worldwide assessment of risk and protection costs. In: *Climate Change: Impact on Coastal Habitation* (Eisma D, ed). London:Lewis Publishers, 137–163.
- IPCC (Intergovernmental Panel on Climate Change). 2001a. Climate Change 2001: Synthesis Report. Cambridge, UK:World Meteorological Organization/United Nations Environment Programme.
- IPCC (Intergovernmental Panel on Climate Change). 2001b. Climate Change 2001: The Scientific Basis: Contribution of Working Group I to the Third Assessment Report. Cambridge, UK:Cambridge University Press.
- Kay D, Prüss A, Corvalan C. 2000. Methodology for Assessment of Environmental Burden of Disease. Report on the ISESE Session on Environmental Burden of Disease, Buffalo, 22 August 2000. WHO/SDE/WSH/00.7. Geneva:World Health Organization.
- Knutson TR, Tuleya RE. 2004. Impact of CO₂-induced warming on simulated hurricane intensity and precipitation: sensitivity to the choice of climate model and convective parameterization. *J Clim* 17:3577–3595.
- Kosatsky T. 2005. The 2003 European heat waves. *Euro Surveill* 10:148–149.
- Kovats RS, Campbell-Lendrum DH, McMichael AJ, Woodward A, Corvalan C. 2001. Early effects of climate change: do they include changes in vector-borne disease? *Philos Trans R Soc Lond B Biol Sci* 356:1057–1068.
- Kuh KG, Campbell-Lendrum DH, Armstrong B, Davies CR. 2003. Malaria in Britain: past, present, and future. *Proc Natl Acad Sci USA* 100:9997–10001.
- Martens WJ. 1998. Human and Climate Change: Modelling the Impacts of Global Warming and Ozone Depletion. London:Earthscan.
- McMichael A, Campbell-Lendrum DH, Corvalan C, Ebi K, Githeko A, Scheraga J, et al., eds. 2003a. *Climate Change and Health: Risks and Responses*. Geneva:World Health Organization.
- McMichael A, Campbell-Lendrum DH, Kovats RS, Edwards S, Wilkinson P, Wilson T, et al. 2004. Climate change. In: *Comparative Quantification of Health Risks: Global and Regional Burden of Disease due to Selected Major Risk Factors* (Ezzati M, Lopez A, Rodgers A, Murray C, eds). Geneva:World Health Organization, 1543–1649.
- McMichael A, Woodruff R, Whetton P, Hennessy P, Nicholls N, Hales S, et al. 2003b. Human Health and Climate Change in Oceania: A Risk Assessment. Canberra:Commonwealth Department of Health and Ageing.
- McMichael AJ, Githeko A. 2001. Human health. In: *Climate Change 2001: Impacts, Adaptation and Vulnerability* (McCarthy JJ, Canziani OF, Leary NA, Dokken DJ, White KS, eds). Cambridge, UK:Cambridge University Press, 451–485.
- Murray CJL. 1994. Quantifying the burden of disease—the technical

- basis for disability-adjusted life years. *Bull World Health Organ* 72:429–445.
- Murray CJL, Lopez AD. 1997. Alternative projections of mortality and disability by cause 1990–2020: Global Burden of Disease Study. *Lancet* 349:1486–1504.
- Nakicenovic N, Alcamo J, Davis G, de Vries, B, Fenhann J, Gaffin S, et al. 2000. IPCC Special Report on Emissions Scenarios. Cambridge, UK:Cambridge University Press.
- National Research Council. 2001. Under the Weather: Climate, Ecosystems, and Infectious Disease. Washington, DC: National Academy Press.
- Nicholls RJ, Hooymans FMJ, Marchand M. 1999. Increasing flood risk and wetland losses due to global sea-level rise: regional and global analyses. *Glob Environ Change* 9:S69–S87.
- OFDA/CRED (Office of U.S. Foreign Disaster Assistance/Center for Research on the Epidemiology of Disasters). 2001. EM-DAT: The International Disaster Database. Brussels: Université Catholique de Louvain. Available: <http://www.cred.be/medat> [accessed July 2002].
- Oreskes N. 2004. Beyond the ivory tower. The scientific consensus on climate change [Letter]. *Science* 306:1686.
- Parmesan C, Yohe G. 2003. A globally coherent fingerprint of climate change impacts across natural systems. *Nature* 421:37–42.
- Parry M, Rosenzweig C, Iglesias A, Fischer G, Livermore M. 1999. Climate change and world food security: a new assessment. *Glob Environ Change* 9:S51–S87.
- Patz JA, Campbell-Lendrum D, Holloway T, Foley JA. 2005. Impact of regional climate change on human health. *Nature* 438:310–317.
- Reiter P. 2001. Climate change and mosquito-borne disease. *Environ Health Perspect* 109:141–161.
- Rogers DJ, Randolph SE. 2000. The global spread of malaria in a future, warmer world. *Science* 289:1763–1766.
- Shindell DT, Rind D, Lerner P. 1998. Increased polar stratospheric ozone losses and delayed eventual recovery owing to increasing greenhouse-gas concentrations. *Nature* 392:589–592.
- Singh RB, Hales S, de Wet N, Raj R, Hearnden M, Weinstein P, et al. 2001. The influence of climate variation and change on diarrheal disease in the Pacific Islands. *Environ Health Perspect* 109:155–159.
- Small J, Goetz SJ, Hay SI. 2003. Climatic suitability for malaria transmission in Africa, 1911–1998. *Proc Natl Acad Sci USA* 100:15341–15345.
- Stott PA, Stone DA, Allen MR. 2003. Human contribution to the European heatwave of 2003. *Nature* 432:610–614.
- Tanser FC, Sharp B, le Sueur D. 2003. Potential effect of climate change on malaria transmission in Africa. *Lancet* 362:1792–1798.
- Thomas CD, Cameron A, Green RE, Bakkenes M, Beaumont LJ, Collingham YC, et al. 2004a. Extinction risk from climate change. *Nature* 427:145–148.
- Thomas CJ, Davies G, Dunn CE. 2004b. Mixed picture for changes in stable malaria distribution with future climate in Africa. *Trends Parasitol* 20:216–220.
- United Nations. 1998. Kyoto Protocol to the United Nations Framework Convention on Climate Change. Available: <http://unfccc.int/resource/docs/convkp/kpeng.pdf> [accessed 4 November 2006].
- WHO. 2002. The World Health Report 2002. Geneva:World Health Organization.
- Woodward A, Hales S, Weinstein P. 1998. Climate change and human health in the Asia Pacific region: who will be most vulnerable? *Climate Res* 11:31–38.
- World Meteorological Organization. 1989. Calculation of Monthly and Annual 30-Year Standard Normal. WCDP-No. 10. WMO-TD/ No. 34. Geneva:World Meteorological Organization.
- Zhou G, Minakawa N, Githeko AK, Yan G. 2004. Association between climate variability and malaria epidemics in the East African highlands. *Proc Natl Acad Sci USA* 101:2375–2380.