


Learning Nominal Automata

Matteo Sammartino

RISE, IST Austria, May 10 2017

Learning Nominal Automata (POPL '17)



Joshua Moerman
Radboud University



Matteo Sammartino
UCL



Alexandra Silva
UCL




Bartek Klin
Warsaw University




Michał Szynwelski
Warsaw University

Active learning



Active learning



No formal specification available? **Learn it!**

L^* algorithm (D.Angluin '87)


Finite alphabet of system's actions A

set of system behaviors is a **regular language** $\mathcal{L} \subseteq A^*$

L^* algorithm (D.Angluin '87)

Finite alphabet of system's actions A


set of system behaviors is a **regular language** $\mathcal{L} \subseteq A^*$



L^* algorithm (D.Angluin '87)

Finite alphabet of system's actions A


set of system behaviors is a **regular language** $\mathcal{L} \subseteq A^*$



L^* algorithm (D.Angluin '87)

Finite alphabet of system's actions A


set of system behaviors is a **regular language** $\mathcal{L} \subseteq A^*$



L^* algorithm (D.Angluin '87)

Finite alphabet of system's actions A

set of system behaviors is a **regular language** $\mathcal{L} \subseteq A^*$



Observation table

		E		
		ϵ	a	aa
S	ϵ	0	0	1
	a	0	1	0
	b	0	0	0

$$\text{row}: S \cup S \cdot A \rightarrow 2^E$$

$$\text{row}(s)(e) = 1 \iff se \in \mathcal{L}$$

$$S, E \subseteq A^* \quad A = \{a, b\}$$

Example

$$\mathcal{L} = \{aa, bb\} \quad A = \{a, b\}$$

$$\mathcal{L} = \{aa, bb\} \quad A = \{a, b\}$$

$S, E \leftarrow \{\epsilon\}$

repeat

while (S, E) is not closed or not consistent

if (S, E) is not closed

 find $s_1 \in S, a \in A$ such that

$\text{row}(s_1a) \neq \text{row}(s)$, for all $s \in S$

$S \leftarrow S \cup \{s_1a\}$

if (S, E) is not consistent

 find $s_1, s_2 \in S, a \in A$, and $e \in E$ such that

$\text{row}(s_1) = \text{row}(s_2)$

 and $\text{row}(s_1a)(e) \neq \text{row}(s_2a)(e)$

$E \leftarrow E \cup \{ae\}$

Submit hypothesis H to the Teacher

if the Teacher replies **no**, with a counter-example t

$S \leftarrow S \cup \text{prefixes}(t)$

until the Teacher replies **yes** to H

return H

$$\mathcal{L} = \{aa, bb\} \quad A = \{a, b\}$$

$$S, E \leftarrow \{\epsilon\}$$

repeat

while (S, E) is not closed or not consistent

if (S, E) is not closed

 find $s_1 \in S, a \in A$ such that

$\text{row}(s_1a) \neq \text{row}(s)$, for all $s \in S$

$S \leftarrow S \cup \{s_1a\}$

if (S, E) is not consistent

 find $s_1, s_2 \in S, a \in A$, and $e \in E$ such that

$\text{row}(s_1) = \text{row}(s_2)$

 and $\text{row}(s_1a)(e) \neq \text{row}(s_2a)(e)$

$E \leftarrow E \cup \{ae\}$

Submit hypothesis H to the Teacher

if the Teacher replies **no**, with a counter-example t

$S \leftarrow S \cup \text{prefixes}(t)$

until the Teacher replies **yes** to H

return H

$$\mathcal{L} = \{aa, bb\} \quad A = \{a, b\}$$

$$S, E \leftarrow \{\epsilon\}$$

repeat

while (S, E) is not closed or not consistent

if (S, E) is not closed

 find $s_1 \in S, a \in A$ such that

$\text{row}(s_1a) \neq \text{row}(s)$, for all $s \in S$

$S \leftarrow S \cup \{s_1a\}$

if (S, E) is not consistent

 find $s_1, s_2 \in S, a \in A$, and $e \in E$ such that

$\text{row}(s_1) = \text{row}(s_2)$

 and $\text{row}(s_1a)(e) \neq \text{row}(s_2a)(e)$

$E \leftarrow E \cup \{ae\}$

Submit hypothesis H to the Teacher

if the Teacher replies **no**, with a counter-example t

$S \leftarrow S \cup \text{prefixes}(t)$

until the Teacher replies **yes** to H

return H

$\text{row} =$

	ϵ
ϵ	0
a	0
b	0

$$\mathcal{L} = \{aa, bb\} \quad A = \{a, b\}$$

$S, E \leftarrow \{\epsilon\}$

repeat

while (S, E) is not closed or not consistent

if (S, E) is not closed

 find $s_1 \in S, a \in A$ such that

$\text{row}(s_1a) \neq \text{row}(s)$, for all $s \in S$

$S \leftarrow S \cup \{s_1a\}$

if (S, E) is not consistent

 find $s_1, s_2 \in S, a \in A$, and $e \in E$ such that

$\text{row}(s_1) = \text{row}(s_2)$

 and $\text{row}(s_1a)(e) \neq \text{row}(s_2a)(e)$

$E \leftarrow E \cup \{ae\}$

Submit hypothesis H to the Teacher

if the Teacher replies **no**, with a counter-example t

$S \leftarrow S \cup \text{prefixes}(t)$

until the Teacher replies **yes** to H

return H

$\text{row} =$

	ϵ
ϵ	0
a	0
b	0

$$\mathcal{L} = \{aa, bb\} \quad A = \{a, b\}$$

$S, E \leftarrow \{\epsilon\}$

repeat

while (S, E) is not closed or not consistent

if (S, E) is not closed

 find $s_1 \in S, a \in A$ such that

$\text{row}(s_1a) \neq \text{row}(s)$, for all $s \in S$

$S \leftarrow S \cup \{s_1a\}$

if (S, E) is not consistent

 find $s_1, s_2 \in S, a \in A$, and $e \in E$ such that

$\text{row}(s_1) = \text{row}(s_2)$

 and $\text{row}(s_1a)(e) \neq \text{row}(s_2a)(e)$

$E \leftarrow E \cup \{ae\}$

Submit hypothesis H to the Teacher

if the Teacher replies **no**, with a counter-example t

$S \leftarrow S \cup \text{prefixes}(t)$

until the Teacher replies **yes** to H

return H

$\text{row} =$

	ϵ
ϵ	0
a	0
b	0

Closed

$$\forall t \in S \cdot A \quad \exists s \in S \quad \text{row}(t) = \text{row}(s)$$

$$\mathcal{L} = \{aa, bb\} \quad A = \{a, b\}$$

$S, E \leftarrow \{\epsilon\}$

repeat

while (S, E) is not closed or not consistent

if (S, E) is not closed

 find $s_1 \in S, a \in A$ such that

$\text{row}(s_1a) \neq \text{row}(s)$, for all $s \in S$

$S \leftarrow S \cup \{s_1a\}$

if (S, E) is not consistent

 find $s_1, s_2 \in S, a \in A$, and $e \in E$ such that

$\text{row}(s_1) = \text{row}(s_2)$

 and $\text{row}(s_1a)(e) \neq \text{row}(s_2a)(e)$

$E \leftarrow E \cup \{ae\}$

Submit hypothesis H to the Teacher

if the Teacher replies **no**, with a counter-example t

$S \leftarrow S \cup \text{prefixes}(t)$

until the Teacher replies **yes** to H

return H

$\text{row} =$

	ϵ
ϵ	0
a	0
b	0

Closed ✓

$\forall t \in S \cdot A \quad \exists s \in S \quad \text{row}(t) = \text{row}(s)$

$$\mathcal{L} = \{aa, bb\} \quad A = \{a, b\}$$

$S, E \leftarrow \{\epsilon\}$

repeat

while (S, E) is not closed or not consistent

if (S, E) is not closed

 find $s_1 \in S, a \in A$ such that

$\text{row}(s_1a) \neq \text{row}(s)$, for all $s \in S$

$S \leftarrow S \cup \{s_1a\}$

if (S, E) is not consistent

 find $s_1, s_2 \in S, a \in A$, and $e \in E$ such that

$\text{row}(s_1) = \text{row}(s_2)$

 and $\text{row}(s_1a)(e) \neq \text{row}(s_2a)(e)$

$E \leftarrow E \cup \{ae\}$

Submit hypothesis H to the Teacher

if the Teacher replies **no**, with a counter-example t

$S \leftarrow S \cup \text{prefixes}(t)$

until the Teacher replies **yes** to H

return H

$\text{row} =$

	ϵ
ϵ	0
a	0
b	0

Closed ✓

$\forall t \in S \cdot A \quad \exists s \in S \quad \text{row}(t) = \text{row}(s)$

Consistent

$\forall s_1, s_2 \in S \quad \text{row}(s_1) = \text{row}(s_2) \implies \forall a \in A \quad \text{row}(s_1a) = \text{row}(s_2a)$

$$\mathcal{L} = \{aa, bb\} \quad A = \{a, b\}$$

$S, E \leftarrow \{\epsilon\}$

repeat

while (S, E) is not closed or not consistent

if (S, E) is not closed

 find $s_1 \in S, a \in A$ such that

$\text{row}(s_1a) \neq \text{row}(s)$, for all $s \in S$

$S \leftarrow S \cup \{s_1a\}$

if (S, E) is not consistent

 find $s_1, s_2 \in S, a \in A$, and $e \in E$ such that

$\text{row}(s_1) = \text{row}(s_2)$

 and $\text{row}(s_1a)(e) \neq \text{row}(s_2a)(e)$

$E \leftarrow E \cup \{ae\}$

Submit hypothesis H to the Teacher

if the Teacher replies **no**, with a counter-example t

$S \leftarrow S \cup \text{prefixes}(t)$

until the Teacher replies **yes** to H

return H

$\text{row} =$

	ϵ
ϵ	0
a	0
b	0

Closed ✓

$\forall t \in S \cdot A \quad \exists s \in S \quad \text{row}(t) = \text{row}(s)$

Consistent ✓

$\forall s_1, s_2 \in S \quad \text{row}(s_1) = \text{row}(s_2) \implies \forall a \in A \quad \text{row}(s_1a) = \text{row}(s_2a)$

$$\mathcal{L} = \{aa, bb\} \quad A = \{a, b\}$$

$S, E \leftarrow \{\epsilon\}$

repeat

while (S, E) is not closed or not consistent

if (S, E) is not closed

 find $s_1 \in S, a \in A$ such that

$\text{row}(s_1a) \neq \text{row}(s)$, for all $s \in S$

$S \leftarrow S \cup \{s_1a\}$

if (S, E) is not consistent

 find $s_1, s_2 \in S, a \in A$, and $e \in E$ such that

$\text{row}(s_1) = \text{row}(s_2)$

 and $\text{row}(s_1a)(e) \neq \text{row}(s_2a)(e)$

$E \leftarrow E \cup \{ae\}$

Submit hypothesis H to the Teacher

if the Teacher replies **no**, with a counter-example t

$S \leftarrow S \cup \text{prefixes}(t)$

until the Teacher replies **yes** to H

return H

$\text{row} =$

	ϵ
ϵ	0
a	0
b	0

Closed ✓

$\forall t \in S \cdot A \quad \exists s \in S \quad \text{row}(t) = \text{row}(s)$

Consistent ✓

$\forall s_1, s_2 \in S \quad \text{row}(s_1) = \text{row}(s_2) \implies \forall a \in A \quad \text{row}(s_1a) = \text{row}(s_2a)$

$$\mathcal{L} = \{aa, bb\} \quad A = \{a, b\}$$

$S, E \leftarrow \{\epsilon\}$

repeat

while (S, E) is not closed or not consistent

if (S, E) is not closed

 find $s_1 \in S, a \in A$ such that

$\text{row}(s_1a) \neq \text{row}(s)$, for all $s \in S$

$S \leftarrow S \cup \{s_1a\}$

if (S, E) is not consistent

 find $s_1, s_2 \in S, a \in A$, and $e \in E$ such that

$\text{row}(s_1) = \text{row}(s_2)$

 and $\text{row}(s_1a)(e) \neq \text{row}(s_2a)(e)$

$E \leftarrow E \cup \{ae\}$

Submit hypothesis H to the Teacher

if the Teacher replies **no**, with a counter-example t

$S \leftarrow S \cup \text{prefixes}(t)$

until the Teacher replies **yes** to H

return H

$\text{row} =$

	ϵ
ϵ	0
a	0
b	0

Closed ✓

$\forall t \in S \cdot A \quad \exists s \in S \quad \text{row}(t) = \text{row}(s)$

Consistent ✓

$\forall s_1, s_2 \in S \quad \text{row}(s_1) = \text{row}(s_2) \implies \forall a \in A \quad \text{row}(s_1a) = \text{row}(s_2a)$

Hypothesis

$$\mathcal{L} = \{aa, bb\} \quad A = \{a, b\}$$

$S, E \leftarrow \{\epsilon\}$

repeat

while (S, E) is not closed or not consistent

if (S, E) is not closed

 find $s_1 \in S, a \in A$ such that

$\text{row}(s_1a) \neq \text{row}(s)$, for all $s \in S$

$S \leftarrow S \cup \{s_1a\}$

if (S, E) is not consistent

 find $s_1, s_2 \in S, a \in A$, and $e \in E$ such that

$\text{row}(s_1) = \text{row}(s_2)$

 and $\text{row}(s_1a)(e) \neq \text{row}(s_2a)(e)$

$E \leftarrow E \cup \{ae\}$

Submit hypothesis H to the Teacher

if the Teacher replies **no**, with a counter-example t

$S \leftarrow S \cup \text{prefixes}(t)$

until the Teacher replies **yes** to H

return H

$\text{row} =$

	ϵ
ϵ	0
a	0
b	0

Closed ✓

$\forall t \in S \cdot A \quad \exists s \in S \quad \text{row}(t) = \text{row}(s)$

Consistent ✓

$\forall s_1, s_2 \in S \quad \text{row}(s_1) = \text{row}(s_2) \implies \forall a \in A \quad \text{row}(s_1a) = \text{row}(s_2a)$

Hypothesis

states =

$\{\text{row}(s) \mid s \in S\}$

$$\mathcal{L} = \{aa, bb\} \quad A = \{a, b\}$$

$S, E \leftarrow \{\epsilon\}$

repeat

while (S, E) is not closed or not consistent

if (S, E) is not closed

 find $s_1 \in S, a \in A$ such that

$\text{row}(s_1a) \neq \text{row}(s)$, for all $s \in S$

$S \leftarrow S \cup \{s_1a\}$

if (S, E) is not consistent

 find $s_1, s_2 \in S, a \in A$, and $e \in E$ such that

$\text{row}(s_1) = \text{row}(s_2)$

 and $\text{row}(s_1a)(e) \neq \text{row}(s_2a)(e)$

$E \leftarrow E \cup \{ae\}$

Submit hypothesis H to the Teacher

if the Teacher replies **no**, with a counter-example t

$S \leftarrow S \cup \text{prefixes}(t)$

until the Teacher replies **yes** to H

return H

$\text{row} =$

	ϵ
ϵ	0
a	0
b	0

Closed ✓

$\forall t \in S \cdot A \quad \exists s \in S \quad \text{row}(t) = \text{row}(s)$

Consistent ✓

$\forall s_1, s_2 \in S \quad \text{row}(s_1) = \text{row}(s_2) \implies \forall a \in A \quad \text{row}(s_1a) = \text{row}(s_2a)$

Hypothesis

$\text{row}(\epsilon)$

states =
 $\{\text{row}(s) \mid s \in S\}$

$$\mathcal{L} = \{aa, bb\} \quad A = \{a, b\}$$

$S, E \leftarrow \{\epsilon\}$

repeat

while (S, E) is not closed or not consistent

if (S, E) is not closed

 find $s_1 \in S, a \in A$ such that

$\text{row}(s_1a) \neq \text{row}(s)$, for all $s \in S$

$S \leftarrow S \cup \{s_1a\}$

if (S, E) is not consistent

 find $s_1, s_2 \in S, a \in A$, and $e \in E$ such that

$\text{row}(s_1) = \text{row}(s_2)$

 and $\text{row}(s_1a)(e) \neq \text{row}(s_2a)(e)$

$E \leftarrow E \cup \{ae\}$

Submit hypothesis H to the Teacher

if the Teacher replies **no**, with a counter-example t

$S \leftarrow S \cup \text{prefixes}(t)$

until the Teacher replies **yes** to H

return H

$\text{row} =$

	ϵ
ϵ	0
a	0
b	0

Closed ✓

$\forall t \in S \cdot A \quad \exists s \in S \quad \text{row}(t) = \text{row}(s)$

Consistent ✓

$\forall s_1, s_2 \in S \quad \text{row}(s_1) = \text{row}(s_2) \implies \forall a \in A \quad \text{row}(s_1a) = \text{row}(s_2a)$

Hypothesis

$\text{row}(\epsilon)$

initial state =
 $\text{row}(\epsilon)$

$$\mathcal{L} = \{aa, bb\} \quad A = \{a, b\}$$

$S, E \leftarrow \{\epsilon\}$

repeat

while (S, E) is not closed or not consistent

if (S, E) is not closed

 find $s_1 \in S, a \in A$ such that

$\text{row}(s_1a) \neq \text{row}(s)$, for all $s \in S$

$S \leftarrow S \cup \{s_1a\}$

if (S, E) is not consistent

 find $s_1, s_2 \in S, a \in A$, and $e \in E$ such that

$\text{row}(s_1) = \text{row}(s_2)$

 and $\text{row}(s_1a)(e) \neq \text{row}(s_2a)(e)$

$E \leftarrow E \cup \{ae\}$

Submit hypothesis H to the Teacher

if the Teacher replies **no**, with a counter-example t

$S \leftarrow S \cup \text{prefixes}(t)$

until the Teacher replies **yes** to H

return H

$\text{row} =$

	ϵ
ϵ	0
a	0
b	0

Closed ✓

$\forall t \in S \cdot A \quad \exists s \in S \quad \text{row}(t) = \text{row}(s)$

Consistent ✓

$\forall s_1, s_2 \in S \quad \text{row}(s_1) = \text{row}(s_2) \implies \forall a \in A \quad \text{row}(s_1a) = \text{row}(s_2a)$

Hypothesis

$\rightarrow \boxed{\text{row}(\epsilon)}$

initial state =
 $\text{row}(\epsilon)$

$$\mathcal{L} = \{aa, bb\} \quad A = \{a, b\}$$

$S, E \leftarrow \{\epsilon\}$

repeat

while (S, E) is not closed or not consistent

if (S, E) is not closed

 find $s_1 \in S, a \in A$ such that

$\text{row}(s_1a) \neq \text{row}(s)$, for all $s \in S$

$S \leftarrow S \cup \{s_1a\}$

if (S, E) is not consistent

 find $s_1, s_2 \in S, a \in A$, and $e \in E$ such that

$\text{row}(s_1) = \text{row}(s_2)$

 and $\text{row}(s_1a)(e) \neq \text{row}(s_2a)(e)$

$E \leftarrow E \cup \{ae\}$

Submit hypothesis H to the Teacher

if the Teacher replies **no**, with a counter-example t

$S \leftarrow S \cup \text{prefixes}(t)$

until the Teacher replies **yes** to H

return H

$\text{row} =$

	ϵ
ϵ	0
a	0
b	0

Closed ✓

$\forall t \in S \cdot A \quad \exists s \in S \quad \text{row}(t) = \text{row}(s)$

Consistent ✓

$\forall s_1, s_2 \in S \quad \text{row}(s_1) = \text{row}(s_2) \implies \forall a \in A \quad \text{row}(s_1a) = \text{row}(s_2a)$

Hypothesis

$\rightarrow \boxed{\text{row}(\epsilon)}$

final states =

$\{\text{row}(s) \mid \text{row}(s)(\epsilon) = 1\}$

$$\mathcal{L} = \{aa, bb\} \quad A = \{a, b\}$$

$S, E \leftarrow \{\epsilon\}$

repeat

while (S, E) is not closed or not consistent

if (S, E) is not closed

 find $s_1 \in S, a \in A$ such that

$\text{row}(s_1a) \neq \text{row}(s)$, for all $s \in S$

$S \leftarrow S \cup \{s_1a\}$

if (S, E) is not consistent

 find $s_1, s_2 \in S, a \in A$, and $e \in E$ such that

$\text{row}(s_1) = \text{row}(s_2)$

 and $\text{row}(s_1a)(e) \neq \text{row}(s_2a)(e)$

$E \leftarrow E \cup \{ae\}$

Submit hypothesis H to the Teacher

if the Teacher replies **no**, with a counter-example t

$S \leftarrow S \cup \text{prefixes}(t)$

until the Teacher replies **yes** to H

return H

$\text{row} =$

	ϵ
ϵ	0
a	0
b	0

Closed ✓

$\forall t \in S \cdot A \quad \exists s \in S \quad \text{row}(t) = \text{row}(s)$

Consistent ✓

$\forall s_1, s_2 \in S \quad \text{row}(s_1) = \text{row}(s_2) \implies \forall a \in A \quad \text{row}(s_1a) = \text{row}(s_2a)$

Hypothesis

$\rightarrow \boxed{\text{row}(\epsilon)}$

transitions =

$\text{row}(s) \xrightarrow{a} \text{row}(sa)$

$$\mathcal{L} = \{aa, bb\} \quad A = \{a, b\}$$

$S, E \leftarrow \{\epsilon\}$

repeat

while (S, E) is not closed or not consistent

if (S, E) is not closed

 find $s_1 \in S, a \in A$ such that

$\text{row}(s_1a) \neq \text{row}(s)$, for all $s \in S$

$S \leftarrow S \cup \{s_1a\}$

if (S, E) is not consistent

 find $s_1, s_2 \in S, a \in A$, and $e \in E$ such that

$\text{row}(s_1) = \text{row}(s_2)$

 and $\text{row}(s_1a)(e) \neq \text{row}(s_2a)(e)$

$E \leftarrow E \cup \{ae\}$

Submit hypothesis H to the Teacher

if the Teacher replies **no**, with a counter-example t

$S \leftarrow S \cup \text{prefixes}(t)$

until the Teacher replies **yes** to H

return H

$\text{row} =$

	ϵ
ϵ	0
a	0
b	0


Closed ✓

$\forall t \in S \cdot A \quad \exists s \in S \quad \text{row}(t) = \text{row}(s)$

Consistent ✓

$\forall s_1, s_2 \in S \quad \text{row}(s_1) = \text{row}(s_2) \implies \forall a \in A \quad \text{row}(s_1a) = \text{row}(s_2a)$

Hypothesis



transitions =

$\text{row}(s) \xrightarrow{a} \text{row}(sa)$

$$\mathcal{L} = \{aa, bb\} \quad A = \{a, b\}$$

$S, E \leftarrow \{\epsilon\}$

repeat

while (S, E) is not closed or not consistent

if (S, E) is not closed

 find $s_1 \in S, a \in A$ such that

$\text{row}(s_1a) \neq \text{row}(s)$, for all $s \in S$

$S \leftarrow S \cup \{s_1a\}$

if (S, E) is not consistent

 find $s_1, s_2 \in S, a \in A$, and $e \in E$ such that

$\text{row}(s_1) = \text{row}(s_2)$

 and $\text{row}(s_1a)(e) \neq \text{row}(s_2a)(e)$

$E \leftarrow E \cup \{ae\}$

Submit hypothesis H to the Teacher

if the Teacher replies **no**, with a counter-example t

$S \leftarrow S \cup \text{prefixes}(t)$

until the Teacher replies **yes** to H

return H

$\text{row} =$

	ϵ
ϵ	0
a	0
b	0


Closed ✓ transitions are well-defined

$\forall t \in S \cdot A \quad \exists s \in S \quad \text{row}(t) = \text{row}(s)$

Consistent ✓

$\forall s_1, s_2 \in S \quad \text{row}(s_1) = \text{row}(s_2) \implies \forall a \in A \quad \text{row}(s_1a) = \text{row}(s_2a)$

Hypothesis



transitions =

$\text{row}(s) \xrightarrow{a} \text{row}(sa)$

$$\mathcal{L} = \{aa, bb\} \quad A = \{a, b\}$$

$S, E \leftarrow \{\epsilon\}$

repeat

while (S, E) is not closed or not consistent

if (S, E) is not closed

 find $s_1 \in S, a \in A$ such that

$\text{row}(s_1a) \neq \text{row}(s)$, for all $s \in S$

$S \leftarrow S \cup \{s_1a\}$

if (S, E) is not consistent

 find $s_1, s_2 \in S, a \in A$, and $e \in E$ such that

$\text{row}(s_1) = \text{row}(s_2)$

 and $\text{row}(s_1a)(e) \neq \text{row}(s_2a)(e)$

$E \leftarrow E \cup \{ae\}$

Submit hypothesis H to the Teacher

if the Teacher replies **no**, with a counter-example t

$S \leftarrow S \cup \text{prefixes}(t)$

until the Teacher replies **yes** to H

return H

$\text{row} =$

	ϵ
ϵ	0
a	0
b	0


Closed ✓ transitions are well-defined

$\forall t \in S \cdot A \quad \exists s \in S \quad \text{row}(t) = \text{row}(s)$

Consistent ✓ determinism

$\forall s_1, s_2 \in S \quad \text{row}(s_1) = \text{row}(s_2) \implies \forall a \in A \quad \text{row}(s_1a) = \text{row}(s_2a)$

Hypothesis



transitions =

$\text{row}(s) \xrightarrow{a} \text{row}(sa)$

$$\mathcal{L} = \{aa, bb\} \quad A = \{a, b\}$$

$$S, E \leftarrow \{\epsilon\}$$

repeat

while (S, E) is not closed or not consistent

if (S, E) is not closed

 find $s_1 \in S, a \in A$ such that

$\text{row}(s_1a) \neq \text{row}(s)$, for all $s \in S$

$S \leftarrow S \cup \{s_1a\}$

if (S, E) is not consistent

 find $s_1, s_2 \in S, a \in A$, and $e \in E$ such that

$\text{row}(s_1) = \text{row}(s_2)$

 and $\text{row}(s_1a)(e) \neq \text{row}(s_2a)(e)$

$E \leftarrow E \cup \{ae\}$

Submit hypothesis H to the Teacher

if the Teacher replies **no**, with a counter-example t

$S \leftarrow S \cup \text{prefixes}(t)$

until the Teacher replies **yes** to H

return H

$\text{row} =$

	ϵ
ϵ	0
a	0
b	0


Closed ✓ transitions are well-defined

$\forall t \in S \cdot A \quad \exists s \in S \quad \text{row}(t) = \text{row}(s)$

Consistent ✓ determinism

$\forall s_1, s_2 \in S \quad \text{row}(s_1) = \text{row}(s_2) \implies \forall a \in A \quad \text{row}(s_1a) = \text{row}(s_2a)$

Hypothesis



transitions =

$\text{row}(s) \xrightarrow{a} \text{row}(sa)$

Counterexample = aa

$$\mathcal{L} = \{aa, bb\} \quad A = \{a, b\}$$

$S, E \leftarrow \{\epsilon\}$

repeat

while (S, E) is not closed or not consistent

if (S, E) is not closed

 find $s_1 \in S, a \in A$ such that

$\text{row}(s_1a) \neq \text{row}(s)$, for all $s \in S$

$S \leftarrow S \cup \{s_1a\}$

if (S, E) is not consistent

 find $s_1, s_2 \in S, a \in A$, and $e \in E$ such that

$\text{row}(s_1) = \text{row}(s_2)$

 and $\text{row}(s_1a)(e) \neq \text{row}(s_2a)(e)$

$E \leftarrow E \cup \{ae\}$

Submit hypothesis H to the Teacher

if the Teacher replies **no**, with a counter-example t

$S \leftarrow S \cup \text{prefixes}(t)$

until the Teacher replies **yes** to H

return H

$\text{row} =$

	ϵ
ϵ	0
a	0
b	0


Closed ✓ transitions are well-defined

$\forall t \in S \cdot A \quad \exists s \in S \quad \text{row}(t) = \text{row}(s)$

Consistent ✓ determinism

$\forall s_1, s_2 \in S \quad \text{row}(s_1) = \text{row}(s_2) \implies \forall a \in A \quad \text{row}(s_1a) = \text{row}(s_2a)$

Hypothesis



transitions =

$\text{row}(s) \xrightarrow{a} \text{row}(sa)$

Counterexample = aa

$$\mathcal{L} = \{aa, bb\} \quad A = \{a, b\}$$

$S, E \leftarrow \{\epsilon\}$

repeat

while (S, E) is not closed or not consistent

if (S, E) is not closed

 find $s_1 \in S, a \in A$ such that

$\text{row}(s_1a) \neq \text{row}(s)$, for all $s \in S$

$S \leftarrow S \cup \{s_1a\}$

if (S, E) is not consistent

 find $s_1, s_2 \in S, a \in A$, and $e \in E$ such that

$\text{row}(s_1) = \text{row}(s_2)$

 and $\text{row}(s_1a)(e) \neq \text{row}(s_2a)(e)$

$E \leftarrow E \cup \{ae\}$

Submit hypothesis H to the Teacher

if the Teacher replies **no**, with a counter-example t

$S \leftarrow S \cup \text{prefixes}(t)$

until the Teacher replies **yes** to H

return H

$\text{row} =$

	ϵ
ϵ	0
a	0
b	0


Closed ✓ transitions are well-defined

$\forall t \in S \cdot A \quad \exists s \in S \quad \text{row}(t) = \text{row}(s)$

Consistent ✓ determinism

$\forall s_1, s_2 \in S \quad \text{row}(s_1) = \text{row}(s_2) \implies \forall a \in A \quad \text{row}(s_1a) = \text{row}(s_2a)$

Hypothesis



transitions =

$\text{row}(s) \xrightarrow{a} \text{row}(sa)$

Counterexample = aa

$$\mathcal{L} = \{aa, bb\} \quad A = \{a, b\}$$

$S, E \leftarrow \{\epsilon\}$

repeat

while (S, E) is not closed or not consistent

if (S, E) is not closed

 find $s_1 \in S, a \in A$ such that

$\text{row}(s_1a) \neq \text{row}(s)$, for all $s \in S$

$S \leftarrow S \cup \{s_1a\}$

if (S, E) is not consistent

 find $s_1, s_2 \in S, a \in A$, and $e \in E$ such that

$\text{row}(s_1) = \text{row}(s_2)$

 and $\text{row}(s_1a)(e) \neq \text{row}(s_2a)(e)$

$E \leftarrow E \cup \{ae\}$

Submit hypothesis H to the Teacher

if the Teacher replies **no**, with a counter-example t

$S \leftarrow S \cup \text{prefixes}(t)$

until the Teacher replies **yes** to H

return H

$row =$

	ϵ
ϵ	0
a	0
aa	1
b	0
ab	0
aaa	0
aab	0

$$\mathcal{L} = \{aa, bb\} \quad A = \{a, b\}$$

$S, E \leftarrow \{\epsilon\}$

repeat

while (S, E) is not closed or not consistent

if (S, E) is not closed

 find $s_1 \in S, a \in A$ such that

$\text{row}(s_1a) \neq \text{row}(s)$, for all $s \in S$

$S \leftarrow S \cup \{s_1a\}$

if (S, E) is not consistent

 find $s_1, s_2 \in S, a \in A$, and $e \in E$ such that

$\text{row}(s_1) = \text{row}(s_2)$

 and $\text{row}(s_1a)(e) \neq \text{row}(s_2a)(e)$

$E \leftarrow E \cup \{ae\}$

Submit hypothesis H to the Teacher

if the Teacher replies **no**, with a counter-example t

$S \leftarrow S \cup \text{prefixes}(t)$

until the Teacher replies **yes** to H

return H

$row =$

	ϵ
ϵ	0
a	0
aa	1
b	0
ab	0
aaa	0
aab	0

$$\mathcal{L} = \{aa, bb\} \quad A = \{a, b\}$$

$S, E \leftarrow \{\epsilon\}$

repeat

while (S, E) is not closed or not consistent

if (S, E) is not closed

 find $s_1 \in S, a \in A$ such that

$\text{row}(s_1a) \neq \text{row}(s)$, for all $s \in S$

$S \leftarrow S \cup \{s_1a\}$

if (S, E) is not consistent

 find $s_1, s_2 \in S, a \in A$, and $e \in E$ such that

$\text{row}(s_1) = \text{row}(s_2)$

 and $\text{row}(s_1a)(e) \neq \text{row}(s_2a)(e)$

$E \leftarrow E \cup \{ae\}$

Submit hypothesis H to the Teacher

if the Teacher replies **no**, with a counter-example t

$S \leftarrow S \cup \text{prefixes}(t)$

until the Teacher replies **yes** to H

return H

$\text{row} =$

	ϵ
ϵ	0
a	0
aa	1
b	0
ab	0
aaa	0
aab	0

Closed

$\forall t \in S \cdot A \quad \exists s \in S \quad \text{row}(t) = \text{row}(s)$

$$\mathcal{L} = \{aa, bb\} \quad A = \{a, b\}$$

$S, E \leftarrow \{\epsilon\}$

repeat

while (S, E) is not closed or not consistent

if (S, E) is not closed

 find $s_1 \in S, a \in A$ such that

$\text{row}(s_1a) \neq \text{row}(s)$, for all $s \in S$

$S \leftarrow S \cup \{s_1a\}$

if (S, E) is not consistent

 find $s_1, s_2 \in S, a \in A$, and $e \in E$ such that

$\text{row}(s_1) = \text{row}(s_2)$

 and $\text{row}(s_1a)(e) \neq \text{row}(s_2a)(e)$

$E \leftarrow E \cup \{ae\}$

Submit hypothesis H to the Teacher

if the Teacher replies **no**, with a counter-example t

$S \leftarrow S \cup \text{prefixes}(t)$

until the Teacher replies **yes** to H

return H

$\text{row} =$

	ϵ
ϵ	0
a	0
aa	1
b	0
ab	0
aaa	0
aab	0

Closed ✓

$\forall t \in S \cdot A \quad \exists s \in S \quad \text{row}(t) = \text{row}(s)$

$$\mathcal{L} = \{aa, bb\} \quad A = \{a, b\}$$

$S, E \leftarrow \{\epsilon\}$

repeat

while (S, E) is not closed or not consistent

if (S, E) is not closed

 find $s_1 \in S, a \in A$ such that

$\text{row}(s_1a) \neq \text{row}(s)$, for all $s \in S$

$S \leftarrow S \cup \{s_1a\}$

if (S, E) is not consistent

 find $s_1, s_2 \in S, a \in A$, and $e \in E$ such that

$\text{row}(s_1) = \text{row}(s_2)$

 and $\text{row}(s_1a)(e) \neq \text{row}(s_2a)(e)$

$E \leftarrow E \cup \{ae\}$

Submit hypothesis H to the Teacher

if the Teacher replies **no**, with a counter-example t

$S \leftarrow S \cup \text{prefixes}(t)$

until the Teacher replies **yes** to H

return H

$\text{row} =$

	ϵ
ϵ	0
a	0
aa	1
b	0
ab	0
aaa	0
aab	0

Closed ✓

$\forall t \in S \cdot A \quad \exists s \in S \quad \text{row}(t) = \text{row}(s)$

Consistent

$\forall s_1, s_2 \in S \quad \text{row}(s_1) = \text{row}(s_2) \implies \forall a \in A \quad \text{row}(s_1a) = \text{row}(s_2a)$

$$\mathcal{L} = \{aa, bb\} \quad A = \{a, b\}$$

$S, E \leftarrow \{\epsilon\}$

repeat

while (S, E) is not closed or not consistent

if (S, E) is not closed

 find $s_1 \in S, a \in A$ such that

$\text{row}(s_1a) \neq \text{row}(s)$, for all $s \in S$

$S \leftarrow S \cup \{s_1a\}$

if (S, E) is not consistent

 find $s_1, s_2 \in S, a \in A$, and $e \in E$ such that

$\text{row}(s_1) = \text{row}(s_2)$

 and $\text{row}(s_1a)(e) \neq \text{row}(s_2a)(e)$

$E \leftarrow E \cup \{ae\}$

Submit hypothesis H to the Teacher

if the Teacher replies **no**, with a counter-example t

$S \leftarrow S \cup \text{prefixes}(t)$

until the Teacher replies **yes** to H

return H

$row =$

	ϵ
ϵ	0
a	0
aa	1
b	0
ab	0
aaa	0
aab	0

Closed ✓

$\forall t \in S \cdot A \quad \exists s \in S \quad \text{row}(t) = \text{row}(s)$

Consistent

$\forall s_1, s_2 \in S \quad \text{row}(s_1) = \text{row}(s_2) \implies \forall a \in A \quad \text{row}(s_1a) = \text{row}(s_2a)$

$$\mathcal{L} = \{aa, bb\} \quad A = \{a, b\}$$

$S, E \leftarrow \{\epsilon\}$

repeat

while (S, E) is not closed or not consistent

if (S, E) is not closed

 find $s_1 \in S, a \in A$ such that

$\text{row}(s_1a) \neq \text{row}(s)$, for all $s \in S$

$S \leftarrow S \cup \{s_1a\}$

if (S, E) is not consistent

 find $s_1, s_2 \in S, a \in A$, and $e \in E$ such that

$\text{row}(s_1) = \text{row}(s_2)$

 and $\text{row}(s_1a)(e) \neq \text{row}(s_2a)(e)$

$E \leftarrow E \cup \{ae\}$

Submit hypothesis H to the Teacher

if the Teacher replies **no**, with a counter-example t

$S \leftarrow S \cup \text{prefixes}(t)$

until the Teacher replies **yes** to H

return H

$\text{row} =$

	ϵ
ϵ	0
a	0
aa	1
b	0
ab	0
aaa	0
aab	0

Closed ✓

$\forall t \in S \cdot A \quad \exists s \in S \quad \text{row}(t) = \text{row}(s)$

Consistent ✗

$\forall s_1, s_2 \in S \quad \text{row}(s_1) = \text{row}(s_2) \implies \forall a \in A \quad \text{row}(s_1a) = \text{row}(s_2a)$

$$\mathcal{L} = \{aa, bb\} \quad A = \{a, b\}$$

$S, E \leftarrow \{\epsilon\}$

repeat

while (S, E) is not closed or not consistent

if (S, E) is not closed

 find $s_1 \in S, a \in A$ such that

$\text{row}(s_1a) \neq \text{row}(s)$, for all $s \in S$

$S \leftarrow S \cup \{s_1a\}$

if (S, E) is not consistent

 find $s_1, s_2 \in S, a \in A$, and $e \in E$ such that

$\text{row}(s_1) = \text{row}(s_2)$

 and $\text{row}(s_1a)(e) \neq \text{row}(s_2a)(e)$

$E \leftarrow E \cup \{ae\}$

Submit hypothesis H to the Teacher

if the Teacher replies **no**, with a counter-example t

$S \leftarrow S \cup \text{prefixes}(t)$

until the Teacher replies **yes** to H

return H

$row =$

	ϵ
ϵ	0
a	0
aa	1
b	0
ab	0
aaa	0
aab	0

Closed ✓

$\forall t \in S \cdot A \quad \exists s \in S \quad \text{row}(t) = \text{row}(s)$

Consistent ✗

$\forall s_1, s_2 \in S \quad \text{row}(s_1) = \text{row}(s_2) \implies \forall a \in A \quad \text{row}(s_1a) = \text{row}(s_2a)$

$$\mathcal{L} = \{aa, bb\} \quad A = \{a, b\}$$

$S, E \leftarrow \{\epsilon\}$

repeat

while (S, E) is not closed or not consistent

if (S, E) is not closed

 find $s_1 \in S, a \in A$ such that

$\text{row}(s_1a) \neq \text{row}(s)$, for all $s \in S$

$S \leftarrow S \cup \{s_1a\}$

if (S, E) is not consistent

 find $s_1, s_2 \in S, a \in A$, and $e \in E$ such that

$\text{row}(s_1) = \text{row}(s_2)$

 and $\text{row}(s_1a)(e) \neq \text{row}(s_2a)(e)$

$E \leftarrow E \cup \{ae\}$

Submit hypothesis H to the Teacher

if the Teacher replies **no**, with a counter-example t

$S \leftarrow S \cup \text{prefixes}(t)$

until the Teacher replies **yes** to H

return H

$row =$

	ϵ
ϵ	0
a	0
aa	1
b	0
ab	0
aaa	0
aab	0

Closed ✓

$\forall t \in S \cdot A \quad \exists s \in S \quad \text{row}(t) = \text{row}(s)$

Consistent ✗

$\forall s_1, s_2 \in S \quad \text{row}(s_1) = \text{row}(s_2) \implies \forall a \in A \quad \text{row}(s_1a) = \text{row}(s_2a)$

$$\mathcal{L} = \{aa, bb\} \quad A = \{a, b\}$$

$S, E \leftarrow \{\epsilon\}$

repeat

while (S, E) is not closed or not consistent

if (S, E) is not closed

 find $s_1 \in S, a \in A$ such that

$\text{row}(s_1a) \neq \text{row}(s)$, for all $s \in S$

$S \leftarrow S \cup \{s_1a\}$

if (S, E) is not consistent

 find $s_1, s_2 \in S, a \in A$, and $e \in E$ such that

$\text{row}(s_1) = \text{row}(s_2)$

 and $\text{row}(s_1a)(e) \neq \text{row}(s_2a)(e)$

$E \leftarrow E \cup \{ae\}$

Submit hypothesis H to the Teacher

if the Teacher replies **no**, with a counter-example t

$S \leftarrow S \cup \text{prefixes}(t)$

until the Teacher replies **yes** to H

return H

$row =$

	ϵ
ϵ	0
a	0
aa	1
b	0
ab	0
aaa	0
aab	0

$s_1 = \epsilon$

$s_2 = a$

a

a

Closed ✓

$\forall t \in S \cdot A \quad \exists s \in S \quad \text{row}(t) = \text{row}(s)$

Consistent ✗

$\forall s_1, s_2 \in S \quad \text{row}(s_1) = \text{row}(s_2) \implies \forall a \in A \quad \text{row}(s_1a) = \text{row}(s_2a)$

$$\mathcal{L} = \{aa, bb\} \quad A = \{a, b\}$$

$S, E \leftarrow \{\epsilon\}$

repeat

while (S, E) is not closed or not consistent

if (S, E) is not closed

 find $s_1 \in S, a \in A$ such that

$\text{row}(s_1a) \neq \text{row}(s), \text{ for all } s \in S$

$S \leftarrow S \cup \{s_1a\}$

if (S, E) is not consistent

 find $s_1, s_2 \in S, a \in A$, and $e \in E$ such that

$\text{row}(s_1) = \text{row}(s_2)$

 and $\text{row}(s_1a)(e) \neq \text{row}(s_2a)(e)$

$E \leftarrow E \cup \{ae\}$

Submit hypothesis H to the Teacher

if the Teacher replies **no**, with a counter-example t

$S \leftarrow S \cup \text{prefixes}(t)$

until the Teacher replies **yes** to H

return H

$row =$

ϵ	0	$s_1 = \epsilon$
ϵ	0	$s_2 = a$
a	0	a
aa	1	
b	0	
ab	0	
aaa	0	
aab	0	

Closed ✓

$\forall t \in S \cdot A \quad \exists s \in S \quad \text{row}(t) = \text{row}(s)$

Consistent ✗

$\forall s_1, s_2 \in S \quad \text{row}(s_1) = \text{row}(s_2) \implies \forall a \in A \quad \text{row}(s_1a) = \text{row}(s_2a)$

$$\mathcal{L} = \{aa, bb\} \quad A = \{a, b\}$$

$$S, E \leftarrow \{\epsilon\}$$

repeat

while (S, E) is not closed or not consistent

if (S, E) is not closed

 find $s_1 \in S, a \in A$ such that

$\text{row}(s_1a) \neq \text{row}(s)$, for all $s \in S$

$S \leftarrow S \cup \{s_1a\}$

if (S, E) is not consistent

 find $s_1, s_2 \in S, a \in A$, and $e \in E$ such that

$\text{row}(s_1) = \text{row}(s_2)$

 and $\text{row}(s_1a)(e) \neq \text{row}(s_2a)(e)$

$E \leftarrow E \cup \{ae\}$

Submit hypothesis H to the Teacher

if the Teacher replies **no**, with a counter-example t

$S \leftarrow S \cup \text{prefixes}(t)$

until the Teacher replies **yes** to H

return H

$row =$

ϵ	0	$s_1 = \epsilon$
ϵ	0	$s_2 = a$
a	0	a
aa	1	
b	0	
ab	0	
aaa	0	
aab	0	

Closed ✓

$$\forall t \in S \cdot A \quad \exists s \in S \quad \text{row}(t) = \text{row}(s)$$

Consistent ✗

$$\forall s_1, s_2 \in S \quad \text{row}(s_1) = \text{row}(s_2) \implies \forall a \in A \quad \text{row}(s_1a) = \text{row}(s_2a)$$

$$\mathcal{L} = \{aa, bb\} \quad A = \{a, b\}$$

$S, E \leftarrow \{\epsilon\}$

repeat

while (S, E) is not closed or not consistent

if (S, E) is not closed

 find $s_1 \in S, a \in A$ such that

$\text{row}(s_1a) \neq \text{row}(s)$, for all $s \in S$

$S \leftarrow S \cup \{s_1a\}$

if (S, E) is not consistent

 find $s_1, s_2 \in S, a \in A$, and $e \in E$ such that

$\text{row}(s_1) = \text{row}(s_2)$

 and $\text{row}(s_1a)(e) \neq \text{row}(s_2a)(e)$

$E \leftarrow E \cup \{ae\}$

Submit hypothesis H to the Teacher

if the Teacher replies **no**, with a counter-example t

$S \leftarrow S \cup \text{prefixes}(t)$

until the Teacher replies **yes** to H

return H

$row =$

	ϵ	a
ϵ	0	0
a	0	1
aa	1	0
b	0	0
ab	0	0
aaa	0	0
aab	0	0

$$\mathcal{L} = \{aa, bb\} \quad A = \{a, b\}$$

$S, E \leftarrow \{\epsilon\}$

repeat

while (S, E) is not closed or not consistent

if (S, E) is not closed

 find $s_1 \in S, a \in A$ such that

$\text{row}(s_1a) \neq \text{row}(s)$, for all $s \in S$

$S \leftarrow S \cup \{s_1a\}$

if (S, E) is not consistent

 find $s_1, s_2 \in S, a \in A$, and $e \in E$ such that

$\text{row}(s_1) = \text{row}(s_2)$

 and $\text{row}(s_1a)(e) \neq \text{row}(s_2a)(e)$

$E \leftarrow E \cup \{ae\}$

Submit hypothesis H to the Teacher

if the Teacher replies **no**, with a counter-example t

$S \leftarrow S \cup \text{prefixes}(t)$

until the Teacher replies **yes** to H

return H

$\text{row} =$

	ϵ	a
ϵ	0	0
a	0	1
aa	1	0
b	0	0
ab	0	0
aaa	0	0
aab	0	0

$$\mathcal{L} = \{aa, bb\} \quad A = \{a, b\}$$

$S, E \leftarrow \{\epsilon\}$

repeat

while (S, E) is not closed or not consistent

if (S, E) is not closed

 find $s_1 \in S, a \in A$ such that

$\text{row}(s_1a) \neq \text{row}(s)$, for all $s \in S$

$S \leftarrow S \cup \{s_1a\}$

if (S, E) is not consistent

 find $s_1, s_2 \in S, a \in A$, and $e \in E$ such that

$\text{row}(s_1) = \text{row}(s_2)$

 and $\text{row}(s_1a)(e) \neq \text{row}(s_2a)(e)$

$E \leftarrow E \cup \{ae\}$

Submit hypothesis H to the Teacher

if the Teacher replies **no**, with a counter-example t

$S \leftarrow S \cup \text{prefixes}(t)$

until the Teacher replies **yes** to H

return H

$\text{row} =$

	ϵ	a
ϵ	0	0
a	0	1
aa	1	0
b	0	0
ab	0	0
aaa	0	0
aab	0	0

Closed ✓

Consistent ✓

$$\mathcal{L} = \{aa, bb\} \quad A = \{a, b\}$$

$S, E \leftarrow \{\epsilon\}$

repeat

while (S, E) is not closed or not consistent

if (S, E) is not closed

 find $s_1 \in S, a \in A$ such that

$\text{row}(s_1a) \neq \text{row}(s)$, for all $s \in S$

$S \leftarrow S \cup \{s_1a\}$

if (S, E) is not consistent

 find $s_1, s_2 \in S, a \in A$, and $e \in E$ such that

$\text{row}(s_1) = \text{row}(s_2)$

 and $\text{row}(s_1a)(e) \neq \text{row}(s_2a)(e)$

$E \leftarrow E \cup \{ae\}$

Submit hypothesis H to the Teacher

if the Teacher replies **no**, with a counter-example t

$S \leftarrow S \cup \text{prefixes}(t)$

until the Teacher replies **yes** to H

return H

$\text{row} =$

	ϵ	a
ϵ	0	0
a	0	1
aa	1	0
b	0	0
ab	0	0
aaa	0	0
aab	0	0

Closed ✓

Consistent ✓

$$\mathcal{L} = \{aa, bb\} \quad A = \{a, b\}$$

$S, E \leftarrow \{\epsilon\}$

repeat

while (S, E) is not closed or not consistent

if (S, E) is not closed

 find $s_1 \in S, a \in A$ such that

$\text{row}(s_1a) \neq \text{row}(s)$, for all $s \in S$

$S \leftarrow S \cup \{s_1a\}$

if (S, E) is not consistent

 find $s_1, s_2 \in S, a \in A$, and $e \in E$ such that

$\text{row}(s_1) = \text{row}(s_2)$

 and $\text{row}(s_1a)(e) \neq \text{row}(s_2a)(e)$

$E \leftarrow E \cup \{ae\}$

Submit hypothesis H to the Teacher

if the Teacher replies **no**, with a counter-example t

$S \leftarrow S \cup \text{prefixes}(t)$

until the Teacher replies **yes** to H

return H


$\text{row} =$

	ϵ	a
ϵ	0	0
a	0	1
aa	1	0
b	0	0
ab	0	0
aaa	0	0
aab	0	0

Closed ✓

Consistent ✓

Hypothesis



$$\mathcal{L} = \{aa, bb\} \quad A = \{a, b\}$$

$S, E \leftarrow \{\epsilon\}$

repeat

while (S, E) is not closed or not consistent

if (S, E) is not closed

 find $s_1 \in S, a \in A$ such that

$\text{row}(s_1a) \neq \text{row}(s)$, for all $s \in S$

$S \leftarrow S \cup \{s_1a\}$

if (S, E) is not consistent

 find $s_1, s_2 \in S, a \in A$, and $e \in E$ such that

$\text{row}(s_1) = \text{row}(s_2)$

 and $\text{row}(s_1a)(e) \neq \text{row}(s_2a)(e)$

$E \leftarrow E \cup \{ae\}$

Submit hypothesis H to the Teacher

if the Teacher replies **no**, with a counter-example t

$S \leftarrow S \cup \text{prefixes}(t)$

until the Teacher replies **yes** to H

return H


$\text{row} =$

	ϵ	a
ϵ	0	0
a	0	1
aa	1	0
b	0	0
ab	0	0
aaa	0	0
aab	0	0

Closed ✓

Consistent ✓

Hypothesis



$$\mathcal{L} = \{aa, bb\} \quad A = \{a, b\}$$

$S, E \leftarrow \{\epsilon\}$

repeat

while (S, E) is not closed or not consistent

if (S, E) is not closed

 find $s_1 \in S, a \in A$ such that

$\text{row}(s_1a) \neq \text{row}(s)$, for all $s \in S$

$S \leftarrow S \cup \{s_1a\}$

if (S, E) is not consistent

 find $s_1, s_2 \in S, a \in A$, and $e \in E$ such that

$\text{row}(s_1) = \text{row}(s_2)$

 and $\text{row}(s_1a)(e) \neq \text{row}(s_2a)(e)$

$E \leftarrow E \cup \{ae\}$

Submit hypothesis H to the Teacher

if the Teacher replies **no**, with a counter-example t

$S \leftarrow S \cup \text{prefixes}(t)$

until the Teacher replies **yes** to H

return H


$\text{row} =$

	ϵ	a
ϵ	0	0
a	0	1
aa	1	0
b	0	0
ab	0	0
aaa	0	0
aab	0	0

Closed ✓

Consistent ✓

Hypothesis



Counterexample = bb

$$\mathcal{L} = \{aa, bb\} \quad A = \{a, b\}$$

$S, E \leftarrow \{\epsilon\}$

repeat

while (S, E) is not closed or not consistent

if (S, E) is not closed

 find $s_1 \in S, a \in A$ such that

$\text{row}(s_1a) \neq \text{row}(s)$, for all $s \in S$

$S \leftarrow S \cup \{s_1a\}$

if (S, E) is not consistent

 find $s_1, s_2 \in S, a \in A$, and $e \in E$ such that

$\text{row}(s_1) = \text{row}(s_2)$

 and $\text{row}(s_1a)(e) \neq \text{row}(s_2a)(e)$

$E \leftarrow E \cup \{ae\}$

Submit hypothesis H to the Teacher

if the Teacher replies **no**, with a counter-example t

$S \leftarrow S \cup \text{prefixes}(t)$

until the Teacher replies **yes** to H

return H

$row =$

	ϵ	a
ϵ	0	0
a	0	1
aa	1	0
b	0	0
bb	1	0
ab	0	0
aaa	0	0
aab	0	0
ba	0	0
bba	0	0
bbb	0	0

$$\mathcal{L} = \{aa, bb\} \quad A = \{a, b\}$$

$S, E \leftarrow \{\epsilon\}$

repeat

while (S, E) is not closed or not consistent

if (S, E) is not closed

 find $s_1 \in S, a \in A$ such that

$\text{row}(s_1a) \neq \text{row}(s)$, for all $s \in S$

$S \leftarrow S \cup \{s_1a\}$

if (S, E) is not consistent

 find $s_1, s_2 \in S, a \in A$, and $e \in E$ such that

$\text{row}(s_1) = \text{row}(s_2)$

and $\text{row}(s_1a)(e) \neq \text{row}(s_2a)(e)$

$E \leftarrow E \cup \{ae\}$

Submit hypothesis H to the Teacher

if the Teacher replies **no**, with a counter-example t

$S \leftarrow S \cup \text{prefixes}(t)$

until the Teacher replies **yes** to H

return H

$row =$

	ϵ	a
ϵ	0	0
a	0	1
aa	1	0
b	0	0
bb	1	0
ab	0	0
aaa	0	0
aab	0	0
ba	0	0
bba	0	0
bbb	0	0

$$\mathcal{L} = \{aa, bb\} \quad A = \{a, b\}$$

$S, E \leftarrow \{\epsilon\}$

repeat

while (S, E) is not closed or not consistent

if (S, E) is not closed

 find $s_1 \in S, a \in A$ such that

$\text{row}(s_1a) \neq \text{row}(s)$, for all $s \in S$

$S \leftarrow S \cup \{s_1a\}$

if (S, E) is not consistent

 find $s_1, s_2 \in S, a \in A$, and $e \in E$ such that

$\text{row}(s_1) = \text{row}(s_2)$

and $\text{row}(s_1a)(e) \neq \text{row}(s_2a)(e)$

$E \leftarrow E \cup \{ae\}$

Submit hypothesis H to the Teacher

if the Teacher replies **no**, with a counter-example t

$S \leftarrow S \cup \text{prefixes}(t)$

until the Teacher replies **yes** to H

return H

$\text{row} =$

	ϵ	a
ϵ	0	0
a	0	1
aa	1	0
b	0	0
bb	1	0
ab	0	0
aaa	0	0
aab	0	0
ba	0	0
bba	0	0
bbb	0	0

Closed ✓

Consistent ✗

$$\mathcal{L} = \{aa, bb\} \quad A = \{a, b\}$$

$S, E \leftarrow \{\epsilon\}$

repeat

while (S, E) is not closed or not consistent

if (S, E) is not closed

 find $s_1 \in S, a \in A$ such that

$\text{row}(s_1a) \neq \text{row}(s)$, for all $s \in S$

$S \leftarrow S \cup \{s_1a\}$

if (S, E) is not consistent

 find $s_1, s_2 \in S, a \in A$, and $e \in E$ such that

$\text{row}(s_1) = \text{row}(s_2)$

 and $\text{row}(s_1a)(e) \neq \text{row}(s_2a)(e)$

$E \leftarrow E \cup \{ae\}$

Submit hypothesis H to the Teacher

if the Teacher replies **no**, with a counter-example t

$S \leftarrow S \cup \text{prefixes}(t)$

until the Teacher replies **yes** to H

return H

$row =$

	ϵ	a
ϵ	0	0
a	0	1
aa	1	0
b	0	0
bb	1	0
ab	0	0
aaa	0	0
aab	0	0
ba	0	0
bba	0	0
bbb	0	0

Closed ✓

Consistent ✗

$$\mathcal{L} = \{aa, bb\} \quad A = \{a, b\}$$

$S, E \leftarrow \{\epsilon\}$

repeat

while (S, E) is not closed or not consistent

if (S, E) is not closed

 find $s_1 \in S, a \in A$ such that

$\text{row}(s_1a) \neq \text{row}(s)$, for all $s \in S$

$S \leftarrow S \cup \{s_1a\}$

if (S, E) is not consistent

 find $s_1, s_2 \in S, a \in A$, and $e \in E$ such that

$\text{row}(s_1) = \text{row}(s_2)$

 and $\text{row}(s_1a)(e) \neq \text{row}(s_2a)(e)$

$E \leftarrow E \cup \{ae\}$

Submit hypothesis H to the Teacher

if the Teacher replies **no**, with a counter-example t

$S \leftarrow S \cup \text{prefixes}(t)$

until the Teacher replies **yes** to H

return H

$row =$

	ϵ	a
ϵ	0	0
a	0	1
aa	1	0
b	0	0
bb	1	0
ab	0	0
aaa	0	0
aab	0	0
ba	0	0
bba	0	0
bbb	0	0

b

b

Closed ✓

Consistent ✗

$$\mathcal{L} = \{aa, bb\} \quad A = \{a, b\}$$

$S, E \leftarrow \{\epsilon\}$

repeat

while (S, E) is not closed or not consistent

if (S, E) is not closed

 find $s_1 \in S, a \in A$ such that

$\text{row}(s_1a) \neq \text{row}(s)$, for all $s \in S$

$S \leftarrow S \cup \{s_1a\}$

if (S, E) is not consistent

 find $s_1, s_2 \in S, a \in A$, and $e \in E$ such that

$\text{row}(s_1) = \text{row}(s_2)$

 and $\text{row}(s_1a)(e) \neq \text{row}(s_2a)(e)$

$E \leftarrow E \cup \{ae\}$

Submit hypothesis H to the Teacher

if the Teacher replies **no**, with a counter-example t

$S \leftarrow S \cup \text{prefixes}(t)$

until the Teacher replies **yes** to H

return H

$row =$

	ϵ	a
ϵ	0	0
a	0	1
aa	1	0
b	0	0
bb	1	0
ab	0	0
aaa	0	0
aab	0	0
ba	0	0
bba	0	0
bbb	0	0

Closed ✓

Consistent ✗

$$\mathcal{L} = \{aa, bb\} \quad A = \{a, b\}$$

$S, E \leftarrow \{\epsilon\}$

repeat

while (S, E) is not closed or not consistent

if (S, E) is not closed

 find $s_1 \in S, a \in A$ such that

$\text{row}(s_1a) \neq \text{row}(s)$, for all $s \in S$

$S \leftarrow S \cup \{s_1a\}$

if (S, E) is not consistent

 find $s_1, s_2 \in S, a \in A$, and $e \in E$ such that

$\text{row}(s_1) = \text{row}(s_2)$

 and $\text{row}(s_1a)(e) \neq \text{row}(s_2a)(e)$

$E \leftarrow E \cup \{ae\}$

Submit hypothesis H to the Teacher

if the Teacher replies **no**, with a counter-example t

$S \leftarrow S \cup \text{prefixes}(t)$

until the Teacher replies **yes** to H

return H

$\text{row} =$

	ϵ	a	b
ϵ	0	0	0
a	0	1	0
aa	1	0	0
b	0	0	1
bb	1	0	0
ab	0	0	0
aaa	0	0	0
aab	0	0	0
ba	0	0	0
bba	0	0	0
bbb	0	0	0

$$\mathcal{L} = \{aa, bb\} \quad A = \{a, b\}$$

$S, E \leftarrow \{\epsilon\}$

repeat

while (S, E) is not closed or not consistent

if (S, E) is not closed

 find $s_1 \in S, a \in A$ such that

$\text{row}(s_1a) \neq \text{row}(s)$, for all $s \in S$

$S \leftarrow S \cup \{s_1a\}$

if (S, E) is not consistent

 find $s_1, s_2 \in S, a \in A$, and $e \in E$ such that

$\text{row}(s_1) = \text{row}(s_2)$

 and $\text{row}(s_1a)(e) \neq \text{row}(s_2a)(e)$

$E \leftarrow E \cup \{ae\}$

Submit hypothesis H to the Teacher

if the Teacher replies **no**, with a counter-example t

$S \leftarrow S \cup \text{prefixes}(t)$

until the Teacher replies **yes** to H

return H

$\text{row} =$

	ϵ	a	b
ϵ	0	0	0
a	0	1	0
aa	1	0	0
b	0	0	1
bb	1	0	0
ab	0	0	0
aaa	0	0	0
aab	0	0	0
ba	0	0	0
bba	0	0	0
bbb	0	0	0

$$\mathcal{L} = \{aa, bb\} \quad A = \{a, b\}$$

$S, E \leftarrow \{\epsilon\}$

repeat

while (S, E) is not closed or not consistent

if (S, E) is not closed

 find $s_1 \in S, a \in A$ such that

$\text{row}(s_1a) \neq \text{row}(s)$, for all $s \in S$

$S \leftarrow S \cup \{s_1a\}$

if (S, E) is not consistent

 find $s_1, s_2 \in S, a \in A$, and $e \in E$ such that

$\text{row}(s_1) = \text{row}(s_2)$

 and $\text{row}(s_1a)(e) \neq \text{row}(s_2a)(e)$

$E \leftarrow E \cup \{ae\}$

Submit hypothesis H to the Teacher

if the Teacher replies **no**, with a counter-example t

$S \leftarrow S \cup \text{prefixes}(t)$

until the Teacher replies **yes** to H

return H

$\text{row} =$

	ϵ	a	b
ϵ	0	0	0
a	0	1	0
aa	1	0	0
b	0	0	1
bb	1	0	0
ab	0	0	0
aaa	0	0	0
aab	0	0	0
ba	0	0	0
bba	0	0	0
bbb	0	0	0

Closed ✓

Consistent ✓

$$\mathcal{L} = \{aa, bb\} \quad A = \{a, b\}$$

$S, E \leftarrow \{\epsilon\}$

repeat

while (S, E) is not closed or not consistent

if (S, E) is not closed

 find $s_1 \in S, a \in A$ such that

$\text{row}(s_1a) \neq \text{row}(s)$, for all $s \in S$

$S \leftarrow S \cup \{s_1a\}$

if (S, E) is not consistent

 find $s_1, s_2 \in S, a \in A$, and $e \in E$ such that

$\text{row}(s_1) = \text{row}(s_2)$

 and $\text{row}(s_1a)(e) \neq \text{row}(s_2a)(e)$

$E \leftarrow E \cup \{ae\}$

Submit hypothesis H to the Teacher

if the Teacher replies **no**, with a counter-example t

$S \leftarrow S \cup \text{prefixes}(t)$

until the Teacher replies **yes** to H

return H

$\text{row} =$

	ϵ	a	b
ϵ	0	0	0
a	0	1	0
aa	1	0	0
b	0	0	1
bb	1	0	0
ab	0	0	0
aaa	0	0	0
aab	0	0	0
ba	0	0	0
bba	0	0	0
bbb	0	0	0

Closed ✓

Consistent ✓

$$\mathcal{L} = \{aa, bb\} \quad A = \{a, b\}$$

$S, E \leftarrow \{\epsilon\}$

repeat

while (S, E) is not closed or not consistent

if (S, E) is not closed

 find $s_1 \in S, a \in A$ such that

$\text{row}(s_1a) \neq \text{row}(s)$, for all $s \in S$

$S \leftarrow S \cup \{s_1a\}$

if (S, E) is not consistent

 find $s_1, s_2 \in S, a \in A$, and $e \in E$ such that

$\text{row}(s_1) = \text{row}(s_2)$

 and $\text{row}(s_1a)(e) \neq \text{row}(s_2a)(e)$

$E \leftarrow E \cup \{ae\}$

Submit hypothesis H to the Teacher

if the Teacher replies **no**, with a counter-example t

$S \leftarrow S \cup \text{prefixes}(t)$

until the Teacher replies **yes** to H

return H


$\text{row} =$

	ϵ	a	b
ϵ	0	0	0
a	0	1	0
aa	1	0	0
b	0	0	1
bb	1	0	0
ab	0	0	0
aaa	0	0	0
aab	0	0	0
ba	0	0	0
bba	0	0	0
bbb	0	0	0

Closed ✓

Consistent ✓

Hypothesis



$$\mathcal{L} = \{aa, bb\} \quad A = \{a, b\}$$

$S, E \leftarrow \{\epsilon\}$

repeat

while (S, E) is not closed or not consistent

if (S, E) is not closed

 find $s_1 \in S, a \in A$ such that

$\text{row}(s_1a) \neq \text{row}(s)$, for all $s \in S$

$S \leftarrow S \cup \{s_1a\}$

if (S, E) is not consistent

 find $s_1, s_2 \in S, a \in A$, and $e \in E$ such that

$\text{row}(s_1) = \text{row}(s_2)$

 and $\text{row}(s_1a)(e) \neq \text{row}(s_2a)(e)$

$E \leftarrow E \cup \{ae\}$

Submit hypothesis H to the Teacher

if the Teacher replies **no**, with a counter-example t

$S \leftarrow S \cup \text{prefixes}(t)$

until the Teacher replies **yes** to H

return H

$\text{row} =$


	ϵ	a	b
ϵ	0	0	0
a	0	1	0
aa	1	0	0
b	0	0	1
bb	1	0	0
ab	0	0	0
aaa	0	0	0
aab	0	0	0
ba	0	0	0
bba	0	0	0
bbb	0	0	0

Closed ✓

Consistent ✓

Hypothesis

Counterexample = $babb$



$$\mathcal{L} = \{aa, bb\} \quad A = \{a, b\}$$

$S, E \leftarrow \{\epsilon\}$

repeat

while (S, E) is not closed or not consistent

if (S, E) is not closed

 find $s_1 \in S, a \in A$ such that

$\text{row}(s_1a) \neq \text{row}(s)$, for all $s \in S$

$S \leftarrow S \cup \{s_1a\}$

if (S, E) is not consistent

 find $s_1, s_2 \in S, a \in A$, and $e \in E$ such that

$\text{row}(s_1) = \text{row}(s_2)$

 and $\text{row}(s_1a)(e) \neq \text{row}(s_2a)(e)$

$E \leftarrow E \cup \{ae\}$

Submit hypothesis H to the Teacher

if the Teacher replies **no**, with a counter-example t

$S \leftarrow S \cup \text{prefixes}(t)$

until the Teacher replies **yes** to H

return H

$$\mathcal{L} = \{aa, bb\} \quad A = \{a, b\}$$

$S, E \leftarrow \{\epsilon\}$

repeat

while (S, E) is not closed or not consistent

if (S, E) is not closed

 find $s_1 \in S, a \in A$ such that

$\text{row}(s_1a) \neq \text{row}(s)$, for all $s \in S$

$S \leftarrow S \cup \{s_1a\}$

if (S, E) is not consistent

 find $s_1, s_2 \in S, a \in A$, and $e \in E$ such that

$\text{row}(s_1) = \text{row}(s_2)$

 and $\text{row}(s_1a)(e) \neq \text{row}(s_2a)(e)$

$E \leftarrow E \cup \{ae\}$

Submit hypothesis H to the Teacher

if the Teacher replies **no**, with a counter-example t

$S \leftarrow S \cup \text{prefixes}(t)$

until the Teacher replies **yes** to H

return H

$$\mathcal{L} = \{aa, bb\} \quad A = \{a, b\}$$

$$S, E \leftarrow \{\epsilon\}$$

repeat

while (S, E) is not closed or not consistent

if (S, E) is not closed

 find $s_1 \in S, a \in A$ such that

$\text{row}(s_1a) \neq \text{row}(s)$, for all $s \in S$

$S \leftarrow S \cup \{s_1a\}$

if (S, E) is not consistent

 find $s_1, s_2 \in S, a \in A$, and $e \in E$ such that

$\text{row}(s_1) = \text{row}(s_2)$

 and $\text{row}(s_1a)(e) \neq \text{row}(s_2a)(e)$

$E \leftarrow E \cup \{ae\}$


Submit hypothesis H to the Teacher

if the Teacher replies **no**, with a counter-example t

$S \leftarrow S \cup \text{prefixes}(t)$

until the Teacher replies **yes** to H

return H



Pros of L^* ...

simple is
beautiful

&

POWERFUL

Applications : Hardware verification, security/network protocols...

Generalizations : Mealy machines, I/O automata, ...

... and shortcomings

L^* learns **control-flow**

What if program model needs to express **data-flow**?

operations on **data values**

comparisons between data values



Automata over infinite alphabets (nominal automata)

Automata over infinite alphabets (nominal automata)

$A = \{a, b, c, d, \dots\}$ **infinite alphabet**


$\mathcal{L} = \{aa, bb, cc, dd, \dots\}$

Automata over infinite alphabets (nominal automata)

$$A = \{a, b, c, d, \dots\}$$

infinite alphabet

$$\mathcal{L} = \{aa, bb, cc, dd, \dots\}$$




infinite automaton

Automata over infinite alphabets (nominal automata)


$$A = \{a, b, c, d, \dots\}$$

infinite alphabet

$$\mathcal{L} = \{aa, bb, cc, dd, \dots\}$$



infinite automaton



but with a finite representation

How to learn them?

How to learn them?

Ad-hoc algorithm? **NO!**

How to learn them?

Ad-hoc algorithm? **NO!**

Challenges:

- table needs to be **infinite**
- code operates on **infinite sets**

$$\boxed{\forall t \in S \cdot A \quad \exists s \in S \quad \text{row}(t) = \text{row}(s).}$$

$$\boxed{\forall s_1, s_2 \in S \quad \text{row}(s_1) = \text{row}(s_2) \implies \forall a \in A \quad \text{row}(s_1 a) = \text{row}(s_2 a)}$$

How to learn them?

Ad-hoc algorithm? **NO!**

Challenges:

- table needs to be **infinite**
- code operates on **infinite sets**

$$\forall t \in S \cdot A \quad \exists s \in S \quad \text{row}(t) = \text{row}(s).$$

Everything is “finitely representable”

$$\forall s_1, s_2 \in S \quad \text{row}(s_1) = \text{row}(s_2) \implies \forall a \in A \quad \text{row}(s_1 a) = \text{row}(s_2 a)$$

Theory of nominal sets

Model of data

\mathbb{D}

Domain of data + relations

$Sym(\mathbb{D})$

group of bijective $\pi: \mathbb{D} \rightarrow \mathbb{D}$ **(permutations)**

E.g.

$(A, =)$ + **bijective functions**

$(\mathbb{Q}, <)$ + **bijective monotonic functions**

Nominal set

$$(X, \cdot)$$

Nominal set

$$(X, \underline{\cdot})$$

group action of $Sym(\mathbb{D})$ on X

$$Sym(\mathbb{D}) \times X \rightarrow X$$

$$\pi \cdot x = x'$$

Nominal set

$$(X, \underline{\cdot})$$

group action of $Sym(\mathbb{D})$ on X

$$Sym(\mathbb{D}) \times X \rightarrow X$$

$$\pi \cdot x = x'$$

with **finite support** $\text{supp}(x) \subseteq \mathbb{D}$

π acts as the identity on $\text{supp}(x) \implies \pi \cdot x = x$

Nominal set

$$(X, \underline{\cdot})$$

group action of $Sym(\mathbb{D})$ on X

$$Sym(\mathbb{D}) \times X \rightarrow X$$

$$\pi \cdot x = x'$$

with **finite support** $\text{supp}(x) \subseteq \mathbb{D}$

π acts as the identity on $\text{supp}(x) \implies \pi \cdot x = x$

Data values “stored” in x


Example: nominal set of strings

$$(A^*, \cdot)$$


$$\pi \cdot abcd = \pi(a)\pi(b)\pi(c)\pi(d)$$

$$\text{supp}(abcd) = \{a, b, c, d\}$$

Nominal automaton



Nominal automaton



Nominal set of states (Q, \cdot)

$$\{q_0\} \cup \{q_a \mid a \in A\} \cup \{q_3\} \cup \{q_4\}$$


$$\pi \cdot q_0 = q_0$$

$$\pi \cdot q_3 = q_3$$

$$\pi \cdot q_4 = q_4$$

$$\pi \cdot q_x = q_{\pi(x)} \quad x \in A$$

Nominal automaton



Nominal set of states (Q, \cdot)

$$\{q_0\} \cup \{q_a \mid a \in A\} \cup \{q_3\} \cup \{q_4\}$$

$$\pi \cdot q_0 = q_0$$

$$\pi \cdot q_3 = q_3$$

$$\pi \cdot q_4 = q_4$$

$$\pi \cdot q_x = q_{\pi(x)} \quad x \in A$$

$$\text{supp}(q_x) = \{x\} \quad x \in A$$

$$\text{supp}(q_0) = \text{supp}(q_3) = \text{supp}(q_4) = \emptyset$$

Equivariant functions


$$f: (X, \cdot) \rightarrow (Y, \cdot)$$

$$f(\pi \cdot x) = \pi \cdot f(x)$$


Nominal automaton

Equivariant transition function

$$\delta: (Q \times A, \cdot) \rightarrow (Q, \cdot)$$



Nominal automaton




Equivariant transition function

$$\delta: (Q \times A, \cdot) \rightarrow (Q, \cdot)$$

$$q_0 \xrightarrow{a} q_a$$

Nominal automaton




Equivariant transition function

$$\delta: (Q \times A, \cdot) \rightarrow (Q, \cdot)$$

$$(a\ b)\ q_0 \xrightarrow{(a\ b)a} q_a$$

Nominal automaton




Equivariant transition function

$$\delta: (Q \times A, \cdot) \rightarrow (Q, \cdot)$$

$$(a\ b)\ q_0 \xrightarrow{(a\ b)a} (a\ b)\ q_a$$

Nominal automaton



Equivariant transition function

$$\delta: (Q \times A, \cdot) \rightarrow (Q, \cdot)$$

$$\begin{array}{c} (a\ b)\ q_0 \xrightarrow{(a\ b)a} (a\ b)q_a \\ = \\ q_0 \xrightarrow{b} q_b \end{array}$$


“closed under permutations”

Orbits

$$\text{orb}(x) = \{\pi \cdot x \mid \pi \in \text{Sym}(\mathbb{D})\}$$

(X, \cdot) **orbit-finite** whenever $\{\text{orb}(x) \mid x \in X\}$ is finite

Nominal automaton



$$\{q_0\} \cup \{q_a \mid a \in A\} \cup \{q_3\} \cup \{q_4\}$$


$$\pi \cdot q_0 = q_0$$

$$\pi \cdot q_3 = q_3$$

$$\pi \cdot q_4 = q_4$$

$$\pi \cdot q_x = q_{\pi(x)} \quad x \in A$$

Nominal automaton



$$\underline{\{q_0\}} \cup \underline{\{q_a \mid a \in A\}} \cup \underline{\{q_3\}} \cup \underline{\{q_4\}}$$


4 orbits

$$\pi \cdot q_0 = q_0$$

$$\pi \cdot q_3 = q_3$$

$$\pi \cdot q_4 = q_4$$


$$\pi \cdot q_x = q_{\pi(x)} \quad x \in A$$



$$\underline{\{q_0\}} \cup \underline{\{q_a \mid a \in A\}} \cup \underline{\{q_3\}} \cup \underline{\{q_4\}}$$

4 orbits

**one representative
per orbit**



Nominal \mathbb{L}^*

S, E

finite sets

$row: S \cup S \times A \rightarrow 2^E$

function

Nominal \mathcal{L}^*

S, E

orbit-finite nominal sets

$row: S \cup S \times A \rightarrow 2^E$

equivariant function

Example

$$\mathcal{L} = \{aa, bb, cc, dd, \dots\} \quad A = \{a, b, c, d, \dots\}$$

$$\mathcal{L} = \{aa, bb, cc, dd, \dots\} \quad A = \{a, b, c, d, \dots\}$$

$S, E \leftarrow \{\epsilon\}$

repeat

while (S, E) is not closed or not consistent

if (S, E) is not closed

 find $s_1 \in S, a \in A$ such that

$\text{row}(s_1a) \neq \text{row}(s)$, for all $s \in S$

$S \leftarrow S \cup \{s_1a\}$

if (S, E) is not consistent

 find $s_1, s_2 \in S, a \in A$, and $e \in E$ such that

$\text{row}(s_1) = \text{row}(s_2)$

 and $\text{row}(s_1a)(e) \neq \text{row}(s_2a)(e)$

$E \leftarrow E \cup \{ae\}$

Submit hypothesis H to the Teacher

if the Teacher replies **no**, with a counter-example t

$S \leftarrow S \cup \text{prefixes}(t)$

until the Teacher replies **yes** to H

return H

$$\mathcal{L} = \{aa, bb, cc, dd, \dots\} \quad A = \{a, b, c, d, \dots\}$$

$$S, E \leftarrow \{\epsilon\}$$

repeat

while (S, E) is not closed or not consistent

if (S, E) is not closed

 find $s_1 \in S, a \in A$ such that

$\text{row}(s_1a) \neq \text{row}(s)$, for all $s \in S$

$S \leftarrow S \cup \{s_1a\}$

if (S, E) is not consistent

 find $s_1, s_2 \in S, a \in A$, and $e \in E$ such that

$\text{row}(s_1) = \text{row}(s_2)$

 and $\text{row}(s_1a)(e) \neq \text{row}(s_2a)(e)$

$E \leftarrow E \cup \{ae\}$

Submit hypothesis H to the Teacher

if the Teacher replies **no**, with a counter-example t

$S \leftarrow S \cup \text{prefixes}(t)$

until the Teacher replies **yes** to H

return H

$$\mathcal{L} = \{aa, bb, cc, dd, \dots\} \quad A = \{a, b, c, d, \dots\}$$

$$S, E \leftarrow \{\epsilon\}$$

repeat

while (S, E) is not closed or not consistent

if (S, E) is not closed

 find $s_1 \in S, a \in A$ such that

$\text{row}(s_1a) \neq \text{row}(s)$, for all $s \in S$

$S \leftarrow S \cup \{s_1a\}$

if (S, E) is not consistent

 find $s_1, s_2 \in S, a \in A$, and $e \in E$ such that

$\text{row}(s_1) = \text{row}(s_2)$

 and $\text{row}(s_1a)(e) \neq \text{row}(s_2a)(e)$

$E \leftarrow E \cup \{ae\}$

Submit hypothesis H to the Teacher

if the Teacher replies **no**, with a counter-example t

$S \leftarrow S \cup \text{prefixes}(t)$

until the Teacher replies **yes** to H

return H

$\text{row} =$

	ϵ
ϵ	0
a	0
b	0
c	0
d	0
...	...

$$\mathcal{L} = \{aa, bb, cc, dd, \dots\} \quad A = \{a, b, c, d, \dots\}$$

$$S, E \leftarrow \{\epsilon\}$$

repeat

while (S, E) is not closed or not consistent

if (S, E) is not closed

 find $s_1 \in S, a \in A$ such that

$\text{row}(s_1a) \neq \text{row}(s)$, for all $s \in S$

$S \leftarrow S \cup \{s_1a\}$

if (S, E) is not consistent

 find $s_1, s_2 \in S, a \in A$, and $e \in E$ such that

$\text{row}(s_1) = \text{row}(s_2)$

 and $\text{row}(s_1a)(e) \neq \text{row}(s_2a)(e)$

$E \leftarrow E \cup \{ae\}$

Submit hypothesis H to the Teacher

if the Teacher replies **no**, with a counter-example t

$S \leftarrow S \cup \text{prefixes}(t)$

until the Teacher replies **yes** to H

return H

$\text{row} =$

	ϵ
ϵ	0
a	0
b	0
c	0
d	0
...	...

infinitely-many rows!

$$\mathcal{L} = \{aa, bb, cc, dd, \dots\} \quad A = \{a, b, c, d, \dots\}$$

$$S, E \leftarrow \{\epsilon\}$$

repeat

while (S, E) is not closed or not consistent

if (S, E) is not closed

 find $s_1 \in S, a \in A$ such that

$\text{row}(s_1a) \neq \text{row}(s)$, for all $s \in S$

$S \leftarrow S \cup \{s_1a\}$

if (S, E) is not consistent

 find $s_1, s_2 \in S, a \in A$, and $e \in E$ such that

$\text{row}(s_1) = \text{row}(s_2)$

 and $\text{row}(s_1a)(e) \neq \text{row}(s_2a)(e)$

$E \leftarrow E \cup \{ae\}$

Submit hypothesis H to the Teacher

if the Teacher replies **no**, with a counter-example t

$S \leftarrow S \cup \text{prefixes}(t)$

until the Teacher replies **yes** to H

return H

$$\text{row} =$$

	ϵ
ϵ	0
a	0
b	0
c	0
d	0
...	...

infinitely-many rows!

$$\text{row}(\pi \cdot s)(\pi \cdot e) = \text{row}(s)(e)$$

$$\mathcal{L} = \{aa, bb, cc, dd, \dots\} \quad A = \{a, b, c, d, \dots\}$$

$$S, E \leftarrow \{\epsilon\}$$

repeat

while (S, E) is not closed or not consistent

if (S, E) is not closed

 find $s_1 \in S, a \in A$ such that

$\text{row}(s_1a) \neq \text{row}(s)$, for all $s \in S$

$S \leftarrow S \cup \{s_1a\}$

if (S, E) is not consistent

 find $s_1, s_2 \in S, a \in A$, and $e \in E$ such that

$\text{row}(s_1) = \text{row}(s_2)$

 and $\text{row}(s_1a)(e) \neq \text{row}(s_2a)(e)$

$E \leftarrow E \cup \{ae\}$

Submit hypothesis H to the Teacher

if the Teacher replies **no**, with a counter-example t

$S \leftarrow S \cup \text{prefixes}(t)$

until the Teacher replies **yes** to H

return H

$$\text{row} =$$

	ϵ
ϵ	0
a	0
b	0
c	0
d	0
...	...

$\xleftarrow{\text{row}((a b)a)((a b)\epsilon)}$
 $\xleftarrow{\text{row}((a c)a)((a c)\epsilon)}$
 $\xleftarrow{\text{row}((a d)a)((a d)\epsilon)}$

infinitely-many rows!

$$\text{row}(\pi \cdot s)(\pi \cdot e) = \text{row}(s)(e)$$

$$\mathcal{L} = \{aa, bb, cc, dd, \dots\} \quad A = \{a, b, c, d, \dots\}$$

$$S, E \leftarrow \{\epsilon\}$$

repeat

while (S, E) is not closed or not consistent

if (S, E) is not closed

 find $s_1 \in S, a \in A$ such that

$\text{row}(s_1a) \neq \text{row}(s)$, for all $s \in S$

$S \leftarrow S \cup \{s_1a\}$

if (S, E) is not consistent

 find $s_1, s_2 \in S, a \in A$, and $e \in E$ such that

$\text{row}(s_1) = \text{row}(s_2)$

 and $\text{row}(s_1a)(e) \neq \text{row}(s_2a)(e)$

$E \leftarrow E \cup \{ae\}$

Submit hypothesis H to the Teacher

if the Teacher replies **no**, with a counter-example t

$S \leftarrow S \cup \text{prefixes}(t)$

until the Teacher replies **yes** to H

return H

$$\text{row} =$$

	ϵ
ϵ	0
a	0
b	0
c	0
d	0
...	...

$\xleftarrow{\text{row}((a b)a)((a b)\epsilon)}$
 $\xleftarrow{\text{row}((a c)a)((a c)\epsilon)}$
 $\xleftarrow{\text{row}((a d)a)((a d)\epsilon)}$

infinitely-many rows!

$$\text{row}(\pi \cdot s)(\pi \cdot e) = \text{row}(s)(e)$$

finite representation by keeping only
orbit representatives of rows

$$\mathcal{L} = \{aa, bb, cc, dd, \dots\} \quad A = \{a, b, c, d, \dots\}$$

$$S, E \leftarrow \{\epsilon\}$$

repeat

while (S, E) is not closed or not consistent

if (S, E) is not closed

 find $s_1 \in S, a \in A$ such that

$\text{row}(s_1a) \neq \text{row}(s)$, for all $s \in S$

$$S \leftarrow S \cup \{s_1a\}$$

if (S, E) is not consistent

 find $s_1, s_2 \in S, a \in A$, and $e \in E$ such that

$\text{row}(s_1) = \text{row}(s_2)$

 and $\text{row}(s_1a)(e) \neq \text{row}(s_2a)(e)$

$$E \leftarrow E \cup \{ae\}$$

Submit hypothesis H to the Teacher

if the Teacher replies **no**, with a counter-example t

$$S \leftarrow S \cup \text{prefixes}(t)$$

until the Teacher replies **yes** to H

return H

$$\text{row} =$$

	ϵ
ϵ	0
a	0

infinitely-many rows!

$$\text{row}(\pi \cdot s)(\pi \cdot e) = \text{row}(s)(e)$$

finite representation by keeping only
orbit representatives of rows

$$\mathcal{L} = \{aa, bb, cc, dd, \dots\} \quad A = \{a, b, c, d, \dots\}$$

$S, E \leftarrow \{\epsilon\}$

repeat

while (S, E) is not closed or not consistent

if (S, E) is not closed

 find $s_1 \in S, a \in A$ such that

$\text{row}(s_1a) \neq \text{row}(s)$, for all $s \in S$

$S \leftarrow S \cup \{s_1a\}$

if (S, E) is not consistent

 find $s_1, s_2 \in S, a \in A$, and $e \in E$ such that

$\text{row}(s_1) = \text{row}(s_2)$

 and $\text{row}(s_1a)(e) \neq \text{row}(s_2a)(e)$

$E \leftarrow E \cup \{ae\}$

Submit hypothesis H to the Teacher

if the Teacher replies **no**, with a counter-example t

$S \leftarrow S \cup \text{prefixes}(t)$

until the Teacher replies **yes** to H

return H

$\text{row} =$

	ϵ
ϵ	0
a	0

$$\mathcal{L} = \{aa, bb, cc, dd, \dots\} \quad A = \{a, b, c, d, \dots\}$$

$S, E \leftarrow \{\epsilon\}$

repeat

while (S, E) is not closed or not consistent

if (S, E) is not closed

 find $s_1 \in S, a \in A$ such that

$\text{row}(s_1a) \neq \text{row}(s)$, for all $s \in S$

$S \leftarrow S \cup \{s_1a\}$

if (S, E) is not consistent

 find $s_1, s_2 \in S, a \in A$, and $e \in E$ such that

$\text{row}(s_1) = \text{row}(s_2)$

 and $\text{row}(s_1a)(e) \neq \text{row}(s_2a)(e)$

$E \leftarrow E \cup \{ae\}$

Submit hypothesis H to the Teacher

if the Teacher replies **no**, with a counter-example t

$S \leftarrow S \cup \text{prefixes}(t)$

until the Teacher replies **yes** to H

return H

$\text{row} =$

	ϵ
ϵ	0
a	0

Closed ✓

Consistent ✓

$$\mathcal{L} = \{aa, bb, cc, dd, \dots\} \quad A = \{a, b, c, d, \dots\}$$

$S, E \leftarrow \{\epsilon\}$

repeat

while (S, E) is not closed or not consistent

if (S, E) is not closed

 find $s_1 \in S, a \in A$ such that

$\text{row}(s_1a) \neq \text{row}(s)$, for all $s \in S$

$S \leftarrow S \cup \{s_1a\}$

if (S, E) is not consistent

 find $s_1, s_2 \in S, a \in A$, and $e \in E$ such that

$\text{row}(s_1) = \text{row}(s_2)$

 and $\text{row}(s_1a)(e) \neq \text{row}(s_2a)(e)$

$E \leftarrow E \cup \{ae\}$

Submit hypothesis H to the Teacher

if the Teacher replies **no**, with a counter-example t

$S \leftarrow S \cup \text{prefixes}(t)$

until the Teacher replies **yes** to H

return H


$\text{row} =$

	ϵ
ϵ	0
a	0

Closed ✓

Consistent ✓

Hypothesis



$$\mathcal{L} = \{aa, bb, cc, dd, \dots\} \quad A = \{a, b, c, d, \dots\}$$

$$S, E \leftarrow \{\epsilon\}$$

repeat

while (S, E) is not closed or not consistent

if (S, E) is not closed

 find $s_1 \in S, a \in A$ such that

$\text{row}(s_1a) \neq \text{row}(s)$, for all $s \in S$

$S \leftarrow S \cup \{s_1a\}$

if (S, E) is not consistent

 find $s_1, s_2 \in S, a \in A$, and $e \in E$ such that

$\text{row}(s_1) = \text{row}(s_2)$

 and $\text{row}(s_1a)(e) \neq \text{row}(s_2a)(e)$

$E \leftarrow E \cup \{ae\}$

Submit hypothesis H to the Teacher

if the Teacher replies **no**, with a counter-example t

$S \leftarrow S \cup \text{prefixes}(t)$

until the Teacher replies **yes** to H

return H


$\text{row} =$

	ϵ
ϵ	0
a	0

Closed ✓

Consistent ✓

Hypothesis



counterexample = aa

$$\mathcal{L} = \{aa, bb, cc, dd, \dots\} \quad A = \{a, b, c, d, \dots\}$$

$$S, E \leftarrow \{\epsilon\}$$

repeat

while (S, E) is not closed or not consistent

if (S, E) is not closed

 find $s_1 \in S, a \in A$ such that

$\text{row}(s_1a) \neq \text{row}(s)$, for all $s \in S$

$S \leftarrow S \cup \{s_1a\}$

if (S, E) is not consistent

 find $s_1, s_2 \in S, a \in A$, and $e \in E$ such that

$\text{row}(s_1) = \text{row}(s_2)$

 and $\text{row}(s_1a)(e) \neq \text{row}(s_2a)(e)$

$E \leftarrow E \cup \{ae\}$

Submit hypothesis H to the Teacher

if the Teacher replies **no**, with a counter-example t

$S \leftarrow S \cup \text{prefixes}(t)$

until the Teacher replies **yes** to H

return H


$\text{row} =$

	ϵ
ϵ	0
a	0

Closed ✓

Consistent ✓

Hypothesis



counterexample = aa

$$\mathcal{L} = \{aa, bb, cc, dd, \dots\} \quad A = \{a, b, c, d, \dots\}$$

$S, E \leftarrow \{\epsilon\}$

repeat

while (S, E) is not closed or not consistent

if (S, E) is not closed

 find $s_1 \in S, a \in A$ such that

$\text{row}(s_1a) \neq \text{row}(s)$, for all $s \in S$

$S \leftarrow S \cup \{s_1a\}$

if (S, E) is not consistent

 find $s_1, s_2 \in S, a \in A$, and $e \in E$ such that

$\text{row}(s_1) = \text{row}(s_2)$

 and $\text{row}(s_1a)(e) \neq \text{row}(s_2a)(e)$

$E \leftarrow E \cup \{ae\}$

Submit hypothesis H to the Teacher

if the Teacher replies **no**, with a counter-example t

$S \leftarrow S \cup \text{prefixes}(t)$

until the Teacher replies **yes** to H

return H


$\text{row} =$

	ϵ
ϵ	0
a	0

Closed ✓

Consistent ✓

Hypothesis



counterexample = aa

$$\mathcal{L} = \{aa, bb, cc, dd, \dots\} \quad A = \{a, b, c, d, \dots\}$$

$$S, E \leftarrow \{\epsilon\}$$

repeat

while (S, E) is not closed or not consistent

if (S, E) is not closed

 find $s_1 \in S, a \in A$ such that

$\text{row}(s_1a) \neq \text{row}(s)$, for all $s \in S$

$$S \leftarrow S \cup \{s_1a\}$$

if (S, E) is not consistent

 find $s_1, s_2 \in S, a \in A$, and $e \in E$ such that

$\text{row}(s_1) = \text{row}(s_2)$

 and $\text{row}(s_1a)(e) \neq \text{row}(s_2a)(e)$

$$E \leftarrow E \cup \{ae\}$$

Submit hypothesis H to the Teacher

if the Teacher replies **no**, with a counter-example t

$$S \leftarrow S \cup \text{orb}(\text{prefixes}(t))$$

until the Teacher replies **yes** to H

return H


$$\text{row} =$$

	ϵ
ϵ	0
a	0

Closed ✓

Consistent ✓

Hypothesis



counterexample = aa

$$\mathcal{L} = \{aa, bb, cc, dd, \dots\} \quad A = \{a, b, c, d, \dots\}$$

$$S, E \leftarrow \{\epsilon\}$$

repeat

while (S, E) is not closed or not consistent

if (S, E) is not closed

 find $s_1 \in S, a \in A$ such that

$\text{row}(s_1a) \neq \text{row}(s)$, for all $s \in S$

$S \leftarrow S \cup \{s_1a\}$

if (S, E) is not consistent

 find $s_1, s_2 \in S, a \in A$, and $e \in E$ such that

$\text{row}(s_1) = \text{row}(s_2)$

 and $\text{row}(s_1a)(e) \neq \text{row}(s_2a)(e)$

$E \leftarrow E \cup \{ae\}$

Submit hypothesis H to the Teacher

if the Teacher replies **no**, with a counter-example t

$S \leftarrow S \cup \text{orb}(\text{prefixes}(t))$

until the Teacher replies **yes** to H

return H

	ϵ
ϵ	0
a	0
aa	1
ab	0
aaa	0
aab	0

$$\text{row} =$$

$$\mathcal{L} = \{aa, bb, cc, dd, \dots\} \quad A = \{a, b, c, d, \dots\}$$

$$S, E \leftarrow \{\epsilon\}$$

repeat

while (S, E) is not closed or not consistent

if (S, E) is not closed

 find $s_1 \in S, a \in A$ such that

$\text{row}(s_1a) \neq \text{row}(s)$, for all $s \in S$

$S \leftarrow S \cup \{s_1a\}$

if (S, E) is not consistent

 find $s_1, s_2 \in S, a \in A$, and $e \in E$ such that

$\text{row}(s_1) = \text{row}(s_2)$

 and $\text{row}(s_1a)(e) \neq \text{row}(s_2a)(e)$

$E \leftarrow E \cup \{ae\}$

Submit hypothesis H to the Teacher

if the Teacher replies **no**, with a counter-example t

$S \leftarrow S \cup \text{orb}(\text{prefixes}(t))$

until the Teacher replies **yes** to H

return H

$$\text{row} =$$

	ϵ
ϵ	0
a	0
aa	1
ab	0
aaa	0
aab	0

$$\mathcal{L} = \{aa, bb, cc, dd, \dots\} \quad A = \{a, b, c, d, \dots\}$$

$$S, E \leftarrow \{\epsilon\}$$

repeat

while (S, E) is not closed or not consistent

if (S, E) is not closed

 find $s_1 \in S, a \in A$ such that

$\text{row}(s_1a) \neq \text{row}(s)$, for all $s \in S$

$S \leftarrow S \cup \{s_1a\}$

if (S, E) is not consistent

 find $s_1, s_2 \in S, a \in A$, and $e \in E$ such that

$\text{row}(s_1) = \text{row}(s_2)$

 and $\text{row}(s_1a)(e) \neq \text{row}(s_2a)(e)$

$E \leftarrow E \cup \{ae\}$

Submit hypothesis H to the Teacher

if the Teacher replies **no**, with a counter-example t

$S \leftarrow S \cup \text{orb}(\text{prefixes}(t))$

until the Teacher replies **yes** to H

return H

$$\text{row} =$$

	ϵ
ϵ	0
a	0
aa	1
ab	0
aaa	0
aab	0

Closed ✓

Consistent ✗

$$\mathcal{L} = \{aa, bb, cc, dd, \dots\} \quad A = \{a, b, c, d, \dots\}$$

$$S, E \leftarrow \{\epsilon\}$$

repeat

while (S, E) is not closed or not consistent

if (S, E) is not closed

 find $s_1 \in S, a \in A$ such that

$\text{row}(s_1a) \neq \text{row}(s)$, for all $s \in S$

$S \leftarrow S \cup \{s_1a\}$

if (S, E) is not consistent

 find $s_1, s_2 \in S, a \in A$, and $e \in E$ such that

$\text{row}(s_1) = \text{row}(s_2)$

 and $\text{row}(s_1a)(e) \neq \text{row}(s_2a)(e)$

$E \leftarrow E \cup \{ae\}$

Submit hypothesis H to the Teacher

if the Teacher replies **no**, with a counter-example t

$S \leftarrow S \cup \text{orb}(\text{prefixes}(t))$

until the Teacher replies **yes** to H

return H

$row =$

	ϵ
ϵ	0
a	0
aa	1
ab	0
aaa	0
aab	0

Closed ✓
Consistent ✗

$$\mathcal{L} = \{aa, bb, cc, dd, \dots\} \quad A = \{a, b, c, d, \dots\}$$

$$S, E \leftarrow \{\epsilon\}$$

repeat

while (S, E) is not closed or not consistent

if (S, E) is not closed

 find $s_1 \in S, a \in A$ such that

$\text{row}(s_1a) \neq \text{row}(s), \text{ for all } s \in S$

$S \leftarrow S \cup \{s_1a\}$

if (S, E) is not consistent

 find $s_1, s_2 \in S, a \in A$, and $e \in E$ such that

$\text{row}(s_1) = \text{row}(s_2)$

 and $\text{row}(s_1a)(e) \neq \text{row}(s_2a)(e)$

$E \leftarrow E \cup \{ae\}$

Submit hypothesis H to the Teacher

if the Teacher replies **no**, with a counter-example t

$S \leftarrow S \cup \text{orb}(\text{prefixes}(t))$

until the Teacher replies **yes** to H

return H

$row =$

	ϵ
ϵ	0
a	0
aa	1
ab	0
aaa	0
aab	0

Closed ✓
Consistent ✗

$$\mathcal{L} = \{aa, bb, cc, dd, \dots\} \quad A = \{a, b, c, d, \dots\}$$

$$S, E \leftarrow \{\epsilon\}$$

repeat

while (S, E) is not closed or not consistent

if (S, E) is not closed

 find $s_1 \in S, a \in A$ such that

$\text{row}(s_1a) \neq \text{row}(s), \text{ for all } s \in S$

$S \leftarrow S \cup \{s_1a\}$

if (S, E) is not consistent

 find $s_1, s_2 \in S, a \in A$, and $e \in E$ such that

$\text{row}(s_1) = \text{row}(s_2)$

 and $\text{row}(s_1a)(e) \neq \text{row}(s_2a)(e)$

$E \leftarrow E \cup \{ae\}$

Submit hypothesis H to the Teacher

if the Teacher replies **no**, with a counter-example t

$S \leftarrow S \cup \text{orb}(\text{prefixes}(t))$

until the Teacher replies **yes** to H

return H

$row =$

	ϵ
ϵ	0
a	0
aa	1
ab	0
aaa	0
aab	0

Closed ✓
Consistent ✗

$$\mathcal{L} = \{aa, bb, cc, dd, \dots\} \quad A = \{a, b, c, d, \dots\}$$

$$S, E \leftarrow \{\epsilon\}$$

repeat

while (S, E) is not closed or not consistent

if (S, E) is not closed

 find $s_1 \in S, a \in A$ such that

$\text{row}(s_1a) \neq \text{row}(s), \text{ for all } s \in S$

$S \leftarrow S \cup \{s_1a\}$

if (S, E) is not consistent

 find $s_1, s_2 \in S, a \in A$, and $e \in E$ such that

$\text{row}(s_1) = \text{row}(s_2)$

 and $\text{row}(s_1a)(e) \neq \text{row}(s_2a)(e)$

$E \leftarrow E \cup \{ae\}$

Submit hypothesis H to the Teacher

if the Teacher replies **no**, with a counter-example t

$S \leftarrow S \cup \text{orb}(\text{prefixes}(t))$

until the Teacher replies **yes** to H

return H

$row =$

	ϵ
ϵ	0
a	0
aa	1
ab	0
aaa	0
aab	0

Closed ✓
Consistent ✗

$$\mathcal{L} = \{aa, bb, cc, dd, \dots\} \quad A = \{a, b, c, d, \dots\}$$

$$S, E \leftarrow \{\epsilon\}$$

repeat

while (S, E) is not closed or not consistent

if (S, E) is not closed

 find $s_1 \in S, a \in A$ such that

$\text{row}(s_1a) \neq \text{row}(s)$, for all $s \in S$

$S \leftarrow S \cup \{s_1a\}$

if (S, E) is not consistent

 find $s_1, s_2 \in S, a \in A$, and $e \in E$ such that

$\text{row}(s_1) = \text{row}(s_2)$

 and $\text{row}(s_1a)(e) \neq \text{row}(s_2a)(e)$

$E \leftarrow E \cup \text{orb}(\{ae\})$

Submit hypothesis H to the Teacher

if the Teacher replies **no**, with a counter-example t

$S \leftarrow S \cup \text{orb}(\text{prefixes}(t))$

until the Teacher replies **yes** to H

return H

$\text{row} =$

	ϵ
ϵ	0
a	0
aa	1
ab	0
aaa	0
aab	0

Closed ✓
Consistent ✗

$$\mathcal{L} = \{aa, bb, cc, dd, \dots\} \quad A = \{a, b, c, d, \dots\}$$

$$S, E \leftarrow \{\epsilon\}$$

repeat

while (S, E) is not closed or not consistent

if (S, E) is not closed

 find $s_1 \in S, a \in A$ such that

$\text{row}(s_1a) \neq \text{row}(s)$, for all $s \in S$

$S \leftarrow S \cup \{s_1a\}$

if (S, E) is not consistent

 find $s_1, s_2 \in S, a \in A$, and $e \in E$ such that

$\text{row}(s_1) = \text{row}(s_2)$

 and $\text{row}(s_1a)(e) \neq \text{row}(s_2a)(e)$

$E \leftarrow E \cup \text{orb}(\{ae\})$

Submit hypothesis H to the Teacher

if the Teacher replies **no**, with a counter-example t

$S \leftarrow S \cup \text{orb}(\text{prefixes}(t))$

until the Teacher replies **yes** to H

return H

$$\text{row} =$$

	ϵ
ϵ	0
a	0
aa	1
ab	0
aaa	0
aab	0

Closed ✓

Consistent ✗

Add all $\pi(a)$ **to** E

$$\mathcal{L} = \{aa, bb, cc, dd, \dots\} \quad A = \{a, b, c, d, \dots\}$$

$S, E \leftarrow \{\epsilon\}$

repeat

while (S, E) is not closed or not consistent

if (S, E) is not closed

 find $s_1 \in S, a \in A$ such that

$\text{row}(s_1a) \neq \text{row}(s)$, for all $s \in S$

$S \leftarrow S \cup \{s_1a\}$

if (S, E) is not consistent

 find $s_1, s_2 \in S, a \in A$, and $e \in E$ such that

$\text{row}(s_1) = \text{row}(s_2)$

 and $\text{row}(s_1a)(e) \neq \text{row}(s_2a)(e)$

$E \leftarrow E \cup \text{orb}(\{ae\})$

Submit hypothesis H to the Teacher

if the Teacher replies **no**, with a counter-example t

$S \leftarrow S \cup \text{orb}(\text{prefixes}(t))$

until the Teacher replies **yes** to H

return H

$\text{row} =$

	ϵ	a	b	c	d	\dots
ϵ	0	0	0	0	0	\dots
a	0	1	0	0	0	\dots
aa	1	0	0	0	0	\dots
ab	0	0	0	0	0	\dots
aaa	0	0	0	0	0	\dots
aab	0	0	0	0	0	\dots

$$\mathcal{L} = \{aa, bb, cc, dd, \dots\} \quad A = \{a, b, c, d, \dots\}$$

$S, E \leftarrow \{\epsilon\}$

repeat

while (S, E) is not closed or not consistent

if (S, E) is not closed

 find $s_1 \in S, a \in A$ such that

$\text{row}(s_1a) \neq \text{row}(s)$, for all $s \in S$

$S \leftarrow S \cup \{s_1a\}$

if (S, E) is not consistent

 find $s_1, s_2 \in S, a \in A$, and $e \in E$ such that

$\text{row}(s_1) = \text{row}(s_2)$

 and $\text{row}(s_1a)(e) \neq \text{row}(s_2a)(e)$

$E \leftarrow E \cup \text{orb}(\{ae\})$

Submit hypothesis H to the Teacher

if the Teacher replies **no**, with a counter-example t

$S \leftarrow S \cup \text{orb}(\text{prefixes}(t))$

until the Teacher replies **yes** to H

return H

$\text{row} =$

	ϵ	a	b	c	d	\dots
ϵ	0	0	0	0	0	\dots
a	0	1	0	0	0	\dots
aa	1	0	0	0	0	\dots
ab	0	0	0	0	0	\dots
aaa	0	0	0	0	0	\dots
aab	0	0	0	0	0	\dots

infinitely-many columns!

$$\mathcal{L} = \{aa, bb, cc, dd, \dots\} \quad A = \{a, b, c, d, \dots\}$$

$$S, E \leftarrow \{\epsilon\}$$

repeat

while (S, E) is not closed or not consistent

if (S, E) is not closed

 find $s_1 \in S, a \in A$ such that

$\text{row}(s_1a) \neq \text{row}(s)$, for all $s \in S$

$S \leftarrow S \cup \{s_1a\}$

if (S, E) is not consistent

 find $s_1, s_2 \in S, a \in A$, and $e \in E$ such that

$\text{row}(s_1) = \text{row}(s_2)$

 and $\text{row}(s_1a)(e) \neq \text{row}(s_2a)(e)$

$E \leftarrow E \cup \text{orb}(\{ae\})$

Submit hypothesis H to the Teacher

if the Teacher replies **no**, with a counter-example t

$S \leftarrow S \cup \text{orb}(\text{prefixes}(t))$


until the Teacher replies **yes** to H

return H

$\text{row} =$

	ϵ	a	b	c	d	\dots
ϵ	0	0	0	0	0	\dots
a	0	1	0	0	0	\dots
aa	1	0	0	0	0	\dots
ab	0	0	0	0	0	\dots
aaa	0	0	0	0	0	\dots
aab	0	0	0	0	0	\dots

$\text{row}(s)(c) = \text{row}(s)(d) = \dots$



infinitely-many columns!

$$\mathcal{L} = \{aa, bb, cc, dd, \dots\} \quad A = \{a, b, c, d, \dots\}$$

$$S, E \leftarrow \{\epsilon\}$$

repeat

while (S, E) is not closed or not consistent

if (S, E) is not closed

 find $s_1 \in S, a \in A$ such that

$\text{row}(s_1a) \neq \text{row}(s)$, for all $s \in S$

$S \leftarrow S \cup \{s_1a\}$

if (S, E) is not consistent

 find $s_1, s_2 \in S, a \in A$, and $e \in E$ such that

$\text{row}(s_1) = \text{row}(s_2)$

 and $\text{row}(s_1a)(e) \neq \text{row}(s_2a)(e)$

$E \leftarrow E \cup \text{orb}(\{ae\})$

Submit hypothesis H to the Teacher

if the Teacher replies **no**, with a counter-example t

$S \leftarrow S \cup \text{orb}(\text{prefixes}(t))$

until the Teacher replies **yes** to H

return H

$\text{row} =$

	ϵ	a	b	c	d	\dots
ϵ	0	0	0	0	0	\dots
a	0	1	0	0	0	\dots
aa	1	0	0	0	0	\dots
ab	0	0	0	0	0	\dots
aaa	0	0	0	0	0	\dots
aab	0	0	0	0	0	\dots

\uparrow \uparrow \uparrow
 $\text{row}(s)(c) = \text{row}(s)(d) = \dots$
 $c, d \notin \text{supp}(\text{row}(s)) \subseteq |s|$

infinitely-many columns!

$$\mathcal{L} = \{aa, bb, cc, dd, \dots\} \quad A = \{a, b, c, d, \dots\}$$

$$S, E \leftarrow \{\epsilon\}$$

repeat

while (S, E) is not closed or not consistent

if (S, E) is not closed

 find $s_1 \in S, a \in A$ such that

$\text{row}(s_1a) \neq \text{row}(s)$, for all $s \in S$

$S \leftarrow S \cup \{s_1a\}$

if (S, E) is not consistent

 find $s_1, s_2 \in S, a \in A$, and $e \in E$ such that

$\text{row}(s_1) = \text{row}(s_2)$

 and $\text{row}(s_1a)(e) \neq \text{row}(s_2a)(e)$

$E \leftarrow E \cup \text{orb}(\{ae\})$

Submit hypothesis H to the Teacher

if the Teacher replies **no**, with a counter-example t

$S \leftarrow S \cup \text{orb}(\text{prefixes}(t))$

until the Teacher replies **yes** to H

return H

$\text{row} =$

	ϵ	a	b	c	d	\dots
ϵ	0	0	0	0	0	\dots
a	0	1	0	0	0	\dots
aa	1	0	0	0	0	\dots
ab	0	0	0	0	0	\dots
aaa	0	0	0	0	0	\dots
aab	0	0	0	0	0	\dots

\uparrow \downarrow \rightarrow
 $\text{row}(s)(c) = \text{row}(s)(d) = \dots$

$c, d \notin \text{supp}(\text{row}(s)) \subseteq |s|$

infinitely-many columns!

we can just keep one column

$c \notin |s|$

$$\mathcal{L} = \{aa, bb, cc, dd, \dots\} \quad A = \{a, b, c, d, \dots\}$$

$$S, E \leftarrow \{\epsilon\}$$

repeat

while (S, E) is not closed or not consistent

if (S, E) is not closed

 find $s_1 \in S, a \in A$ such that

$\text{row}(s_1a) \neq \text{row}(s)$, for all $s \in S$

$S \leftarrow S \cup \{s_1a\}$

if (S, E) is not consistent

 find $s_1, s_2 \in S, a \in A$, and $e \in E$ such that

$\text{row}(s_1) = \text{row}(s_2)$

 and $\text{row}(s_1a)(e) \neq \text{row}(s_2a)(e)$

$E \leftarrow E \cup \text{orb}(\{ae\})$

Submit hypothesis H to the Teacher

if the Teacher replies **no**, with a counter-example t

$S \leftarrow S \cup \text{orb}(\text{prefixes}(t))$

until the Teacher replies **yes** to H

return H

$\text{row} =$

	ϵ	a	b	c
ϵ	0	0	0	0
a	0	1	0	0
aa	1	0	0	0
ab	0	0	0	0
aaa	0	0	0	0
aab	0	0	0	0

infinitely-many columns!

we can just keep one column

$c \notin |s|$

Nominal L^*

$S, E \leftarrow \{\epsilon\}$

repeat

while (S, E) is not closed or not consistent

if (S, E) is not closed

 find $s_1 \in S, a \in A$ such that

$\text{row}(s_1a) \neq \text{row}(s)$, for all $s \in S$

$S \leftarrow S \cup \text{orb}(\{s_1a\})$

if (S, E) is not consistent

 find $s_1, s_2 \in S, a \in A$, and $e \in E$ such that

$\text{row}(s_1) = \text{row}(s_2)$

 and $\text{row}(s_1a)(e) \neq \text{row}(s_2a)(e)$

$E \leftarrow E \cup \text{orb}(\{ae\})$

 Submit hypothesis H to the Teacher

if the Teacher replies **no**, with a counter-example t

$S \leftarrow S \cup \text{orb}(\text{prefixes}(t))$

until the Teacher replies **yes** to H

return H

we only changed 3 lines

Implementation

N λ

Extension of Haskell with **nominal data-types** and **primitives**

atoms : Set Atoms - - the infinite (but orbit-finite) set of all atoms

Internal representation as **FO formulae**, resolved via **Z3**

<https://www.mimuw.edu.pl/~szynwelski/nlambda/>


Correctness from scratch?

Correctness from scratch?

Not really

Set-based proofs as **guidelines**

L^* enjoys a nice **category-theoretic** generalization



Categorical Automata Learning Framework

calf-project.org

The general picture

(finite) sets  (orbit-finite) nominal sets

functions  equivariant functions

(change category from **Set** to **Nom**)

The general picture

(finite) sets  (orbit-finite) nominal sets

functions  equivariant functions

(change category from **Set** to **Nom**)


Nominal automata theory

Mikolaj Bojanczyk, Bartek Klin, Slawomir Lasota:

Automata with Group Actions. LICS 2011

The general picture

(finite) sets



(orbit-finite) nominal sets

functions



equivariant functions

(change category from **Set** to **Nom**)

Nominal automata theory

Mikolaj Bojanczyk, Bartek Klin, Slawomir Lasota:
Automata with Group Actions. LICS 2011

Nominal Programming languages

Eryk Kopczynski, Szymon Torunczyk:
LOIS: syntax and semantics. POPL 2017

The general picture

(finite) sets → (orbit-finite) nominal sets

functions → equivariant functions

(change category from **Set** to **Nom**)

Nominal L^*

Nominal automata theory


Nominal Programming languages

Mikolaj Bojanczyk, Bartek Klin, Slawomir Lasota:
Automata with Group Actions. LICS 2011

Eryk Kopczynski, Szymon Torunczyk:
LOIS: syntax and semantics. POPL 2017

The general picture

(finite) sets



(orbit-finite) nominal sets

functions



equivariant functions

(change category from **Set** to **Nom**)

Nominal L^{*}

Works with any
(suitable) data domain

Nominal automata theory

Nominal Programming languages

Mikolaj Bojanczyk, Bartek Klin, Slawomir Lasota:
Automata with Group Actions. LICS 2011

Eryk Kopczynski, Szymon Torunczyk:
LOIS: syntax and semantics. POPL 2017

In the paper...

- Nominal L^*
- Variations, Nominal NL^*
- Implementation
- Experimental results

<https://github.com/Jaxan/nominal-lstar>



Ongoing and future work

- **Library & tool**



Maverick Chardet
ENS Lyon

- Applications:
 - Specification mining
 - Network verification, with **amazon**
 - Verification of cryptographic protocols
 - Ransomware detection

Demo