

# Magritte: a new Multidimensional Accelerated General-purpose Radiative Transfer code

F. De Ceuster<sup>1,2\*</sup>, J. Yates<sup>1</sup>, J. Hetherington<sup>3</sup>, P.A. Boyle<sup>4</sup>, L. Decin<sup>2</sup>, S. Viti<sup>1</sup> and T.G.

<sup>1</sup>*Department of Physics and Astronomy, University College London, Gower Place, London, WC1E 6BT, UK*

<sup>2</sup>*Department of Physics and Astronomy, Institute of Astronomy, KU Leuven, Celestijnenlaan 200D, 3001 Leuven, Belgium*

<sup>3</sup>*Research Software Development Group, University College London, Podium Building, 1 Eversholt Street, London, NW12DN, UK*

<sup>4</sup>*School of Physics and Astronomy, The University of Edinburgh, Edinburgh EH9 3FD, UK*

<sup>5</sup>*Department of Astronomy and Physics, University of Florida, Gainesville, FL 32611, USA*

<sup>6</sup>*Max-Planck-Institut für Extraterrestrische Physik, Giessenbachstrasse 1, D-85748 Garching, Germany*

Last updated 2011 May 22; in original form 2013 September 5

## ABSTRACT

The In this paper we present the basis for a new multidimensional accelerated general-purpose radiative transfer code, called Magritte.

**Key words:** radiative transfer, astrochemistry, methods: numerical

## 1 INTRODUCTION

The Radiative Transfer problem is one of the oldest and

New C++ code based on 3D-PDR Bisbas et al. (2012) improved performance. separate modules for chemistry, radiative transfer and thermal balance.

The paper is organized as follows. In section 2 we present the computational scheme and the different modules. Section 3 describes the different benchmarks that were done to and to compare its performance with 3D-PDR.

## 2 COMPUTATIONAL SCHEME

Although Magritte is based on 3D-PDR, the whole code has been rewritten from scratch.


### 2.1 Data structures and memory layout

Magritte assumes an unstructured grid of cells

Much better memory scaling and improved overall performance.

Plot of memory use as function of the number of grid cells, Magritte vs. 3D-PDR.

In: density, velocity and radiation field



**Figure 1.** Scheme of different modules.

\* Contact e-mail: [frederik.deceuster@kuleuven.be](mailto:frederik.deceuster@kuleuven.be)

## 2.2 Modules

### 2.2.1 Ray-tracing

### 2.2.2 Chemistry

### 2.2.3 Radiative transfer

Lines plus continuum.

Acceleration methods Ng-acceleration and Rybicki-Hummer acceleration scheme [Rybicki & Hummer \(1991\)](#)

Later versions will be able to treat multiple dust scattering

### 2.2.4 Thermal balance

Magritte can self-consistently determine the temperature assuming local thermal balance, i.e. equal heating and cooling rates for each cell. (Argument on time scales?)

## 2.3 Parallelization strategy

## 3 BENCHMARKS

## 4 APPLICATIONS

The modular character of Magritte

## 5 CONCLUSIONS

We have presented Magritte: a new multidimensional accelerated general

Once all modules are finished and extensively tested, the source code for Magritte and its separate modules will be made freely available on [github.com/Magritte-code](https://github.com/Magritte-code).

## ACKNOWLEDGEMENTS

FDC is supported by the EPSRC iCASE studentship programme, Intel Corporation and Cray Inc.

## REFERENCES

- Bisbas T. G., Bell T. A., Viti S., Yates J., Barlow M. J., 2012, [Mon. Not. R. Astron. Soc.](#), 427, 2100  
Rybicki G. B., Hummer D. G., 1991, *Astron. Astrophys.*, 245, 171

This paper has been typeset from a  $\text{\TeX}/\text{\LaTeX}$  file prepared by the author.