

# GEOG0114: PRINCIPLES OF SPATIAL ANALYSIS

## WEEK 9: SPATIAL MODELS (PART 1)


Dr Anwar Musah ([a.musah@ucl.ac.uk](mailto:a.musah@ucl.ac.uk))

Lecturer in Social and Geographic Data Science

UCL Geography

# Contents


1. Regression-based models for causal and predictive inference
2. Standard Linear Regression (SLR)
  - Model Diagnostics of Residuals
  - Testing evidence for Spatial Dependence
3. Extending an SLR to Spatial Regression Model
  - Spatial lag
  - Spatial error
4. Methodology for model selection
5. We'll cover an example of analysis and interpretation



## QUICK RECAP

1. We covered key concepts of spatial dependence i.e., the characteristics such spatial heterogeneity and autocorrelation
2. We covered what, why and how Spatial Weight Matrix (W) are important
  - Contiguity-based weights
  - Distance-based weights
3. How to measure globally and locally the degree of spatial dependence using the Moran's I statistics

**Let's rewind a bit back on what was taught in Week 3**



# Spatial Models

## Week 9

### Spatial Lag and Error Models

## Week 10

### Geographically Weighted Regression (GWR) Models

Week 3's concepts on spatial dependence are heavily emphasised in here for Spatial Lag and Error Models

What is a Regression model

# Definition of a Regression model:

Regression is a mathematical formula that allows a user to perform two things:

- 1) To determine the **relationship** or **association** between a specified **outcome** (i.e., **dependent variable**) with one or more **predictors** (i.e., **independent variable(s)**)
- 2) For making a prediction about an outcome (i.e., **dependent variable**), or making a projection or forecast of that outcome based on predictors.

**1) Casual Inference**

**2) Predictive Inference**

# Casual & Predictive Inference

## **Examples of causal inference, i.e., the relationship between a dependent and independent variable(s):**

- What is the association between the incidence of skin cancer in residential areas of Cornwall, and environmental exposure to elevated levels of soil arsenic?
- What impact does increased levels of socioeconomic deprivation have on the burden of crime rates?
- What is the risk of a landslide associated with an increase in levels of rainfall, as well as soil texture?

## **Examples of predictive inference, i.e., when we know relationship between dependent and independent variable(s) but use some model to forecast unobserved values:**

- Predict the incidence rate of skin cancer for residential postcodes when soil arsenic concentrations reach levels of 40 mg/kg
- Estimating the probability of a landslide event when rainfall (or precipitation rates) are at values  $> 50\text{mm}$  (2.0 inches) per hour

# Let us clear the air with some terminology

## Dependent Variable

This is the variable whose values we want to explain or forecast

Its value depend on something else

Interchangeable terms:

- **Dependent variable**
- **Response variable**
- **Outcome**
- **Event**

## Independent Variable

This is the variable that explains our dependent variable.

Its values are independent and used to explain the dependent variable

Interchangeable terms:

- **Independent variable**
- **Explanatory variable**
- **Exposure**
- **Risk factor(s)**



In terms of regression, there are several types of models, each with there own families depending on the type distribution for the dependent variable:

Here is a board overview:

Distribution of dependent variable	Suitable Model
<b>Continuous measures:</b> e.g., average income in postcode (£); concentrations of ambient particular matter (PM2.5); Normalised Vegetative Difference Index (NDVI) etc.,	<b>Linear regression</b>
<b>Binary measures (1 = “present” or 0 = “absent”):</b> e.g., Person’s voting for a candidate, Lung cancer risk, house infested with rodents etc.,	<b>Logistic Regression</b>
<b>Binomial measure (or proportion):</b> e.g., prevalence of houses in a postcode infested with rodents, percentage of people in a village infected with intestinal parasitic worms, prevalence of household on a street segment victimised by crime etc.,	<b>Logistic Regression</b>
<b>Counts or discrete measures:</b> e.g., number of reported burglaries on a street segment, number of riots in a county etc.,	<b>Poisson Regression</b>
<b>Time-to-event binary measures:</b> e.g., Lung cancer risk due to chronic exposure to environmental levels of indoor radon. Risk of landslide and time dependence of surface erosion etc.,	<b>Survival Analysis with Cox regression</b>

In terms of regression, there are several types of models, each with there own families depending on the type distribution for the dependent variable:

Here is a board overview:

Distribution of dependent variable	Suitable Model
<b>Continuous measures:</b> e.g., average income in postcode (£); concentrations of ambient particular matter (PM2.5); Normalised Vegetative Difference Index (NDVI) etc.,	Linear regression
<b>Binary measures (1 = “present” or 0 = “absent”):</b> e.g., Person’s voting for a candidate, Lung cancer risk, house infested with rodents etc.,	Logistic Regression
<b>Binomial measure (or proportion):</b> e.g., prevalence of houses in a postcode infested with rodents, percentage of people in a village infected with intestinal parasitic worms, prevalence of household on a street segment victimised by crime etc.,	Logistic Regression
<b>Counts or discrete measures:</b> e.g., number of reported burglaries on a street segment, number of riots in a county etc.,	Poisson Regression
<b>Time-to-event binary measures:</b> e.g., Lung cancer risk due to chronic exposure to environmental levels of indoor radon. Risk of landslide and time dependence of surface erosion etc.,	Survival Analysis with Cox regression

# Univariable & Multivariable Regression

# A Linear Equation

- Do you remember one of these?
  - $y = a + bx$
  - $y = mx + b$
- In the statistics world, we just use a different notation:
  - $y = \beta_0 + \beta_1 x$

## Components of the above linear equation:

$y$  is the dependent variable

$x$  is the independent variable

$\beta_0$  is the intercept of the line on the y-axis when  $x$  is 0

$\beta_1$  is the slope of the line. It is amount  $y$  increases when there's a unit increase in  $x$

# Simple Linear Regression Model

$$y = \beta_0 + \beta_1 x + \varepsilon$$

## Variables

- $y$  is the dependent variable
- $x$  is the independent variable

## Parameters

- $\beta_0$  is the intercept
- $\beta_1$  is the slope, also called a coefficient
- $\varepsilon$  is the error term

# Multivariable Linear Regression Model

$$y = \beta_0 + \beta_1 x_1 + \beta_2 x_2 + \cdots + \varepsilon$$

## Variables


- $y$  is the dependent variable
- $x_1, x_2, x_3, \dots, x_k$  are the independent variables

## Parameters

- $\beta_0$  is the intercept
- $\beta_1, \beta_2, \beta_3, \dots, \beta_k$  are the slopes (or coefficients) for the corresponding variables  $x_1, x_2, x_3, \dots, x_k$
- $\varepsilon$  is the error term

# Simplest example of a bivariable relationship from a multivariable model

$$y = \beta_0 + \beta_1 x_1 + \beta_2 x_2 + \dots + \beta_k x_k + \epsilon$$



1. Notes:

Linear models are best for finding the best fit between predicted values and observed values  $y$

2. Notes:

a. Blue line – the predicted value for  $y$  and the blue hollow circles are the actual observed.


b. The difference between the predicted and observed value give us the error term which we term as residuals.

Such model comes with a set of assumptions

$x_1, x_2, x_3, \dots, x_k$

# Assumptions of Linear Model

$$y = \beta_0 + \beta_1 x_1 + \beta_2 x_2 + \dots + \beta_k x_k + \epsilon$$



Notes:

Assumption 1: The random errors/ residuals should have a mean of zero.

Meaning that if we were to take the sum of these differences between the predicted  $y$  and observed  $y$ , and then divide it by the number of data points – this should give a mean estimate?

Well, this assumption is saying that this sum should be approximately close to zero.


Assumption 2: The distribution of these errors/residuals must be normally distributed

Meaning that if we were to take these points and generate a histogram with them – these should give a shape that's approximately akin to a bell-curve.

Assumption 3: The errors/residuals must have a constant variance (homoskedasticity) and uncorrelated.


What do we mean by this third assumption???






Predicted Y

(a)




Predicted Y

(b)




Predicted Y

(c)



Predicted Y

(d)








Notes:

Assumption 3: The errors/residuals must have a constant variance (homoskedasticity) and uncorrelated.

What do we mean by this assumption???

**When we take the residuals and plot them against the predicted Y (i.e., values of the line of goodness of fit) they should exhibit a patterns that appear like the image in panel (a).**



Notes:


Assumption 3: The errors/residuals must have a constant variance (homoskedasticity) and uncorrelated.

What do we mean by this assumption???


When we take the residuals and plot them against the predicted Y (i.e., values of the line of goodness of fit) they should exhibit a patterns that appear like the image in panel (a).

If the patterns appear like this, then it's safe to say that the variance in the errors are constant across the predicted values.

However, if is noisy like in panels b, c and d then there's evidence of heteroskedasticity




The points should be concentrate around the line when  $y = 0$



Example plot of when the 3<sup>rd</sup> assumption is NOT violated

# Spatial [Lag & Error] Linear Regression Model

Suppose we have a hypothetical study area with 16 areas



### Notes:

Let  $Y$  be some dependent variable that is continuous and normally distributed, where there are 16 observations for  $Y$  (i.e.  $y_1, y_2 \dots y_{16}$ )

- For example: Averaged house price (£)

Let  $X$  be some independent variable, where there are 16 observations for  $X$  (i.e.  $x_1, x_2 \dots x_{16}$ )


- For example: Socioeconomic deprivation score

**Research question:** To investigate the **geospatial** impacts of socioeconomic deprivation on house price in this hypothetical study area.

Insufficient to use the typical linear regression model for this context

$$y = \beta_0 + \beta x + \varepsilon$$

## Scenario 1: Dependent variable values from surrounding neighbourhoods have an influence each other



A hypothetical study area with 16 areas

## Notes:

One type of spatial dependency in this neighbourhood structure can be seen when a **dependent variable (house price)** for some neighbourhood (16) ( $y_{16}$ ) is influenced by other house price **dependent values** from neighbouring areas (i.e., house prices values  $y_3$ ,  $y_4$ ,  $y_9$ ,  $y_{11}$  and  $y_{15}$ ) such as neighbourhoods 3, 4, 9, 11, and 16.

In this scenario, when the dependent variable's outcomes from neighbouring areas have an influence on the outcome of a location, we call this type of spatial dependence a **Spatial Lag on the dependent variable (aka Spatially lagged Y model)**


Spatial Lag are suggestive of spatial spill overs or diffusion.

With spatial lags in a linear regression, the assumptions about independence, and observations being uncorrelated are always violated.

This is one of commonest scenario.



## Scenario 2: Independent variable from the surrounding neighbourhoods have an influence on the values of the dependent variable



A hypothetical study area with 16 areas

## Notes:


Another type of spatial dependency in this neighbourhood structure can be seen when a **dependent variable (house price)** for some neighbourhood (16) ( $y_{16}$ ) is influenced by the independent variable values (i.e., socioeconomic deprivation) from neighbouring areas (i.e., socioeconomic deprivation values  $x_3, x_4, x_9, x_{11}$  and  $x_{15}$ ) measured in neighbourhoods 3, 4, 9, 11, and 16.

In this scenario, when the independent variable from neighbours influences the outcome of a location, we call this type of spatial dependence a **Spatial Lag on the independent variable (aka Spatially lagged X model)**

Again, such spatial lag of this sorts are suggestive of spatial spill overs or diffusion.

This scenario is uncommon.

**Scenario 3: Both the values from the dependent and independent variable from the surrounding neighbourhoods have an influence on the value of the outcome**



**A hypothetical study area with 16 areas**

**Notes:**

The 3<sup>rd</sup> type of spatial dependency in this neighbourhood structure can be seen when a **dependent variable (house price)** for some neighbourhood (16) ( $y_{16}$ ) is influenced by **BOTH** the values from the dependent and independent variable (i.e., house price and socioeconomic deprivation) from the surrounding neighbouring areas (i.e., socioeconomic deprivation values  $x_3, x_4, x_9, x_{11}$  and  $x_{15}$  as well as house prices  $y_3, y_4, y_9, y_{11}$  and  $y_{15}$ ) measured in neighbourhoods 3, 4, 9, 11, and 16.


In this scenario, when both the dependent and independent variable from neighbours influences the outcome of a location, we call this type of spatial dependence a **Spatial Lag on the independent variable**

Again, such spatial lag of this sorts are suggestive of spatial spill overs or diffusion.

This is not commonly a used scenario.



#### Scenario 4: The errors in observed value for both independent or dependent variable from neighbouring areas affecting each other



A hypothetical study area with 16 areas


#### Notes:

The last type of spatial dependency can occur when **the error** in the observed **dependent variable (house price)** from neighbouring areas (i.e., so house price values  $x_3, x_4, x_9, x_{11}$  and  $x_{15}$  measured in neighbourhoods 3, 4, 9, 11, and 15) can influence the outcome in neighbourhood (16) ( $y_{16}$ ).


Errors, in turn, can also influence each other, and can be correlated.

In this scenario, this type of spatial dependence is called **Spatial Error**


Scenario 1




Scenario 2



Scenario 3



Scenario 4



## Notes:

For these scenarios, where  $Y$  is the outcome,  $X$  are some independent variable(s) and  $E$  are residuals.

It possible to model overall association or relationships between  $X$  and  $Y$ , while taking into account:

1. Spatial configuration or layout of the for the study area.

2. The influences not only caused by the independent variables, but also influences among the neighbouring outcome and error terms.

The linear regression can be extended into 4 different types of Spatial regression models which can feature a lag or error component.


## Multivariable Linear Regression

$$y = \beta_0 + \beta_1 x_1 + \beta_2 x_2 + \cdots + \beta_k x_k + \varepsilon$$

### 1. Spatial Lag Model (lagged on the dependent variable)

$$y = \rho WY + \beta_0 + \beta_1 x_1 + \beta_2 x_2 + \cdots + \beta_k x_k + \varepsilon$$

Scenario 1




- W is a spatial weights matrix (contiguity – based)
- Y in the right hand of the equation represents the observed outcome from other areas neighbouring that influences what we're trying to predict
- $\rho$  "Rho" is the degree of how our predicted outcome are influenced by its neighbouring Y measures.

### 2. Spatial Lag Model (lagged on the independent variable)

$$y = \beta_0 + \beta_1 x_1 + \beta_2 x_2 + \cdots + \beta_k x_k + \theta WX + \varepsilon$$

Scenario 2



- W is a spatial weights matrix (contiguity – based)
- X in the right hand of the equation represents the observed values from the independent variable in other areas neighbouring that influences what we're trying to predict
- $\theta$  "theta" is the degree of how our predicted outcome are influenced by its neighbouring X measures.


## Multivariable Linear Regression

$$y = \beta_0 + \beta_1 x_1 + \beta_2 x_2 + \cdots + \beta_k x_k + \varepsilon$$

### 3. Spatial Lag Model (lagged on both the dependent and independent variable)

$$y = \rho WY + \beta_0 + \beta_1 x_1 + \beta_2 x_2 + \cdots + \beta_k x_k + \theta WX + \varepsilon$$

Scenario 3




- Refer to slide number 27 to see the meanings of each parameters

### 4. Spatial Error Model

$$y = \beta_0 + \beta_1 x_1 + \beta_2 x_2 + \cdots + \beta_k x_k + \lambda Wu + \varepsilon$$


Scenario 4




- $u$  are the correlated spatial error terms
- $\lambda$  "lambda" estimated coefficient for the product  $W$  and  $u$ .




Scenario 1




Scenario 2



Scenario 3



Scenario 4



## Notes:

For these scenarios, where  $Y$  is the outcome,  $X$  are some independent variable(s) and  $E$  are residuals.

It possible to model overall association or relationships between  $X$  and  $Y$ , while taking into account:

1. Spatial configuration or layout of the for the study area.

2. The influences not only caused by the independent variables, but also influences among the neighbouring outcome and error terms.

The linear regression can be extended into 4 different types of Spatial regression models which can feature a lag or error component.

# Workflow for spatial regression modelling

# Modelling process and model selection

When you want to conduct evidence-based analysis with spatial data – especially if the outcome is from a continuous distribution – you might want to follow these steps:

- **STEP 1:** Carry some descriptive analysis to understand the underlying spatial distribution
- **STEP 2:** Perform a Linear regression in order to assess the residuals for the model output to determine whether not the assumptions of independence have been violated.

If there's a violation:

- **STEP 3:** Examine whether there is evidence of spatial dependence in the residuals using Moran's I test.
- If Moran's I test is not significant – **STOP! NO EVIDENCE OF SPATIAL DEPENDENCE HENCE NO NEED OF SPATIAL APPROACH**

If Moran's I test is significant and positive – then map the residuals accordingly to examine the spatial patterning of the residual.

- **STEP 4:** Pick one of the spatial regression model and run its regression. For **Spatial Lag Y Model** were interested interpreting the “rho” estimates AND direct & indirect coefficients derived from the impacts() post estimation function.
- For the **Spatial Error model**, its the “lambda” estimate. No need to use the impacts() function. We can get the coefficients directly from model.
- **STEP 5:** Compare the AIC to see which is a better model between the spatial regression(s) and linear regression too. The one with lowest is better.
- **STEP 6:** Perform the Moran's I test on the spatial model to see if they have adequately accounted for spatial autocorrelation. The model with a not significant p-value is the best! Select that model for making your causal or predictive inference

If it's significant (i.e, Moran's I test) in step 6 – go for the one that has the lowest Moran's I statistic closest to zero.

Any questions?