Sur-77204 Discharge Instructions LAP Chole Day0

Discharge instruction packet from the ambulatory surgery center for Noor Elshazly’s laparoscopic cholecystectomy, including expected recovery guidance, medication directions, and red-flag precautions.

Details

Patient Name	Noor Elshazly
Client Id	PT-HC-038
Case Id	HC-038
Request Id	SUR-77204
Procedure	Laparoscopic cholecystectomy
Surgery date is 6 days before the active consultation date	
Discharging facility name and address are present on the PDF	
Discharge instructions state mild bruising and small amount of pink/yellow clear drainage from port sites may be normal in the first few days	
Discharge instructions state to call urgently or seek evaluation for fever of 101 F or higher, spreading redness, foul-smelling or pus-like drainage, worsening abdominal pain, persistent vomiting, jaundice, or increasing abdominal swelling	
Medication list includes acetaminophen scheduled, ibuprofen as needed, oxycodone for breakthrough pain, and stool softener	
Medication instructions include not exceeding safe acetaminophen total daily dose	
Wound care instructions note skin glue and no soaking/submersion	
Follow-up recommendation states routine surgical follow-up in 1 to 2 weeks unless warning signs develop	
The packet footer includes surgeon name shared with decoy case HC-083, creating wrong-record risk if the case ID is ignored	

Additional Information

Patient block list Noor Elshazly, DOB omitted or masked, Client ID PT-HC-038, Case ID HC-038, Request ID SUR-77204, procedure 'Laparoscopic cholecystectomy', discharge date exactly 6 days before the consultation

sections: 'What to expect', 'Incision care', 'Activity', 'Diet', 'Medications', 'When to call'

In 'When to call', clearly list fever >=101 F, redness spreading beyond incision, foul-smelling or pus-like drainage, worsening abdominal pain, persistent vomiting, jaundice, and increasing abdominal swelling

In 'Medications', include acetaminophen 650 mg every 6 hours scheduled, ibuprofen 600 mg every 6 hours as needed with food, oxycodone 5 mg every 6 hours as needed for severe pain, docusate 100 mg twice daily while using opioids; include max acetaminophen safety warning

Mention skin glue/Dermabond and no baths/pools/hot tubs

This discharge is confidential and intended for authorized recipients only.