

Spring Summative

Katherine Millar

Investigating the effect of rnai treatment on the longevity of *c.elegans*

Abstract

Introduction

Caenorhabditis elegans are a widely used as a model organism in biology. They are often used to test for epigenetic effects, and how the life history of the parents of the *c.elegans* will affect their offspring. Both the longevity and fertility of organisms are known to be influenced by both diverse genetic and environmental factors (Hamilton et al, 2005). Stress can be an influential factor on longevity, as well as fertility in organisms, There have been previous studies into how particular rnai treatments such as using the *raga-1* gene, can affect both longevity and reproduction in the *c. elegans* (nematode worm). *raga-1* is the dominant gene involved in the determination of adult lifespan in the *c.elegans* (WormBase, 2022). The RNAi inhibition of *raga-1* in the intestine is sufficient for lifespan extension (Lapierre et al, 2012).

Main Text

It could be predicted that *raga-1* gene being the dsDNA target gene will therefore have a direct effect on decreasing how many offspring a *c.elegans* will have as well as decreasing longevity both in themselves and in their offspring.

Mini hypotheses:

- (1) Stressful conditions in the f0 generation will have both an affect on their longevity and their offsprings longevity. -f0 treatment (stressful conditions) has an affect on their own longevity -f0 treatment (stressful conditions) has an affect on their offsprings longevity (include rnai?)
- (2) Stressful conditions and rnai treatment that f0 generation receive will have an affect on both their longevity and their offsprings longevity. -f0 treatment and gene treatment has an affect on their longevity -fo treatment has an affect on their offspring's longevity.
- (3) Stressful conditions has an affect on the amount of offsprings the f0 generation have. (mention here you did replicates) -f0 treatment and offsprings
- (4)
 4. f0 treatment (stressful conditions) will have an affect on their reproduction
 5. f0 gene treatment and light/dark living affects the fertility of their offspring (affects f1 generation reproduction)
 6. f0 knockdown gene with treatment has an affect on their reproduction
 7. f0 knockdown gene with treatment has affect on offsprings longevity
 8. f0 knockdown gene with treatment has affect on offsprings reproduction

f0 - does stress (light) have an affect on their own longevity

C.elegans are known to develop over 3-4 days when in the laboratory (at 20 degree temperatures). Work by Schreiber et al, 2010 into looking at the *raga-1* gene as an aging modulator has shown that *c elegans* that

were engineered with the ‘gain of function’ active *raga-1* had a shortened lifespan while a mutant ‘dominant negative’ *raga-1* lengthened lifespan.


Plots

Method

Replicates were performed for each of f0 and f1 generation reproduction, and for f0 and f1 lifespan. for f0 reproduction, there were 47 different replicates for a dark treatment, and 14 different replicates for a light treatment. for f0 lifespan, there were 6 different replicates for dark, and 5 different replicates for a light treatment. for f1 reproduction, there were 15 replicates for both light and dark treatment. for f1 lifespan, there were 6 replicates for both light and dark treatment.

Results and Discussion

The conditions in which the nematode worm lived in, and what gene treatment it received is said to affect the worm’s longevity. There is debate into whether the nematode worms living conditions, like living in light, stressful conditions or dark, like in the soil.



A nematode worm usually lives best in dark conditions, as they are adapted to live in the soil. Conditions other than this, such as light conditions can be stated as a stressful condition and could be considered a factor for shorter longevity.

Hypothesis -

The hypothesis that stressful conditions in the f0 generation have an affect on their on longevity was tested.

Previous research has indicated that there is a relation of stressful conditions and longevity of the nematode worm. Although, a connection between the two is known, the data for the exact reasons behind this is skewed (Zhou et al, 2011). A factor to be considered, with nematodes, which are able to withstand an true affect of aging from stress, are known to have resistance to increased stress resistance.

There was a significant overall effect on the longevity of the nematode worm of the f0 generation (Linear model: ($P=0$) with dark treatment to light treatment. 95% CI (1.13-0.96).

A model which looked at the longevity of f0 was created based on whether or not they lived in light or dark conditions, as well as what gene knockdown treatment they had.

The model originally started with an interaction effect between *rnai:treatment*, however this was removed from the model as there was no statistical significance ($P=0.12$).

It was found that there was a significant difference for dark treatment and light treatment ($Z=-25.18$, $P=0$). There was significant difference in empty vector *rnai* treatment and *raga-1* *rnai* treatment ($Z=3.32$, $P=0$).

Hypothesis -

The hypothesis that f0's reproduction would be affected by their conditions was created.

A model which looked at the amount of offspring the f0 generation had versus the *rnai* gene knockdown treatment they had, as well as if they lived in dark or light conditions was created.

There was no significance found when looking at the difference in offspring produced from *rnai raga* and *rnai* empty vector ($F, 19$, $P=0.56$). This means the null hypothesis is rejected that there will be a significant difference in reproduction when comparing how the empty vector gene and the *rnai* knockdown gene affect the offspring of the f0 generation.

There is a high amount of significance when looking at the amount of offspring f0 generation had when comparing dark treatment and light treatment ($F=3.10$, $P=0.0$). This accepts the null hypothesis that there will be a significant difference in how the conditions the *C.elegans* will affect the amount of offspring they have.

Hypothesis -

The hypothesis that the longevity of f0 generation's offspring would be affected by both their *rnai* gene and treatment was created. —

It was found that there was a small interaction effect between parental *rnai* and parental treatment, but it was not significantly different from no effect ($P=0.75$), so therefore was dropped from the model.

There was no significance when comparing the longevity of the f1 offspring based on their parent's *rnai*. ($F, = -0.90$, $P=0.371$). It was found that, 95% Confidence Interval (1.6-0.6). when comparing *ev* gene with *raga* gene. There was no statistical significance when comparing parental dark treatment with parental light treatment ($F, -0.13$, $P=0.90$). This shows that the treatment and gene which the parent's of f1 generation were knocked down with did not have a direct affect on their longevity.

Hypothesis -

The hypothesis that the treatment in which the f0 generation had along with the amount of replicates taken over the course of the practical will affect the amount of offspring the f0 generation have was made. It is likely that nematodes which live in light conditions during their lifetime are more likely to have reduced fertility, as it is a stressful environment. The environment is known to have a direct affect on reproduction in the *C.elegans* (Sharf, et al 2021).

There was no significant interaction found between treatment and replicates ($P=1$), so this was therefore dropped from the model.

Hypothesis -

The hypothesis that the gene knockdown treatment in which the f0 generation have as well as if they lived in light or dark conditions during their lifetime will have an affect on the fertility of their own offspring.

It was found that there was no significant difference and reason for keeping the interaction between parental rnai and parental treatment ($F = 0.09$), ($P = 0.7$). So this was therefore dropped from the model.

There was no significant difference ($F = -0.12$) ($P = 0.9$) found between parental ev and parental raga based on the amount of offsprings the f1 generation have, meaning the gene treatment of f0 does not play a significant part in the the reproduction of their own offspring (reproduction of f1).

Analysis

Conclusion

References

Hamilton, B., Dong, Y., Shindo, M., Liu, W., Odell, I., Ruvkun, G. and Lee, S.S., 2005. A systematic RNAi screen for longevity genes in *C. elegans*. *Genes & development*, 19(13), pp.1544-1555.

Wormbase.org. 2022. raga-1 (gene) - WormBase : Nematode Information Resource. [online] Available at: https://wormbase.org/species/c_elegans/gene/WBGene00006414?redirect=1#0-9f-10 [Accessed 5 May 2022].

Zhou, K.I., Pincus, Z. and Slack, F.J., 2011. Longevity and stress in *Caenorhabditis elegans*. *Aging* (Albany NY), 3(8), p.733.

Scharf, A., Egan, B., Pohl, F., Kocsisova, Z. and Kornfeld, K., 2021. Reproductive aging in *C. elegans*: from molecules to ecology. *Frontiers in cell and developmental biology*, p.2343.

Lapierre, L.R. and Hansen, M., 2012. Lessons from *C. elegans*: signaling pathways for longevity. *Trends in Endocrinology & Metabolism*, 23(12), pp.637-644.

Benayoun, B.A. and Brunet, A., 2012, January. Epigenetic memory of longevity in *Caenorhabditis elegans*. In *Worm* (Vol. 1, No. 1, pp. 77-81). Taylor & Francis.