

Design of “pseudo-natural product” inhibitors of DNA gyrase

Christopher Marriott¹, Marco Cominetti¹, Andrew Beekman¹, Anthony Maxwell² & Mark Searcey¹

¹School of Pharmacy, University of East Anglia, Norwich Research Park, Norwich, NR4 7TJ, United Kingdom


²Department of Biological Chemistry, John Innes Centre, Norwich Research Park, Norwich, NR4 7UH, United Kingdom

1. Introduction



- Antimicrobial resistance poses a huge threat to society, with the predicted number of deaths owing to resistance estimated to rise from 700,000 in 2014 to 10,000,000 by 2050.¹ New compounds must therefore be found to combat resistance.
- DNA gyrase is an essential topoisomerase enzyme found in bacteria which catalyses negative supercoiling of closed-circular double-stranded DNA.
- Previous work has found that certain members of the simocyclinone class of natural products, found in *Streptomyces* strains present in soil, inhibit the binding of DNA gyrase to DNA.² However, the structures of these natural products make these compounds difficult to synthesise.

2. DNA gyrase

- DNA gyrase is a type IIA topoisomerase enzyme.




- It is the only topoisomerase capable of catalysing the introduction of negative supercoils into bacterial DNA
- Inhibited by the simocyclinones (e.g. SD8) which bind to GyrA




3. Problems and aims

- SD8 is unable to penetrate cell membranes and is difficult to synthesise
- Aim is to design “pseudo-natural products” based on SD8 *in silico* and to test these compounds against DNA gyrase and cells




4. Linker design

- Examined use of terminal azidoalkyl acids to replace tetraene and D-olivose regions.
- Azides can react with alkynes to form triazoles via Click chemistry
- Generated a series of alkyl-triazolo linked compounds *in silico* which were then docked into GyrA




- Compound (2) had highest docking score so 10-azidodecanoic acid to be used for future *in silico* design steps.



5. *In silico* hybrid testing

- Libraries of amines (to replace aminocoumarin) and terminal alkynes (to replace angucyclinone) prepared from Aldrich Market Select
- Libraries used to generate hybrids which were docked into GyrA, top 5 scorers taken forward for synthesis



Conclusions and Future Work

- DNA gyrase is an essential bacterial enzyme, and so is a good target for novel antimicrobials
- A series of fragments to replace various components of SD8 have been identified *in silico* to show potential for binding to DNA gyrase
- Synthesis of SD8-fragment hybrids is currently underway, starting with SD8-alkyne hybrids
- Hybrids will then be used in a supercoiling assay to determine activity, following which the best angucyclinone-replacing alkyne will be taken forward to study the aminocoumarin-replacing amines

References and acknowledgements

- J. O'Neill, *Antimicrobial Resistance: Tackling a crisis for the health and wealth of nations*, 2014.
- J. Shimana, H.-P. Fiedler, I. Groth, R. Submuth, W. Beil, M. Walker and A. Zeeck, *J. Antibiot. (Tokyo)*, 2000, **53**, 779–787.
- M. J. Buttner, M. Schäfer, D. M. Lawson and A. Maxwell, *FEMS Microbiol. Rev.*, 2018, **42**, 100–112

Central image adapted from PDB ID 4CKL

This work was supported by the Biotechnology and Biological Sciences Research Council Norwich Research Park Biosciences Doctoral Training Partnership [Grant number BB/M011216/1]