

Machine Learning in Python

Supervised Learning - Regression and Evaluation

Cristian A. Marocico, A. Emin Tatar

Center for Information Technology
University of Groningen

Wednesday, July 2nd 2025

Outline

- 1 Introduction to Regression
- 2 Simple Linear Regression
- 3 Evaluation Metrics for Regression
- 4 Robust Regression
- 5 Multiple Linear Regression
- 6 Regularized Regression
- 7 Nonlinearities in Regression

Introduction to Regression

Regression

Definition

Introduction to Regression

Regression

Definition

Regression is a statistical method used to model the relationship between a dependent variable and one or more independent variables.

Simple Linear Regression

Simple Linear Regression

Definition

Simple Linear Regression

Simple Linear Regression

Definition

Simple Linear Regression is a method to model the relationship between two variables by fitting a linear equation to observed data.

Simple Linear Regression

Simple Linear Regression

Definition

Simple Linear Regression is a method to model the relationship between two variables by fitting a linear equation to observed data. Mathematically:

$$y = \beta_0 + \beta_1 x + \epsilon$$

where:

- y is the dependent variable (response).
- x is the independent variable (predictor).
- β_0 is the y-intercept (constant term).
- β_1 is the slope of the line (coefficient).
- ϵ is the error term (residuals).

Simple Linear Regression

A Simple Linear Regression Machine Learning model will learn the coefficients β_0 and β_1 from the training data to minimize the difference between the predicted values and the actual values.

Assumptions of Simple Linear Regression

- Linearity: The relationship between the independent and dependent variable is linear.

Assumptions of Simple Linear Regression

- Linearity: The relationship between the independent and dependent variable is linear.
- Independence: Observations are independent of each other.

Assumptions of Simple Linear Regression

- Linearity: The relationship between the independent and dependent variable is linear.
- Independence: Observations are independent of each other.
- Homoscedasticity: Constant variance of the error terms.

Assumptions of Simple Linear Regression

- Linearity: The relationship between the independent and dependent variable is linear.
- Independence: Observations are independent of each other.
- Homoscedasticity: Constant variance of the error terms.
- Normality: The residuals (errors) of the model are normally distributed.

Evaluation Metrics for Regression

Common Metrics for Regression

Definition

Evaluation Metrics for Regression

Common Metrics for Regression

Definition

Common metrics to evaluate regression models include:

- Mean Absolute Error (MAE)
- Mean Squared Error (MSE)
- Root Mean Squared Error (RMSE)
- R-squared (R^2)
- Adjusted R-squared

Mean Absolute Error (MAE)

Mean Absolute Error (MAE)

Definition

Mean Absolute Error (MAE)

Mean Absolute Error (MAE)

Definition

Mean Absolute Error (MAE) is the average of the absolute differences between the predicted and the actual values:

$$\text{MAE} = \frac{1}{n} \sum_{i=1}^n |y_i - \hat{y}_i|$$

Mean Absolute Error (MAE)

Mean Absolute Error (MAE)

Definition

Mean Absolute Error (MAE) is the average of the absolute differences between the predicted and the actual values:

$$\text{MAE} = \frac{1}{n} \sum_{i=1}^n |y_i - \hat{y}_i|$$

where:

- n is the number of observations.
- y_i is the actual value.
- \hat{y}_i is the predicted value.

Mean Absolute Error (MAE)

Mean Absolute Error (MAE)

Definition

Mean Absolute Error (MAE) is the average of the absolute differences between the predicted and the actual values:

$$\text{MAE} = \frac{1}{n} \sum_{i=1}^n |y_i - \hat{y}_i|$$

where:

- n is the number of observations.
- y_i is the actual value.
- \hat{y}_i is the predicted value.

MAE is a linear score, which can be used when all errors are equally important; it is also less sensitive to outliers compared to MSE.

Mean Squared Error (MSE)

Mean Squared Error (MSE)

Definition

Mean Squared Error (MSE)

Mean Squared Error (MSE)

Definition

Mean Squared Error (MSE) is the average of the squared differences between the predicted and the actual values:

$$\text{MSE} = \frac{1}{n} \sum_{i=1}^n (y_i - \hat{y}_i)^2$$

Mean Squared Error (MSE)

Mean Squared Error (MSE)

Definition

Mean Squared Error (MSE) is the average of the squared differences between the predicted and the actual values:

$$\text{MSE} = \frac{1}{n} \sum_{i=1}^n (y_i - \hat{y}_i)^2$$

where:

- n is the number of observations.
- y_i is the actual value.
- \hat{y}_i is the predicted value.

Mean Squared Error (MSE)

Mean Squared Error (MSE)

Definition

Mean Squared Error (MSE) is the average of the squared differences between the predicted and the actual values:

$$\text{MSE} = \frac{1}{n} \sum_{i=1}^n (y_i - \hat{y}_i)^2$$

where:

- n is the number of observations.
- y_i is the actual value.
- \hat{y}_i is the predicted value.

MSE is more sensitive to outliers than MAE because it squares the errors, which can disproportionately affect the metric if there are large errors; however, it is useful when larger errors are more significant.

Root Mean Squared Error (RMSE)

Root Mean Squared Error (RMSE)

Definition

Root Mean Squared Error (RMSE)

Root Mean Squared Error (RMSE)

Definition

Root Mean Squared Error (RMSE) is the square root of the average of the squared differences between the predicted and the actual values:

$$\text{RMSE} = \sqrt{\text{MSE}} = \sqrt{\frac{1}{n} \sum_{i=1}^n (y_i - \hat{y}_i)^2}$$

Root Mean Squared Error (RMSE)

Root Mean Squared Error (RMSE)

Definition

Root Mean Squared Error (RMSE) is the square root of the average of the squared differences between the predicted and the actual values:

$$\text{RMSE} = \sqrt{\text{MSE}} = \sqrt{\frac{1}{n} \sum_{i=1}^n (y_i - \hat{y}_i)^2}$$

where:

- n is the number of observations.
- y_i is the actual value.
- \hat{y}_i is the predicted value.

Root Mean Squared Error (RMSE)

Root Mean Squared Error (RMSE)

Definition

Root Mean Squared Error (RMSE) is the square root of the average of the squared differences between the predicted and the actual values:

$$\text{RMSE} = \sqrt{\text{MSE}} = \sqrt{\frac{1}{n} \sum_{i=1}^n (y_i - \hat{y}_i)^2}$$

where:

- n is the number of observations.
- y_i is the actual value.
- \hat{y}_i is the predicted value.

RMSE is in the same units as the dependent variable, making it interpretable; it is also sensitive to outliers, similar to MSE.

R-squared (R^2)

R-squared (R^2)

Definition

R-squared (R^2)

R-squared (R^2)

Definition

R-squared (R^2) is a statistical measure that represents the proportion of the variance for a dependent variable that's explained by an independent variable or variables in a regression model:

$$R^2 = 1 - \frac{SS_{\text{res}}}{SS_{\text{tot}}}$$

where:

- SS_{res} is the sum of squares of residuals (errors).
- SS_{tot} is the total sum of squares (variance of the dependent variable).

R-squared (R^2)

R-squared (R^2)

Definition

R-squared (R^2) is a statistical measure that represents the proportion of the variance for a dependent variable that's explained by an independent variable or variables in a regression model:

$$R^2 = 1 - \frac{SS_{\text{res}}}{SS_{\text{tot}}}$$

where:

- SS_{res} is the sum of squares of residuals (errors).
- SS_{tot} is the total sum of squares (variance of the dependent variable).

R^2 values range from 0 to 1, where:

- 0 indicates that the model does not explain any of the variability of the response data around its mean.
- 1 indicates that the model explains all the variability of the response data around its mean.

Adjusted R-squared

Adjusted R-squared

Definition

Adjusted R-squared

Adjusted R-squared

Definition

Adjusted R-squared adjusts the R^2 value for the number of predictors in the model, providing a more accurate measure when comparing models with different numbers of predictors:

$$\text{Adjusted } R^2 = 1 - (1 - R^2) \frac{n - 1}{n - p - 1}$$

where:

- n is the number of observations.
- p is the number of predictors in the model.

Adjusted R-squared

Adjusted R-squared

Definition

Adjusted R-squared adjusts the R^2 value for the number of predictors in the model, providing a more accurate measure when comparing models with different numbers of predictors:

$$\text{Adjusted } R^2 = 1 - (1 - R^2) \frac{n - 1}{n - p - 1}$$

where:

- n is the number of observations.
- p is the number of predictors in the model.

Adjusted R^2 can be negative, which indicates that the model is worse than a horizontal line (mean of the dependent variable); it is useful for comparing models with different numbers of predictors.

Robust Regression

Robust Regression

Definition

Robust Regression

Robust Regression

Definition

Robust Regression is a type of regression analysis designed to be less sensitive to outliers in the data. It provides a more reliable estimate of the relationship between variables when the data contains outliers or violations of assumptions.

Types of Robust Regression

Types of Robust Regression

Definition

There are several types of robust regression techniques, including:

Types of Robust Regression

Types of Robust Regression

Definition

There are several types of robust regression techniques, including:

- Huber Regression

Types of Robust Regression

Types of Robust Regression

Definition

There are several types of robust regression techniques, including:

- Huber Regression
- RANSAC (RANDOM Sample Consensus)

Types of Robust Regression

Types of Robust Regression

Definition

There are several types of robust regression techniques, including:

- Huber Regression
- RANSAC (RANDOM Sample Consensus)
- Least Trimmed Squares (LTS)

Types of Robust Regression

Types of Robust Regression

Definition

There are several types of robust regression techniques, including:

- Huber Regression
- RANSAC (RANDOM Sample Consensus)
- Least Trimmed Squares (LTS)
- Theil-Sen Estimator

Types of Robust Regression

Types of Robust Regression

Definition

There are several types of robust regression techniques, including:

- Huber Regression
- RANSAC (RANDOM Sample Consensus)
- Least Trimmed Squares (LTS)
- Theil-Sen Estimator
- Quantile Regression

Huber Regression

Huber Regression

Definition

Huber Regression

Huber Regression

Definition

Huber Regression is a robust regression technique that uses a loss function that is quadratic for small residuals and linear for large residuals. This makes it less sensitive to outliers compared to traditional least squares regression.

Huber Regression

Huber Regression

Definition

Huber Regression is a robust regression technique that uses a loss function that is quadratic for small residuals and linear for large residuals. This makes it less sensitive to outliers compared to traditional least squares regression. Mathematically, the Huber loss function is defined as:

$$L(y, \hat{y}) = \begin{cases} \frac{1}{2}(y - \hat{y})^2 & \text{if } |y - \hat{y}| \leq \delta \\ \delta|y - \hat{y}| - \frac{1}{2}\delta^2 & \text{if } |y - \hat{y}| > \delta \end{cases}$$

where δ is a threshold that determines the point at which the loss function transitions from quadratic to linear.

RANSAC (RANdom SAmple Consensus)

RANSAC (RANdom SAmple Consensus)

Definition

RANSAC (RANDOM SAMPLE CONSENSUS)

RANSAC (RANDOM SAMPLE CONSENSUS)

Definition

RANSAC (RANDOM SAMPLE CONSENSUS) is an iterative method used to estimate parameters of a mathematical model from a dataset that contains outliers. It works by randomly selecting a subset of the data, fitting a model to this subset, and then determining how many points from the entire dataset fit this model well.

RANSAC (RANDOM Sample Consensus)

RANSAC (RANDOM Sample Consensus)

Definition

RANSAC (RANDOM Sample Consensus) is an iterative method used to estimate parameters of a mathematical model from a dataset that contains outliers. It works by randomly selecting a subset of the data, fitting a model to this subset, and then determining how many points from the entire dataset fit this model well. The RANSAC algorithm consists of the following steps:

- 1 Randomly select a subset of the data points.
- 2 Fit a model to this subset.
- 3 Determine the inliers (points that fit the model well) and outliers (points that do not fit the model).
- 4 Repeat steps 1-3 for a specified number of iterations or until a satisfactory model is found.
- 5 Select the model with the highest number of inliers as the final model.

Multiple Linear Regression

Multiple Linear Regression

Definition

Multiple Linear Regression

Multiple Linear Regression

Definition

Multiple Linear Regression is an extension of simple linear regression that models the relationship between a dependent variable and multiple independent variables. It is used when there are two or more predictors.

Multiple Linear Regression

Multiple Linear Regression

Definition

Multiple Linear Regression is an extension of simple linear regression that models the relationship between a dependent variable and multiple independent variables. It is used when there are two or more predictors. Mathematically, it is represented as:

$$y = \beta_0 + \beta_1 x_1 + \beta_2 x_2 + \dots + \beta_p x_p + \epsilon$$

where:

- y is the dependent variable.
- x_1, x_2, \dots, x_p are the independent variables (predictors).
- β_0 is the y-intercept (constant term).
- $\beta_1, \beta_2, \dots, \beta_p$ are the coefficients (slopes) for each independent variable.
- ϵ is the error term (residuals).

Assumptions of Multiple Linear Regression

Assumptions of Multiple Linear Regression

Definition

Assumptions of Multiple Linear Regression

Assumptions of Multiple Linear Regression

Definition

The assumptions of multiple linear regression are similar to those of simple linear regression:

- **Linearity:** The relationship between the independent variables and the dependent variable is linear.
- **Independence:** Observations are independent of each other.
- **Homoscedasticity:** Constant variance of the error terms.
- **Normality:** The residuals (errors) of the model are normally distributed.
- **No multicollinearity:** The independent variables are not highly correlated with each other.

How do Machine Learning Models Learn?

How do Machine Learning Models Learn?

Definition

How do Machine Learning Models Learn?

How do Machine Learning Models Learn?

Definition

Machine Learning models learn by adjusting their parameters (coefficients) to minimize the difference between the predicted values and the actual values in the training data, also known as the **cost function** or **loss function**.

How do Machine Learning Models Learn?

How do Machine Learning Models Learn?

Definition

Machine Learning models learn by adjusting their parameters (coefficients) to minimize the difference between the predicted values and the actual values in the training data, also known as the **cost function** or **loss function**. This process is often done using optimization algorithms that iteratively update the parameters based on the loss function, which measures the error of the predictions.

Cost Function

Cost Function	Definition

Cost Function

Cost Function

Definition

The **cost function** (or **loss function**) is a mathematical function that quantifies the difference between the predicted values and the actual values in the training data. It is used to guide the learning process of the model.

Cost Function

Cost Function

Definition

The **cost function** (or **loss function**) is a mathematical function that quantifies the difference between the predicted values and the actual values in the training data. It is used to guide the learning process of the model. Common cost functions for regression include:

- Mean Squared Error (MSE)
- Mean Absolute Error (MAE)

Cost Function: Mean Squared Error (MSE)

Cost Function: Mean Squared Error (MSE)

Definition

Cost Function: Mean Squared Error (MSE)

Cost Function: Mean Squared Error (MSE)

Definition

Mean Squared Error (MSE) is a common cost function used in regression tasks. It measures the average of the squares of the errors, which are the differences between the predicted values and the actual values:

$$J_{\text{MSE}}(w) = \frac{1}{2m} \sum_{i=1}^m (f(x_i, w) - y_i)^2$$

where:

- m is the number of observations.
- $f(x_i, w)$ is the predicted value for the i -th observation given the input features x_i and model parameters w .
- y_i is the actual value for the i -th observation.

Cost Function: Mean Squared Error (MSE)

Cost Function: Mean Squared Error (MSE)

Definition

Mean Squared Error (MSE) is a common cost function used in regression tasks. It measures the average of the squares of the errors, which are the differences between the predicted values and the actual values:

$$J_{\text{MSE}}(w) = \frac{1}{2m} \sum_{i=1}^m (f(x_i, w) - y_i)^2$$

where:

- m is the number of observations.
- $f(x_i, w)$ is the predicted value for the i -th observation given the input features x_i and model parameters w .
- y_i is the actual value for the i -th observation.

MSE is sensitive to outliers, and is, therefore, useful when larger errors are more significant.

Cost Function: Mean Absolute Error (MAE)

Cost Function: Mean Absolute Error (MAE)

Definition

Cost Function: Mean Absolute Error (MAE)

Cost Function: Mean Absolute Error (MAE)

Definition

Mean Absolute Error (MAE) is another common cost function used in regression tasks. It measures the average of the absolute differences between the predicted values and the actual values:

$$J_{\text{MAE}}(w) = \frac{1}{m} \sum_{i=1}^m |f(x_i, w) - y_i|$$

where:

- m is the number of observations.
- $f(x_i, w)$ is the predicted value for the i -th observation given the input features x_i and model parameters w .
- y_i is the actual value for the i -th observation.

Cost Function: Mean Absolute Error (MAE)

Cost Function: Mean Absolute Error (MAE)

Definition

Mean Absolute Error (MAE) is another common cost function used in regression tasks. It measures the average of the absolute differences between the predicted values and the actual values:

$$J_{\text{MAE}}(w) = \frac{1}{m} \sum_{i=1}^m |f(x_i, w) - y_i|$$

where:

- m is the number of observations.
- $f(x_i, w)$ is the predicted value for the i -th observation given the input features x_i and model parameters w .
- y_i is the actual value for the i -th observation.

MAE is less sensitive to outliers compared to MSE; useful when all errors are equally important.

Gradient Descent

Gradient Descent

Definition

Gradient Descent

Gradient Descent

Definition

Gradient Descent is an optimization algorithm used to minimize the cost function by iteratively updating the model parameters in the direction of the steepest descent of the cost function.

Gradient Descent

Gradient Descent

Definition

Gradient Descent is an optimization algorithm used to minimize the cost function by iteratively updating the model parameters in the direction of the steepest descent of the cost function. The update rule for gradient descent is:

$$w := w - \alpha \nabla J(w)$$

where:

- w is the vector of model parameters (coefficients).
- α is the learning rate, which controls the step size of each update.
- $\nabla J(w)$ is the gradient of the cost function with respect to the parameters w .

Gradient Descent

Gradient Descent

Definition

Gradient Descent is an optimization algorithm used to minimize the cost function by iteratively updating the model parameters in the direction of the steepest descent of the cost function. The update rule for gradient descent is:

$$w := w - \alpha \nabla J(w)$$

where:

- w is the vector of model parameters (coefficients).
- α is the learning rate, which controls the step size of each update.
- $\nabla J(w)$ is the gradient of the cost function with respect to the parameters w .

Gradient descent continues until convergence, which is when the change in the cost function is below a certain threshold or a maximum number of iterations is reached.

Gradient Descent Algorithm

Gradient Descent Algorithm

Definition

Gradient Descent Algorithm

Gradient Descent Algorithm

Definition

The **Gradient Descent Algorithm** consists of the following steps:

- 1 Initialize the model parameters (coefficients) randomly or to zero.

Gradient Descent Algorithm

Gradient Descent Algorithm

Definition

The **Gradient Descent Algorithm** consists of the following steps:

- 1 Initialize the model parameters (coefficients) randomly or to zero.
- 2 Compute the predicted values using the current parameters.

Gradient Descent Algorithm

Gradient Descent Algorithm

Definition

The **Gradient Descent Algorithm** consists of the following steps:

- ➊ Initialize the model parameters (coefficients) randomly or to zero.
- ➋ Compute the predicted values using the current parameters.
- ➌ Calculate the cost function (loss) using the predicted values and actual values.

Gradient Descent Algorithm

Gradient Descent Algorithm

Definition

The **Gradient Descent Algorithm** consists of the following steps:

- 1 Initialize the model parameters (coefficients) randomly or to zero.
- 2 Compute the predicted values using the current parameters.
- 3 Calculate the cost function (loss) using the predicted values and actual values.
- 4 Compute the gradient of the cost function with respect to the parameters.

Gradient Descent Algorithm

Gradient Descent Algorithm

Definition

The **Gradient Descent Algorithm** consists of the following steps:

- 1 Initialize the model parameters (coefficients) randomly or to zero.
- 2 Compute the predicted values using the current parameters.
- 3 Calculate the cost function (loss) using the predicted values and actual values.
- 4 Compute the gradient of the cost function with respect to the parameters.
- 5 Update the parameters using the gradient descent update rule.

Gradient Descent Algorithm

Gradient Descent Algorithm

Definition

The **Gradient Descent Algorithm** consists of the following steps:

- 1 Initialize the model parameters (coefficients) randomly or to zero.
- 2 Compute the predicted values using the current parameters.
- 3 Calculate the cost function (loss) using the predicted values and actual values.
- 4 Compute the gradient of the cost function with respect to the parameters.
- 5 Update the parameters using the gradient descent update rule.
- 6 Repeat steps 2-5 until convergence (i.e., the change in the cost function is below a threshold or a maximum number of iterations is reached).

Overfitting and Underfitting

Overfitting and Underfitting

Definition

Overfitting and Underfitting

Overfitting and Underfitting

Definition

Overfitting occurs when a model learns the noise in the training data instead of the underlying pattern, resulting in poor generalization to unseen data. It typically happens when the model is too complex relative to the amount of training data.

Overfitting and Underfitting

Overfitting and Underfitting

Definition

Overfitting occurs when a model learns the noise in the training data instead of the underlying pattern, resulting in poor generalization to unseen data. It typically happens when the model is too complex relative to the amount of training data.

Underfitting occurs when a model is too simple to capture the underlying pattern in the data, resulting in poor performance on both training and unseen data. It typically happens when the model is not complex enough relative to the data.

Overfitting and Underfitting

Overfitting and Underfitting

Definition

Overfitting occurs when a model learns the noise in the training data instead of the underlying pattern, resulting in poor generalization to unseen data. It typically happens when the model is too complex relative to the amount of training data.

Underfitting occurs when a model is too simple to capture the underlying pattern in the data, resulting in poor performance on both training and unseen data. It typically happens when the model is not complex enough relative to the data.

To mitigate overfitting, techniques such as **regularization**, **cross-validation**, and **pruning** can be used. To address underfitting, more complex models or additional features can be considered.

Regularization

Regularization

Definition

Regularization

Regularization

Definition

Regularization is a technique used in machine learning to prevent overfitting by adding a penalty term to the cost function. This penalty discourages overly complex models by constraining the size of the model parameters (coefficients).

Regularization

Regularization

Definition

Regularization is a technique used in machine learning to prevent overfitting by adding a penalty term to the cost function. This penalty discourages overly complex models by constraining the size of the model parameters (coefficients).

Types of Regularized Regression

Types of Regularized Regression

Definition

There are two common types of regularized regression:

- Ridge Regression (L2 Regularization)
- Lasso Regression (L1 Regularization)

Lasso Regression (L1 Regularization)

Lasso Regression (L1 Regularization)

Definition

Lasso Regression (L1 Regularization)

Lasso Regression (L1 Regularization)

Definition

Lasso Regression adds a penalty equal to the absolute value of the magnitude of coefficients to the loss function. The objective function for lasso regression is:

$$J_{\text{Lasso}}(w) = \frac{1}{2m} \sum_{i=1}^m (f(x_i, w) - y_i)^2 + \lambda \sum_{j=1}^p |w_j|$$

where:

- λ is the regularization parameter that controls the strength of the penalty.
- w_j are the coefficients of the independent variables.

Lasso Regression (L1 Regularization)

Lasso Regression (L1 Regularization)

Definition

Lasso Regression adds a penalty equal to the absolute value of the magnitude of coefficients to the loss function. The objective function for lasso regression is:

$$J_{\text{Lasso}}(w) = \frac{1}{2m} \sum_{i=1}^m (f(x_i, w) - y_i)^2 + \lambda \sum_{j=1}^p |w_j|$$

where:

- λ is the regularization parameter that controls the strength of the penalty.
- w_j are the coefficients of the independent variables.

Lasso regression can shrink some coefficients to exactly zero, effectively performing variable selection and resulting in a simpler model.

Ridge Regression (L2 Regularization)

Ridge Regression (L2 Regularization)

Definition

Ridge Regression (L2 Regularization)

Ridge Regression (L2 Regularization)

Definition

Ridge Regression adds a penalty equal to the square of the magnitude of coefficients to the loss function. The objective function for ridge regression is:

$$J_{\text{Ridge}}(w) = \frac{1}{2m} \sum_{i=1}^m (f(x_i, w) - y_i)^2 + \frac{\lambda}{2m} \sum_{j=1}^p w_j^2$$

where:

- λ is the regularization parameter that controls the strength of the penalty.
- w_j are the coefficients of the independent variables.

Ridge Regression (L2 Regularization)

Ridge Regression (L2 Regularization)

Definition

Ridge Regression adds a penalty equal to the square of the magnitude of coefficients to the loss function. The objective function for ridge regression is:

$$J_{\text{Ridge}}(w) = \frac{1}{2m} \sum_{i=1}^m (f(x_i, w) - y_i)^2 + \frac{\lambda}{2m} \sum_{j=1}^p w_j^2$$

where:

- λ is the regularization parameter that controls the strength of the penalty.
- w_j are the coefficients of the independent variables.

Ridge regression helps to reduce model complexity and multicollinearity by shrinking the coefficients towards zero, but it does not set any coefficients exactly to zero.

Nonlinearities in Regression

Nonlinearities in Regression

Definition

Nonlinearities in Regression

Nonlinearities in Regression

Definition

Nonlinearities in Regression refers to the presence of nonlinear relationships between the independent and dependent variables in a regression model. In such cases, a simple linear regression model may not adequately capture the underlying relationship, leading to poor predictions and insights.

Nonlinearities in Regression

Nonlinearities in Regression

Definition

Nonlinearities in Regression refers to the presence of nonlinear relationships between the independent and dependent variables in a regression model. In such cases, a simple linear regression model may not adequately capture the underlying relationship, leading to poor predictions and insights. Nonlinear regression models can help address this limitation by allowing for more flexible relationships between the independent and dependent variables.

Types of Nonlinear Regression Models

Types of Nonlinear Regression Models

Definition

There are several types of nonlinear regression models, including:

- **Polynomial Regression:** Fits a polynomial equation to the data, allowing for curves and bends in the relationship between the independent and dependent variables. The degree of the polynomial can be adjusted to capture more complex relationships.

Types of Nonlinear Regression Models

Types of Nonlinear Regression Models

Definition

There are several types of nonlinear regression models, including:

- **Polynomial Regression:** Fits a polynomial equation to the data, allowing for curves and bends in the relationship between the independent and dependent variables. The degree of the polynomial can be adjusted to capture more complex relationships.
- **Exponential Regression:** Fits an exponential function to the data, which is useful for modeling relationships where the rate of change increases or decreases exponentially.

Types of Nonlinear Regression Models

Types of Nonlinear Regression Models

Definition

There are several types of nonlinear regression models, including:

- **Polynomial Regression:** Fits a polynomial equation to the data, allowing for curves and bends in the relationship between the independent and dependent variables. The degree of the polynomial can be adjusted to capture more complex relationships.
- **Exponential Regression:** Fits an exponential function to the data, which is useful for modeling relationships where the rate of change increases or decreases exponentially.
- **Logarithmic Regression:** Fits a logarithmic function to the data, which is useful for modeling relationships where the rate of change decreases as the independent variable increases.