

Machine Learning in Python

Supervised Learning - Regression and Evaluation

Cristian A. Marocico, A. Emin Tatar

Center for Information Technology
University of Groningen

Wednesday, July 2nd 2025

Outline

- 1 Introduction to Regression
- 2 Simple Linear Regression
- 3 Evaluation Metrics for Regression

Introduction to Regression

Regression

Definition

Introduction to Regression

Regression

Definition

Regression is a statistical method used to model the relationship between a dependent variable and one or more independent variables.

Simple Linear Regression

Simple Linear Regression

Definition

Simple Linear Regression

Simple Linear Regression

Definition

Simple Linear Regression is a method to model the relationship between two variables by fitting a linear equation to observed data.

Simple Linear Regression

Simple Linear Regression

Definition

Simple Linear Regression is a method to model the relationship between two variables by fitting a linear equation to observed data. Mathematically:

$$y = \beta_0 + \beta_1 x + \epsilon$$

where:

- y is the dependent variable (response).
- x is the independent variable (predictor).
- β_0 is the y-intercept (constant term).
- β_1 is the slope of the line (coefficient).
- ϵ is the error term (residuals).

Simple Linear Regression

A Simple Linear Regression Machine Learning model will learn the coefficients β_0 and β_1 from the training data to minimize the difference between the predicted values and the actual values.

Assumptions of Simple Linear Regression

- Linearity: The relationship between the independent and dependent variable is linear.

Assumptions of Simple Linear Regression

- Linearity: The relationship between the independent and dependent variable is linear.
- Independence: Observations are independent of each other.

Assumptions of Simple Linear Regression

- Linearity: The relationship between the independent and dependent variable is linear.
- Independence: Observations are independent of each other.
- Homoscedasticity: Constant variance of the error terms.

Assumptions of Simple Linear Regression

- Linearity: The relationship between the independent and dependent variable is linear.
- Independence: Observations are independent of each other.
- Homoscedasticity: Constant variance of the error terms.
- Normality: The residuals (errors) of the model are normally distributed.

Evaluation Metrics for Regression

Common Metrics for Regression

Definition

Evaluation Metrics for Regression

Common Metrics for Regression

Definition

Common metrics to evaluate regression models include:

- Mean Absolute Error (MAE)
- Mean Squared Error (MSE)
- Root Mean Squared Error (RMSE)
- R-squared (R^2)
- Adjusted R-squared

Mean Absolute Error (MAE)

Mean Absolute Error (MAE)

Definition

Mean Absolute Error (MAE)

Mean Absolute Error (MAE)

Definition

Mean Absolute Error (MAE) is the average of the absolute differences between the predicted and the actual values:

$$\text{MAE} = \frac{1}{n} \sum_{i=1}^n |y_i - \hat{y}_i|$$

Mean Absolute Error (MAE)

Mean Absolute Error (MAE)

Definition

Mean Absolute Error (MAE) is the average of the absolute differences between the predicted and the actual values:

$$\text{MAE} = \frac{1}{n} \sum_{i=1}^n |y_i - \hat{y}_i|$$

where:

- n is the number of observations.
- y_i is the actual value.
- \hat{y}_i is the predicted value.

Mean Absolute Error (MAE)

Mean Absolute Error (MAE)

Definition

Mean Absolute Error (MAE) is the average of the absolute differences between the predicted and the actual values:

$$\text{MAE} = \frac{1}{n} \sum_{i=1}^n |y_i - \hat{y}_i|$$

where:

- n is the number of observations.
- y_i is the actual value.
- \hat{y}_i is the predicted value.

MAE is a linear score, which can be used when all errors are equally important; it is also less sensitive to outliers compared to MSE.

Mean Squared Error (MSE)

Mean Squared Error (MSE)

Definition

Mean Squared Error (MSE)

Mean Squared Error (MSE)

Definition

Mean Squared Error (MSE) is the average of the squared differences between the predicted and the actual values:

$$\text{MSE} = \frac{1}{n} \sum_{i=1}^n (y_i - \hat{y}_i)^2$$

Mean Squared Error (MSE)

Mean Squared Error (MSE)

Definition

Mean Squared Error (MSE) is the average of the squared differences between the predicted and the actual values:

$$\text{MSE} = \frac{1}{n} \sum_{i=1}^n (y_i - \hat{y}_i)^2$$

where:

- n is the number of observations.
- y_i is the actual value.
- \hat{y}_i is the predicted value.

Mean Squared Error (MSE)

Mean Squared Error (MSE)

Definition

Mean Squared Error (MSE) is the average of the squared differences between the predicted and the actual values:

$$\text{MSE} = \frac{1}{n} \sum_{i=1}^n (y_i - \hat{y}_i)^2$$

where:

- n is the number of observations.
- y_i is the actual value.
- \hat{y}_i is the predicted value.

MSE is more sensitive to outliers than MAE because it squares the errors, which can disproportionately affect the metric if there are large errors; however, it is useful when larger errors are more significant.

Root Mean Squared Error (RMSE)

Root Mean Squared Error (RMSE)

Definition

Root Mean Squared Error (RMSE)

Root Mean Squared Error (RMSE)

Definition

Root Mean Squared Error (RMSE) is the square root of the average of the squared differences between the predicted and the actual values:

$$\text{RMSE} = \sqrt{\text{MSE}} = \sqrt{\frac{1}{n} \sum_{i=1}^n (y_i - \hat{y}_i)^2}$$

Root Mean Squared Error (RMSE)

Root Mean Squared Error (RMSE)

Definition

Root Mean Squared Error (RMSE) is the square root of the average of the squared differences between the predicted and the actual values:

$$\text{RMSE} = \sqrt{\text{MSE}} = \sqrt{\frac{1}{n} \sum_{i=1}^n (y_i - \hat{y}_i)^2}$$

where:

- n is the number of observations.
- y_i is the actual value.
- \hat{y}_i is the predicted value.

Root Mean Squared Error (RMSE)

Root Mean Squared Error (RMSE)

Definition

Root Mean Squared Error (RMSE) is the square root of the average of the squared differences between the predicted and the actual values:

$$\text{RMSE} = \sqrt{\text{MSE}} = \sqrt{\frac{1}{n} \sum_{i=1}^n (y_i - \hat{y}_i)^2}$$

where:

- n is the number of observations.
- y_i is the actual value.
- \hat{y}_i is the predicted value.

RMSE is in the same units as the dependent variable, making it interpretable; it is also sensitive to outliers, similar to MSE.

R-squared (R^2)

R-squared (R^2)

Definition

R-squared (R^2)

R-squared (R^2)

Definition

R-squared (R^2) is a statistical measure that represents the proportion of the variance for a dependent variable that's explained by an independent variable or variables in a regression model:

$$R^2 = 1 - \frac{SS_{\text{res}}}{SS_{\text{tot}}}$$

where:

- SS_{res} is the sum of squares of residuals (errors).
- SS_{tot} is the total sum of squares (variance of the dependent variable).

R-squared (R^2)

R-squared (R^2)

Definition

R-squared (R^2) is a statistical measure that represents the proportion of the variance for a dependent variable that's explained by an independent variable or variables in a regression model:

$$R^2 = 1 - \frac{SS_{\text{res}}}{SS_{\text{tot}}}$$

where:

- SS_{res} is the sum of squares of residuals (errors).
- SS_{tot} is the total sum of squares (variance of the dependent variable).

R^2 values range from 0 to 1, where:

- 0 indicates that the model does not explain any of the variability of the response data around its mean.
- 1 indicates that the model explains all the variability of the response data around its mean.

Adjusted R-squared

Adjusted R-squared

Definition

Adjusted R-squared

Adjusted R-squared

Definition

Adjusted R-squared adjusts the R^2 value for the number of predictors in the model, providing a more accurate measure when comparing models with different numbers of predictors:

$$\text{Adjusted } R^2 = 1 - (1 - R^2) \frac{n - 1}{n - p - 1}$$

where:

- n is the number of observations.
- p is the number of predictors in the model.

Adjusted R-squared

Adjusted R-squared

Definition

Adjusted R-squared adjusts the R^2 value for the number of predictors in the model, providing a more accurate measure when comparing models with different numbers of predictors:

$$\text{Adjusted } R^2 = 1 - (1 - R^2) \frac{n - 1}{n - p - 1}$$

where:

- n is the number of observations.
- p is the number of predictors in the model.

Adjusted R^2 can be negative, which indicates that the model is worse than a horizontal line (mean of the dependent variable); it is useful for comparing models with different numbers of predictors.