Cancer Incidence AAPC Values - Men

■ 20-49 years ■ 50+ years

		P Value	P Value
		20-49	50+
Prostate*^		<0.0001	<0.0001
Thyroid*		<0.0001	<0.0001
MultipleMyeloma*		<0.0001	<0.0001
Kidney*		<0.0001	<0.0001
Gallbladder*		0.012	<0.0001
Colorectum*^	_ _	<0.0001	0.55
Pancreas*		<0.0001	<0.0001
Leukaemia*		<0.0001	0.044
Liver		0.29	<0.0001
Oral*	- - -	<0.0001	<0.0001
Hodgkin Lymphoma*	- 	0.00014	< 0.0001
Melanoma		0.066	<0.0001
Testis*	- 	0.00021	<0.0001
Brain and CNS*^		0.0002	0.26
Non-Hodgkin Lymphoma	— — ——————————————————————————————————	0.76	<0.0001
Kaposi Sarcoma		0.96	<0.0001
Oesophagus		0.068	<0.0001
Lung		0.00053	<0.0001
Larynx		0.016	<0.0001
Stomach	<u> </u>	0.00074	<0.0001
Bladder	-4 -3 -2 -1 0 1 2 3 4 5 6	<0.0001	0.00094

 $^{^{\}star}$ – AAPC for 20–49 is significantly greater than 0

^{^ –} AAPC for 20–49 is significantly greater than 0 and the AAPC for 50+ is not significantly greater than 0 or does not overlap with the confidence interval of the AAPC for 20–49