

See discussions, stats, and author profiles for this publication at: <https://www.researchgate.net/publication/322505910>

Remarks on: Swanson's Hamiltonian [J.Math.Phys 45(2);585(2004)]

Negative Results · January 2018

DOI: 10.13140/RG.2.2.22605.54246

CITATIONS

0

READS

91

2 authors, including:



Biswanath Rath

Maharaja Sriram Chandra Bhanja Deo University, Takatpur ,Baripada ,Mayurbhanj ,Odisha ,India

151 PUBLICATIONS 211 CITATIONS

[SEE PROFILE](#)

Some of the authors of this publication are also working on these related projects:



Applied mathematics [View project](#)



Applied Mathematics [View project](#)

Remarks on: Swanson's Hamiltonian

[J.Math.Phys **45**(**2**);585(2004)]

Biswanath Rath; Pravanjan Mallick and Prachiprava

Mohapatra

Department of Physics, North Orissa University, Takatpur,

Baripada -757003, Odisha, INDIA

(e.mail:biswanathrath10@gmail.com)

We find that the Hamiltonian $H = wa^\dagger a + \alpha a^2 + \beta (a^\dagger)^2$ satisfying the condition $w^2 - 4\alpha\beta \gg 0$ can admit negative spectra , which is in contradiction to all real positive eigenvalues (as stated in the abstract).

PACS:03.65.Ge

Key words-Negative eigenvalue , matrix diagonalisation method .

I.Introduction

Nearly 14 years back M.S.Swanson [1] proposed a non-hermitian quadratic Hamiltonian

$$H = wa^\dagger + \alpha a^2 + \beta (a^\dagger)^2 \quad (1)$$

with $\alpha \neq \beta$ and suggested that the spectra becomes positive and real provided

$$w^2 - 4\alpha\beta \gg 0 \quad (2)$$




Figure 1: $H = 64a^\dagger a + 12a^2 + 20(a^\dagger)^2$

We feel that the above Hamiltonian is a different manifestation of Harmonic oscillator . As such the construction deals with non-hermitian quantum mechanics with real non-diagonal terms . By curiosity we performed matrix diagonalisation calculation to check the validity of pure real positive spectra satisfying the above condition. To our surprise the spectra can admit real negative energy in addition to positive energy.

II. Method and Results

Here we use simple matrix diagonalisation method [2] to find eigenvalues . For some values of $w; \alpha$ and β we summarise the value of negative energy as given below.

$$E_0 = -0.5 \rightarrow w = 8; \alpha = 1.5; \beta = 2.5$$

$$E_0 = -1 \rightarrow w = 16; \alpha = 3; \beta = 5$$

$$E_0 = -2 \rightarrow w = 32; \alpha = 6; \beta = 10$$

$$E_0 = -4 \rightarrow w = 64; \alpha = 12; \beta = 20$$

For the interest of the reader we plot the first two eigenvalues corresponding to $w = 64, \alpha = 12, \beta = 20$ in fig-1.

III. Conclusion

Our findings contradict the conclusion presented earlier[1] .One can verify our numerical findings using analytical method [3]. We believe that the non-hermitian Hamiltonian can admit negative energy in addition to positive energy . Probably a slight modified form of Swanson's Hamiltonian can also reflect all pure real negative eigenvalues [4] satisfying the condition in Eq(2).

References

- [1] M.S.Swanson ; J.Math.Phys **45(2)**,585(2004).
- [2] B.Rath;P.Mallick and P.K.samal :The African Review of Physics**10:0007**,(2015).
- [3] H.B.Zhang,G.Y.Jiang and G.C.Wang J.Math.Phys**56**,072103(2015)(Use Eq(62;52) to verify our numerical results).
- [4] B.Rath and P.Mallick,Open.Phys.**14**,492(2016).