

the 1990s, the incidence of *S. flexneri* infections in the United Kingdom has increased, and the incidence of *S. flexneri* infection in the United States has increased in the 1980s and 1990s [10].

There is a paucity of data on the incidence of *S. flexneri* infection in the United Kingdom. In the 1980s, *S. flexneri* was the second most commonly isolated serotype of *Shigella* from patients with shigellosis in the United Kingdom [11]. In the 1990s, *S. flexneri* was the most commonly isolated serotype of *Shigella* from patients with shigellosis in the United Kingdom [12]. In the 1980s, *S. flexneri* was the second most commonly isolated serotype of *Shigella* from patients with shigellosis in the United States [13]. In the 1990s, *S. flexneri* was the most commonly isolated serotype of *Shigella* from patients with shigellosis in the United States [14].

The purpose of this study was to determine the incidence of *S. flexneri* infection in the United Kingdom in the 1990s. The study was conducted in the United Kingdom, where the incidence of *S. flexneri* infection has increased in the 1990s.

The study was conducted in the United Kingdom, where the incidence of *S. flexneri* infection has increased in the 1990s. The study was conducted in the United Kingdom, where the incidence of *S. flexneri* infection has increased in the 1990s.

The study was conducted in the United Kingdom, where the incidence of *S. flexneri* infection has increased in the 1990s. The study was conducted in the United Kingdom, where the incidence of *S. flexneri* infection has increased in the 1990s.

The study was conducted in the United Kingdom, where the incidence of *S. flexneri* infection has increased in the 1990s. The study was conducted in the United Kingdom, where the incidence of *S. flexneri* infection has increased in the 1990s.

The study was conducted in the United Kingdom, where the incidence of *S. flexneri* infection has increased in the 1990s. The study was conducted in the United Kingdom, where the incidence of *S. flexneri* infection has increased in the 1990s.

The study was conducted in the United Kingdom, where the incidence of *S. flexneri* infection has increased in the 1990s. The study was conducted in the United Kingdom, where the incidence of *S. flexneri* infection has increased in the 1990s.

The study was conducted in the United Kingdom, where the incidence of *S. flexneri* infection has increased in the 1990s. The study was conducted in the United Kingdom, where the incidence of *S. flexneri* infection has increased in the 1990s.

The study was conducted in the United Kingdom, where the incidence of *S. flexneri* infection has increased in the 1990s. The study was conducted in the United Kingdom, where the incidence of *S. flexneri* infection has increased in the 1990s.

The study was conducted in the United Kingdom, where the incidence of *S. flexneri* infection has increased in the 1990s. The study was conducted in the United Kingdom, where the incidence of *S. flexneri* infection has increased in the 1990s.