Descriptives for model cohort

Cohort Definition

Xilin Chen

2021-06-29

Contents

Table1		 1
Case level		
Surgeon level		
By top 10 procedu	res	1

Table1

Case level

Characteristic	**Failed**, $N = 19,286$	**Passed**, $N = 768,132$	**p-value**
flg_male	9,342 (48%)	323,150 (42%)	< 0.001
age_at_admit	75 (70, 81)	76 (70, 82)	< 0.001
AHRQ_score	8 (2, 17)	9 (1, 19)	0.019
race_white	16,833 (87%)	670,902 (87%)	0.8
ses			< 0.001
high_ses	6,780 (35%)	346,373 (45%)	
low_ses	12,506 (65%)	421,759 (55%)	
emergent_admit	7,777 (40%)	384,372 (50%)	< 0.001
had_assist_surg	207 (1.1%)	5,569 (0.7%)	< 0.001

Surgeon level

Characteristic	**Failed**, $N = 817$	**Passed**, $N = 37,473$	**p-value**
surgeon_yearly_load	18 (10, 33)	15 (7, 27)	< 0.001

By top 10 procedures

procedure		percentage_in_pass_or_fail_grp
Small bowel resection with anastomosis (open)		7.0%
Partial colectomy with anastomosis (open)		6.0%
Partial colectomy with ileocolostomy (open)		4.0%
Partial colectomy with cecostomy or colostomy (open)	33023	4.0%
Splenic flexure takedown (in addition to primary procedure)		4.0%
Lysis of adhesions (open)		4.0%
Ventral Hernia Repair W/Mesh		3.0%
Ventral hernia repair (open)		3.0%
Cholecystectomy (open)		3.0%
Partial colectomy with coloproctostomy (open)	21897	3.0%