




B530-A




B585-A




B695-A




B780-A




R670-A




R780-A




Blue_LaserDelay




Red_LaserDelay




FSC-A




FSC-H




FSC-W




SSC-A




SSC-H




SSC-W




B530-A_Gain




B585-A_Gain




B695-A_Gain




B780-A_Gain




R670-A_Gain




R780-A_Gain




FSC-A_Gain




SSC-A_Gain




Blue_LaserDelay




Red_LaserDelay




Blue_AreaScalingFactor




Red_AreaScalingFactor




B530-A-% rCV




B585-A-% rCV




B695-A-% rCV




B780-A-% rCV




R670-A-% rCV




R780-A-% rCV




FSC-A-% rCV




FSC-H-% rCV



FSC-W-% rCV



SSC-A-% rCV



The graph displays the daily number of COVID-19 cases in the United States. The y-axis is on a logarithmic scale, with major ticks at 100,000 and 1,000,000. The x-axis spans from October to April. A red dashed vertical line is positioned at the beginning of the data series in early November. The data shows a period of fluctuation between approximately 100,000 and 500,000 cases from November to early January. A massive spike occurs in early January, peaking at over 1,000,000 cases. Following this peak, the number of cases drops sharply and remains relatively low, fluctuating between 100,000 and 200,000 through April.

The graph displays the daily count of COVID-19 cases in the United States. The y-axis is labeled with values 0, 10, 15, and 20. The x-axis is labeled with months: Oct, Jan, and Apr. A red dashed vertical line is positioned at approximately January 20, 2020. The data shows a period of low activity from late September to early January, followed by a rapid increase in cases, peaking at over 20 cases in early February. After this peak, the number of cases drops sharply and remains relatively low, with minor fluctuations, through April.