




B530-A




B585-A




B695-A




B780-A




R670-A




R780-A




Blue_LaserDelay




Red_LaserDelay




FSC-A




FSC-H




FSC-W




SSC-A




SSC-H




SSC-W




B530-A_Gain




B585-A_Gain




B695-A_Gain




B780-A_Gain



R670-A_Gain



R780-A_Gain



FSC-A_Gain






Figure 1 is a scatter plot showing the time series of the difference between the estimated and true values of the parameter α . The y-axis ranges from -0.050 to 0.050. The x-axis shows months from October to April. Two vertical dashed red lines are drawn at approximately October 15 and November 15. The data points are clustered around zero, with a slight increase in variance after the second dashed line.

Scatter plot showing monthly mean precipitation (mm) for the period 1979-2019. The y-axis represents precipitation in mm, ranging from 29 to 32. The x-axis represents months, with labels for Oct, Jan, and Apr. Two vertical dashed red lines are present at approximately July and August. The data points show a seasonal cycle, with higher precipitation (around 32.5 mm) occurring from October to June, and lower precipitation (around 28.5 mm) occurring from July to April.


Blue_AreaScalingFactor




Red_AreaScalingFactor




B530-A-% rCV




B585-A-% rCV




B695-A-% rCV




B780-A-% rCV




R670-A-% rCV




R780-A-% rCV




FSC-A-% rCV




FSC-H-% rCV




FSC-W-% rCV



SSC-A-% rCV



SSC-H-% rCV



SSC-W-% rCV

