




B530-A




B585-A




B695-A




B780-A




R670-A




R780-A




Blue_LaserDelay




Red_LaserDelay




FSC-A




FSC-H




FSC-W




SSC-A




SSC-H




SSC-W




B530-A_Gain




B585-A_Gain




B695-A_Gain




B780-A_Gain




R670-A_Gain




R780-A_Gain




FSC-A_Gain




SSC-A_Gain




Blue_LaserDelay




Red_LaserDelay




Blue_AreaScalingFactor



Red_AreaScalingFactor



B530-A-% rCV




The graph displays the daily number of COVID-19 cases in the United States from October to April. The y-axis represents the number of cases, and the x-axis represents time. A blue line with circular markers shows the daily case counts. A vertical dashed red line marks the date of the first case in Washington state (January 15, 2020). The graph shows a significant increase in cases starting in late October, peaking in early January, followed by a sharp decline and then a period of low, fluctuating case counts.


The graph displays the percentage of reads with coverage variation (%rCV) over time. The y-axis represents %rCV, ranging from 5 to 20. The x-axis shows months from October to April. A vertical dashed red line is positioned at approximately November 15. Before this line, the %rCV is highly variable, with values ranging from approximately 9 to 21. After the line, the %rCV drops significantly, remaining mostly between 5 and 7, with a sharp spike to 20 in late March.

The graph displays the percentage of replicates with a coefficient of variation greater than 10% (%rCV) over a period from October to April. The y-axis represents %rCV, ranging from 0 to 25. The x-axis shows months: Oct, Jan, and Apr. A vertical dashed red line is positioned at approximately November 15, indicating a significant change in the data. Before this line, the %rCV values are generally higher, fluctuating between approximately 14 and 26. After the line, the values are mostly lower, fluctuating between approximately 6 and 13, with a notable spike in late March.


R670-A-% rCV



R780-A-% rCV



FSC-A-% rCV



The graph displays the daily number of COVID-19 cases in the United States from October to April. The x-axis represents time, with labels for October, January, and April. The y-axis represents the number of cases, with a scale from 0 to 100,000. A vertical dashed red line is positioned at the end of December, indicating the start of the epidemic curve. The data shows a significant increase in cases starting in late December, peaking in early January at approximately 100,000 cases, and then declining. There is a second, smaller peak in cases around mid-March, reaching approximately 20,000 cases, before declining again.

The graph displays the daily number of COVID-19 cases in the Netherlands. The x-axis is labeled with months: Oct, Jan, and Apr. The y-axis is labeled with case counts: 5, 10, and 15. A red dashed vertical line is positioned at approximately November 25th. The data shows a first wave peaking at about 12 cases in late November, followed by a decline to around 2 cases by early January. A second wave begins in late January, peaking at 15 cases in late March, and then declining again by April.

The graph displays the daily count of COVID-19 cases in the United States. The y-axis is labeled with values 7.5, 10.0, 12.5, and 15.0, representing the number of cases. The x-axis is labeled with months: Oct, Jan, and Apr. A red dashed vertical line is positioned at the beginning of the data series in early November. The data shows a period of fluctuation between approximately 6.5 and 11.5 cases from late October to late November. Following the red line, there is a rapid ascent to a peak of about 15.0 cases in early January. This is followed by a steep decline to around 7.0 cases by mid-January, and then a period of relative stability, with cases generally staying between 6.5 and 8.5 from February through April.

The graph displays the daily count of COVID-19 cases in the United States. The data begins in late October at approximately 7 cases. It shows a period of fluctuation with a peak of about 14 cases in early January. A major spike occurs in early February, reaching over 20 cases. Following this, the case count drops sharply and remains relatively low, fluctuating between 2 and 8 cases through April.