Check fitted flocker model

Specify model to check:

```
spec <- params$spec #final_species_eco_sorted[6] # 1
spec
```

```
[1] "Lazuli Bunting"
```

```
buffer_km <- params$buffer_km
buffer_km
```

```
[1] 750
```

```
fold <- params$fold
# fold

results_dir <- params$results_dir
results_dir
```

```
[1] "M:/Documents/DEBTs/analysis/Schifferle_BBS_occupancy_models_2023/results/full_model"
```

Load fitted model:

```
load(file = file.path(results_dir,
                      paste0("out_", spec, "_fm_buffer", buffer_km, ".RData"))) # xx adjus
#print(out)

posterior <- brms::as_draws(out)
```

Check whether MCMC worked:

Do we have divergent transitions?

```
[1] "divergent transitions: 0"
```

Explorations regarding divergent transitions:

...

Is effective number of MCMC samples large enough:

The larger the ratio of n_eff to N (number of samples), the better.




```
[1] "effective sample size too small"
```

```
[1] "n_eff ratio bulk ESS:"
```

```
   Min. 1st Qu.  Median    Mean 3rd Qu.    Max. 
0.09984 0.51714 0.64311 0.74803 0.85688 2.01462 
```

```
[1] "n_eff ratio tail ESS:"

   Min. 1st Qu.  Median    Mean 3rd Qu.    Max. 
 0.2068  0.5993  0.6628  0.6519  0.7173  0.8717 

[1] "effective sample size in bulk or tail too small"
```

Do chains mix well:

```
No divergences to plot.
No divergences to plot.
```



No divergences to plot.


b_colo_Intercept
b_colo_bio1
b_colo_bio2
b_colo_bio3
b_colo_bio7
b_colo_bio14
b_colo_bio15
b_colo_pr_spring
b_colo_pr_summer
b_colo_pr_autumn
b_colo_pr_winter
b_colo_Ibio1E2
b_colo_Ibio2E2
b_colo_Ibio3E2
b_colo_Ibio7E2
b_colo_Ibio14E2
b_colo_Ibio15E2
b_colo_Ipr_springE2
b_colo_Ipr_summerE2
b_colo_Ipr_autumnE2
b_colo_Ipr_winterE2
_colo_sum_annual_crop
b_colo_secdf
b_colo_pastr
b_colo_urban
olo_Isum_annual_crop
b_colo_IsecdfE2
b_colo_IpastrE2
b_colo_IurbanE2
Chain
1
2
3
4
Post−warmup iteration



b_ex_Intercept
b_ex_bio1
b_ex_bio2
b_ex_bio3
b_ex_bio7
b_ex_bio14
b_ex_bio15
b_ex_pr_spring
b_ex_pr_summer
b_ex_pr_autumn
b_ex_pr_winter
b_ex_Ibio1E2
b_ex_Ibio2E2
b_ex_Ibio3E2
b_ex_Ibio7E2
b_ex_Ibio14E2
b_ex_Ibio15E2
b_ex_Ipr_springE2
b_ex_Ipr_summerE2
b_ex_Ipr_autumnE2
b_ex_Ipr_winterE2
)_ex_sum_annual_crops
b_ex_secdf
b_ex_pastr
b_ex_urban
ex_Isum_annual_crops
b_ex_IsecdfE2
b_ex_IpastrE2
b_ex_IurbanE2
Chain
1
2
3
4
Post−warmup iteration

Are R-hat values fine:

```
   Min. 1st Qu.  Median    Mean 3rd Qu.    Max. 
 0.9991  0.9999  1.0002  1.0009  1.0010  1.0136 
```



Print model posteriors

```
[1] "parameters for which uncertainty interval doesn't overlap zero:"

[1] "b_occ_Intercept"                 "b_occ_bio1_3yrs"                
[3] "b_occ_pr_summer_3yrs"            "b_occ_Ibio3_3yrsE2"             
[5] "b_occ_Ipr_winter_3yrsE2"         "b_occ_Isum_annual_crops_3yrsE2"

[1] "b_colo_bio1"                 "b_colo_pr_summer"           
[3] "b_colo_Ibio3E2"              "b_colo_Ipr_autumnE2"        
[5] "b_colo_sum_annual_crops"     "b_colo_secdf"               
[7] "b_colo_Isum_annual_cropsE2" "b_colo_IpastrE2"            
[9] "b_colo_IurbanE2"            

[1] "b_ex_bio1"              "b_ex_Ibio1E2"           "b_ex_sum_annual_crops"
[4] "b_ex_IsecdfE2"          "b_ex_IpastrE2"          "b_ex_IurbanE2"        
```



b_occ_bio1_3yrs
b_occ_bio2_3yrs
b_occ_bio3_3yrs
b_occ_bio7_3yrs
b_occ_bio14_3yrs
b_occ_bio15_3yrs
b_occ_pr_spring_3yrs
b_occ_pr_summer_3yrs
b_occ_pr_autumn_3yrs
b_occ_pr_winter_3yrs
b_occ_Ibio1_3yrsE2
b_occ_Ibio2_3yrsE2
b_occ_Ibio3_3yrsE2
b_occ_Ibio7_3yrsE2
b_occ_Ibio14_3yrsE2
b_occ_Ibio15_3yrsE2
b_occ_Ipr_spring_3yrsE2
b_occ_Ipr_summer_3yrsE2
b_occ_Ipr_autumn_3yrsE2
b_occ_Ipr_winter_3yrsE2
b_occ_sum_annual_crops_3yrs
b_occ_secdf_3yrs
b_occ_pastr_3yrs
b_occ_urban_3yrs
b_occ_Isum_annual_crops_3yrsE2
b_occ_Isecdf_3yrsE2
b_occ_Ipastr_3yrsE2
b_occ_Iurban_3yrsE2
−5.0
−2.5
0.0
2.5


b_colo_bio1
b_colo_bio2
b_colo_bio3
b_colo_bio7
b_colo_bio14
b_colo_bio15
b_colo_pr_spring
b_colo_pr_summer
b_colo_pr_autumn
b_colo_pr_winter
b_colo_Ibio1E2
b_colo_Ibio2E2
b_colo_Ibio3E2
b_colo_Ibio7E2
b_colo_Ibio14E2
b_colo_Ibio15E2
b_colo_Ipr_springE2
b_colo_Ipr_summerE2
b_colo_Ipr_autumnE2
b_colo_Ipr_winterE2
b_colo_sum_annual_crops
b_colo_secdf
b_colo_pastr
b_colo_urban
b_colo_Isum_annual_cropsE2
b_colo_IsecdfE2
b_colo_IpastrE2
b_colo_IurbanE2
−5.0
−2.5
0.0
2.5


b_ex_bio1
b_ex_bio2
b_ex_bio3
b_ex_bio7
b_ex_bio14
b_ex_bio15
b_ex_pr_spring
b_ex_pr_summer
b_ex_pr_autumn
b_ex_pr_winter
b_ex_Ibio1E2
b_ex_Ibio2E2
b_ex_Ibio3E2
b_ex_Ibio7E2
b_ex_Ibio14E2
b_ex_Ibio15E2
b_ex_Ipr_springE2
b_ex_Ipr_summerE2
b_ex_Ipr_autumnE2
b_ex_Ipr_winterE2
b_ex_sum_annual_crops
b_ex_secdf
b_ex_pastr
b_ex_urban
b_ex_Isum_annual_cropsE2
b_ex_IsecdfE2
b_ex_IpastrE2
b_ex_IurbanE2
−6
−3
0
3