

Check fitted flocker model

Specify model to check:

```
spec <- params$spec #final_species_eco_sorted[6] # 1
spec
```

```
[1] "Blue-winged Warbler"
```

```
buffer_km <- params$buffer_km
buffer_km
```

```
[1] 750
```

```
fold <- params$fold
# fold

results_dir <- params$results_dir
results_dir
```

```
[1] "M:/Documents/DEBTs/analysis/Schifferle_BBS_occupancy_models_2023/results/full_model"
```

Load fitted model:

```
load(file = file.path(results_dir,
                       paste0("out_", spec, "_fm_buffer", buffer_km, ".RData"))) # xx adjust
#print(out)

posterior <- brms::as_draws(out)
```

Check whether MCMC worked:

Do we have divergent transitions?


```
[1] "divergent transitions: 0"
```

Explorations regarding divergent transitions:

...

Is effective number of MCMC samples large enough:

The larger the ratio of n_{eff} to N (number of samples), the better.



```
[1] "effective sample size too small"
```

```
[1] "n_eff ratio bulk ESS:"
```

Min.	1st Qu.	Median	Mean	3rd Qu.	Max.
0.2931	0.6917	0.8205	0.8810	0.9661	2.0300

```
[1] "n_eff ratio tail ESS:"
```

	Min.	1st Qu.	Median	Mean	3rd Qu.	Max.
	0.07764	0.69401	0.74334	0.72974	0.78552	0.87557

```
[1] "effective sample size in bulk or tail too small"
```

Do chains mix well:

No divergences to plot.

No divergences to plot.




No divergences to plot.





Are R-hat values fine:

Min.	1st Qu.	Median	Mean	3rd Qu.	Max.
0.9993	0.9996	1.0001	1.0004	1.0005	1.0198



Print model posteriors

```
[1] "parameters for which uncertainty interval doesn't overlap zero:"
```

```
[1] "b_occ_Ibio1_3yrsE2" "b_occ_secdf_3yrs" "b_occ_Isecdf_3yrsE2"
```

```
[1] "b_colo_bio1" "b_colo_pr_summer" "b_colo_Ibio1E2" "b_colo_Ibio7E2"
```

```
[5] "b_colo_secdf" "b_colo_pastr" "b_colo_urban" "b_colo_IpastrE2"
```

```
[1] "b_ex_bio1" "b_ex_Ibio1E2" "b_ex_secdf" "b_ex_IsecdfE2"
```

