

Check fitted flocker model

Specify model to check:

```
spec <- params$spec #final_species_eco_sorted[6] # 1
spec
```

```
[1] "Golden Eagle"
```

```
buffer_km <- params$buffer_km
buffer_km
```

```
[1] 750
```

```
fold <- params$fold
# fold

results_dir <- params$results_dir
results_dir
```

```
[1] "M:/Documents/DEBTs/analysis/Schifferle_BBS_occupancy_models_2023/results/full_model"
```

Load fitted model:

```
load(file = file.path(results_dir,
                       paste0("out_", spec, "_fm_buffer", buffer_km, ".RData"))) # xx adjust
#print(out)

posterior <- brms::as_draws(out)
```

Check whether MCMC worked:

Do we have divergent transitions?


```
[1] "divergent transitions: 0"
```

Explorations regarding divergent transitions:

...

Is effective number of MCMC samples large enough:

The larger the ratio of n_{eff} to N (number of samples), the better.



```
[1] "effective sample size too small"
```

Warning: Parts of the model have not converged (some Rhats are > 1.05). Be careful when analysing the results! We recommend running more iterations and/or setting stronger priors.

```
[1] "n_eff ratio bulk ESS:"
```

Min.	1st Qu.	Median	Mean	3rd Qu.	Max.
0.004975	0.015733	0.036190	0.046602	0.062960	0.201719

[1] "n_eff ratio tail ESS:"

Min.	1st Qu.	Median	Mean	3rd Qu.	Max.
0.005392	0.052317	0.108455	0.149588	0.224167	0.522231

[1] "effective sample size in bulk or tail too small"


Do chains mix well:

No divergences to plot.

No divergences to plot.



No divergences to plot.




Post-warmup iteration



Are R-hat values fine:

Min.	1st Qu.	Median	Mean	3rd Qu.	Max.
1.000	1.010	1.025	1.038	1.052	1.154

```
[1] "R-hat > 1.05 for 26 parameters."
```



Print model posteriors

```
[1] "parameters for which uncertainty interval doesn't overlap zero:"
```

```
character(0)
```

```
[1] "b_colo_Ibio1E2" "b_colo_Ibio2E2"
```

```
[1] "b_ex_Ibio1E2"
```

