

Check fitted flocker model

Specify model to check:

```
spec <- params$spec #final_species_eco_sorted[6] # 1
spec
```

```
[1] "Pine Warbler"
```

```
buffer_km <- params$buffer_km
buffer_km
```

```
[1] 750
```

```
fold <- params$fold
# fold

results_dir <- params$results_dir
results_dir
```

```
[1] "M:/Documents/DEBTs/analysis/Schifferle_BBS_occupancy_models_2023/results/full_model"
```

Load fitted model:

```
load(file = file.path(results_dir,
                       paste0("out_", spec, "_fm_buffer", buffer_km, ".RData"))) # xx adjust
#print(out)

posterior <- brms::as_draws(out)
```

Check whether MCMC worked:

Do we have divergent transitions?


```
[1] "divergent transitions: 0"
```

Explorations regarding divergent transitions:

...

Is effective number of MCMC samples large enough:

The larger the ratio of n_{eff} to N (number of samples), the better.



```
[1] "effective sample size fine"
```

```
[1] "n_eff ratio bulk ESS:"
```

Min.	1st Qu.	Median	Mean	3rd Qu.	Max.
0.6203	1.0584	1.2414	1.3164	1.4498	3.2180

```
[1] "n_eff ratio tail ESS:"
```

Min.	1st Qu.	Median	Mean	3rd Qu.	Max.
0.5609	0.7132	0.7632	0.7541	0.7973	0.9459

```
[1] "effective sample size in bulk and tail fine"
```


Do chains mix well:


No divergences to plot.

No divergences to plot.



No divergences to plot.





Are R-hat values fine:

Min.	1st Qu.	Median	Mean	3rd Qu.	Max.
0.9991	0.9994	0.9997	0.9997	0.9999	1.0015

```
[1] "All R-hat values < 1.01"
```

Print model posteriors

```
[1] "parameters for which uncertainty interval doesn't overlap zero:"
```

```
[1] "b_occ_Intercept" "b_occ_Ibio14_3yrsE2"
```

```
[3] "b_occ_sum_annual_crops_3yrs" "b_occ_secdf_3yrs"
```

```
[5] "b_occ_Isecdf_3yrsE2"
```

```
[1] "b_colo_pr_autumn" "b_colo_sum_annual_crops"
```

```
[3] "b_colo_secdf" "b_colo_pastr"
```

```
[5] "b_colo_IpastrE2"
```

```
[1] "b_ex_pastr" "b_ex_Isum_annual_cropsE2"
```

