

Diabetes Part 2



Talent Management Department

صيدليـة المتحـدة
United Pharmacy

Diabetes drugs classification



*** (DPP-4) ➡** Dipeptidyl peptidase-4
**** SLGT-2 ➡** Sodium-dependent glucose transporter-2

Today we will learn about

06 Insulins

07 GLP-1 Analogues

08 Combinations

صيدلية المتحدة
United Pharmacy

Insulins

Insulin is a hormone replacement therapy used in Type 1 and advanced Type 2 Diabetes Mellitus to control blood glucose.

1. Rapid-Acting Insulin ✅
- Onset: 10–20 min
- Duration: 3–5 hrs
- Use: Taken just before meals.

2. Short-Acting Insulin ✅
- Onset: 30 min
- Duration: 4–6 hrs
- Use: 30 min before meals.

3. Intermediate-Acting Insulin ✅
- Onset: 1–2 hrs
- Duration: 12–16 hrs
- Use: Basal insulin (background control).

4. Long-Acting Insulin ✅
- Onset: 1–3 hrs
- Duration: Up to more than 24 hrs
- Use: Once daily for basal control.

5. Premixed Insulin ✅
- Combo of short/intermediate or rapid/ intermediate.
- Provides both meal-time and basal coverage.

Insulins

6

Short Acting

- **ACTRAPID®**
- **HUMULIN R®**

Intermediate Acting

- **INSULATARD®**
- **HUMULIN N®**
- **INSULATARD HM®**

Ultrashort Acting

- **HUMALOG®** | *Insulin Lispro*
- **APIDRA®** | *Insulin Glulisine*
- **NOVORAPID®, TRUVELOG®** | *Insulin Aspart*

Long Acting

- **RYZODEG® , TRESIBA®** | *Insulin Degludec*
- **LEVEMIR®** | *Insulin Detemir*
- **BASAGLAR® , VIVARO® , LANTUS®, TOUJEO®** | *Insulin Glargine*

Premixed

- **MIXTARD® 70/30 – 50/50**
- **HUMULIN® 70/30**
- **NOVOMIX® 70/30 – 50/50** | *Aspart + Aspart protamine*
- **HUMALOG MIX® 75/25 – 50/50** | *Lispro + lispro protamine*

GLP-1 Analogues

GLP-1 receptor agonists (also called incretin mimetics) are injectable drugs used in Type 2 Diabetes to improve blood sugar control and may be used also as weight loss support.

Mechanism of Action ✅

- They Increase Insulin secretion.
- Decrease Glucagon release.
- Slow gastric emptying rate.
- Increase satiety → ↓ appetite and weight.

Important Notes: ✅

- But may cause Nausea, vomiting, Diarrhea & Pancreatitis.
- Have a Blackbox warning of causing medullary thyroid carcinoma, must be avoided with patient with high risk or past history.
- They can be combined with Long-acting Insulin.
- They are better to have dose titration (Gradually increasing the dose) to avoid GI symptoms.

GLP-1 Analogues

GLP-1 Analogues

- **BYETTA®** | *Exenatide*
- **VICTOZA®** | *Liraglutide*
- **TRULICITY®** | *Dulaglutide*
- **OZEMPIC®** | *Semaglutide*
- **MOUNJARO®** | *Tirzepatide*
- **SAXENDA®** | *Liraglutide*
- **WEGOVY®** | *Semaglutide*
- **RYBELSUS®** | *Semaglutide*
  (The Only Oral Member)

Combinations

- **SOLIQUA®** | *Lixisenatide & Insulin Glargine*
- **XULTOPHY®** | *Liraglutide & Insulin Degludec*

الآن حان دور الأنشطة
اضغط على الزر للبدء



ابدأ النشاط