

Top13 Paracompactness, Complete Metric Spaces

KYB

Thrn, it's a Fact

mathrnfact@gmail.com

June 10, 2020

41. Paracompactness

Theorems and definitions

Exercises

41.

Paracompactness

Theorems and
definitions

Exercises

43. Complete Metric Spaces

Theorems and definitions

Exercises

43. Complete
Metric Spaces

Theorems and
definitions

Exercises

Recall

Definition

A collection \mathcal{A} of subsets of X is said to be locally finite in X if every point of X has a neighborhood that intersects only finitely many elements of \mathcal{A} .

Definition

Let \mathcal{A} be a collection of subsets of X . A collection \mathcal{B} of subsets of X is said to be a refinement of \mathcal{A} if for each element B of \mathcal{B} , there is an element A of \mathcal{A} containing B .

41.
ParacompactnessTheorems and
definitions

Exercises

43. Complete
Metric SpacesTheorems and
definitions

Exercises

Definition (Paracompact)

X is paracompact if every open covering \mathcal{A} of X has a locally finite open refinement \mathcal{B} that covers X .

41.

Paracompactness

Theorems and
definitions

Exercises

43. Complete
Metric SpacesTheorems and
definitions

Exercises

Theorem (41.1)

Every paracompact Hausdorff space X is normal.

Theorem (41.2)

Every closed subspace of a paracompact space is paracompact.

Note

- ▶ A paracompact subspace of a Hausdorff space X need not be closed in X .
- ▶ A subspace of a paracompact space need not be paracompact.

Lemma (41.3)

Let X be regular. TFAE:

Every open covering of X has a refinement that is:

- (1) An open covering of X and countably locally finite.
- (2) A covering of X and locally finite.
- (3) A closed covering of X and locally finite.
- (4) An open covering of X and locally finite.

Theorem (41.4)

Every metrizable space is paracompact.

Theorem (41.4)

Every regular Lindelöf space is paracompact.

Note

- ▶ The product of two paracompact spaces need not be paracompact.
- ▶ \mathbb{R}^ω is paracompact in both the product and uniform topologies.
- ▶ It is not known whether \mathbb{R}^ω is paracompact in the box topology.
- ▶ The product space \mathbb{R}^J is not paracompact if J is uncountable.

41.

Paracompactness

Theorems and
definitions

Exercises

43. Complete

Metric Spaces

Theorems and
definitions

Exercises

Definition

Let $\{U_\alpha\}$ be an indexed open covering of X . An indexed family of continuous functions $\phi_\alpha : X \rightarrow [0, 1]$ is said to be a partition of unity on X dominated by $\{U_\alpha\}$ if

1. $\text{supp}(\phi_\alpha) \subset U_\alpha$ for each α
2. $\{\text{supp}(\phi_\alpha)\}$ is locally finite
3. $\sum \phi_\alpha(x) = 1$ for each x .

Lemma (41.6 Shrinking lemma)

Let X be a paracompact Hausdorff space; let $\{U_\alpha\}_{\alpha \in J}$ be an indexed family of open sets covering X . Then there exists a locally finite indexed family $\{V_\alpha\}_{\alpha \in J}$ of open sets covering X such that $\bar{V}_\alpha \subset U_\alpha$ for each α .

Theorem (41.7)

Let X be a paracompact Hausdorff space; let $\{U_\alpha\}_{\alpha \in J}$ be an indexed family of open sets covering X . Then there exists a partition of unity on X dominated by $\{U_\alpha\}$.

41.

Paracompactness

Theorems and
definitions

Exercises

43. Complete

Metric Spaces

Theorems and
definitions

Exercises

Partitions of unity are most often used in mathematics to "patch together" functions that are defined locally so as to obtain a function that is defined globally.

Theorem (41.8)

Let X be a paracompact Hausdorff space; let \mathcal{C} be a collection of subsets of X ; for each $C \in \mathcal{C}$, let ϵ_C be a positive number. If \mathcal{C} is locally finite, there is a continuous function $f : X \rightarrow \mathbb{R}$ such that $f(x) > 0$ for all x , and $f(x) \leq \epsilon_C$ for $x \in C$.

41.

Paracompactness

Theorems and
definitions

Exercises

43. Complete
Metric SpacesTheorems and
definitions

Exercises

Ex41.1

Give an example to show that if X is paracompact, it does not follow that for every open covering \mathcal{A} of X , there is a locally finite subcollection of \mathcal{A} that covers X .

Proof.

In \mathbb{R} , $\mathcal{A} = \{(-\infty, n) : n \in \mathbb{Z}\}$.



41. Paracompactness

Theorems and
definitions

Exercises

43. Complete
Metric SpacesTheorems and
definitions

Exercises

Ex41.2(a)

Show that the product of a paracompact space and a compact space is paracompact.

Proof

Main idea : Given open cover \mathcal{A} , find refinement

$\{U_\alpha \times V_{\alpha,i} : i = 1, \dots, n_\alpha\}$ where $\bigcup V_{\alpha,i} = Y$.

Step1

Fix $x \in X$. Then we can find A_α 's in \mathcal{A} such that $\{x\} \times Y \subset \bigcup A_\alpha$. The trick is that

we can find $U_\alpha \subset X, V_\alpha \subset Y$ such that $\{x\} \times Y \subset \bigcup U_\alpha \times V_\alpha$
and $U_\alpha \times V_\alpha \subset A_\alpha$ for some $A_\alpha \in \mathcal{A}$.

Then by the compactness of Y , we can find U_i, V_i for $i = 1, \dots, n$ such that $\{x\} \times Y \subset \bigcup_1^n U_i \times V_i$. By the tube lemma, we can find $W_x \subset X$ such that

$$\{x\} \times Y \subset W_x \times Y \subset \bigcup_1^n U_i \times V_i$$

41.

Paracompactness

Theorems and
definitions

Exercises

43. Complete

Metric Spaces

Theorems and
definitions

Exercises

Step2

For each $x \in X$, we can find $\{W_x\}$ in Step1 that covers X . By the paracompactness, we can find a locally finite open refinement $\mathcal{B} = \{R_\alpha\}$ that covers X . Then for each x find open O_x that intersects only finitely many elements in \mathcal{B} . Now for each α , there is x_α such that $R_\alpha \subset W_{x_\alpha}$. Then define $\mathcal{C} = \{R_\alpha \times V_{x_\alpha, i}\}$

Step3

Claim) \mathcal{C} is a locally finite open refinement of \mathcal{A} .

- ▶ (refinement) $R_\alpha \times V_{x_\alpha, i} \subset W_{x_\alpha} \times V_{x_\alpha, i} \subset U_{x_\alpha, i} \times V_{x_\alpha, i} \subset A_{x_\alpha, i}$
- ▶ (open cover) Trivial.
- ▶ (locally finite)

Let $(x, y) \in X \times Y$. Then O_x intersects only finitely many R_α 's, say $R_{\alpha_1}, \dots, R_{\alpha_k}$. Then $O_x \times Y$ is the desired open neighborhood.

41.

Paracompactness

Theorems and
definitions

Exercises

43. Complete
Metric SpacesTheorems and
definitions

Exercises

Ex.41.5 Expansion lemma

Let $\{B_\alpha\}_{\alpha \in J}$ be a locally finite indexed family of subsets of the paracompact Hausdorff space X . Then there is a locally finite indexed family $\{U_\alpha\}$ of open sets in X such that $B_\alpha \subset U_\alpha$ for each α .

41.

Paracompactness

Theorems and
definitions

Exercises

43. Complete
Metric SpacesTheorems and
definitions

Exercises

Definition (Cauchy sequence)

Let (X, d) be a metric space. A sequence (x_n) of points of X is said to be a Cauchy sequence in (X, d) if it has the property that given $\epsilon > 0$, there is an integer N such that

$$d(x_n, x_m) < \epsilon \quad \text{whenever } n, m \geq N$$

The metric space (X, d) is said to be complete if every Cauchy sequence in X converges.

41.

Paracompactness

Theorems and
definitions

Exercises

43. Complete
Metric SpacesTheorems and
definitions

Exercises

Lemma (43.1)

A metric space X is complete if every Cauchy sequence in X has a convergent subsequence.

Proof

Given $\epsilon > 0$, choose N large enough that $d(x_n, x_m) < \epsilon/2$ for all $n, m \geq N$. Then choose an i large enough that $n_i \geq N$ and $d(x_{n_i}, x) < \epsilon/2$. Then for $n \geq N$,

$$d(x_n, x) \leq d(x_n, x_{n_i}) + d(x_{n_i}, x) < \epsilon.$$

41.

Paracompactness

Theorems and
definitions

Exercises

43. Complete
Metric SpacesTheorems and
definitions

Exercises

Lemma (43.3)

Let X be the product space $X = \prod X_\alpha$; let x_n be a sequence of points of X . Then $x_n \rightarrow x$ if and only if $\pi_\alpha(x_n) \rightarrow \pi_\alpha(x)$.

41.

Paracompactness

Theorems and
definitions

Exercises

43. Complete
Metric SpacesTheorems and
definitions

Exercises

Definition

Let (Y, d) be a metric space; let $\bar{d}(a, b) = \min\{d(a, b), 1\}$ be the standard bounded metric on Y derived from d . If $x = (x_\alpha)_{\alpha \in J}$ and $y = (y_\alpha)_{\alpha \in J}$ are points of Y^J , let

$$\bar{\rho}(x, y) = \sup\{\bar{d}(x_\alpha, y_\alpha) : \alpha \in J\}.$$

Theorem (43.5)

If Y is complete d , then the space Y^J is complete in $\bar{\rho}$.

Definition

$$\mathcal{C}(X, Y) \subset Y^X$$

is the set of all continuous functions from X to Y .

$$\mathcal{B}(X, Y) \subset Y^X$$

is the set of all bounded functions from X to Y .

Definition

Let M be a Euclidean space (\mathbb{R}^n) or smooth n -manifold. Then we define \mathcal{C}^k be the set of all function f such that f has n th derivative $f^{(n)}$ and $f^{(n)}$ is continuous.

Then

$$\mathcal{C} = \mathcal{C}^{(0)} \supset \mathcal{C}^{(1)} \supset \dots$$

Define $\mathcal{C}^\infty = \bigcap \mathcal{C}^n$ the set of all smooth functions.

Theorem (43.6)

X : top'l. (Y, d) metric.

- ▶ $\mathcal{C}(X, Y)$ is closed in Y^X under $\bar{\rho}$.
- ▶ $\mathcal{B}(X, Y)$ is closed in Y^X under $\bar{\rho}$.

Therefore, if Y is complete, these spaces are complete in $\bar{\rho}$.

Definition

(Y, d) metric. We can define another metric on $\mathcal{B}(X, Y)$ by

$$\rho(f, g) = \sup\{d(f(x), g(x)) \mid x \in X\}$$

For $f, g \in \mathcal{B}(X, Y)$, $\rho(\bar{f}, g) = \min\{\rho(f, g), 1\}$

41.

Paracompactness

Theorems and
definitions

Exercises

43. Complete
Metric SpacesTheorems and
definitions

Exercises

Ex43.4

Show that the metric space (X, d) is complete if and only if for every nested sequence $A_1 \supset A_2 \supset \cdots$ of nonempty closed sets of X such that $\text{diam} A_n \rightarrow 0$, the intersection of the sets A_n is nonempty.

41.

Paracompactness

Theorems and
definitions

Exercises

43. Complete
Metric SpacesTheorems and
definitions

Exercises

Ex43.5

If (X, d) is a metric space, recall that a map $f : X \rightarrow X$ is called a contraction if there is a number $\alpha < 1$ such that

$$d(f(x), f(y)) \leq \alpha d(x, y)$$

for all $x, y \in X$. Show that if f is a contraction of a complete metric space, then there is a unique point $x \in X$ such that $f(x) = x$.

41.
ParacompactnessTheorems and
definitions

Exercises

43. Complete
Metric SpacesTheorems and
definitions

Exercises

Ex43.8

Define $e : X \times \mathcal{C}(X, Y) \rightarrow Y$ be the equation $e(x, f) = f(x)$. We call e the evaluation map. Show that if d is a metric for Y and $\mathcal{C}(X, Y)$ has the corresponding uniform topology, then e is continuous.

41.

Paracompactness

Theorems and
definitions

Exercises

43. Complete
Metric SpacesTheorems and
definitions

Exercises

Ex43.9 Another completion

(X, d) :metric space. Let \tilde{X} be the set of all Cauchy sequences $x = (x_n)_n$. Define $x \sim y$ if $d(x_n, y_n) \rightarrow 0$. Let $[x]$ be the equivalence class of x ; let $Y = \tilde{X}/\sim$. Define a metric D on Y by

$$D([x], [y]) = \lim d(x_n, y_n)$$

Then Y is complete and there is an isometric imbedding $h : X \rightarrow Y$

In the set theory, there are two way to construct real number. One is 'Dedekind cut' and the other is 'completion'.

41.

Paracompactness

Theorems and
definitions

Exercises

43. Complete
Metric SpacesTheorems and
definitions

Exercises

(a)

Show that \sim is an equivalence relation, and show that D is a well-defined metric

41.

Paracompactness

Theorems and
definitions

Exercises

43. Complete
Metric SpacesTheorems and
definitions

Exercises

(b)

Define $h : X \rightarrow Y$ by letting $h(x)$ be the equivalence class of the constant sequence (x, x, \dots) . Show that h is an isometric imbedding.

41.

Paracompactness

Theorems and
definitions

Exercises

43. Complete
Metric SpacesTheorems and
definitions

Exercises

(c)

Show that $h(X)$ is dense in Y ; indeed, given $x = (x_n)_n \in \tilde{X}$, show the $h(x_n)$ of point of Y converges to the point $[x]$.

41.

Paracompactness

Theorems and
definitions

Exercises

43. Complete
Metric SpacesTheorems and
definitions

Exercises

(d)

Show that if A is a dense subset of a metric space (Z, ρ) , and if every Cauchy sequence in A converges in Z , then Z is complete.

41.

Paracompactness

Theorems and
definitions

Exercises

43. Complete

Metric Spaces

Theorems and
definitions

Exercises

(e)

Show that (Y, D) is complete.

41.

Paracompactness

Theorems and
definitions

Exercises

43. Complete
Metric SpacesTheorems and
definitions

Exercises

Ex43.10 Uniqueness of the completion

Let $h : X \rightarrow Y$ and $h' : X \rightarrow Y'$ be isometric imbeddings of the metric space (X, d) in the complete metric spaces (Y, D) and (Y', D') , respectively. Then there is an isometry of $(h(X), D)$ with $(h'(X), D')$ that equals $h'h^{-1}$ in the subspace $h(X)$.

41.

Paracompactness

Theorems and
definitions

Exercises

43. Complete
Metric SpacesTheorems and
definitions

Exercises

Bump Function

Consider $f: \mathbb{R} \rightarrow \mathbb{R}$ such that

$$f(x) = \begin{cases} e^{-1/x} & x > 0 \\ 0 & x \leq 0 \end{cases}$$

Prove that f is smooth.

41.

Paracompactness

Theorems and
definitions

Exercises

43. Complete

Metric Spaces

Theorems and
definitions

Exercises

Bump function

Given $r_1 < r_2$, define $h : \mathbb{R} \rightarrow \mathbb{R}$ such that

$$h(x) = \frac{f(r_2 - x)}{f(r_2 - x) + f(x - r_1)}.$$

Prove that h is smooth.

41.

Paracompactness

Theorems and
definitions

Exercises

43. Complete

Metric Spaces

Theorems and
definitions

Exercises

Bump function

Given any real number r_1 and r_2 such that $r_1 < r_2$, define $H : \mathbb{R}^n \rightarrow \mathbb{R}$ such that

$$H(x) = h(|x|).$$

Then

$$\begin{aligned} H &\equiv 1 && \text{on } \overline{B}(0, r_1) \\ 0 < H < 1 && \text{for all } x \in B(0, r_2) - \overline{B}(0, r_1) \text{ .} \\ H &\equiv 0 && \text{on } \mathbb{R}^n - B(0, r_2) \end{aligned}$$

and H is smooth.

41.

Paracompactness

Theorems and
definitions

Exercises

43. Complete

Metric Spaces

Theorems and
definitions

Exercises