

LA4 Linear Operator, Matrix

KYB

Thrn, it's a Fact

mathrnfact@gmail.com

February 16, 2021

Overview

Ch2. Fields and vector spaces

Ch3. Linear Operators

3.1 Linear Operators

3.2 More Properties of linear operators

3.3 Isomorphic Vector Spaces

3.4 Linear Operator Equations

Ex 2.6.11, 2.7.13

Let F be a finite field with $|F| = q$. Consider $\dim(\mathcal{P}_n(F))$.

(1) If $n \leq q - 1$, then $\{1, x, \dots, x^n\}$ is linearly independent. So $\dim(\mathcal{P}_n(F)) = n + 1$.

Proof.

Suppose $a_0 + a_1x + \dots + a_nx^n = 0$. Since every nonzero polynomial has at most “ n ” roots, $a_0 = a_1 = \dots = a_n = 0$. □

Ex 2.6.11, 2.7.13

Let F be a finite field with $|F| = q$. Consider $\dim(\mathcal{P}_n(F))$.

(2) If $n \geq q$, then $\{1, x, \dots, x^{q-1}\}$ is linearly independent. So $\dim(\mathcal{P}_n(F)) \geq q$. In fact, $\dim(\mathcal{P}_n(F)) = q$.

Proof.

Suppose $F = \{\alpha_1, \dots, \alpha_q\}$. Then any function $f : F \rightarrow F$ is determined by

$$(f(\alpha_1), \dots, f(\alpha_q)) \in F^q.$$

Thus there are at most $|F^q|$ many polynomials. On the other hand, $\dim(\mathcal{P}_n(F)) \geq q$ implies there are at least $|F^q|$ many polynomials. Hence $\dim(\mathcal{P}_n(F)) = q$.

(Note that every $v \in F^q$ can be identified with a function $v : F \rightarrow F$ by $v(\alpha_i) = v_i$).



Linear Operator

$$L(\alpha x + \beta y) = \alpha L(x) + \beta L(y).$$

Example

1. $C^1(\mathbb{R}) \rightarrow C^0(\mathbb{R})$ by $f \mapsto \frac{df}{dx}$.
2. $C^0[0, 1] \rightarrow \mathbb{R}$ by $f \mapsto \int_0^1 f(x)dx$.

$m \times n$ matrix, $F^{m \times n}$

$$\begin{array}{c}
 \rightarrow \\
 \rightarrow \\
 m \quad \vdots \\
 \rightarrow
 \end{array}
 \begin{array}{c}
 \downarrow \quad \downarrow \quad \downarrow \\
 \left[\begin{array}{ccc}
 A_{11} & \cdots & A_{1n} \\
 A_{21} & \ddots & \vdots \\
 \vdots & \ddots & \vdots \\
 A_{m1} & \cdots & A_{mn}
 \end{array} \right]
 \end{array}
 \begin{array}{c}
 n \\
 \\
 \\
 \\
 \end{array}
 = A,$$

$A_{ij} \in \mathbb{R}.$

$$A_j = \begin{bmatrix} A_{1j} \\ \vdots \\ A_{mj} \end{bmatrix} = (A_{1j}, \dots, A_{mj}) \in F^m, \quad r_i = [A_{i1} | \cdots | A_{in}]$$

$m \times n$ matrix, $F^{m \times n}$

$$Ax = \begin{bmatrix} A_{11} & \cdots & A_{1n} \\ \vdots & \ddots & \vdots \\ A_{m1} & \cdots & A_{mn} \end{bmatrix} \begin{bmatrix} x_1 \\ \vdots \\ x_n \end{bmatrix} = x_1 A_1 + \cdots + x_n A_n$$

$$\implies Ax = \sum_{i=1}^n x_i A_i, \quad (Ax)_i = \sum_{j=1}^n A_{ij} x_j$$

For $A \in F^{m \times n}$, $B \in F^{n \times l}$,

$$AB = [AB_1 | \cdots | AB_l] \in F^{m \times l}.$$

$$A^T = F^{n \times m}, \text{ and } (A^T)_{ij} = A_{ji}.$$

Ex 3.1.8

Let $L : \mathcal{P}_n \rightarrow \mathcal{P}_{2n-1}$ be defined by $L(p) = pp'$. Is L linear or nonlinear?.

Ex 3.1.10

(a) $A \in \mathbb{R}^{3 \times 3}$, $b \in \mathbb{R}^4$,

$$A = \begin{bmatrix} 1 & -2 & -4 \\ 5 & -11 & -15 \\ -2 & 6 & -1 \end{bmatrix}, b = \begin{bmatrix} 4 \\ 23 \\ -14 \end{bmatrix}.$$

Ex 3.1.10

(b) $A \in \mathbb{Z}^{3 \times 3}$, $b \in \mathbb{Z}^3$,

$$A = \begin{bmatrix} 2 & 1 & 2 \\ 1 & 0 & 1 \\ 1 & 2 & 0 \end{bmatrix}, b = \begin{bmatrix} 0 \\ 1 \\ 2 \end{bmatrix}.$$

Ex 3.1.12

Give a formula for $(AB)_{ij}$, assuming $A \in F^{m \times n}$, $B \in F^{n \times p}$.

Theorem

Let X and U be vector spaces over a field F .

- 1. If $L : X \rightarrow U$ is linear and $\alpha \in F$, then αL is also linear.*
- 2. If $L, M : X \rightarrow U$ are linear, then so is $L + M$.*

Remark

- Let $\mathcal{L}(X, U)$ be the set of all linear operator from X to U . Then it is a vector space over F .*
- For a vector space X , $\mathcal{L}(X, F)$ is called the (algebraic) “dual space” of X , and write $X^* = \mathcal{L}(X, F)$.*

The Matrix of a Linear Operator on Euclidean Spaces

Let $L : F^n \rightarrow F^m$ be linear. We can find a matrix $A \in F^{m \times n}$ such that $L(x) = Ax$ as follows:

$$L(e_i) = a_{i1}e_1 + \cdots + a_{im}e_m$$

$$A = \begin{bmatrix} a_{11} & \cdots & a_{1n} \\ a_{21} & \ddots & \vdots \\ \vdots & & \vdots \\ a_{m1} & \cdots & a_{mn} \end{bmatrix}$$

Then $L(e_i) = A_i$ and

$$L\left(\sum_{i=1}^n \alpha_i e_i\right) = \sum_{i=1}^n \alpha_i L(e_i) = \sum_{i=1}^n \alpha_i A_i = A \begin{bmatrix} \alpha_1 \\ \vdots \\ \alpha_n \end{bmatrix} = Ax$$

Hence, $L(x) = Ax$.

Associated Matrix with Bases

Let $L : X \rightarrow U$ be linear and let $\mathcal{X} = \{x_1, \dots, x_n\}$, $\mathcal{U} = \{u_1, \dots, u_m\}$ be bases for X and U , respectively. Then

$$L(x) = L\left(\sum_{i=1}^n \alpha_i x_i\right) = \sum_{i=1}^n \alpha_i L(x_i)$$

and

$$L(x_i) = \sum_{j=1}^m \beta_{ij} u_j.$$

Associated Matrix with Bases

Let $E_X : X \rightarrow F^n$ and $E_U : U \rightarrow F^m$ by $E_X(x_i) = e_i$, $E_U(u_j) = e_j$. Then

$$\begin{array}{ccccc}
 x_i & X & \xrightarrow{L} & U & u_j \\
 \downarrow & \downarrow E_X & & \downarrow E_U & \downarrow \\
 e_i & F^n & \xrightarrow{A} & F^m & e_j
 \end{array}$$

So

$$L = E_Y^{-1} \circ A \circ E_X$$

Associated Matrix with Bases

Main Idea: $(X, \mathcal{X}), (U, \mathcal{U}), \dim X = n, \dim U = m$.

$$L \left(\sum_1^n \alpha_i x_i \right) = \sum_1^m \beta_j u_j \implies F^n \longrightarrow F^m$$
$$\begin{bmatrix} \alpha_1 \\ \vdots \\ \alpha_n \end{bmatrix} \mapsto \begin{bmatrix} \beta_1 \\ \vdots \\ \beta_m \end{bmatrix}$$

Then there is a unique $A \in F^{m \times n}$ such that

$$A \begin{bmatrix} \alpha_1 \\ \vdots \\ \alpha_n \end{bmatrix} = \begin{bmatrix} \beta_1 \\ \vdots \\ \beta_m \end{bmatrix}$$

Write $A = [L]_{\mathcal{X}, \mathcal{U}}$. A depends on the choices of bases

Remark

Every linear map is fully determined by basis elements. If for all $x_i \in \mathcal{B}$ $L(x_i) = y_i$, then for all $x \in X$, $L(x)$ is determined uniquely.

$$[L]_{\mathcal{X}, \mathcal{U}} = [L(x_1) | \cdots | L(x_n)] .$$

Ex 3.2.1

Let A be an $m \times n$ matrix with real entries, $n > m$. Prove that $Ax = 0$ has a nontrivial solution $x \in \mathbb{R}^n$.

Ex 3.2.2

Let $R : \mathbb{R}^2 \rightarrow \mathbb{R}^2$ be the rotation of angle θ about the origin.

- (a) Give a geometric argument that R is linear.
- (b) Find the matrix A such that $R(x) = Ax$ for all $x \in \mathbb{R}^2$.

Ex 3.2.9, Convolution

Let $x \in \mathbb{R}^N$ be denoted as $x = (x_0, x_1, \dots, x_{N-1})$. Given $x, y \in \mathbb{R}^N$, the convolution of x and y is the vector $x * y \in \mathbb{R}^N$ defined by

$$(x * y)_n = \sum_{m=0}^{N-1} x_m y_{n-m}$$

In this formula, if $n - m < 0$, we take $y_{n-m} = y_{N+n-m}$. Prove that if $y \in \mathbb{R}^N$ is fixed, then the mapping $x \mapsto x * y$ is linear. Find the matrix representing this operator.

Definition

Let X and Y be any sets and let $f : X \rightarrow Y$ be a function.

1. f is injective if and only if for all $x_1, x_2 \in X$, $f(x_1) = f(x_2)$ implies $x_1 = x_2$.
2. f is surjective if and only if for each $y \in Y$, there exists $x \in X$ such that $f(x) = y$.
3. If f is both injective and surjective, then f is called bijective.

Remark

- ▶ $f : X \rightarrow Y$ is bijective if and only if there is $g : Y \rightarrow X$ such that $f \circ g(y) = y$ and $g \circ f(x) = x$ for all $x \in X$, $y \in Y$. Define $g = f^{-1}$ and we say “ f is invertible”.
- ▶ If $L : X \rightarrow U$ is an invertible linear operator, then L^{-1} is again linear. In this case, we say 1) “ L is an isomorphism” and 2) X is isomorphic to U .

Theorem

Suppose X and Y are n -dimensional vector spaces over a field F . Then $X \cong Y$.

Remark (Notation)

$$x \in X \implies [x]_{\mathcal{X}} := E_X(x), u \in U \implies [u]_{\mathcal{U}} := E_U(u)$$
$$A = [L]_{\mathcal{X}, \mathcal{U}}, [L]_{\mathcal{X}, \mathcal{U}}[x]_{\mathcal{X}} = [L(x)]_{\mathcal{U}}.$$

$$\begin{array}{ccc} F^n & \xrightarrow{A} & F^m \\ E_X \uparrow & & \uparrow E_U \\ X & \xrightarrow{L} & U \end{array}$$

Remark

$$\mathcal{X} = \{x_1, \dots, x_n\}, \mathcal{U} = \{u_1, \dots, u_m\}.$$

$$(1) [x_i]_{\mathcal{X}} = E_X(x_i) = e_i^n, [u_j]_{\mathcal{U}} = E_U(u_j) = e_j^m.$$

$$(2) [L]_{\mathcal{X}, \mathcal{U}} = A = [A_1 | \dots | A_n].$$

$$A_i = Ae_i = A[x_i]_{\mathcal{X}} = [L]_{\mathcal{X}, \mathcal{U}}[x_i]_{\mathcal{X}} = [L(x_i)]_{\mathcal{U}}.$$

Hence,

$$[L]_{\mathcal{X}, \mathcal{U}} = [[L(x_1)]_{\mathcal{U}} | \dots | [L(x_n)]_{\mathcal{U}}]$$

Ex 3.3.7

Let $L : \mathbb{R}^2 \rightarrow \mathbb{R}^2$ be defined by $L(x) = Ax$, where

$$A = \begin{bmatrix} 1 & 1 \\ 1 & 1 \end{bmatrix}.$$

Let \mathcal{S} be the standard basis, and let $\mathcal{X} = \{(1, 1), (1, 2)\}$. Find $[L]_{\mathcal{X}, \mathcal{X}}$.

Ex 3.3.12

Let $\mathcal{X} = \{(1, 0, 0), (1, 1, 0), (1, 1, 1)\} \subset \mathbb{Z}_2^3$.

(b) Find $[x]_{\mathcal{X}}$ for arbitrary vector x in \mathbb{Z}_2^3 .

Ex 3.3.13

Let $\mathcal{X} = \{(1, 0, 0), (1, 1, 0), (1, 1, 1)\} \subset \mathbb{Z}_2^3$, let $A \in \mathbb{Z}_2^{3 \times 3}$ be defined by

$$A = \begin{bmatrix} 1 & 1 & 0 \\ 1 & 0 & 1 \\ 0 & 1 & 1 \end{bmatrix},$$

and define $L : \mathbb{Z}_2^3 \rightarrow \mathbb{Z}_2^3$ by $L(x) = Ax$. Find $[L]_{\mathcal{X}, \mathcal{X}}$.

Ex 3.3.17

$$F^{m \times n} \cong F^{mn}.$$

Ex 3.3.5

Let X , Y , and Z be sets, and suppose $f : X \rightarrow Y$, $g : Y \rightarrow Z$ are given functions.

(a) f and $g \circ f$ invertible $\implies g$ invertible?

Injective:

Surjective:

(b) g and $g \circ f$ invertible $\implies f$ invertible?

Injective:

Surjective:

(c) $g \circ f$ invertible $\implies f, g$ invertible?

Definition

Let X, U be vector spaces over a field F , and let $L : X \rightarrow U$ be linear.

- ▶ $\ker(L) = \{x \in X : L(x) = 0\}$.
- ▶ $\mathcal{R}(L) = \{L(x) : x \in X\} = \{u \in U : L(x) = u \text{ for some } x \in X\}$.

Theorem

$\ker(L)$ is a subspace of X and $\mathcal{R}(L)$ is a subspace of U .

Remark

- (1) L is injective if and only if $\ker(L) = \{0\}$.
- (2) L is surjective if and only if $\mathcal{R}(L) = U$.
- (3) If X and U are both n -dimensional, then

$$\begin{aligned} L \text{ is injective} &\iff L \text{ is surjective} \iff L \text{ is bijective} \\ &\iff \ker(L) = \{0\} \iff \mathcal{R}(L) = U \end{aligned}$$

The End